

1 Quantification of Optical Coherence 2 Tomography Angiography in Age and Age- 3 related Macular Degeneration using Vessel 4 Density analysis

5
6 **Ehsan Vaghefi, PhD^{1,2}, Sophie Hill, MD³, Hannah M Kersten, PhD¹, David
7 Squirrell, FRANZCO³**

8 1- School of Optometry and Vision Science, University of Auckland, New
9 Zealand

10 2- Auckland Bioengineering Institute, University of Auckland, New Zealand

11 3- Department of Ophthalmology, University of Auckland, New Zealand

12

13 * Corresponding Author

14 Dr Ehsan Vaghefi

15 85 Park Rd, Grafton

16 School of Optometry and Vision Science

17 University of Auckland, Auckland

18 New Zealand

19 E: e.vaghefi@auckland.ac.nz

20 P: +64 9 9231036

21 This work was funded by Faculty Research Development fund, The University of

22 Auckland.

23 None of the authors of this manuscript has a proprietary interest to declare.

24 **Keywords:** Age-related Macular Degeneration, Optical Coherence Tomography

25 Angiography, Retinal Blood Flow, Vessel Density

26

27 Word Count: 3768

28

29 **Brief Summary**

30 In this manuscript, we have studied the potential of retinal vessel density as

31 measured by optical coherence tomography angiography (OCT-A), as a biomarker

32 for detection of high-risk of developing exudative age-related macular degeneration

33 (AMD).

34

35 **Abstract**

36 **Purpose**

37 To determine whether vessel density (VD) as measured by optical coherence tomography
38 angiography (OCT-A) provide insights into retinal and choriocapillaris vascular changes with
39 ageing and intermediate dry age related macular degeneration (AMD).

40 **Methods**

41 Seventy-five participants were recruited into three cohorts; young healthy (YH) group, old healthy
42 (OH) and those at high-risk for exudative AMD. Raw OCT and OCT-A data from TOPCON DRI
43 OCT Triton were exported using Topcon IMAGENET 6.0 software, and 3D datasets were
44 analysed to determine retinal thickness and vessel density.

45 **Results**

46 Central macular thickness measurements revealed a trend of overall retinal thinning with
47 increasing age. VD through the full thickness of the retina was highest in ETDRS sector 4 (the
48 inferior macula) in all the cohorts. Mean VD was significantly higher in the deep capillary plexus
49 than the superficial capillary plexus in all ETDRS sectors in all cohorts but there was no
50 significant difference noted between groups. Choriocapillaris VD was significantly lower in all
51 ETDRS sectors in the in the AMD group compared with the YH and the OH groups.

52 **Conclusions**

53 Retinal vessel density maps, derived from the retinal plexi are not reliable biomarkers for
54 assessing the ageing macular. Our non-proprietary analysis of the vascular density of the
55 choriocapillaris revealed a significant drop off of VD with age and disease but further work is
56 required to corroborate this finding. If repeatable, choriocapillaris VD may provide a non-invasive
57 biomarker of healthy ageing and disease.

58 **Introduction**

59 Currently we lack a reliable, non-invasive vascular biomarker to monitor both the “healthy”
60 ageing and disease progression of age related macular degeneration (AMD). Optical
61 coherence tomography angiography (OCT-A) represents a novel, non-invasive, dye-less
62 retinal vascular imaging technique that can be rapidly acquired during a clinical consultation.
63 OCT-A is a progression of OCT, using repeated B-scans to detect motion contrast¹. Based
64 on the assumption that this represents the movement of erythrocytes travelling through the
65 retinal blood vessels, the retinal vasculature can therefore be visualised.² Previous studies
66 have shown the value of this imaging technique in the evaluation of glaucoma, diabetic
67 retinopathy (DR), retinal vein occlusions and AMD³.

68 The laminar structure of the retina lends itself to segmentation analysis. Histologically, the
69 retina has four retinal capillary plexuses, within the macular only three of these are
70 considered; the superficial, deep and intermediate capillary plexuses^{4, 5}. When the macular
71 vasculature is imaged using OCT-A the vascular layers are segmented automatically by the
72 software. This results in the merging of the intermediate with the deep capillary plexus³. The
73 regions of interest identified on the OCT-A therefore include the superficial capillary plexus
74 (SCP) and the deep capillary plexus (DCP)³. The SCP consists of the vasculature within the
75 retinal nerve fibre layer (RNFL) and ganglion cell layer (GCL). The DCP represents the
76 vascular plexuses present at two locations: These plexuses are at the border of the inner
77 nuclear layer and the outer plexiform layer, and the border of the inner plexiform layer and
78 inner nuclear layer⁶.

79 OCT-A images can be subjectively appraised for the presence of disease or post processed
80 to produce quantitative data that can be evaluated objectively. A number of studies^{7, 8} have
81 used perfusion indices such as vessel density and flow index⁹ as a method of quantitatively
82 analysing OCT-A images. Vessel density is defined as the “percentage area occupied by
83 vessels in the segmented area”⁹. Flow index is defined as “the average decorrelation values
84 in the segmented area”⁹. It has been proposed that these indices could be of value in

85 monitoring disease progression in AMD and DR^{7, 10, 11}. When measuring vessel density, both
 86 vessel length and diameter need to be considered. Poor perfusion can also result in vessel
 87 dilation, therefore vessel density may not give complete information about the retinal
 88 vascular status¹².

89 There are a wide range of OCT-A devices available, each using a variety of algorithms to
 90 capture retinal images. These include OCT-based optical micro-angiography (OMAG); split-
 91 spectrum amplitude decorrelation angiography (SSADA); OCT angiography ratio analysis
 92 (OCT-ARA); speckle variance; phase variance; and correlation mapping¹³. Specific OCT-A
 93 models such as the AngioVue (Optovue, Inc., Fremont, CA, USA) have inbuilt software for
 94 perfusion indices calculation¹⁴. However, the majority of studies have applied post image
 95 analysis in a variety of methods after the images are exported from the OCT-A instrument².

96 This heterogeneity in the image capture and analysis means that data calculated using
 97 different algorithms are not directly comparable, due to systemic difference and poor
 98 agreement². Therefore, external normative databases are unreliable and longitudinal
 99 monitoring of disease using vessel density requires scans to be performed using the same
 100 instrument, scan location and algorithm².

101 The aims of this study are to compare the macular vessel density values for; young healthy,
 102 old healthy, and patients with intermediate dry age related macular degeneration at high risk
 103 of progression to exudative AMD, and to determine if these perfusion indices could usefully
 104 serve as a biomarker for both “healthy” ageing and the identification of disease risk.

105

106

107

108

109

Materials and Methods

Seventy-five participants were recruited through Auckland Optometry Clinic and Milford Eye Clinic, Auckland, New Zealand. All participants provided written informed consent prior to imaging. Ethics approval (#018241) from the University of Auckland Human Participants Ethics Committee was obtained for this study. This research adhered to the tenets of the Declaration of Helsinki.

Participants were divided into three groups, young healthy (YH) group, old healthy (OH) group and the high-risk intermediate AMD group. Twenty participants were recruited into the YH group, twenty-one were recruited into the OH group, and thirty-four recruited into the AMD group. A comprehensive eye exam was conducted on each participant prior to the OCT and OCT-A scans in order to determine high contrast BCVA and the ocular health status of the fundus. Patients with any posterior eye disease that could potentially affect the choroidal or retinal vasculature including, but not limited to glaucoma, polypoidal choroidal vasculopathy, DR, hypertensive retinopathy, and high myopia ($\geq 6D$), were excluded from the study. Patients with any history of neurological disorders were also excluded. The Young Healthy group consisted entirely of individuals between the ages of 20 and 26 and a best corrected visual acuity of $\geq 6/9$ in the eye under test. The “Old Healthy” group consisted of individuals over the age of 55 years and who had and a best corrected visual acuity of $\geq 6/9$ in the eye under test. The “AMD group” consisted entirely of patients with high risk intermediate dry AMD. This being diagnosed if the individual had at least two of the following risk factors; reticular pseudodrusen, established neovascular AMD in the fellow eye, confluent soft drusen with accompanying changes within the retinal pigment epithelium. In order to ensure that all patients in the “AMD cohort” could maintain fixation during OCT-A imaging, only those patients with a best-corrected visual acuity (BCVA) of 6/15 or better were enrolled. The mean age of the participants in the YH group, OH and AMD participants were 23 ± 3 , 65 ± 10 and 75 ± 8 years respectively. Only one eye of each patient was used for the analysis, and if the patient had both eyes scanned, the OCT-A scan that was the better quality (assessed subjectively by the clinical grader) of the two was used. Mean best-corrected visual acuity for the YH, OH and AMD group were 6/6, 6/9, and 6/12 respectively.

The ocular health of all participants was assessed at Auckland Optometry Clinic, by a registered optometrist, prior to enrolment in the study. The macular status of patients enrolled into the AMD group was assessed separately by an experienced retinal specialist (DS).

Data collection protocol

Participants were dilated with 1.0% tropicamide if the pupils were deemed to be too small for adequate OCT scans. Intraocular pressures were measured before and after dilation by iCare tonometry. OCT/OCT-A scans, fundus photography and clinical measurements were all performed at the University of Auckland Optometry Clinic, in a single session.

SS-OCT-A device and scanning protocol

The swept source OCT-A device (Topcon DRI OCT Triton, Topcon Corporation, Japan) was used to obtain the OCT and OCT-A scans. A macular line and 3x3mm² OCT, and 3x3mm² OCT-A scan were performed on each participant. All scans were centered on the fovea, and retinal layers were identified using the IMAGENET 6.0 automated layer detection tool so that the superficial (SCP) and deep (DCP) capillary plexuses could be evaluated. The quality of the generated layers were checked manually. Scan quality was evaluated at the time of acquisition, and repeated if required.

Quantitative analysis of OCT-A Images

Raw OCT and OCT-A data were exported using Topcon IMAGENet 6.0 software. Since OCT-A images are only qualitative, we chose to normalize the OCT-A datasets, prior to further analysis. These datasets were then correlated with vessel density in each retinal sector. OCT-A 3D datasets were analysed in three different ways (outlined below), to investigate the retinal and choroidal thickness and vessel density (VD).

1. ETDRS-specific

The nine Early Treatment Diabetic Retinopathy Study (ETDRS) sectors were computationally generated and centred manually by the clinical examiner, on the fovea (using enface OCT-A) for

each participant. In some instances, the macular region was not in the centre of the enface 3x3mm² OCT-A scan. Hence, after manual adjustment of the ETDRS central positioning on the fovea, the outer ETDRS regions (i.e. 6-9) could have been positioned outside of the imaging area. Therefore, in this study in which we were interested in assessing the role vessel density may play as a potential biomarker for ageing and disease of the central macular, we only used ETDRS regions 1 to 5 for consistency in our analysis. In this process, the OCT-A signal is extracted and saved for each ETDRS region, superimposed on every OCT-A enface layer. In other words, all the 3x3mm² OCT-A layers through the retinal thickness are combined to create an OCT-A volume dataset, 920 of them per 3x3mm² scan, using the Topcon DRI OCT. Finally, the ETDRS regions are manually applied to each of the 920 enface OCT-A images and OCT-A normalized data per region are extracted (see Figure 1). This method has been recently published¹⁵. The averaged normalized vessel density (VD) of the full thickness retina and choriocapillaris of each ETDRS region was then extracted from every participant for further analysis. Normalisation was performed by converting the raw data to percentage of VD in all ETDRS regions, throughout the thickness of the retina.

2. SCP, DCP and Choriocapillaris

Using the same ETDRS regions as above, the VD was extracted from layer 1 and 2 for the SCP calculation, layers 3 and 4 for the DCP and layer 8 for the choriocapillaris. Averaged normalized VD from the SCP, DCP and choriocapillaris was then extracted for each participant, in the three groups.

3. Retinal and choroidal thickness

In addition to OCT-A data, we also recorded the foveal thickness and subfoveal choroidal thickness from the OCT B-scans.

Statistical analysis

MATLAB programming software and custom-written code was used to import and analyse the data, as detailed above. Statistical analysis was performed using MATLAB Statistics Toolbox.

196 Due to unequal sample sizes between the sample groups, we performed the two sample non-
197 parametric Kolmogorov and Smirnov test, which is one of the most general nonparametric
198 methods for comparing two samples. The Kolmogorov and Smirnov method assumes that the
199 data in both groups follow Gaussian distributions. Significant differences between groups were
200 defined as $D < 0.565$ for the Kolmogorov and Smirnov test.
201

Results

Mean age, central macular thickness, and foveal avascular zone diameter.

Seventy-five participants were included in this study of which 20 were in the Young Healthy group (23 ± 3 years old), 21 in the Old Healthy group, (65 ± 10 years old) and 34 in the AMD group (75 ± 8 years old). OCT data was segmented by IMAGEnet 6.0 software, and the mean central macular thickness measurements for the three participating cohorts was extracted: YH 217.4 ± 9.28 μm , OH 209.8 ± 6.13 μm , AMD 174.2 ± 20.3 μm . Horizontal diameter of the foveal avascular zone (FAZ) was also measured from each OCT and mean FAZ measurements were calculated: YH 672.5 ± 73.4 μm , OH 577.7 ± 98.8 μm , AMD 639 ± 126.2 μm .

Vessel density measurements: Full retinal thickness.

ETDRS Sectors	Foveal % (1)	Superior % (2)	Nasal % (3)	Inferior % (4)	Temporal % (5)
Young Healthy n=20	7.5 ± 6.6	18.1 ± 7.3	16.0 ± 7.0	20.5 ± 8.3	18.2 ± 8.1
Old	14.7 ± 9.8	21.1 ± 8.3	20.8 ± 9.9	21.5 ± 11	21.1 ± 9.6

Table 1: Mean Vessel Density from full retinal thickness per ETDRS Sector, for each group.

Healthy n=21					
AMD n=34	29.6±19.8	48.5±31.4	43.8±31.9	47.7±27.9	48.6±24.8

ETDRS Sectors	Foveal % (1)	Superior % (2)	Nasal % (3)	Inferior % (4)	Temporal % (5)
--------------------------	---------------------	-----------------------	--------------------	-----------------------	-----------------------

214

215 Table 1 shows the mean VD throughout the **full retinal thickness**, for each ETDRS sector. In
216 all groups the highest VD was found in sector 4, the inferior macula. Lowest vessel density in all
217 groups was in sector 1 the foveal/parafoveal region. A statistical difference between the vessel
218 density in Sector 1 was found between all the groups, with the vessel density being higher in the
219 AMD cohort (29.6±19.8) compared to the Young Healthy (7.5±6.6) and Old healthy groups
220 (14.7±9.68). Analysis of the vessel density of ETDRS sectors 2-5, revealed that there was a
221 statistical difference in the VD between the AMD group and both the YH and OH groups; but not
222 between the YH and OH groups.

223 **Vessel density measurements: SCP Segment.**

224 *Table 2: Mean Vessel Density in superficial capillary plexus (SCP) defined in sectors based on the*
225 *ETDRS chart (Foveal, Superior, Nasal, Inferior, and Temporal).*

Young Healthy		27.0±12.9		62.3±23.1		60.0±20.1		64.8±21.5		59.9±21.2	
ETDRS n=20		Foveal % (1)		Superior % (2)		Nasal % (3)		Inferior % (4)		Temporal % (5)	
Sectors											
Old											
Young Healthy		35.9±20.3		57.1±19.9		56.3±22.6		56.6 ± 22.1		54.1±24.8	
Healthy n=21		19.8±3.3		25.0±4.7		24.5±4.2		27.8±4.8		25.5±4.8	
AMD n=20											
Old n=34		43.8±31.6		55.0±44		53.9±41.1		62.0±50.1		55.6±40.6	
Healthy n=21		21.3±3.9		25.9±3.5		25.7 ± 4.1		26.7±4.9		24.6±5.0	

226

227 Table 2 shows the mean VD of the SCP segment within each ETDRS sector in each group.

228 Although, there was a trend towards the mean vessel density derived from the SCP in all ETDRS

229 sectors being higher in the AMD cohort compared to the 2 healthy cohorts, no significant

230 difference was found in the VD's recorded in any of the 5 ETDRS sectors between the 3 cohorts.

231 **Vessel density measurements: DCP Segment.**

232 *Table 3: Mean Vessel Density in deep capillary plexus (DCP) defined in sectors based on the*
 233 *ETDRS chart (Foveal, Superior, Nasal, Inferior, and Temporal).*

AMD	25.4 ±7.9	30.4±10.5	30.7±11.1	32.5±11.0	32.0±8.5
ETDRS	Foveal ratio	Superior ratio	Nasal ratio	Inferior ratio	Temporal ratio

234

235 Table 3 shows the mean VD of the DCP segment within each ETDRS sector for all groups.

236 The mean VD was found to be significantly higher (<0.01) in the DCP than the SCP in all ETDRS

237 sectors and in all groups. Although, there was a trend towards the mean vessel density derived

238 from the DCP in ETDRS sector 1 being higher in the AMD cohort compared to the 2 “healthy”

239 cohorts, the difference was not statistically significant. There was a trend for the mean vessel

240 density derived from the DCP in the remaining ETDRS sectors; 2-5, to be higher in the YH cohort

241 compared to the other 2 cohorts, this difference was not statistically significant.

242

243 **Vessel density measurements: Choriocapillaris Segment.**

244

245 *Table 4: Mean Vessel Density in the choriocapillaris, based on the ETDRS chart (Foveal, Superior,*
246 *Nasal, Inferior, and Temporal)*

S Sector s	(1)	(2)	(3)	(4)	(5)
Young Health y n=20	97 ± 10.5	86.2 ± 10.7	90.8 ± 11	103.8 ± 9.8	103 ± 11.3
Old Health y n=21	97 ± 4.5	85.5 ± 8.5	83.9 ± 9.5	86.3 ± 10.5	80.1 ± 12.8
AMD n=34	71.4 ± 24.3	68.1 ± 23.8	69.7 ± 24.4	68.4 ± 22.7	73.1 ± 19.4

247

248 Table 4 shows the mean VD of the Choriocapillaris segment within each ETDRS sector for all
 249 groups. In contrast to the measurements recorded in the retinal segments, statistically
 250 significantly lower mean VD's were recorded in the choriocapillaris segment layer in the AMD
 251 group across all ETDRS sectors compared to the two healthy groups. There was no significant
 252 difference in the VD's measured in each of the ETDRS sectors between YH and the OH groups.

Discussion

Although a number of changes within the ageing macular have previously been reported, a reliable, non-invasive biomarker of both “healthy” macular ageing and disease risk remains elusive. Retinal thickness, both at the fovea and extra-foveal regions, has previously been shown to differ significantly between individuals with early AMD and age-matched controls with the retina being thinner in individuals with disease¹⁶. The results of our study also revealed that there was a trend for the overall retinal thickness to thin with age but this difference was not statistically significant between cohorts. It is also recognised that the choroid tends to thin with age; with the nasal area being thinnest, and the subfoveal region being thickest¹⁷. In addition to thinning with “normal ageing,” Choroidal thinning has also been demonstrated in individuals with early AMD compared to age matched controls^{16, 18-20} however, the significance of this observation is controversial as both the extent and the pattern of thinning is significant in some²⁰ but not all studies^{16,18,19}. The histological evidence regarding choroidal thickness is also contradictory with some studies showing reduction in late AMD²¹ and others showing no change¹⁹; moreover, the changes in the vasculature of the choriocapillaris in early AMD that have been reported from histological studies^{19, 21} appear to be independent of choroidal thickness.

Similarly inconclusive observations have been reported with respect to the width of the Foveal avascular zone with age. Whilst some studies report no difference with age^{22, 23} others have found that the FAZ enlarges with increasing age^{24, 25}. Where reported, the FA at the level of the SCP has been found to be significantly smaller in >60 year olds compared to younger study groups¹⁴, but the significance of this difference is unknown and furthermore no difference was observed when the FAZ at the level of the DCP was analysed¹⁴. Recently, a study studying the FAZ in patients with intermediate non-exudative AMD found no significant difference in the width of the FAZ between patients with non-exudative AMD and healthy controls²⁶. We likewise found there to be no statically significant difference in the diameter of the FAZ between any of the 3 cohorts studied.

Where objective measures of macular anatomy have failed, OCTA may provide an opportunity to develop a reliable, non-invasive vascular biomarker of the ageing macular. The proprietary

OCTA software now has a vessel density function which, in brief uses the proportion of bright pixels to the proportion of dark pixels to derive a measure of vascular density. In the first instance we reviewed the mean vessel density percentage throughout the **full** retinal thickness, for each ETDRS sector. We then used the **segmentation** function of the enface OCTA to explore the relationship of the vessel densities within each of the ETDRS sectors 1-5 in each of the 3 vascular zones of the central macular; (SCP, DCP, Choriocapillaris) across the 3 cohorts; YH, OH and AMD.

As one might expect the VD's derived from the relatively avascular ETDRS sector 1 were, in the **full** thickness maps, significantly reduced compared to the VD's derived from ETDRS sectors 2-5 in the YH, OH, and AMD cohorts. However, and curiously, the VD recorded in the AMD cohort was significantly higher than in the two "Healthy" cohorts, and whilst there was a trend for the VD's derived from the YH cohort to be reduced compared to the OH cohort, this difference was not significant. Aside the marked difference between the AMD cohort and the other two "healthy" cohorts, no other clinically relevant or statistically significant patterns were noted in the VD's derived from the full thickness retinal maps.

Overall, and when the retinal vascular complexes were analysed as separate segments, the highest vessel densities were found in sectors 2 and 4; the superior and inferior macula. Again the VDs derived from ETDRS sector 1 was greater in the AMD cohort compared to the two "Healthy" cohorts. Whilst there was a trend for the VD's derived from the OH cohort to be lower compared to the YH cohort, this difference was not significant. The spatial variation in VD's that we observed; a higher mean VD in the inferior and superior EDTRS sectors and a lower mean VD in the temporal and nasal sectors, in both the deep and superficial capillary plexi, is consistent with published data from previous studies.^{14, 22, 23} Like others^{14, 27}, we also observed that in all ETDRS sectors and in all 3 study groups, the mean VD derived from the DCP was significantly higher than that derived from the SCP.

The association between age and vessel density remains unclear, with reports of statistically significant lower mean vessel density in SCP, DCP and full thickness of retina in each ETDRS sector in patients >60 years old compared to younger cohorts;¹⁴, however, others who have studied this have found no such correlation²². On the whole, our results do not further our

understanding of VD changes with age, being similarly inconclusive. On the basis of these data, one has to conclude that without further work, VD's derived from the retinal segments are not a reliable biomarker for "healthy" ageing.

One consistent feature of the VD's derived from the retinal segments in the present study was the finding that the VD's recorded from ETDRS sector 1, and indeed in many other ETDRS sectors in the AMD cohort, were as a group consistently higher than those obtained from the "healthy" cohorts. One possible explanation for this surprising result is the technical challenges of obtaining high quality OCT-A scans from individuals with even modest visual impairment as accurate central fixation during the OCT-A capture can be challenging. Many of the patients in our AMD cohort struggled to maintain fixation during the image acquisition stage, despite the majority recording a Snellen acuity of 6/12. . Increased acquisition time and the resultant compromise in image quality means there is a higher likelihood of projection and banding artefacts.²⁸ Motion artefacts can also lead to the duplication of inner retinal vascular structures²⁹. As vessel density is derived from the percentage of the area occupied by bright pixels in a segmented area, errors such as vessel duplication and banding artefacts could be incorrectly interpreted during computational analysis as increased vessel density within the image. Such an error would be magnified in ETDRS sector 1 where the effects of these banding and motion artefacts would be more pronounced in what should be avascular tissue.

Although the proprietary OCTA software to assess VD is designed to measure the inner retinal plexi, a vascular bed which has a very different architecture to the choriocapillaris³⁰ we were curious to know whether the principle of VD analysis could be used to assess the vascularity of the choriocapillaris. We therefore extracted the raw data, and then used a manual analysis method previously published in the literature¹⁵, to assess the "Vascular density" of the choriocapillaris segment. Even allowing for the influence of banding and motion artefacts, issues which should increase VD measurements, we observed a significant reduction in the VD in both the AMD and OH cohorts compared to the YH cohort with the reduction in VD being most marked in the AMD cohort (Table 4). This contrasts sharply with the data derived from the retinal plexi which broadly showed no difference between groups. Although interesting, one has to interpret these findings with caution. It is possible that the observed reduction in VD is simply a

result of increased projection artefact and as VD has, to the best of our knowledge, not previously been used to assess the choriocapillaris, it is not possible to corroborate these findings. However, a global reduction in choriocapillaris density, associated with drusen on OCT-A in dry AMD has been demonstrated previously.⁶ It is also recognised that the inner choroid and the choriocapillaris disproportionately thin with age and disease³¹. It is feasible that a thinning choriocapillaris would be less vascular and other groups have hypothesised that choroidal perfusion changes on the OCT-A, may predict disease progression⁶. Debate about whether this is related to actual flow impairment⁶ or projection artefact “masking” the choriocapillaris^{29, 32} due to large drusen or pigment epithelial detachments attenuating the OCT-A signal is still ongoing. Without further corroborative work, the importance of our finding that there is a progressive reduction in choriocapillaris VD in age and disease is difficult to assess. Furthermore, because our old healthy group and AMD cohorts are not aged matched, it is not possible to comment on whether any observed reduction is simply an ageing phenomenon or a marker of disease. Nevertheless, if reproducible, such a measure has the potential to provide a non-invasive biomarker of both “healthy” macular ageing and disease risk. Further limitations of this study include a small sample size, which might reduce the generalisability of our findings and the interpretation of the statistical significance of our data. Furthermore, the Kolmogorov and Smirnov statistical test assumes the Gaussian distribution of the tested data.

Conclusion

OCT-A mean retinal vessel density results for normal subjects are in line with the existing literature. Challenges of image acquisition and the significant influence of image artefact on post-acquisition analysis of patients with mild visual loss due to intermediate age related macular degeneration further limits the value of VD measurements in this patient group. Traditional measures of vessel density, derived from the retinal plexi are therefore unreliable biomarkers for both “healthy” ageing and quantifying risk of developing AMD. Our non-proprietary analysis of the vascular density of the choriocapillaris revealed a significant drop off of VD with age and

368 disease but further work is required to corroborate this finding. If repeatable choriocapillaris VD
369 may provide a non-invasive biomarker of healthy ageing and disease.

370

371 **Acknowledgments**

372 The authors wish to acknowledge the assistance by Mr Diego Sonderegger, from Device
373 Technologies®.

374


References

1. Spaide RF, Fujimoto JG, Waheed NK, Sadda SR, Staurengi G. Optical coherence tomography angiography. *Prog Retin Eye Res* 2018;64:1-55.
2. Rabiolo A, Gelormini F, Sacconi R, et al. Comparison of methods to quantify macular and peripapillary vessel density in optical coherence tomography angiography. *PLoS One* 2018;13:e0205773.
3. Spaide RF, Klancnik JM, Jr., Cooney MJ. Retinal vascular layers imaged by fluorescein angiography and optical coherence tomography angiography. *JAMA Ophthalmol* 2015;133:45-50.
4. Tan PE, Yu PK, Balaratnasingam C, et al. Quantitative confocal imaging of the retinal microvasculature in the human retina. *Invest Ophthalmol Vis Sci* 2012;53:5728-5736.
5. Gariano RF, Iruela-Arispe ML, Hendrickson AE. Vascular development in primate retina: comparison of laminar plexus formation in monkey and human. *Invest Ophthalmol Vis Sci* 1994;35:3442-3455.
6. de Carlo TE, Romano A, Waheed NK, Duker JS. A review of optical coherence tomography angiography (OCTA). *Int J Retina Vitreous* 2015;1:5.
7. Reif R, Qin J, An L, Zhi Z, Dziennis S, Wang R. Quantifying optical microangiography images obtained from a spectral domain optical coherence tomography system. *Int J Biomed Imaging* 2012;2012:509783.
8. Jia Y, Bailey ST, Hwang TS, et al. Quantitative optical coherence tomography angiography of vascular abnormalities in the living human eye. *Proc Natl Acad Sci U S A* 2015;112:E2395-2402.
9. Chalam KV, Sambhav K. Optical Coherence Tomography Angiography in Retinal Diseases. *J Ophthalmic Vis Res* 2016;11:84-92.
10. Pichi F, Nucci P, Baynes K, Lowder CY, Srivastava SK. Sustained-release dexamethasone intravitreal implant in juvenile idiopathic arthritis-related uveitis. *Int Ophthalmol* 2017;37:221-228.
11. Agemy SA, Sripsema NK, Shah CM, et al. Retinal Vascular Perfusion Density Mapping Using Optical Coherence Tomography Angiography in Normals and Diabetic Retinopathy Patients. *Retina* 2015;35:2353-2363.
12. Chu Z, Lin J, Gao C, et al. Quantitative assessment of the retinal microvasculature using optical coherence tomography angiography. *J Biomed Opt* 2016;21:66008.
13. Zhang A ZQ, Chen CL, Wang RK. Methods and algorithms for optical coherence tomography-based angiography: a review and comparison. *J Biomed Opt* 2015;20:100901.
14. Coscas F, Sellam A, Glacet-Bernard A, et al. Normative Data for Vascular Density in Superficial and Deep Capillary Plexuses of Healthy Adults Assessed by Optical Coherence Tomography Angiography. *Invest Ophthalmol Vis Sci* 2016;57:OCT211-223.
15. Hirano T, Chanwimol K, Weichsel J, Tepelus T, Sadda S. Distinct Retinal Capillary Plexuses in Normal Eyes as Observed in Optical Coherence Tomography Angiography Axial Profile Analysis. *Sci Rep* 2018;8:9380.
16. Wood A, Binns A, Margrain T, et al. Retinal and choroidal thickness in early age-related macular degeneration. *Am J Ophthalmol* 2011;152:1030-1038 e1032.
17. MANJUNATH V TM, FUJIMOTO JG, DUKER JS. Choroidal Thickness in Normal Eyes Measured Using Cirrus HD Optical Coherence Tomography. *Am J Ophthalmol* 2010;150:325-329.
18. Chung SE, Kang SW, Lee JH, Kim YT. Choroidal thickness in polypoidal choroidal vasculopathy and exudative age-related macular degeneration. *Ophthalmology* 2011;118:840-845.

19. Ramrattan RS, van der Schaft TL, Mooy CM, de Bruijn WC, Mulder PG, de Jong PT. Morphometric analysis of Bruch's membrane, the choriocapillaris, and the choroid in aging. *Invest Ophthalmol Vis Sci* 1994;35:2857-2864.
20. Sigler EJ, Randolph JC. Comparison of macular choroidal thickness among patients older than age 65 with early atrophic age-related macular degeneration and normals. *Invest Ophthalmol Vis Sci* 2013;54:6307-6313.
21. Sarks SH. Ageing and degeneration in the macular region: a clinico-pathological study. *Br J Ophthalmol* 1976;60:324-341.
22. Rosen RK, B; Mo, S; Geyman, L; Phillips, E; Carroll, J; Weitz, R; Chui, TYP. Age-Related Variations in Foveal Avascular Zone Geometry and Vessel Density - an Optical Coherence Tomography Angiography (OCTA) study. *Invest Ophthalmol Vis Sci* September 2016;57:5501.
23. Gadde SG, Anegondi N, Bhanushali D, et al. Quantification of Vessel Density in Retinal Optical Coherence Tomography Angiography Images Using Local Fractal Dimension. *Invest Ophthalmol Vis Sci* 2016;57:246-252.
24. Iafe NA, Phasukkijwatana N, Chen X, Sarraf D. Retinal Capillary Density and Foveal Avascular Zone Area Are Age-Dependent: Quantitative Analysis Using Optical Coherence Tomography Angiography. *Invest Ophthalmol Vis Sci* 2016;57:5780-5787.
25. Falavarjani KG, Shenazandi H, Naseri D, et al. Foveal Avascular Zone and Vessel Density in Healthy Subjects: An Optical Coherence Tomography Angiography Study. *J Ophthalmic Vis Res* 2018;13:260-265.
26. Stavrev V, Sivkova N, Koleva-Georgieva D. Quantitative Assessment of Foveal Avascular Zone in Patients with Early and Intermediate Nonexudative Age-Related Macular Degeneration Using Optical Coherence Tomography-Angiography %J Open Journal of Ophthalmology. 2018;Vol.08No.03:7.
27. Shahlaee A, Samara WA, Hsu J, et al. In Vivo Assessment of Macular Vascular Density in Healthy Human Eyes Using Optical Coherence Tomography Angiography. *Am J Ophthalmol* 2016;165:39-46.
28. Al-Sheikh M, Ghasemi Falavarjani K, Akil H, Sadda SR. Impact of image quality on OCT angiography based quantitative measurements. *Int J Retina Vitreous* 2017;3:13.
29. Chen FK, Viljoen RD, Bukowska DM. Classification of image artefacts in optical coherence tomography angiography of the choroid in macular diseases. *Clin Exp Ophthalmol* 2016;44:388-399.
30. Spaide RF. Choriocapillaris flow features follow a power law distribution: implications for characterization and mechanisms of disease progression. *American journal of ophthalmology* 2016;170:58-67.
31. Esmaeelpour M, Kajic V, Zabihian B, et al. Choroidal Haller's and Sattler's layer thickness measurement using 3-dimensional 1060-nm optical coherence tomography. *PloS one* 2014;9:e99690.
32. Spaide RF, Fujimoto JG, Waheed NK. Image Artifacts in Optical Coherence Tomography Angiography. *Retina* 2015;35:2163-2180.


466 **Figure Legend**

467 **Figure 1: Retinal layers were automatically identified by IMAGENET 6 software (A) the SCP and**
468 **DCP were then isolated from layers 1 and 2, and 3 and 4 respectively. ETDRS regions 1-9 are**
469 **superimposed onto the enface OCT-A images and manually adjusted so that sector 1 was**
470 **centred on the fovea. In this study we selected sectors 1-5 (B).**

A

NFL: Nerve Fiber Layer
 GCL: Ganglion Cell Layer
 IPL: Inner Plexiform Layer
 INL: Inner Nuclear Layer
 OPL: Outer Plexiform Layer

ONL: Outer Nuclear Layer
 IS: Inner Segment
 OS: Outer Segment
 RPE: Retinal Pigment Epithelium

B

mean OCT - A ETDRS 1