1 Optimizing the deployment of ultra-low volume and indoor

2 residual spraying for dengue outbreak response

3 Authors

- 4 Sean M. Cavany¹, Guido España¹, Alun L. Lloyd², Lance A. Waller³, Uriel Kitron⁴, Helvio Astete⁵,
- 5 William H. Elson⁶, Gonzalo M. Vazquez-Prokopec⁴, Thomas W. Scott⁶, Amy C. Morrison⁷, Robert C.
- 6 Reiner Jr.⁸, T. Alex Perkins¹

7 Affiliations

- 8 1. Department of Biological Sciences & Eck Institute of Global Health, University of Notre Dame
- 9 2. Department of Mathematics & Biomathematics Graduate Program, North Carolina State
- 10 University
- 11 3. Department of Biostatistics and Bioinformatics, Rollins School of Public Health, Emory

12 University

- 13 4. Department of Environmental Sciences, Emory University
- 14 5. U.S. Naval Medical Research Unit Six, Lima
- 15 6. Department of Entomology and Nematology, University of California, Davis
- 16 7. Department of Pathology, Microbiology, and Immunology, School of Veterinary Medicine,
- 17 University of California, Davis
- 18 8. Institute of Health Metrics and Evaluation, University of Washington

19 Funding

- 20 SMC, GFCE, GMVP, ACM, TWS, RCR, and TAP were supported by grant P01AI098670 (TWS, PI)
- 21 from the National Institutes of Health, National Institute for Allergy and Infectious Disease
- 22 (<u>https://www.niaid.nih.gov</u>). The funders had no role in the study design, data collection and analysis,

23 decision to publish, or preparation of the manuscript.

24 Competing interests

25 The authors have declared that no competing interests exist.

26 Abstract (word limit: 300)

Recent years have seen rising incidence of dengue and large outbreaks of Zika and chikungunya, which 27 28 are all caused by viruses transmitted by *Aedes aegypti* mosquitoes. In most settings, the primary 29 intervention against Aedes-transmitted viruses is vector control, such as indoor, ultra-low volume 30 (ULV) spraying. Targeted indoor residual spraying (TIRS) has the potential to more effectively impact 31 Aedes-borne diseases, but its implementation requires careful planning and evaluation. The optimal 32 time to deploy these interventions and their relative epidemiological effects are not well understood, 33 however. We used an agent-based model of dengue virus transmission calibrated to data from Iquitos, 34 Peru to assess the epidemiological effects of these interventions under differing strategies for deploying them. Specifically, we compared strategies where spray application was initiated when incidence rose 35 36 above a threshold based on incidence in recent years to strategies where spraying occurred at the same 37 time(s) each year. In the absence of spraying, the model predicted 361,000 infections [inter-quartile 38 range (IQR): 347,000 – 383,000] in the period 2000-2010. The ULV strategy with the fewest median 39 infections was spraying twice yearly, in March and October, which led to a median of 172,000 infections [IQR: 158,000 – 183,000] over the 11-year study period, a 52% reduction from baseline. 40 41 Compared to spraying once yearly in September, the best threshold-based strategy utilizing ULV had 42 fewer median infections (254,000 vs. 261,000), but required more spraying (351 vs. 274 days). For 43 TIRS, the best strategy was threshold-based, which led to the fewest infections of all strategies tested 44 (9,900; [IQR: 8,720 – 11,400], a 94% reduction), and required fewer days spraying than the equivalent 45 ULV strategy (280). Although spraying twice each year is likely to avert the most infections, our

46 results indicate that a threshold-based strategy can become an alternative to better balance the

47 translation of spraying effort into impact, particularly if used with a residual insecticide.

48 Author Summary (150-200)

49 Over half of the world's population is at risk of infection by dengue virus (DENV) from *Aedes* aegypti 50 *mosquitoes.* While most infected people experience mild or asymptomatic infections, dengue can cause 51 severe symptoms, such as hemorrhage, shock, and death. A vaccine against dengue exists, but it can 52 increase the risk of severe disease in people who have not been previously infected by one of the four 53 DENV serotypes. In many places, therefore, the best currently available way to prevent outbreaks is by 54 controlling the mosquito population. Our study used a simulation model to explore alternative 55 strategies for deploying insecticide in the city of Iquitos in the Peruvian Amazon. Our simulations closely matched empirical patterns from studies of dengue's ecology and epidemiology in Iquitos, such 56 57 as mosquito population dynamics, human household structure, demography, human and mosquito 58 movement, and virus transmission. Our results indicate that an insecticide that has a long-lasting, 59 residual effect will have the biggest impact on reducing DENV transmission. For non-residual insecticides, we find that it is best to begin spraving close to the start of the dengue transmission 60 61 season, as mosquito populations can rebound quickly and resume previous levels of transmission.

62 Introduction

Dengue incidence is rising [1]. Current estimates indicate that over half of the world's population is at risk of dengue virus infection (DENV) [2]. The last decade has also seen explosive outbreaks of Zika and chikungunya viruses, which are transmitted by *Aedes aegypti*, too. Because the only licensed dengue vaccine is contraindicated in individuals without prior DENV exposure [3], and there are no therapeutic options for Zika and chikungunya, the only intervention available to address these diseases in most settings is vector control. The most common method for controlling adult *Ae. aegypti* is ultralow volume (ULV) spraying, defined as a treatment with minimum effective volume of the active ingredient [4–6]. It can be implemented outdoors by plane [7] or trucks [8], or indoors by handheld devices. Indoor application is considered most effective, because *Ae. aegypti* lives primarily inside human habitations [4,9]. In Iquitos, Peru, where our study was focused, ULV is the most commonly used method and has been repeatedly applied city-wide in response to past *Aedes*-transmitted virus outbreaks [10].

75 Apart from the Western hemisphere-wide Ae. aegypti control program, which focused on 76 yellow fever prevention during the 1950s and 1960s [11], there have been two vector control programs that have successfully controlled dengue: Cuba, which used ULV spraying complemented by larval 77 78 source reduction [12], and Singapore, which utilized larval source reduction and community 79 engagement [13]. A 2010 systematic review found five studies on indoor ULV [4], with generally high 80 (up to 100%) mosquito mortality effects that were sustained for only about one month [14–17]. An 81 exception was a study in Thailand, which found a sustained drop in *Ae. aeaypti* landing rates out to six 82 months [18]. A more recent 2016 systematic review found no randomized controlled trials assessing 83 the impact of ULV spraying. A recent study in Iquitos reported that city-wide indoor ULV spraying reduced the Ae. aegypti population by 60%, but effects were only sustained for a short period, also 84 85 about one month [9]. The sum of all this evidence indicates that indoor ULV spraving is effective at 86 reducing adult numbers in the short term [14–16], but with mixed evidence on its impact on virus transmission and disease [4,19,20] and a lack of information on best practices for how to deploy ULV 87 at a city level [19,21]. 88

Traditionally, indoor residual spraying (IRS) has been widely used against malaria, but has not been recommended for control of *Aedes*-borne diseases [4]. In recent years, however, there has been increased interest in utilizing IRS to combat *Aedes*-borne diseases. A 2016 systematic review found no

92 evidence of an effect of IRS on DENV infection risk [19], though only two studies were included in 93 that analysis. A more recent study using contact tracing reported a large epidemiological effect, 94 reducing the probability of future transmission by 86-96% in Cairns, Australia [22]. A study of IRS in 95 Iquitos found more than 80% mosquito mortality in 24 hours for eight weeks after spraying [10]. Recent work to develop targeted IRS (TIRS), where insecticide is sprayed only where Ae. aegypti are 96 likely to rest, led to gains in speed of application, without significant declines in effectiveness [23]. 97 Increased speed coupled with the relatively small size of Iquitos makes it a feasible location to 98 99 undertake city-wide TIRS spraying. The primary drawback is that, there are no published details on 100 best practices for undertaking this approach.

101 Field trials to measure effectiveness and compare different strategies are logistically 102 challenging and in some cases prohibitively expensive due to the complex interplay of mosquito 103 population dynamics, seasonal dynamics, human movement, and fine-scale heterogeneities [19,24–32]. 104 Mathematical modeling can be helpful in multifaceted cases like this for predicting the best 105 intervention strategies. Additionally, the rebound in adult mosquito abundance following spraving [9,16,19], due to immature emergence and movement, and feedbacks caused by reduced egg-laying due 106 107 to increased adult mortality, mean that it is important to capture mosquito population dynamics when modeling vector control strategies. For example, models have been used for many diseases to analyze 108 109 causes of outbreaks and to help optimize response strategies; increasingly, this is happening in real-110 time during outbreaks [33]. Examples include diphtheria among Rohingva refugees [34], where realtime modeling informed resource allocation and transmission mechanisms; the 2013-16 west African 111 112 Ebola outbreak [35.36]: optimum vaccination strategies in response to measles outbreaks [37–40]: and 113 seasonal malaria prophylaxis [41,42]. Recent modeling studies evaluated the impact on dengue of outdoor, truck-mounted ULV spraving in Porto Alegre, Brazil and IRS in Merida, Mexico [8,43]. The 114

former reported that 24% of cases were averted and the latter found that IRS strategies initiated early in the transmission season were generally superior to those initiated late in the season.

117 One challenge associated with ULV and IRS campaigns is determining the criteria for initiating such a 118 response. Brady et al. [44] discussed a variety of ways to determine when a dengue outbreak is 119 occurring, predominantly based on comparison of current incidence from patterns in recent years. An 120 alternative to initiating an intervention when a threshold has been exceeded would be to start the 121 intervention at the same time each year in an effort to prevent transmission from reaching outbreak 122 levels. Hladish et al., considered campaign start date for IRS, finding that deploying IRS four months before the seasonal peak produced the greatest impact on infections [43]. Several studies of malaria 123 124 also found that IRS timing was important [42,45], and one study assessed ULV timing in relation to 125 *Triatoma dimidiata*, the vector for Chagas disease [46]. Few studies have compared alternative 126 methods for initiating outbreak response though, and none, to our knowledge, did so for the impact of 127 indoor ULV on dengue. An added complication is the characteristic variation in seasonal patterns of 128 DENV transmission [20], which along with the aforementioned complex interplay of heterogeneities 129 can result in vector control strategies with the biggest impact on mosquitoes not necessarily 130 corresponding to the biggest reduction in human infections.

To address these challenges, in this study we used a transmission model to investigate the optimal application of indoor ULV or TIRS for dengue control in Iquitos, Peru. Because the timing of DENV transmission seasons can vary considerably across years, we sought criteria that were optimal in the sense of being robust across multiple years, rather than optimizing outbreak response for a single outbreak year. We compared several possible threshold-based strategies, based on a variety of outbreak definitions, to strategies in which insecticide was sprayed regularly on the same date, either once or twice a year, starting at different times each year.

138 Methods

139 Study area and synthetic location generation

140 Our model was calibrated to data from Iquitos, which has a population of about 450,000 people in the Peruvian Amazon [47,48] and where all four DENV serotypes are endemic. Our analysis focuses on 141 the period 2000-2010, in which DENV-3 and DENV-4 were introduced (in 2001 and 2008, 142 respectively). Locations and coordinates of almost half (40,839/92,891) of the locations in the city were 143 collected during surveys conducted as part of prospective cohort studies [49]. For the remaining 144 locations, we randomly assigned them to ministry of health zones, so that the total number within each 145 146 zone matched that recorded in past citywide spraying campaigns [9]. The location type (e.g., home, shop, etc.) of each of these new locations was randomly assigned so that the final distribution of 147 location types matched that from the aforementioned surveys. Their positions were distributed using 148 149 the rSSI algorithm in the spatstat package in R [50,51], so that they were evenly distributed, and at least 5 m separated each location. 150

151 Model overview

152 We simulated outbreak response strategies using an established agent-based model of dengue virus 153 dynamics in Iquitos. This model has been shown to accurately recreate the dynamics of all four DENV 154 serotypes in Iquitos, and has previously been used to answer questions relating to DENV vaccination [47.48.52]. Human agents in the model move according to realistic movement patterns in Iguitos [53]. 155 Household composition and demographic patterns match those seen in Iquitos and Peru as a whole. 156 157 respectively. Mosquito agents move with fixed probability of 0.3 to a nearby location [54] and have a 158 propensity to bite that depends on temperature, the host's body size, and whether it is the mosquito's first bite. Four stages of mosquito development are explicitly modeled (eggs, larvae, pupae, and female 159 160 adults), with density-dependent mortality occurring in the larval stage. Mosquito population dynamics

161 were calibrated, via an additional mortality rate acting on pupae and larvae, so that adult female abundance matched a spatiotemporal estimate of abundance in Iquitos [49]. The model assumed that all 162 163 four DENV serotypes can be transmitted when either a mosquito or human is infectious, the other 164 susceptible, and the mosquito takes a bloodmeal. Transmission occurs with probability 0.9 from 165 mosquitoes to humans and a time-varying probability from humans to mosquitoes [55,56]. Following infection with one DENV serotype, human agents exhibit permanent immunity to that serotype and 166 temporary immunity to the other serotypes for a period of 686 days on average [57]. The rate of 167 168 introduction of each DENV serotype into the population was calibrated so that serotype-specific 169 incidence of infection matched that predicted for Iquitos in a previous study [58]. All features of the 170 model have been thoroughly described in a prior publication [47], and further details are described in 171 the Supplementary Material.

172 Hypothetical spraying protocol

173 We set in place an outbreak response intervention based on a zonal spraving strategy. Spraving takes 174 place on Monday through Saturday. There are 35 Ministry of Health zones in Iquitos, and the outbreak response sprays these 35 zones in batches of a fixed number, until all zones are sprayed (Figure 1). 175 176 After a period of time, another cycle of spraying the 35 zones in batches is initiated, and this process is 177 repeated until a fixed number of cycles have been completed. The number of houses to spray per day is 178 limited by a maximum number of houses that can be sprayed each day. The probability that occupants will be at home and allow the outbreak response team to spray is represented by a compliance 179 probability. The form of vector control is assumed to be an adult insecticide that increases the baseline 180 181 mortality of mosquitoes by a fixed hazard, called thoroughness. For ultra-low volume (ULV) spraying, 182 the increase in mortality decays exponentially following spraying with a half-life of one day. The most 183 realistic parameterization, based on records of past city-wide spraying campaigns, involved attempting

to spray 11,000 houses per day on average, of which ~6,800 (62%) were compliant. Six or seven zones were sprayed simultaneously so that each zone was sprayed three times over a 3-4 week period. There was no training period, no waiting periods between spray cycles, and no repeat visits to houses. We calibrated the intervention thoroughness such that the intervention campaign would generate an approximately 60% drop in total city-wide mosquito abundance following spraying, consistent with empirical estimates for Iquitos by Gunning et al. [9].



191 Figure 1: Map of Iquitos showing the boundaries of the Ministry of Health zones, which are numbered
192 1-35. Inset shows location of Iquitos in Peru [59].

193 We also simulated city-wide TIRS spraying. In this case, we changed the number of houses 194 spraved per day, the thoroughness (i.e., the increase in mosquito mortality), and the residuality of the 195 insecticide. Based on the estimate that it takes 5-6 times longer to spray a house using TIRS compared 196 to ULV (~3 minutes vs. ~15 minutes) [23], we used 2,000 as an upper limit on the number of houses sprayed daily for TIRS. We calibrated the thoroughness and the residuality so that the 24-hour 197 mortality matched that observed for TIRS in Dunbar et al. [23]. This led to a function which had an 198 increase in mosquito mortality of nine deaths/mosquito-day (i.e., increased the daily risk to close to 1), 199 200 which decayed exponentially after 90 days following treatment, with a half-life of 11 days (Figure 2). 201 In the TIRS scenario, each campaign consisted of just one city-wide cycle, compared to three spray 202 cycles for ULV campaigns.



Figure 2: Mortality over time following TIRS spraying. Circles represent data from Dunbar et al. [23],
line represents mortality function fitted by least squares (R²=0.995).

205 Experiments

206 We considered three ways in which spraying could be initiated: when incidence exceeds a threshold,

207 once yearly, or twice yearly. In the threshold-based strategies, spraying is initiated when the weekly or 208 monthly incidence rises one or two standard deviations above the mean incidence from the 209 corresponding week or month from the previous five years (henceforth, adaptive threshold strategies), 210 or when weekly or monthly incidence rises above a fixed threshold (henceforth, fixed threshold strategies) [44]. This leads to a total of four possible adaptive threshold strategies. Note that for the 211 purposes of initiating threshold strategies, incidence represents cases that are symptomatic, whereas in 212 the results, we generally report the number of infections a particular strategy leads to, irrespective of 213 214 symptoms. The yearly and twice yearly strategies begin at the same time(s) each year. We tested yearly 215 spraying starting in each month (12 strategies) and twice-yearly spraying in each pair of months (66 strategies) (Table 1). Due to the residual effect of TIRS and the longer roll-out of the campaign we did 216 not explore twice-yearly strategies for this intervention. We compared the number of infections 217 218 predicted under each of these initiation strategies to the number predicted had there been no spraying 219 over the years 2000-2010. To focus on the effect of the strategy for initiating spraying, we sprayed the 220 Ministry of Health zones in the same order in each simulation. For the same reason, we used the same 221 time series of DENV introduction in each simulation; namely, the trajectory associated with the highest 222 likelihood following the calibration procedure. We chose the number of simulations so that in the 223 absence of spraying, the change in the coefficient of variation of the number of human infections as 224 new simulations were added was less than 0.1% (about 400 simulations) [60]. For model outputs we 225 present the median and the interquartile range (IQR). We use the IQR as the model is highly stochastic 226 and this measure of dispersion is robust to the presence of outliers.

Strategy	Adaptive threshold	Fixed threshold	Yearly	Twice yearly
Ultra-low volume	Based on mean and standard deviation from recent years.	Vary threshold between 1 and 1,000 per month, and 1 and	Start at the beginning of each month. 12 strategies, 400	All pairs of months. 66 strategies, 400 simulations each.
Indoor residual spraying	4 strategies, 400 simulations each.	230 per week. 2,000 simulations.	simulations each.	N/A

227 Table 1: Summary of simulation experiments.

228 Sensitivity analysis

229 For each of the optimum adaptive threshold, vearly, and, in the case of ULV, twice vearly strategies, 230 we undertook a global sensitivity analysis of the total number of human infections and mosquito 231 abundance. For each of the parameters governing spraving (thoroughness, delay between cycles, for 232 ULV only, and compliance), we selected a range of plausible values using the sampling approach of Saltelli et al. [61], and simulated the best outbreak response strategies for each of these [62]. We then 233 234 decomposed the variance in the output into first and higher order effects of the sampled parameters 235 using the SALib package in Python [63]. Because the data used to parameterize the TIRS strategy were 236 from a controlled experiment, we also reduced the thoroughness in the TIRS adaptive threshold 237 strategy to the value used for ULV spraying (1.5) and to half this value (0.75), while keeping all other parameters the same. Finally, we simulated the adaptive threshold strategies for both IRS and ULV in 238 239 scenarios where (a) only 10% of cases were reported, and (b) there was a lag of 2 weeks between 240 infection and notification.

241 **Results**

Unsurprisingly, TIRS was able to prevent more cases overall than ULV (Table 2). The best adaptive
threshold strategies for TIRS started more quickly following a rise in incidence than the best ULV
strategies and earlier in the year for the best yearly strategy. Due to its higher efficacy and long-lasting
effect, TIRS had an order of magnitude greater impact than ULV on the number of infections predicted.

		None	Adaptive threshold	Yearly	Twice yearly
ULV	Best strategy	N/A	When monthly incidence is 2σ above mean	October	March & October
	Number of infections, 2000-2010 (1,000s) [IQR]	361 [347, 383]	254 [210, 277]	261 [250, 277]	172 [158, 183]

-					
	Days spent spraying [IQR]	0	351 [341, 371]	274 [272, 277]	549 [545, 552]
TIRS	Best strategy	N/A	When weekly incidence is 1σ above mean	September	N/A
	Number of infections, 2000-2010 (1,000s) [IQR]	361 [347, 383]	9.90 [8.72, 11.3]	11.8 [9.75, 14.1]	N/A
	Days spent spraying [IQR]	0	280 [175, 315]	385 [385, 385]	N/A

Table 2: Summary of results and predicted number of infections. N/A denotes not applicable, numbers
in cells represent the number of infections over the 11 year period in 1,000s, and the numbers in
brackets represent the inter-quartile range (IQR).

249 Ultra-low volume spraying

250 In the absence of spraying, the model predicted a median of 361,000 infections [IQR: 347,000 –

251 383,000] across the four serotypes in Iquitos in the period 2000-2010. The adaptive threshold strategy

252 performed best when the incidence was monitored on a monthly basis and when spraying was initiated

when incidence exceeded the mean plus one standard deviation from the last five years (254,000

infections; IQR: 210,000 – 277,000). The difference in the impact on incidence between the adaptive

threshold strategies was small (Figure 3a). When spraying was initiated yearly, our model predicted

that starting in September would lead to the fewest cases (261,000; IQR: 250,000 – 277,000), although

spraying in October produced slightly higher, but similar results (262,000; IQR: 253,000 – 275,000)

258 (Figure 3b). In the case of yearly spraying, timing was important; we saw large differences between the

259 best and worst strategies. It is also worth noting that the yearly strategy which led to the lowest average

260 mosquito abundance was spraying in November, not September (Supplementary Figure 1). The best

261 strategy for spraying twice yearly was spraying in September and November (172,000; IQR: 158,000 –

262 183,000) (Figure 3c). Generally, undertaking the first spray in August or September (typically, just

- 263 before the dengue season) and the second in October or November (typically, near the start of the
- 264 dengue season) led to the fewest cases (Figure 3c). In this case, the strategy that led to the fewest cases

- also led to the lowest average mosquito abundance: spraying in September and November
- 266 (Supplementary Figure 1).



Figure 3: Predicted human infections following city-wide ULV spraying. (a) Comparison of adaptive
threshold strategies for initiating spraying; spraying began when the monthly or weekly incidence was

269 one or two standard deviations above the mean for that period from the last five years, as shown on the 270 x-axis. (b) Comparison of yearly city-wide spraying strategies, beginning on the first day of the shown 271 month. (c) Comparison of the median predicted cases for twice yearly spraying strategies, beginning 272 on the first days of the shown month. Darker colors correspond to fewer cases, and the diagonal shows 273 yearly spraying strategies. (d) Comparison of the best strategies in each category: adaptive threshold corresponds to starting when monthly incidence was more than one standard deviation above the 274 275 mean, once yearly corresponds to spraying in September, twice yearly corresponds to spraying in 276 September and November.

277 Comparing the best strategies, spraying twice yearly (in October and March) averted the most 278 infections, but required the most spraying campaigns: 22 campaigns of spraying, spending a median of 279 549 days spraying in total over 11 years. The best adaptive threshold strategy typically led to fewer 280 cases than the best yearly strategy, but required more spray campaigns (a median of 14 [IQR: 14 - 15] 281 compared to 11 for a yearly strategy). In seasons with a large outbreak (2000-01, 2001-02, 2002-03, 282 2008-09, and 2009-10), the adaptive threshold strategy typically performed better than the yearly strategy (Figure 4, Supplementary Figure 2). In years without a large outbreak, the adaptive threshold 283 284 strategy performed worst, even worse than not spraving at all, because herd immunity was reduced from previous years of spraying, while no spraying happened in that year because the threshold was not 285 286 met (See, for example, the 2003-2004 and 2006-2007 seasons, Supplementary Figure 2).



287 Figure 4: Time-series of incidence of human infections for the best ULV strategy in each category. In

each plot, the green lines represent predictions without spraying, and purple represents the given
strategy. The line represents the median of all 400 simulations and shading represents the interquartile range. The vertical dotted lines represent the start of city-wide ULV campaigns (not displayed
for the threshold strategy, as in this case campaigns start at differing times depending on incidence).

292 When we initiated spraying with a fixed threshold (i.e., one that does not depend on the mean 293 and standard deviation from recent years), then the number of infections had a nonlinear relationship with the magnitude of the threshold (Figure 5). Counterintuitively, we observed higher numbers of 294 295 infections at lower thresholds for spraying than at higher thresholds, even though for low thresholds we sprayed more (Figure 5, top and bottom rows). We saw what appeared to be two regimes: declining 296 297 numbers of infections as we increased the threshold for spraying until about 400 infections per month 298 (or 130 per week), and a more modest increase in the number of infections as we increased the 299 threshold above that number. To explore this pattern, we stratified the simulations into those for which 300 the threshold was below 400 cases per month (or 130 cases per week), and those where the threshold was above this (Figure 6). When the threshold was low, spraving often occurred too soon. before 301 outbreaks began in earnest, and, because we were limited to two spray campaigns per year, we 302 303 effectively used up our guota by the time the outbreak occurred (Figure 6: top row). At higher 304 thresholds, spray campaigns more closely corresponded to times when transmission was ongoing and, 305 consequently, the subsequent incidence was lower even though fewer days spent spraving were 306 required (Figure 6; bottom row).



Figure 5: Results when initiating ULV spraying according to a fixed threshold. The top row shows the number of days spent spraying over the 11 year period, the middle row the mean mosquito abundance, and the bottom row the total number of dengue infections. The left column represents when the threshold is monitored on a monthly basis, and the right column when it is monitored on a weekly basis. Each point represents one model simulation, and the line represents predictions by a fitted



312 generalized additive model.

Figure 6: Comparison of frequency of ULV spraying and human incidence with different fixed
thresholds to initiate spraying. In all panels, lilac bars represent the proportion of simulations which
were undertaking spraying on the given day, the green line represents median incident infections, and
the green shading represents the interquartile range. The left column shows when thresholds were

317 monitored monthly, and the right column when they were monitored weekly. In the top row, the

318 threshold was low (400 and below for monthly, 130 and below for weekly), and in the bottom row, the

319 threshold was high (above 400 for monthly, and above 130 for weekly).

320 Targeted indoor residual spraying

321 The strategy for beginning city-wide TIRS that led to fewest cases was to begin when the weekly 322 incidence was one standard deviation above the mean, resulting in 9.900 infections (IOR: 8.720 – 11,300). This strategy reacted more quickly than the best adaptive threshold ULV strategy (i.e., 323 324 monitoring incidence monthly). The result was many fewer infections (17-fold) than the best ULV 325 strategy, which led to a median of 172,000 infections. Because spraving the whole city once with TIRS 326 took longer than spraving the whole city three times with ULV (35 days vs 25 days), the yearly strategy 327 required spraying for more days for TIRS than for ULV. When using an adaptive threshold strategy though, fewer days were spent spraying than any other strategy tested (median of 280 days), despite it 328 329 also having the largest impact on number of infections (Table 2). The best yearly strategy was to begin 330 spraving each September, which resulted in 11.800 cases (IOR: 9,750 – 14,100). There were smaller 331 differences between the yearly TIRS strategies than for ULV spraying, particularly for the yearly 332 strategies (compare Figure 7 to Figure 3). Overall, all TIRS strategies had a much larger impact on the number of infections than did ULV strategies, and were able to almost completely avert some 333 334 outbreaks in later years (Figure 8). This is because repeated applications of IRS almost eliminate Ae. *aegypti* from Iquitos (Supplementary Figure 3). When initiating TIRS after incidence exceeded a fixed 335 336 threshold, the number of predicted cases increased approximately linearly with the threshold (Figure 9).



Figure 7: Predicted human infections following city-wide TIRS spraying. (a) Comparison of adaptive threshold strategies for initiating spraying – spraying began when the monthly or weekly incidence was one or two standard deviations above the mean for that period from the last five years, as shown on the x-axis. (b) Comparison of yearly city-wide spraying strategies, began on the first day of the indicated month. (c) and (d) Comparison of the best strategies in each category: adaptive threshold

- 342 corresponds to starting when weekly incidence is more than one standard deviation above the mean,
- 343 and once yearly corresponds to spraying in September. Panels (c) and (d) display the same results,
- 344 *except that (d) excludes the no spraying scenario.*



Figure 8: Time-series of incidence of human infections for the best TIRS strategy in each category. In each plot the purple lines represent the predictions without spraying, and the green represents the given strategy. The line represents the median of all 400 simulations and the shading represents the inter-quartile range. The vertical dotted lines represent the start of city-wide TIRS campaigns (not displayed for the threshold strategy as in this case campaigns start at differing times depending on

350 incidence).



Figure 9: Results when initiating TIRS according to a fixed threshold. The top row shows the number
of days spent spraying over the 11-year period, the middle row the mean mosquito abundance, and the
bottom row the total number of human DENV infections. The first column represents when the

threshold was monitored on a monthly basis, and the second column when it was monitored on a
weekly basis. Each point represents one model simulation, and the line represents a fitted generalized
additive model.

357 Sensitivity analysis

358 Keeping all other parameters the same, we jointly varied the thoroughness of spraving (i.e., the increase in the daily mortality rate), the compliance of houses, and, in the case of ULV strategies, the delay 359 between spray cycles. For the best adaptive threshold strategy, surveillance effort did not have a large 360 361 effect on the predicted number of cases. This is because our threshold is based on incidence from recent vears, so if only a proportion of cases are notified, then the threshold will be that proportion of its value 362 363 if all cases were notified, and the time at which spraying starts will be similar. Increasing the thoroughness of the spraying (or, equivalently, the efficacy of the treatment) leads to fewer cases 364 averted for small values of thoroughness (Figure 10, left column). However, for values of thoroughness 365 366 above 3 (a daily mortality risk of about 95%), increasing the thoroughness further does not lead to further gains. The probability that a household complies with spraying has a strong negative 367 relationship with the number of infections predicted for all strategies (Figure 10, right column). 368

369 In the case of TIRS, reducing the thoroughness of spraving did not have a strong effect on the number of infections (Supplementary Figure 4). Even for low thoroughness of 0.5, all TIRS 370 371 implementations lead to fewer than 80,000 infections, a more than 4.5-fold reduction from the baseline 372 number of infections (361,000). For the adaptive threshold strategy, reduced household compliance did not have a large effect on the number of infections above a compliance of 0.4 (much below the 373 observed compliance of 0.7). Compliance had a stronger impact on the number of infections for the 374 yearly strategy, in which improved compliance generally led to fewer cases. For both TIRS and ULV 375 376 spraying campaigns, the variance in compliance determined more than half of the overall variance in

the number of infections in all cases except for the adaptive threshold TIRS strategy (Supplementary
Figure 5). In the case of adaptive threshold TIRS, there was a substantial interaction between
compliance and the thoroughness of the spraying, with the interaction between these terms determining
for 39% of the variance in the output.

381 When we reduced thoroughness of TIRS to 1.5 (i.e., the same as that used in ULV spraying), then the best reactive threshold strategy was the same as when the thoroughness was 9: spraying when 382 383 weekly incidence is more than one standard deviation above the mean. Moreover, it had only a small impact on the number of infections we predicted: 12,800 infections (IQR: 12,000 – 14,100) compared 384 to 9,900 in the baseline case. When we reduced thoroughness further (to 0.75), the best threshold 385 386 strategy remained the same, and the impact of the number of infections was still large: we predicted 14,300 infections (IQR: 13,500 – 15,700). In the case of the once yearly strategy, reducing the 387 thoroughness to 1.5 or 0.75 led to predictions of 35,800 (IQR: 32,500 – 38,500) and 45,300 (IQR: 388 41,900 – 49,700), respectively, and changed the best month to spray to August (compared to September 389 390 in the baseline case).

391 In the baseline case, we assumed that 100% of symptomatic cases were notified when 392 calculating whether the threshold for response was reached. However, as the adaptive threshold was itself based on past incidence of notified cases, it indirectly included reporting rate, and so a fixed level 393 under-reporting should not greatly affect response timing, beyond increasing stochasticity. We tested 394 this logic by decreasing the reporting rate to 10%. In this case we observed 248,000 (IQR: 197,000 – 395 (275,000) and (6,370) (IOR: (5,030 - 7,620)) infections for city-wide ULV and IRS campaigns, 396 397 respectively (recall this is the total number of infections and so is unaffected by reporting rate). These values were both below the baseline median prediction for full reporting. The best strategies in both 398 399 instances remained the same as in the baseline, full-reporting case. If surveillance of cases lagged by

400 two weeks, the best adaptive strategies remained the same for both IRS and ULV. For TIRS we

401 observed slightly more infections than when there was no lag (12,100, IQR: 10,300 – 14,500), whereas

402 for ULV we actually observed many fewer cases compared to when there was no lag (164,000, IQR:

403 156,000 - 172,000).



405 Figure 10: Sensitivity analysis jointly varying the thoroughness of ULV spraying (i.e., the increase in mosquito mortality rate on the day of spraying) (left column), the delay, in days, between cycles 406 (middle column), and the compliance of households. The first row shows the results using the best 407 408 adaptive threshold strategy (starting spraying when the monthly incidence is more than one standard deviation above the mean for that month for the last five years), the second row using the best yearly 409 strategy (spraying in September) and the bottom row the best twice yearly strategy (spraying in 410 September and November). Each point represents a model simulation, and the line represents a fitted 411 412 generalized additive model. Vertical dashed lines show the value used in the baseline simulation.

413 **Discussion**

414 We used an agent-based model of DENV transmission in Iquitos, Peru, to compare strategies for 415 initiating city-wide spraying with either ULV or TIRS. None of the city-wide ULV spraying strategies 416 were able to prevent outbreaks, but the best strategies reduced the total number of infections over an 417 eleven-year period by around a half. Strategies that used TIRS were able to almost completely eliminate *Ae. aeaypti* from Iquitos, and so prevent an order of magnitude more infections than ULV. 418 The best strategy for ULV spraying was to spray twice per year, in September and November. Spraying 419 420 yearly in September prevented slightly fewer infections, but required spraying slightly less, than the 421 best adaptive threshold strategy. The yearly and twice yearly strategies also tended to lead to fewer infections than the adaptive threshold strategies in those years when there was not a large outbreak. The 422 best strategy tested for TIRS was an adaptive threshold one, which had the biggest impact on the 423 numbers of infections of all strategies tested. Moreover, it also required the fewest days spent spraying 424 425 compared to all other strategies (280 days over 11 years).

When considering initiation of a city-wide ULV campaign, two factors stood out as optimizing the
impact of outbreak response: (1) begin spraying when the monthly incidence is one standard deviation

428 above the mean, and (2) use a relatively high fixed threshold (for Iquitos: ~400 cases/month) for 429 initiating outbreak response. Taken together, these observations indicate that after an initial increase in 430 incidence, not reacting too quickly can result in a more effective city-wide ULV response. This makes 431 sense, due to the short-term effect of ULV spraying and the capacity of *Ae. aeqypti* populations to 432 rebound rapidly. If we instead consider TIRS, our results indicate that timing is less important, due to the residual effect of the insecticide. Moreover, reductions in vector abundances and numbers of 433 434 infections would be even further reduced with longer lasting insecticides, such as those with 150-day 435 effects that are now becoming available [43].

Our sensitivity analysis indicated that the level of surveillance effort did not have a strong effect on the 436 437 predicted number of cases, due to the fact that the adaptive threshold calculation inherently captures 438 this under-reporting, if it occurs at a constant rate. This would not be the case, however, if the rate of 439 under-reporting changed over time. For instance, if reporting rate increased through time, then 440 thresholds would be based on a smaller proportion of cases than the current year's incidence, which would lead to us spraving too soon due to an artificially small threshold. This could happen if, for 441 example, reporting of DENV infections became more frequent as awareness of symptoms grew among 442 443 the public and/or clinicians. In addition, a lag in reporting did not have a big impact on our model's results about the number of infections prevented by a city-wide TIRS campaign. This was not the case 444 445 for ULV, though. In that case, a lag of two weeks actually led to fewer infections, implying that, if there is not an inherent lag in reporting, it may be worthwhile to wait once the threshold has passed 446 before beginning a city-wide ULV campaign. 447

448 Reassuringly, the impact of a city-wide TIRS campaign was robust to more pessimistic assumptions 449 about the thoroughness with which the insecticide is sprayed (or its efficacy) and the compliance of 450 households, in addition to under-reporting and lags in reporting. This means that, even when the

451	increase in mosquito mortality caused by TIRS at baseline was an order of magnitude below that
452	observed by Dunbar et al. [23], or when half as many houses were treated as observed in city-wide
453	ULV campaigns, then the effect of a city-wide IRS campaign was not greatly impacted.
454	Because the seasonality of DENV transmission is highly irregular in Iquitos [20], a characteristic of
455	DENV transmission in general, yearly strategies can be expected to perform very well in some years
456	(e.g., 2002-03) but poorly in others (e.g., 2001-02), especially if spraying occurs too soon. This implies
457	that caution should be taken to not overinterpret our result that September seemed to be the best month
458	to initiate ULV spraying. September was largely best because application at that time strongly
459	mitigated the 2002-2003 season, which was specific to the particular importation patterns that sparked
460	local transmission during the 2000-2010 period of our analysis. Generalizing across ULV strategies
461	though, it seems that strategies that initiate just before or early in the season perform best.
462	Although we have compared results of city-wide ULV spraying with results of city-wide TIRS, there
463	are some caveats to this comparison. First, we parameterized ULV spraying using a study of actual
464	city-wide spraying campaigns in Iquitos, while we parameterized TIRS using a controlled study from a
465	different country. It is possible that in reality TIRS may have lower effectiveness, although we saw in
466	our sensitivity analysis that if we reduced TIRS to the level of thoroughness used for ULV, the impact
467	on the number of infections was still much greater for TIRS. Secondly, it may not be feasible to
468	undertake city-wide TIRS campaigns, due to the greater cost and time commitment associated with
469	city-wide spray campaigns. On the other hand, our results show that, in the long-term, we would
470	actually need to spray less using TIRS due to the reduced case-load and large reduction in mosquito
471	abundance.

472 Our prediction that the optimal indoor ULV strategy could lead to a reduction of infections by up to473 around 50% exceeds an estimate from a recent modeling study from Porto Alegre, Brazil, which

474 reported that outdoor truck-mounted ULV spraying reduced the number of secondary infections by around 25% [8]. At the same time, our estimate is lower than the 85% reduction in cases due to outdoor 475 476 ULV spraying reported by Wahid et al. [64] in Malakar, Indonesia, although ULV was applied there in 477 conjunction with other interventions (reactive ULV, larviciding, and larval source reduction) [64]. A 478 previous modeling study of IRS spraying in Merida, Mexico found that proactive strategies (i.e., before the season) outperformed reactive strategies [43]. That is commensurate with our results, as each of our 479 threshold strategies and the best performing yearly strategies all involve spraving before the season. 480 481 The observation of Hladish et al. [43] that campaigns that start after the peak in incidence can still have 482 a large effect is consistent with our result that the month when we started TIRS was less important than 483 for ULV. Our prediction of a 97% reduction in the number of infections for repeated TIRS campaigns with about 70% coverage each year is greater than that found by Haldish et al. [43] (79% reduction in 484 485 caseloads over 5 years with 75% coverage in Merida, Mexico) and Vazquez-Prokopec et al. [22] (86% 486 reduction in transmission in treated houses in Cairns, Australia). Comparing to Hladish et al. [43], our 487 higher predicted impact may be because our model incorporates a more detailed entomological 488 component with immature stages and spatial heterogeneity. This means that feedbacks caused by fewer 489 mosquitoes laying fewer eggs, and stochastic local fade-out of adult mosquitoes, allow the Ae. aegypti 490 population to be reduced to very low numbers after several years of TIRS application.

For all ULV strategies, the approach that led to the fewest cases was not necessarily the same as the strategy that reduced mosquito abundance the most, reemphasizing the point made by previous studies of the importance of measuring epidemiological endpoints when assessing vector control [65,66]. This difference is likely due to DENV transmission being the result of a complex interplay of factors, not simply a direct, positive relationship with *Ae. aegypti* abundance. The best strategy also differed by year. In years with low incidence, the adaptive threshold strategies performed poorly, because a

497 response was not triggered.

498 A limitation of our study is that there are few published results on the impact of ULV or TIRS on 499 dengue incidence with which to validate our model [65,66]. This is mitigated somewhat by our model's 500 incorporation of mosquito population dynamics that match observed patterns from Iquitos [49], as well 501 as matching ULV mortality effects to a study carried out in Iquitos and IRS effects to a controlled study in Mexico [9,10]. A second limitation is that, because our model reproduces the seasonal patterns 502 503 observed in the period 2000-2010, our results may be somewhat specific to DENV transmission and 504 mosquito population patterns at that particular place and time. While our results regarding threshold spraying strategies are likely to be robust to this this concern, our predictions for when regular spraying 505 506 should begin may be less robust, and could, for example, differ if importation rates peak at different 507 points in the transmission season. On the other hand, grounding of our model in data from Iquitos, 508 which is an extremely well-characterized site for DENV transmission and Ae. aegypti population 509 dynamics, is a notable strength. Another strength is our model's level of detail, which is something that 510 enables us to capture the interplay of two important feedbacks in mosquito population dynamics 511 following spraying: (1) density-dependent mortality in the larval stage causing the population to rebound and (2) reduced egg-laying by adults. We are also able to capture local mosquito population 512 513 depletion to zero due to demographic stochasticity and subsequent population rebounding due to 514 mosquito movement.

515 Our results indicate that the city-wide ULV and TIRS campaigns would have reduced the number of 516 DENV infections in Iquitos by up to half relative to the baseline scenario that we modeled. Although a 517 well-timed campaign could be expected to mitigate transmission in a particular season, it would be 518 difficult to prevent an outbreak altogether using ULV or TIRS. Similarly, selecting a single strategy 519 that consistently mitigated outbreaks across multiple years proved to be difficult. For example, our

adaptive threshold strategies performed well during the 2001-2002 transmission season, but poorly in 2002-2003. The opposite was true of the yearly strategy. With indoor ULV spraying, the best strategy was with a fixed threshold of around 400 infections per month. A downside of this approach is that it requires accurate, timely, and potentially expensive surveillance. More field work is needed to better understand the feasibility and effectiveness of city-wide TIRS, including its spatially targeted application in combination with ULV. We predict, however, that city-wide TIRS, if feasible, will have a greater impact than ULV without asking significantly more from surveillance.

527 Disclaimer

528 The views expressed in this article are those of the authors and do not necessarily reflect the official

529 policies or positions of the Department of the Navy, Department of Defense, nor the U.S. Government.

530 Ethical considerations

The study protocol was approved by the Naval Medical Research Unit No. 6 (NAMRU-6) Institutional
Review Board (IRB) (protocol #NAMRU6.2014.0028), in compliance with all applicable Federal
regulations governing the protection of human subjects. IRB relying agreements were established
between NAMRU-6, the University of California, Davis, Tulane University, Emory University and
Notre Dame University. The protocol was reviewed and approved by the Loreto Regional Health
Department, which oversees health research in Iquitos. This study represents historical data analysis
using data without personal identifiers.

538 Author contributions

Contributor Role	Contributors
Conceptualization	GVP, TWS, RCR, TAP
Data Curation	HA, ACM
Formal Analysis	SMC, GFCE, RCR, TAP; all authors assisted with interpretation of results.

Contributor Role	Contributors		
Funding Acquisition	TWS, TAP		
Investigation	SMC, GFCE, TAP		
Methodology	SMC, GFCE, TAP		
Project Administration	GVP, TWS, TAP		
Resources	ТАР		
Software	SMC, GFCE, RCR, TAP		
Supervision	ТАР		
Validation	SMC		
Visualization	SMC, TAP		
Writing – Original Draft Preparation	SMC		
Writing – Review & Editing	All authors		

539 Copyright Statement

- 540 ACM and HA were employees of the United States government. This work was prepared as part of
- their official duties. Title 17 U.S.C. 105 provides that `copyright protection under this title is not
- available for any work of the United States Government.' Title 17 U.S.C. 101 defines a U.S.
- 543 Government work as work prepared by a military service member or employee of the U.S. Government
- 544 as part of that person's official duties.

545 Acknowledgments

546 We thank the residents of Iquitos for their participation in this study. We greatly appreciate the support

- 547 of the Loreto Regional Health Department, including Drs. Hugo Rodriguez-Ferruci, Christian Carey,
- 548 Carlos Alvarez, Hernan Silva, and Lic. Wilma Casanova Rojas who all facilitated our work in Iquitos.
- 549 We thank the NAMRU-6 Virology and Emerging Infections Department (VEID) and Entomology
- 550 Department leadership who provided institutional support, IRB guidance and support supervising field
- staff during the years 2000-2010 when the data used in these models was collected. We also appreciate

- 552 the careful commentary and advice provided by the NAMRU-6 IRB and Research Administration
- 553 Program for the duration of this study. We thank the NAMRU-6 VEID field teams provided who daily
- support through duration of the project and without whom the capture of acute dengue cases would not
- base been possible. In particular we thank Gabriela Vasquez de la Torre for her administrative support
- 556 for the project.

557 **References**

- 1. Campos NBD, Morais MHF, Ceolin APR, Cunha M da CM, Nicolino RR, Schultes OL, et al. Twenty-Two years of dengue fever (1996-2017): an epidemiological study in a Brazilian city. Int J Environ Health Res. 2019; 1–10. doi:10.1080/09603123.2019.1656801
- 2. Messina JP, Brady OJ, Golding N, Kraemer MUG, Wint GRW, Ray SE, et al. The current and future global distribution and population at risk of dengue. Nature Microbiology. 2019; 1. doi:10.1038/s41564-019-0476-8
- 3. Sridhar S, Luedtke A, Langevin E, Zhu M, Bonaparte M, Machabert T, et al. Effect of Dengue Serostatus on Dengue Vaccine Safety and Efficacy. N Engl J Med. 2018;379: 327–340. doi:10.1056/NEJMoa1800820
- 4. Esu E, Lenhart A, Smith L, Horstick O. Effectiveness of peridomestic space spraying with insecticide on dengue transmission; systematic review. Tropical Medicine & International Health. 2010;15: 619–631. doi:10.1111/j.1365-3156.2010.02489.x
- 5. Bonds J a. S. Ultra-low-volume space sprays in mosquito control: a critical review. Medical and Veterinary Entomology. 2012;26: 121–130. doi:10.1111/j.1365-2915.2011.00992.x
- 6. Boubidi SC, Roiz D, Rossignol M, Chandre F, Benoit R, Raselli M, et al. Efficacy of ULV and thermal aerosols of deltamethrin for control of Aedes albopictus in Nice, France. Parasites & Vectors. 2016;9: 597. doi:10.1186/s13071-016-1881-y
- 7. Breidenbaugh MS, Haagsma KA, Wojcik GM, De Szalay FA. Efficacy of aerial spray applications using fuselage booms on Air Force C-130H aircraft against mosquitoes and biting midges. J Am Mosq Control Assoc. 2009;25: 467–473. doi:10.2987/08-5892.1
- 8. Marini G, Guzzetta G, Marques Toledo CA, Teixeira M, Rosà R, Merler S. Effectiveness of Ultra-Low Volume insecticide spraying to prevent dengue in a non-endemic metropolitan area of Brazil. PLoS Comput Biol. 2019;15: e1006831. doi:10.1371/journal.pcbi.1006831
- Gunning CE, Okamoto KW, Astete H, Vasquez GM, Erhardt E, Del Aguila C, et al. Efficacy of Aedes aegypti control by indoor Ultra Low Volume (ULV) insecticide spraying in Iquitos, Peru. Barrera R, editor. PLOS Neglected Tropical Diseases. 2018;12: e0006378. doi:10.1371/journal.pntd.0006378

- 10. Paredes-Esquivel C, Lenhart A, del Río R, Leza MM, Estrugo M, Chalco E, et al. The impact of indoor residual spraying of deltamethrin on dengue vector populations in the Peruvian Amazon. Acta Tropica. 2016;154: 139–144. doi:10.1016/j.actatropica.2015.10.020
- 11. Schliessman, DJ, Calheiros, LB. A review of the status of yellow fever and Aedes aegypti eradication programs in the Americas. [Internet]. 1974. Available: http://archive.org/details/cbarchive_116599_areviewofthestatusofyellowfeve1974
- 12. Armada Gessa JA, Figueredo González R. Application of environmental management principles in the program for eradication of Aedes (Stegomyia) aegypti (Linneus, 1762) in the Republic of Cuba, 1984. Bull Pan Am Health Organ. 1986;20: 186–193.
- 13. Ooi E-E, Goh K-T, Gubler DJ. Dengue prevention and 35 years of vector control in Singapore. Emerging Infect Dis. 2006;12: 887–893. doi:10.3201/10.3201/eid1206.051210
- 14. Perich MJ, Rocha N O, Castro A L, Alfaro A W, Platt KB, Solano T, et al. Evaluation of the efficacy of lambda-cyhalothrin applied by three spray application methods for emergency control of Aedes aegypti in Costa Rica. J Am Mosq Control Assoc. 2003;19: 58–62.
- 15. Perich MJ, Sherman C, Burge R, Gill E, Quintana M, Wirtz RA. Evaluation of the efficacy of lambda-cyhalothrin applied as ultra-low volume and thermal fog for emergency control of Aedes aegypti in Honduras. J Am Mosq Control Assoc. 2001;17: 221–224.
- 16. Koenraadt CJM, Aldstadt J, Kijchalao U, Kengluecha A, Jones JW, Scott TW. Spatial and Temporal Patterns in the Recovery of Aedes aegypti (Diptera: Culicidae) Populations After Insecticide Treatment. J Med Entomol. 2007;44: 65–71. doi:10.1093/jmedent/41.5.65
- 17. Castro M, Quintana N, Quiñones P ML. Evaluating two pyrethroids in dengue vector control in Putumayo, Colombia. Revista de Salud Pública. 2007;9: 106–116.
- 18. Pant CP, Mathis HL, Nelson MJ, Phanthumachinda B. A large-scale field trial of ultra-low-volume fenitrothion applied by a portable mist blower for the control of Aedes aegypti. Bull World Health Organ. 1974;51: 409–415.
- 19. Bowman LR, Donegan S, McCall PJ. Is Dengue Vector Control Deficient in Effectiveness or Evidence?: Systematic Review and Meta-analysis. PLOS Neglected Tropical Diseases. 2016;10: e0004551. doi:10.1371/journal.pntd.0004551
- 20. Stoddard ST, Wearing HJ, Reiner RC, Morrison AC, Astete H, Vilcarromero S, et al. Long-Term and Seasonal Dynamics of Dengue in Iquitos, Peru. Barrera R, editor. PLoS Neglected Tropical Diseases. 2014;8: e3003. doi:10.1371/journal.pntd.0003003
- 21. Achee NL, Gould F, Perkins TA, Reiner RC, Morrison AC, Ritchie SA, et al. A Critical Assessment of Vector Control for Dengue Prevention. Halstead SB, editor. PLOS Neglected Tropical Diseases. 2015;9: e0003655. doi:10.1371/journal.pntd.0003655
- 22. Vazquez-Prokopec GM, Montgomery BL, Horne P, Clennon JA, Ritchie SA. Combining contact tracing with targeted indoor residual spraying significantly reduces dengue transmission. Sci Adv.

2017;3: e1602024. doi:10.1126/sciadv.1602024

- 23. Dunbar MW, Correa-Morales F, Dzul-Manzanilla F, Medina-Barreiro A, Bibiano-Marín W, Morales-Ríos E, et al. Efficacy of novel indoor residual spraying methods targeting pyrethroid-resistant Aedes aegypti within experimental houses. PLOS Neglected Tropical Diseases. 2019;13: e0007203. doi:10.1371/journal.pntd.0007203
- 24. Stoddard ST, Forshey BM, Morrison AC, Paz-Soldan VA, Vazquez-Prokopec GM, Astete H, et al. House-to-house human movement drives dengue virus transmission. Proceedings of the National Academy of Sciences. 2013;110: 994–999.
- 25. Cuong HQ, Vu NT, Cazelles B, Boni MF, Thai KTD, Rabaa MA, et al. Spatiotemporal Dynamics of Dengue Epidemics, Southern Vietnam. Emerg Infect Dis. 2013;19: 945–953. doi:10.3201/eid1906.121323
- 26. Yoon I-K, Getis A, Aldstadt J, Rothman AL, Tannitisupawong D, Koenraadt CJM, et al. Fine Scale Spatiotemporal Clustering of Dengue Virus Transmission in Children and Aedes aegypti in Rural Thai Villages. PLOS Neglected Tropical Diseases. 2012;6: e1730. doi:10.1371/journal.pntd.0001730
- 27. Peterson AT, Martínez-Campos C, Nakazawa Y, Martínez-Meyer E. Time-specific ecological niche modeling predicts spatial dynamics of vector insects and human dengue cases. Trans R Soc Trop Med Hyg. 2005;99: 647–655. doi:10.1016/j.trstmh.2005.02.004
- 28. Getis A, Morrison AC, Gray K, Scott TW. Characteristics of the spatial pattern of the dengue vector, Aedes aegypti, in Iquitos, Peru. Perspectives on Spatial Data Analysis. Springer; 2010. pp. 203–225.
- 29. LaCon G, Morrison AC, Astete H, Stoddard ST, Paz-Soldan VA, Elder JP, et al. Shifting patterns of Aedes aegypti fine scale spatial clustering in Iquitos, Peru. PLoS Negl Trop Dis. 2014;8: e3038. doi:10.1371/journal.pntd.0003038
- Liebman KA, Stoddard ST, Morrison AC, Rocha C, Minnick S, Sihuincha M, et al. Spatial Dimensions of Dengue Virus Transmission across Interepidemic and Epidemic Periods in Iquitos, Peru (1999–2003). Kent Crockett RJ, editor. PLoS Neglected Tropical Diseases. 2012;6: e1472. doi:10.1371/journal.pntd.0001472
- 31. Liebman KA, Stoddard ST, Reiner RC, Perkins TA, Astete H, Sihuincha M, et al. Determinants of Heterogeneous Blood Feeding Patterns by Aedes aegypti in Iquitos, Peru. Barrera R, editor. PLoS Neglected Tropical Diseases. 2014;8: e2702. doi:10.1371/journal.pntd.0002702
- 32. Perkins TA, Scott TW, Menach AL, Smith DL. Heterogeneity, Mixing, and the Spatial Scales of Mosquito-Borne Pathogen Transmission. PLOS Computational Biology. 2013;9: e1003327. doi:10.1371/journal.pcbi.1003327
- 33. Polonsky Jonathan A., Baidjoe Amrish, Kamvar Zhian N., Cori Anne, Durski Kara, Edmunds W. John, et al. Outbreak analytics: a developing data science for informing the response to emerging pathogens. Philosophical Transactions of the Royal Society B: Biological Sciences. 2019;374:

20180276. doi:10.1098/rstb.2018.0276

- Finger F, Funk S, White K, Siddiqui MR, Edmunds WJ, Kucharski AJ. Real-time analysis of the diphtheria outbreak in forcibly displaced Myanmar nationals in Bangladesh. BMC Med. 2019;17: 58. doi:10.1186/s12916-019-1288-7
- 35. Funk S, Camacho A, Kucharski AJ, Lowe R, Eggo RM, Edmunds WJ. Assessing the performance of real-time epidemic forecasts: A case study of Ebola in the Western Area region of Sierra Leone, 2014-15. PLoS Comput Biol. 2019;15: e1006785. doi:10.1371/journal.pcbi.1006785
- 36. Kucharski AJ, Camacho A, Flasche S, Glover RE, Edmunds WJ, Funk S. Measuring the impact of Ebola control measures in Sierra Leone. PNAS. 2015;112: 14366–14371. doi:10.1073/pnas.1508814112
- 37. Tao Y, Shea K, Ferrari M. Logistical constraints lead to an intermediate optimum in outbreak response vaccination. PLoS Comput Biol. 2018;14: e1006161. doi:10.1371/journal.pcbi.1006161
- 38. Gastañaduy PA, Funk S, Paul P, Tatham L, Fisher N, Budd J, et al. Impact of Public Health Responses During a Measles Outbreak in an Amish Community in Ohio: Modeling the Dynamics of Transmission. Am J Epidemiol. 2018;187: 2002–2010. doi:10.1093/aje/kwy082
- 39. Thompson KM, Kisjes KH. Modeling Measles Transmission in the North American Amish and Options for Outbreak Response. Risk Anal. 2016;36: 1404–1417. doi:10.1111/risa.12440
- 40. Grais R.F, Conlan A.J.K, Ferrari M.J, Djibo A, Le Menach A, Bjørnstad O.N, et al. Time is of the essence: exploring a measles outbreak response vaccination in Niamey, Niger. Journal of The Royal Society Interface. 2008;5: 67–74. doi:10.1098/rsif.2007.1038
- 41. Gao D, Amza A, Nassirou B, Kadri B, Sippl-Swezey N, Liu F, et al. Optimal seasonal timing of oral azithromycin for malaria. Am J Trop Med Hyg. 2014;91: 936–942. doi:10.4269/ajtmh.13-0474
- 42. Selvaraj P, Wenger EA, Gerardin J. Seasonality and heterogeneity of malaria transmission determine success of interventions in high-endemic settings: a modeling study. BMC Infect Dis. 2018;18: 413. doi:10.1186/s12879-018-3319-y
- 43. Hladish TJ, Pearson CAB, Patricia Rojas D, Gomez-Dantes H, Halloran ME, Vazquez-Prokopec GM, et al. Forecasting the effectiveness of indoor residual spraying for reducing dengue burden. PLoS Negl Trop Dis. 2018;12: e0006570. doi:10.1371/journal.pntd.0006570
- 44. Brady OJ, Smith DL, Scott TW, Hay SI. Dengue disease outbreak definitions are implicitly variable. Epidemics. 2015;11: 92–102. doi:10.1016/j.epidem.2015.03.002
- 45. Worrall E, Connor SJ, Thomson MC. A model to simulate the impact of timing, coverage and transmission intensity on the effectiveness of indoor residual spraying (IRS) for malaria control. Tropical Medicine & International Health. 2007;12: 75–88. doi:10.1111/j.1365-3156.2006.01772.x

- 46. Barbu C, Dumonteil E, Gourbière S. Optimization of Control Strategies for Non-Domiciliated Triatoma dimidiata, Chagas Disease Vector in the Yucatán Peninsula, Mexico. PLOS Neglected Tropical Diseases. 2009;3: e416. doi:10.1371/journal.pntd.0000416
- 47. Perkins TA, Reiner RC, España G, Ten Bosch QA, Verma A, Liebman KA, et al. An agent-based model of dengue virus transmission shows how uncertainty about breakthrough infections influences vaccination impact projections. PLoS Comput Biol. 2019;15: e1006710. doi:10.1371/journal.pcbi.1006710
- 48. España G, Hogea C, Guignard A, Ten Bosch QA, Morrison AC, Smith DL, et al. Biased efficacy estimates in phase-III dengue vaccine trials due to heterogeneous exposure and differential detectability of primary infections across trial arms. PLoS ONE. 2019;14: e0210041. doi:10.1371/journal.pone.0210041
- 49. Reiner RC, Stoddard ST, Vazquez-Prokopec GM, Astete H, Perkins TA, Sihuincha M, et al. Estimating the impact of city-wide Aedes aegypti population control: An observational study in Iquitos, Peru. PLoS Negl Trop Dis. 2019;13: e0007255. doi:10.1371/journal.pntd.0007255
- 50. Baddeley A, Rubak RT and E. spatstat: Spatial Point Pattern Analysis, Model-Fitting, Simulation, Tests [Internet]. 2019. Available: https://CRAN.R-project.org/package=spatstat
- 51. R Core Team. R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing; 2018.
- 52. Flasche S, Jit M, Rodríguez-Barraquer I, Coudeville L, Recker M, Koelle K, et al. The Long-Term Safety, Public Health Impact, and Cost-Effectiveness of Routine Vaccination with a Recombinant, Live-Attenuated Dengue Vaccine (Dengvaxia): A Model Comparison Study. PLOS Medicine. 2016;13: e1002181. doi:10.1371/journal.pmed.1002181
- 53. Perkins TA, Garcia AJ, Paz-Soldán VA, Stoddard ST, Reiner RC, Vazquez-Prokopec G, et al. Theory and data for simulating fine-scale human movement in an urban environment. J R Soc Interface. 2014;11. doi:10.1098/rsif.2014.0642
- 54. Magori K, Legros M, Puente ME, Focks DA, Scott TW, Lloyd AL, et al. Skeeter Buster: a stochastic, spatially explicit modeling tool for studying Aedes aegypti population replacement and population suppression strategies. PLoS Negl Trop Dis. 2009;3: e508. doi:10.1371/journal.pntd.0000508
- 55. Ellis AM, Garcia AJ, Focks DA, Morrison AC, Scott TW. Parameterization and Sensitivity Analysis of a Complex Simulation Model for Mosquito Population Dynamics, Dengue Transmission, and Their Control. Am J Trop Med Hyg. 2011;85: 257–264. doi:10.4269/ajtmh.2011.10-0516
- 56. Nishiura H, Halstead SB. Natural History of Dengue Virus (DENV)—1 and DENV—4 Infections: Reanalysis of Classic Studies. J Infect Dis. 2007;195: 1007–1013. doi:10.1086/511825
- 57. Reich Nicholas G., Shrestha Sourya, King Aaron A., Rohani Pejman, Lessler Justin, Kalayanarooj Siripen, et al. Interactions between serotypes of dengue highlight epidemiological impact of cross-

immunity. Journal of The Royal Society Interface. 2013;10: 20130414. doi:10.1098/rsif.2013.0414

- 58. Reiner RC, Stoddard ST, Forshey BM, King AA, Ellis AM, Lloyd AL, et al. Time-varying, serotype-specific force of infection of dengue virus. Proc Natl Acad Sci USA. 2014;111: E2694-2702. doi:10.1073/pnas.1314933111
- 59. Huhsunqu. Peru [Internet]. wikimedia.org; 2009. Available: https://commons.wikimedia.org/wiki/ File:Peru_location_map.svg
- 60. Lee J-S, Filatova T, Ligmann-Zielinska A, Hassani-Mahmooei B, Stonedahl F, Lorscheid I, et al. The Complexities of Agent-Based Modeling Output Analysis. JASSS. 2015;18: 4.
- 61. Saltelli A, Annoni P, Azzini I, Campolongo F, Ratto M, Tarantola S. Variance based sensitivity analysis of model output. Design and estimator for the total sensitivity index. Computer Physics Communications. 2010;181: 259–270. doi:10.1016/j.cpc.2009.09.018
- 62. Wu J, Dhingra R, Gambhir M, Remais JV. Sensitivity analysis of infectious disease models: methods, advances and their application. J R Soc Interface. 2013;10. doi:10.1098/rsif.2012.1018
- 63. Herman J, Usher W. SALib: An open-source Python library for Sensitivity Analysis. JOSS. 2017;2: 97. doi:10.21105/joss.00097
- 64. Wahid I, Ishak H, Hafid A, Fajri M, Sidjal S, Nurdin A, et al. Integrated vector management with additional pre-transmission season thermal fogging is associated with a reduction in dengue incidence in Makassar, Indonesia: Results of an 8-year observational study. PLOS Neglected Tropical Diseases. 2019;13: e0007606. doi:10.1371/journal.pntd.0007606
- 65. Bowman LR, Runge-Ranzinger S, McCall PJ. Assessing the Relationship between Vector Indices and Dengue Transmission: A Systematic Review of the Evidence. PLOS Neglected Tropical Diseases. 2014;8: e2848. doi:10.1371/journal.pntd.0002848
- 66. Reiner RC, Achee N, Barrera R, Burkot TR, Chadee DD, Devine GJ, et al. Quantifying the Epidemiological Impact of Vector Control on Dengue. Harris E, editor. PLOS Neglected Tropical Diseases. 2016;10: e0004588. doi:10.1371/journal.pntd.0004588
- 67. Paz-Soldan VA, Stoddard ST, Vazquez-Prokopec G, Morrison AC, Elder JP, Kitron U, et al. Assessing and maximizing the acceptability of global positioning system device use for studying the role of human movement in dengue virus transmission in Iquitos, Peru. Am J Trop Med Hyg. 2010;82: 723–730. doi:10.4269/ajtmh.2010.09-0496
- 68. Paz-Soldan VA, Reiner RC, Morrison AC, Stoddard ST, Kitron U, Scott TW, et al. Strengths and weaknesses of Global Positioning System (GPS) data-loggers and semi-structured interviews for capturing fine-scale human mobility: findings from Iquitos, Peru. PLoS Negl Trop Dis. 2014;8: e2888. doi:10.1371/journal.pntd.0002888
- 69. Otero M, Solari HG, Schweigmann N. A stochastic population dynamics model for Aedes aegypti: formulation and application to a city with temperate climate. Bull Math Biol. 2006;68:

1945-1974. doi:10.1007/s11538-006-9067-y

- 70. Focks DA, Haile DG, Daniels E, Mount GA. Dynamic Life Table Model for Aedes aegypti (Diptera: Culicidae): Analysis of the Literature and Model Development. J Med Entomol. 1993;30: 1003–1017. doi:10.1093/jmedent/30.6.1003
- 71. Ten Bosch QA, Clapham HE, Lambrechts L, Duong V, Buchy P, Althouse BM, et al. Contributions from the silent majority dominate dengue virus transmission. PLoS Pathog. 2018;14: e1006965. doi:10.1371/journal.ppat.1006965
- 72. Chan M, Johansson MA. The incubation periods of Dengue viruses. PLoS ONE. 2012;7: e50972. doi:10.1371/journal.pone.0050972
- 73. Clapham HE, Cummings DAT, Johansson MA. Immune status alters the probability of apparent illness due to dengue virus infection: Evidence from a pooled analysis across multiple cohort and cluster studies. PLoS Negl Trop Dis. 2017;11: e0005926. doi:10.1371/journal.pntd.0005926

Supplementary Text S1. Agent-based model of dengue virus transmission.

An ABM consists of "agents" (in this case, people and mosquitoes) who interact with each other in a 561 shared environment. The environment of our model was represented by climate conditions and a set of 562 locations, such as houses, schools, parks, cemeteries, and churches. Climate conditions affected 563 564 mosquito biting frequency, survival, and incubation time of DENV. Locations represented all 92,891 565 buildings in the city. For about 38,835 of these locations, mainly in the central portion of the city, we 566 had data on exact coordinates and location type. For the remainder, we randomly distributed the 567 locations and randomly assigned a location type. Agents represented approximately 450,000 individuals that live in this area [47]. 568

569 Our synthetic population realistically portrayed the population of Iquitos in terms of demographic

570 characteristics of how people are distributed across houses and over time. Specifically, the

571 demographic profiles of the modeled households were consistent with survey data collected during a

572 previous study [31]. The population-wide sex and age distributions were consistent with U.N. estimates

573 for Peru. To represent population changes in time, we simulated human births and deaths that match

574 those estimated by the U.N. for Iquitos, while simultaneously preserving realistic household

575 compositions by placing newborn children in houses with appropriately aged mothers as determined by

576 U.N. estimates of age-specific fertility of Peru [47]. For each person, we simulated daily human

577 movement patterns with a model previously described by Perkins et al. [53], which was fitted to data

578 from retrospective, semi-structured interviews of residents of Iquitos [67,68].

We modeled three immature mosquito stages (eggs, larvae, and pupae) deterministically at each
location. Development and mortality rates for each of these stages, along with adult mortality rates and
the rate at which pupae emerged as adults, varied daily, and are based on empirically derived

temperature-dependent relationships and weather data from Iquitos [54,69]. Additionally, larval density dependent mortality and an additional rate of death in both the larval and pupal stages are calibrated so that adult mosquito abundance matches the spatio-temporal heterogeneity predicted by a previous study from Iquitos [49]. Adult mosquitoes blood-fed on humans who were co-located with a mosquito during its time of blood-feeding at rates informed by empirical relationships with temperature [69,70]. Each day, mosquitoes moved to adjacent locations based on a probability of 0.3, consistent with another agent-based model of *Ae. aegypti* population dynamics in Iquitos [54].

The transmission of DENV to humans, and to mosquitoes, occurred through mosquito bites. The 589 probability of transmission from humans to mosquitoes was determined by the viremia levels of the 590 591 infecting human at the time of the bite [71]. After completion of a temperature-dependent extrinsic 592 incubation period [72], infectious mosquitoes transmitted DENV to susceptible humans with a fixed 593 probability of 1.0 [55]. Infected humans became infectious and, with a probability informed by 594 empirical studies [73], developed symptoms following a latency period linked to the timing of peak 595 viremia [72]. After recovering from infection, humans gained permanent immunity to the infecting 596 serotype and temporary immunity against heterotypic infections for a period of time. The duration of 597 temporary immunity was exponentially distributed across people with a mean of 686 days, as estimated 598 by a previous model-based analyses of serotype-specific dengue incidence time series [57]. We 599 assigned the initial level of population immunity to each serotype in the population based on estimates 600 by Reiner et al. [58].



602

Supplementary Figure 1: Predicted average mosquito abundance following city-wide ULV spraying.
(a) Comparison of reactive strategies for initiating spraying; spraying began when the monthly or
weekly incidence was one or two standard deviations above the mean for that period from the last five
years, as shown on the x-axis. (b) Comparison of yearly city-wide spraying strategies, beginning on the
first day of the shown month. (c) Comparison of the median predicted cases for twice yearly spraying

- 608 strategies, beginning on the first days of the shown month. Darker colors correspond to fewer cases,
- 609 and the diagonal shows yearly spraying strategies. (d) Comparison of the best strategies in each
- 610 category: adaptive threshold corresponds to starting when weekly incidence was more than two
- 611 standard deviations above the mean, once yearly corresponds to spraying in November, twice yearly
- 612 corresponds to spraying in September and November.



- 614 Supplementary Figure 2: Predicted number of infections following city-wide ULV spraying, by season.
- 615 Each figure compares the best strategies in each category, for that season.



Supplementary Figure 3: Time-series of mosquito abundance for the best TIRS strategy in each
category. In each plot the purple lines represent the predictions without spraying, and the green
represents the given strategy. The line represents the median of all 400 simulations and the shading
represents the inter-quartile range.



Supplementary Figure 4: The effect of varying household compliance and thoroughness of spraying on the median number of infections for the best adaptive threshold strategy with TIRS (spraying when monthly incidence exceeds one standard deviation above the mean). The vertical line shows the value used in the baseline simulations. The solid line represents a fitted multivariable generalized additive model.



Supplementary Figure 5: pie charts showing the proportion of variance in the output that is explained
by variance in the sampled input parameters. Higher-order terms include interactions between

- 629 parameters; in the case of TIRS this is just the interaction between compliance and thoroughness as
- 630 only two parameters were varied.