From frail to fit after allogeneic hematopoietic cell transplantation:

Scope of the problem and a strength-based solution

Jason Sweetnam^{1*}, Eric Twohey^{2*}, Sasha Skendzel³, John Stingle³, Mukta Arora³, Austin Baraki⁴, Qing Cao⁵, Sonya Grillo⁶, Shernan G. Holtan³

- 1. Life Time Fitness, Minneapolis, MN
- 2. University of Minnesota Medical School
- 3. University of Minnesota Blood and Marrow Transplant Program, University of Minnesota, Minneapolis, MN
- 4. Department of Medicine, University of Texas Health Sciences Center, San Antonio, TX
- 5. Bioinformatics and Statistics Core, University of Minnesota
- 6. Best Practices Integrated Informatics Core, University of Minnesota

*Contributed equally

Running title: Frail to fit

Corresponding author:

Shernan G. Holtan, MD Associate Professor of Medicine Blood and Marrow Transplant Program University of Minnesota 420 Delaware Street SE, MMC 480 Minneapolis, MN 55455 Phone: (612) 301-1095 Fax: (612) 625-6919 Email: <u>sgholtan@umn.edu</u>

Abstract: 312 words Manuscript: 2,228 words Tables: 5 Figures: 4 References: 39

Key words: Allogeneic hematopoietic cell transplantation, GVHD, malnutrition, fatigue, weakness, sarcopenia, frailty, strength training

Disclaimers: The authors of this manuscript have no relevant conflicts of interest to disclose.

Abstract

Background: Frailty is a common but underdiagnosed syndrome among long-term survivors of allogeneic hematopoietic cell transplantation (HCT). Conditions such as malnutrition, fatigue, and weakness may suggest frailty despite patients not receiving a formal diagnosis. Furthermore, the optimal interventions to overcome frailty in long-term survivors of allogeneic HCT is not yet established.

Patients and Methods: This study consists of a retrospective and prospective component. First, we completed a retrospective review using diagnosis data from the electronic medical record to estimate the prevalence of components of frailty in 1077 recipients of allogeneic HCT through 5 years post-transplant. Second, we developed a community-based pilot study of strength training for long-term survivors of allogeneic HCT that addressed several common barriers exercise engagement and adherence. Four allogeneic HCT recipients (2 males, 2 females) and 4 controls (2 males, 2 females) completed the strength training pilot study, consisting of a baseline assessment, 10 weeks of personalized and supervised strength programming at least once weekly as a group, and an end-of-study assessment to measure progress in strength, body composition, and a standardized measure of self-efficacy.

<u>Results</u>: Despite a lack of a formal diagnosis, approximately 80% of HCT recipients in this series received a diagnosis of a component of frailty (weakness, fatigue, or malnutrition), and over 1/3 of HCT recipients had symptoms extending beyond 1 year. Over the course of the strength training pilot study, both pre-frail/frail allogeneic HCT recipients and healthy controls doubled their total strength, lost body fat, and gained muscle mass. Self-efficacy improved from baseline in allogeneic HCT recipients after the strength training pilot program.

Discussion: Based upon the prevalence of frailty-related diagnoses among long-term survivors of HCT, a comprehensive mitigation strategy needs to be developed for this population. A community-based strength training program that includes a personalized component, a group setting, and caregiver/partner involvement appears feasible and overcomes several established barriers to exercise engagement and adherence.

Introduction

Frailty is a clinical syndrome of diminished physiologic reserve and low resistance to stressors affecting approximately 10% of the hematopoietic cell transplant (HCT) population.(*1*) Frailty can manifest as weakness, low physical activity, exercise intolerance, unintentional weight loss, self-reported exhaustion, and low-self efficacy.(*2*, *3*) While its manifestations are common and undoubtedly recognized as expected or "normal" to some extent, frailty is likely underdiagnosed in HCT recipients. Conditions such as malnutrition, fatigue, and weakness may suggest frailty despite patients not receiving a formal diagnosis. Because aspects of frailty are so common after HCT, its root causes, including potential anabolic resistance (*4*), may not be adequately addressed by physicians specializing in HCT. Therefore, there is an unmet need to both increase the recognition of frailty by the medical team and to design and test interventions to mitigate it.

The optimal interventions to prevent or mitigate frailty in the allogeneic HCT population are not yet known. However, strength-based training programs directly address multiple domains of frailty and have shown success in other clinical settings, including communitydwelling elderly (5), those residing in long-term care facilities (6), and after lung transplantation (7). We have 2 goals outlined in this paper: (1) estimate the prevalence of components of frailty in 1077 recipients of allogeneic HCT through 5 years post-transplant using diagnosis data from the electronic medical record, and (2) perform a community-based pilot study of strength training in long-term survivors of allogeneic HCT. The results herein are useful to inform the development of a larger strength-focused longitudinal training program to reverse frailty and improve quality of life after allogeneic HCT.

Patients and Methods

We performed this study in two parts. We first performed a retrospective review to estimate frailty from medical diagnosis codes abstracted from the electronic medical record. We subsequently performed a pilot study to improve frailty in cancer survivors. We describe details of each cohort below.

Retrospective Cohort

In order to estimate the prevalence of frailty through 5 years post-HCT, we retrieved both International Classification of Diseases (ICD)9 and ICD10 codes for malnutrition, fatigue, and weakness from our electronic medical record system spanning 3 post-transplant periods: Day 0 – 100, Day 101 – 365, and 1 – 5 years. Patients (N=1077) in this cohort were adults age 18 years or greater undergoing HCT at the University of Minnesota between 2000-2016. We adjusted for mortality by taking into consideration the death rate in each time period. A patient could be counted for a diagnosis of malnutrition, fatigue, and/or weakness in the time period if they survived at all during that time period.

Strength Training Pilot Study

We designed a pilot study of community-based strength training designed to address frailty. Participants (N=4) were required to be at least 18 months post-HCT, meet criteria for pre-frailty or frailty (all had low energy expenditure and self-reported exhaustion), and have a healthy exercise partner (spouse or other relative or friend) who could serve as a control (N=4). Participants in our pilot study received an assessment by a National Academy of Sports Medicine (NASM)-certified trainer prior to design of a personalized strength training program during week 1. During weeks 2-11, each participant completed their personalized program at least weekly as a group under the supervision of the trainer on Saturday mornings, and they were encouraged but not required to complete additional training sessions as individuals during

the other days of the week. These additional training sessions were unsupervised but followed the same protocol as the supervised sessions. Each participants' personalized strength training program consisted of three lower body exercises, four upper body exercises, three core exercises, and one aerobic test (example provided in Figure 1). Prescriptions for all exercises were determined by the number of repetitions at a given weight required to approach muscular failure in the initial assessment. The large majority of exercises were performed on Free Motion® machines. These machines have a rotating cable system so that individuals with different body types and proportions can use the same machines without being limited to a fixed motor pattern or range of motion.

During the course of the study, specific exercises and their loads were progressed based on the abilities of each participant. In assessing the progress of individuals during week 12 (final week of study), the estimate of improvement was based on the change in relative strength. This metric consists of changes in amount of weight lifted (absolute strength) and the progress in the amount of repetitions completed (work capacity). All of the participants' workouts (both those with and without a trainer) were recorded and collected, and the final results were calculated by comparing subject's initial assessments with their final workouts. To assess progress over the trial period, we compared relative strength in each of the muscle groups comparing the initial and final sessions using descriptive statistics (average, range) without statistical tests due to the small sample size.

Body Composition

We recorded body composition, including weight, percentage of body fat, and weight of muscle mass, at the initial and final sessions using an InBody® bioelectrical impedance device. Participants were counseled that the goal of the study was not to lose weight but rather to gain muscle mass. No formal dietary intervention was given, but participants learned about healthy

eating habits with a particular focus on eating for muscular growth from a registered dietician once per month in a group setting.

Self-Efficacy Assessment

The primary endpoint of the strength training pilot study was self-efficacy as determined by a 6-question survey adapted from the Stanford Patient Education Resource Center (8). Participants gave a numeric response on a scale of 1 (not confident at all) to 10 (totally confident) on questions regarding confidence to deal with fatigue, physical discomfort, emotional distress, managing health problems, reducing dependence on the healthcare system, and taking steps to improve health beyond medications.

Ethical Considerations

The University of Minnesota Institutional Review Board approved both the retrospective medical record review and the strength training pilot study. The strength training pilot study is registered on ClinicalTrials.gov as NCT03609203.

Results

Estimation of frailty from diagnosis codes

Demographics of the retrospective cohort are detailed in Table 1. The median age of this cohort was 52 years, with a slight male predominance (58.6% of patients). Table 2 lists the overall frequency of diagnoses of weakness, malnutrition, and fatigue over the period of 2000-2016, and Figure 2 shows how these diagnoses overlap in individual patients across early, intermediate, and late post-HCT recovery. The majority (79.3%) of adult allogeneic HCT recipients had a diagnosis of malnutrition (Table 2), weakness (Table 3), and/or fatigue (Table 4) at some time post-HCT. Only 179 patients (21.7%) had none of these diagnoses at any time point in their post-transplant recovery. Across the 3 time periods, fatigue was the most frequent

diagnosis, with 37% diagnosed with fatigue prior to day 100, 28% diagnosed between days 101 and 365, and 33% diagnosed between years 1-5. The frequency of malnutrition diagnoses decreased over time, with 22% prior to day 100, 18% between days 101-365, and 16% between years 1-5. In contrast, the frequency of muscle weakness diagnoses increased over time, with 6% diagnosed prior to day 100, 9% between days 101 - 365, and 11% between years 1-5. Patients with all 3 diagnoses ranged from 1.8% in the first 100 days, to 3.9% between days 101 - 365, to 4.2% between years 1-5.

Strength Training Pilot Study Outcomes

Four adult allogeneic HCT recipients (2 males, 2 females) and 4 controls (2 males, 2 females) completed the strength training pilot study, consisting of a baseline assessment, 10 weeks of personalized and supervised strength programming at least once weekly as a group, and an end-of-study assessment to measure progress. Participant engagement was excellent, there were no serious adverse events related to strength training. The average number of strength training sessions completed over 10 weeks by participant was 27 (range 17-34). Study participants doubled their strength on average after 10 weeks, as demonstrated by improvements of 194% in upper body exercises, and 199% in lower body exercises.

While this was not designed as a weight loss study, most participants lost fat mass. Average weight loss was 3.0 kilograms over 10 weeks, which reflected in an average change in body fat of 2.54% as assessed by bioelectrical impedance among study participants (Figure 3B). The average muscle gain as assessed by bioelectrical impedance in allogeneic HCT recipients was 1.4 kilograms throughout the study, with no net change in the control group (Figure 3C). However, one female outlier the healthy control group lost a total of 10.8 kg during the study.

We measured self-efficacy via 6-item survey at the start of the study and at its completion. Baseline self-efficacy was lower in recipients of allogeneic HCT than controls

(Figure 4), but increased by an average of 10 points (mean 36.75 to 46.75) in allogeneic HCT recipients over the course of the study.

Discussion

Frailty is a commonly observed syndrome in the adult HCT population, although it is typically not diagnosed formally. Frailty does not have its own ICD-10 code; rather, the code R54 refers to "age-related physical debility" that includes frailty, but excludes sarcopenia and cognitive decline. Despite not receiving a formal diagnosis of frailty, approximately 80% of HCT recipients in this series received a diagnosis of a component of frailty (weakness, fatigue, or malnutrition), and over 1/3 of HCT recipients reported symptoms extending beyond 1 year. Sarcopenia affects 50% of HCT recipients even prior to HCT leading to an increased risk of late effects, poor quality of life, and early mortality.(*9*) Knowing this, and observing the frequency of frailty-related diagnoses in our series, a comprehensive mitigation strategy needs to be developed for long-term HCT survivors.

Recipients of HCT can carry several risk factors for frailty including advanced age, polypharmacy, lack of regular exercise, isolation, malnutrition, and unintentional weight loss (1, 10). Risks traditionally associated with life-threatening complications following SCT have been substantially mitigated by innovations in supportive care.(11) As a result, there has been a considerable increase in the number of SCT recipients living with the systemic effects of cancer and related treatment as a chronic condition.(12) In response, health organizations are now recommending survivorship interventions founded on the theoretical principles of the Chronic Care Model with a focus on programs targeting chronic-disease self-management.(13) Improving patient self-management behavior in the survivorship period has been the focus of multiple chronic disease programs.(14-17) Exercise has well-established beneficial effects including improved quality-of-life and a reduction in untoward cancer-related physical symptoms (18) and should thus be a key component of such self-management programs.

Perhaps the strongest predictor of health-related self-management behavior is selfefficacy.(19) As defined by Bandura, self-efficacy is the perceived ability or confidence to achieve a goal, which can be influenced by motivation and task mastery.(20) Increased selfefficacy among cancer survivors appears to have a relationship with exercise participation and adherence, with higher levels of self-efficacy associated with more exercise behaviors and vice versa.(19, 21) Understanding the critical role of this phenomenon in promoting and sustaining positive behavior change, we used the validated Chronic Disease Self-Management 6-Item scale to assess patient self-efficacy in this pilot study.(8) The scale measures several common domains of chronic disease including the management of symptoms and health related activities. Our pilot study found improvements in self-efficacy after the strength training intervention, but the overall level of self-efficacy still did not match that of healthy control participants. Self-efficacy may have continued to improve further with a longer study duration.

Our pilot study is one among many other larger studies that have investigated the role of exercise in recovery from HCT (Table 5). The results from previous interventions may be variable if the training stimulus was not personalized; either too small to induce muscular growth, or too large, putting individuals at risk of injury, excess fatigue, or pain. Factors that enhanced the feasibility of our pilot study is that the intervention was personalized and occurred in a commercial health club. While the participants were required to meet at least once weekly as a group at one of the club's locations, the participants were able to exercise at a club closer to their home (if applicable) during the week. The required group participation also enhanced the feasibility and reduced expense, allowing the trainer to supervise 8 participants at once during the weekly sessions. The group participation also likely enhanced the sense of community among participants, as some remarked during the study that they felt less isolated than they had felt prior to starting the program. Finally, the partnering of healthy controls (caregiver, spouse, or other relative or friend) with the HCT recipients may have enhanced adherence to the program. We also wanted to involve caregivers in the intervention recognizing

the impact of the HCT process on their lives.(22) We did not want to give them yet one more task to add to their list. Furthermore, involving the caregiver could impact the entire family dynamic toward adopting healthier lifestyle behaviors despite many perceived barriers when a loved one is going through the allogeneic HCT process.

In conclusion, frailty is a clinically recognized but underdiagnosed syndrome after allogeneic HCT. A personalized strength training intervention, conducted along with a caregiver in a community setting, can help address some of the root causes of frailty, increasing lean muscle mass and improving self-efficacy. Larger studies are needed to refine our communitybased strength training program for allogeneic HCT survivors and their caregivers prior to widespread implementation.

Acknowledgement of Research Support:

This study was funded by the University Of Minnesota Department Of Medicine Women's Early Research Career program (SGH) and Marrow on the Move, University of Minnesota (SGH).

 Table 1. Retrospective cohort patient demographics.

	N=1077
AGE	
N of Observed	1077
Median(Min-Max)	52.2(18.1-75.1)
SEX	
Male	631(58.6%)
Female	446(41.4%)
DISEASE	
Aplastic anemia	14(1.3%)
Metabolic disorder	2(0.2%)
Acute lymphoblastic leukemia	148(13.7%)
Acute myeloid leukemia	439(40.8%)
Chronic myeloid leukemia	41(3.8%)
Other Leukemia	13(1.2%)
Myelodysplasia	132(12.3%)
Non-Hodgkin lymphoma	140(13.0%)
Hodgkin lymphoma	45(4.2%)
Myeloproliferative neoplasms	19(1.8%)
Other Malignancy	14(1.3%)
Multiple myeloma	41(3.8%)
Chronic lymphocytic leukemia	29(2.7%)
CONDITIONING	
Myeloablative	386 (35.8%)
Reduced Intensity/Non- myeloablative	686 (63.7%)
Other	5 (<1%)
GRAFT SOURCE	
Marrow	102 (9.5%)

PBSC	459 (42.5%)
Cord	517 (48.0%)
Acute graft-versus-host disease II- IV by day 180	446 (41.4%)
Acute graft-versus-host disease III- IV by day 180	215 (20.0%)
Chronic graft-versus-host disease by day 365	268 (24.9%)

Diagnosis Code	Code Type	Number of times diagnosed	Description
R63.0	ICD10	523	Anorexia
783.0	ICD9	443	Anorexia
783.21	ICD9	236	Loss of weight
R62.7	ICD10	163	Adult failure to thrive
R63.0	ICD10	150	Anorexia
783.7	ICD9	120	Adult failure to thrive
783.21	ICD9	110	Loss of weight
783.7	ICD9	98	Adult failure to thrive
783.0	ICD9	92	Anorexia
			Feeding difficulties and
783.3	ICD9	91	mismanagement
R63.3	ICD10	55	Feeding difficulties
R62.7	ICD10	55	Adult failure to thrive
700.4		45	Body mass index (BMI) 19 or
208.1		45	
/ 03./ DC4		30	FAILURE TO THRIVE-ADULT
R04		29	
799.4	ICD9	28	Moderate protein-calorie
E44.0	ICD10	25	malnutrition
263.0	ICD9	16	Malnutrition of moderate degree
R63.3	ICD10	16	Feeding difficulties
V85.0		16	Body Mass Index less than 19,
F44 1		10	Mild protein-calorie malnutrition
			Feeding difficulties and
783.3	ICD9	9	mismanagement
			Moderate protein-calorie
E44.0	ICD10	6	malnutrition
263.1	ICD9	6	Malnutrition of mild degree
263.0	ICD9	4	Malnutrition of moderate degree
700.4			Body mass index (BMI) 19 or
<u>268.1</u>	ICD10	4	less, adult
R62.51		3	Failure to thrive (child)
E44.1		2	IVIIIG protein-calorie mainutrition
/83.41		2	
700.4		1	
799.4		1	
K64	ICD10	1	Cachexia

Table 2. Frequency of malnutrition diagnoses (raw data) abstracted from electronic medical record.

Table 3. Frequency of weakness diagnoses (raw data) abstracted from electronic medical record.

Diagnosis Code	Code Type	Number of times diagnosed	Description
728.87	ICD9	445	Muscle weakness (generalized)
728.87	ICD9	377	MUSCLE WEAKNESS-GENERAL
M62.81	ICD10	223	Muscle weakness (generalized)
M62.81	ICD10	123	Muscle weakness (generalized)
728.87	ICD9	118	Muscle weakness (generalized)
M62.50	ICD10	59	Muscle wasting and atrophy, not elsewhere classified, unspecified site
M62.50	ICD10	34	Muscle wasting and atrophy, not elsewhere classified, unspecified site
M62.59	ICD10	12	Muscle wasting and atrophy, not elsewhere classified, multiple sites
M62.562	ICD10	12	Muscle wasting and atrophy, not elsewhere classified, left lower leg
M62.561	ICD10	10	Muscle wasting and atrophy, not elsewhere classified, right lower leg
M62.58	ICD10	9	Muscle wasting and atrophy, not elsewhere classified, other site
M62.552	ICD10	5	Muscle wasting and atrophy, not elsewhere classified, left thigh
M62.569	ICD10	2	Muscle wasting and atrophy, not elsewhere classified, unspecified lower leg

Diagnosis Code	Code Type	Number of times diagnosed	Description
780.79	ICD9	2488	Other malaise and fatigue
780.79	ICD9	1398	Other malaise and fatigue
780.79	ICD9	817	MALAISE AND FATIGUE NEC
R53.83	ICD10	699	Other fatigue
R53.83	ICD10	340	Other fatigue
R53.82	ICD10	25	Chronic fatigue, unspecified
780.79	ICD9	10	MALAISE AND FATIGUE NEC
R53.82	ICD10	6	Chronic fatigue, unspecified
780.54	ICD9	5	Hypersomnia, unspecified
780.54	ICD9	5	Hypersomnia, unspecified
780.53	ICD9	4	HYPERSOM W SLP APNEA NOS
780.71	ICD9	3	Chronic fatigue syndrome
780.71	ICD9	3	Chronic fatigue syndrome
G47.10	ICD10	2	Hypersomnia, unspecified
780.54	ICD9	2	HYPERSOMNIA NOS
780.79	ICD9	1	MALAISE AND FATIGUE NEC
G47.10	ICD10	1	Hypersomnia, unspecified
780.54	ICD9	1	HYPERSOMNIA NOS

Table 5. Prospective studies involving strength training in blood and marrow transplantation.

Study Design	N	Age	Interventio n	Primary Endpoints	Results of Primary Endpoints	Secondary endpoints	Results of Secondary Endpoints	Reference s
Randomized pilot study	19	Adult (ages 20-67)	Strength	Physical activity, muscle strength, fatigue, health status perceptions, quality of life	Significant decrease in fatigue with exercise group, no significant effects on all other primary outcomes	Not applicable	Not applicable	(23)
Non- randomized	30	Adolescent s and adults	Strength	Muscle protein status and turnover- nitrogen balance, creatinine, 3- methylhistidine	Muscle protein sparing effect of exercise, significant decrease in creatinine excretion in controls only suggested muscle protein loss associated w/ inactivity	Arm muscle area	Correlated with energy but not protein intake	(24)

[1				1		
Randomized	53	Pediatric (ages 5-17)	Aerobic and Strength	6-minute walk test (6MWT)	Intervention unfit group increased 6MWT, while unfit and fit control groups decreased.	Exercise frequency and intensity	Intervention fit group presented higher exercise frequency and intensity compared to intervention unfit	(25)
Randomized	42	Adult (ages 18-60)	Aerobic and Strength	VO2max	Significant increase	Muscle strength, functional performance, physical activity level, QOL, fatigue, psychological well-being and clinical outcomes	Significant effect on muscle strength: chest press, leg extension, right elbow flexor, right knee flexor, right knee extensor, and functional performance (stair test)	(26)
Randomized	18 7	Adults (ages 25- 76)	Aerobic and strength in the context of Total Therapy for multiple myeloma	Fatigue, nighttime sleep, aerobic capacity	No statistical significance in nighttime sleep, 6MWT, or fatigue between groups	Not applicable	Not applicable	(27)

Randomized	61	Adults (ages 22- 70)	Aerobic and Strength	Performance (50ft walk, 6MWT, functional reach, sit-to-stand, uniped stance)	No statistical significance found between supervised vs. unsupervised exercise groups	Brief Fatigue Inventory	No statistical significance	(28)
Randomized	13 5	Adults (ages 32- 74)	Aerobic and Strength	# of transfusions during transplantation, # of attempts and total number of days of stem cell collection, time to recovery after transplant, response to intensive therapy	Recovery and treatment response not significantly different. Exercise group had significantly fewer RBC transfusions and fewer attempts at stem cell collection	Not applicable	Not applicable	(29)
Randomized	24	Adults (ages 42- 74)	Aerobic and Strength	Feasibility of home exercise program (injuries, adherence)	No injuries in either exercise or control group	Lean Body Weight, muscle strength, fatigue, mood disturbances, sleep time	Increased lean body weight in exercise group (+0.40kg /month vs 0.44kg /month in control)	(30)

Randomized	42	Adults (ages 18- 65)	Aerobic and Strength	Severity of 5 symptoms clusters: gastrointestinal, cognitive, affective, functional, mucositis	Intervention group mean symptom severity scores were significantly lower in all categories except for affective	Symptom burden	Confirmed that patients undergoing allo- HCT experience a considerable symptom burden	(31)
Randomized	13 1	Adult (ages 18-75)	Aerobic and Strength	Physical performance and self-reported physical function	Statistically significant increase in knee extension, walking speed, and 6MWT were found in the exercise group. Grip strength was non- significant	Body composition, Quantified walking activity, self- reported physical activity, fatigue and QOL	No statistical significance	(32)
Multicenter randomized	10 5	Adult (ages 18-71)	Aerobic and strength	Fatigue	Up to 15% improvement in exercise group vs up to 28% worsening in control	Physical functioning, anxiety, global distress, 6MWT	Exercise group was found to have significantly higher physical functioning, improved anxiety and 6MWT. Decreased global distress	(33)

Randomized	67	Adult (ages 19-73)	Strength	Muscle strength, fatigue, functional ability	Fatigue: significant improvement in intervention group Functional ability: significant increase in timed stair climb & timed up and go Muscle strength: no significant increase, but trend toward improving muscle strength in intervention group	Cross-section area of R. femoris, objective physical activity, quality of life	Cross-section of R. femoris: no significance, in quality of life: no significant difference Physical activity: significant increase in objective physical activity in intervention group	(34)
Multicenter randomized	10 9	Adults (ages 19- 67)	Aerobic and strength	Physical Fitness: VO2peak, Wpeak, grip strength, 30s chair stand test	No significant benefits in intervention group on physical fitness and fatigue	BMI, sum of 4 skinfolds, max isometric torque of M. quadriceps, HRQOL, symptoms scales of EORTC Myeloma module, distress via HADS questionnaire, subjective and objective physical activity	No significant effects found in secondary outcome measures	(35)

Randomized	20	Pediatric (ages 4-16)	Aerobic and strength	Immune cell recovery	No difference between groups	Body composition (weight, BMI, Body fat, FFM)	Significant effect across all anthropometric measures (i.e. intervention group increased BMI)	(36)
Randomized	12	Adults (ages 16- 64)	Aerobic and strength	QOL (CARES questionnaire)	Improved QOL in exercise group when compared to pre-transplant and non- exercise group	Peak aerobic capacity	Peak aerobic capacity and QOL were correlated (p<.05)	(37)
Prospective, repeated measures design	10	Adults (ages 48- 71)	Strength	Feasibility of exercise based on week to week exercise frequency of participates	Over 6 week period, number of patients exercising 3+ times a week went from 2/10 to 6/10	Strength (hand grip strength, time needed to stand up, and 30 second chair to stand test). General Health Perceptions via EORTC QLQ-C30	Strength: hand grip strength (32.66+/-12.34 to 28.18 +/- 12.42); time needed to stand 4.63+/-1.65 to 4.05+/-1.5; 7.6+/-3.1 to 8.22+/-3.42; QOL: 23.51+/- 3.65 to 24+/- 3.88	(38)
Prospective feasibility study	16	Adults (ages 60- 70)	Aerobic and Strength	More than 50% of patients complete more than 50% of scheduled exercise days during hospitalization	only 15% of scheduled sessions missed by patients	Strength: isometric knee extensor strength, trunk strength; Endurance: 6MWT; QOL; Fatigue; Depression	Equal lower extremity strength, negative trend in endurance capacity, mixed regard in trunk strength	(39)

Figures

Figure 1. Example strength training prescription and training log.

Workout Session #1					Da	ate:				
Stretching	Perform Before Every Exercise Session									
Foam Roll Legs (Front, Side)										
Foam Roll Calves										
Kneeling Hip Flexor Stretch										
Banded Lateral Walking										
Warm up	Get your body and mind ready for exercise by using a piece of cardio equipment that feels good to you for 5 minutes: stationary bike, tread mill, or elliptical									
	Seat Height and	Set	Set#1 Set#2		#2	#2 Se		Average		
Lower Body	Machine Settings	Weight	Reps	Weight	Reps	Weight	Reps	Weight	Reps	
Free Motion Squat Machine							_			
Quad Extension Machine										
Seated Hamstring Curl										
	Seat Height and	Set	:#1	Set#2		Set #3		Average		
Upper Body	Machine Settings	Weight	Reps	Weight	Reps	Weight	Reps	Weight	Reps	
Free Motion Chest Press										
Free Motion Overhead Press										
Free Motion Row					L					
Free Motion Lat Pull-Down			l		ļ					
			L							
		Set	:#1	Set #2		Set #3		Average		
Glutes and Core	Notes:		Reps		Reps		Reps		Reps	
Glute Bridge	Hold At Top									
Dead Bug	Keep Abs Flexed		L		L			'	L	
Plank	Record In Seconds									
			<u> </u>		<u> </u>				-	
1 Minute Challenge	Max Meters in 1 Minute on Rope Climb Machine. Distance Completed Today:				Last Week:					
This program is personalized for you.	The goal is to complete betwee repetitions with excellent form become MORE capable.	n 10 - 15 repeti at your assigne	tions on each d weight. Eve	⊧exercise with e ∍ry week we are	xce∥entform. trγing to do a	. Let Jason knov LITT LE bit more	w if you can eas e to encourage	ily do more tha our body to ada	n 15 1pt, grow, and	

Figure 2. Number of patients with overlapping diagnoses of malnutrition, fatigue, and muscle weakness through 5 years post-allogeneic hematopoietic cell transplantation.



Figure 3. Comparison of percent change in (A) percent total strength compared to baseline, (B) percent body fat, (C) and lean body weight in kilograms between allogeneic hematopoietic cell transplant recipients (Allo, N=4) versus healthy controls (N=4).



Figure 4. Self-efficacy at the start of the strength training intervention and at its completion (maximum score = 60 points) comparing recipients of allogeneic hematopoietic cell transplantation (gray) to controls (black).



REFERENCES

- 1. M. Arora *et al.*, Physiologic Frailty in Nonelderly Hematopoietic Cell Transplantation Patients: Results From the Bone Marrow Transplant Survivor Study. *JAMA Oncol* **2**, 1277-1286 (2016).
- 2. L. P. Fried *et al.*, Frailty in older adults: evidence for a phenotype. *J Gerontol A Biol Sci Med Sci* **56**, M146-156 (2001).
- 3. M. D. Hladek *et al.*, High coping self-efficacy associated with lower odds of prefrailty/frailty in older adults with chronic disease. *Aging Ment Health*, 1-7 (2019).
- 4. M. P. Engelen, B. S. van der Meij, N. E. Deutz, Protein anabolic resistance in cancer: does it really exist? *Curr Opin Clin Nutr Metab Care* **19**, 39-47 (2016).
- 5. E. F. Binder *et al.*, Effects of exercise training on frailty in community-dwelling older adults: results of a randomized, controlled trial. *J Am Geriatr Soc* **50**, 1921-1928 (2002).
- 6. C. B. Ferreira *et al.*, Effects of a 12-Week Exercise Training Program on Physical Function in Institutionalized Frail Elderly. *J Aging Res* **2018**, 7218102 (2018).
- 7. D. Langer *et al.*, Exercise training after lung transplantation improves participation in daily activity: a randomized controlled trial. *Am J Transplant* **12**, 1584-1592 (2012).
- 8. K. R. Lorig, D. S. Sobel, P. L. Ritter, D. Laurent, M. Hobbs, Effect of a self-management program on patients with chronic disease. *Eff Clin Pract* **4**, 256-262 (2001).
- 9. S. Morishita *et al.*, Prevalence of sarcopenia and relevance of body composition, physiological function, fatigue, and health-related quality of life in patients before allogeneic hematopoietic stem cell transplantation. *Support Care Cancer* **20**, 3161-3168 (2012).
- 10. M. Pamukcuoglu *et al.*, Hematopoietic Cell Transplant-Related Toxicities and Mortality in Frail Recipients. *Biol Blood Marrow Transplant*, (2019).
- 11. T. A. Gooley *et al.*, Reduced mortality after allogeneic hematopoietic-cell transplantation. *N Engl J Med* **363**, 2091-2101 (2010).
- 12. S. Hashmi, P. Carpenter, N. Khera, A. Tichelli, B. N. Savani, Lost in transition: the essential need for long-term follow-up clinic for blood and marrow transplantation survivors. *Biol Blood Marrow Transplant* **21**, 225-232 (2015).
- 13. E. H. Wagner, Chronic disease management: what will it take to improve care for chronic illness? *Eff Clin Pract* **1**, 2-4 (1998).
- 14. T. M. Damush, A. Perkins, K. Miller, The implementation of an oncologist referred, exercise self-management program for older breast cancer survivors. *Psychooncology* **15**, 884-890 (2006).
- 15. S. Lawn *et al.*, Is self-management feasible and acceptable for addressing nutrition and physical activity needs of cancer survivors? *Health Expect* **18**, 3358-3373 (2015).
- 16. R. McCorkle *et al.*, Self-management: Enabling and empowering patients living with cancer as a chronic illness. *CA Cancer J Clin* **61**, 50-62 (2011).
- 17. B. C. Risendal *et al.*, Meeting the challenge of cancer survivorship in public health: results from the evaluation of the chronic disease self-management program for cancer survivors. *Front Public Health* **2**, 214 (2014).
- 18. S. I. Mishra *et al.*, Exercise interventions on health-related quality of life for people with cancer during active treatment. *Clin Otolaryngol* **37**, 390-392 (2012).
- 19. S. D. Neupert, M. E. Lachman, S. B. Whitbourne, Exercise self-efficacy and control beliefs: effects on exercise behavior after an exercise intervention for older adults. *J Aging Phys Act* **17**, 1-16 (2009).
- 20. A. Bandura, Health promotion by social cognitive means. *Health Educ Behav* **31**, 143-164 (2004).

- 21. S. S. Daniali, F. M. Darani, A. A. Eslami, M. Mazaheri, Relationship between Selfefficacy and Physical Activity, Medication Adherence in Chronic Disease Patients. *Adv Biomed Res* **6**, 63 (2017).
- 22. A. J. Applebaum *et al.*, A scoping review of caregiver burden during allogeneic HSCT: lessons learned and future directions. *Bone Marrow Transplant* **51**, 1416-1422 (2016).
- 23. E. D. Hacker *et al.*, Strength training following hematopoietic stem cell transplantation. *Cancer Nurs* **34**, 238-249 (2011).
- 24. B. A. Cunningham *et al.*, Effects of resistive exercise on skeletal muscle in marrow transplant recipients receiving total parenteral nutrition. *JPEN J Parenter Enteral Nutr* **10**, 558-563 (1986).
- 25. S. Wallek *et al.*, Impact of the initial fitness level on the effects of a structured exercise therapy during pediatric stem cell transplantation. *Pediatr Blood Cancer* **65**, (2018).
- 26. M. Jarden, M. T. Baadsgaard, D. J. Hovgaard, E. Boesen, L. Adamsen, A randomized trial on the effect of a multimodal intervention on physical capacity, functional performance and quality of life in adult patients undergoing allogeneic SCT. *Bone Marrow Transplant* **43**, 725-737 (2009).
- 27. E. A. Coleman *et al.*, Effects of exercise on fatigue, sleep, and performance: a randomized trial. *Oncol Nurs Forum* **39**, 468-477 (2012).
- 28. M. L. Shelton *et al.*, A randomized control trial of a supervised versus a self-directed exercise program for allogeneic stem cell transplant patients. *Psychooncology* **18**, 353-359 (2009).
- 29. E. A. Coleman *et al.*, Effects of exercise in combination with epoetin alfa during highdose chemotherapy and autologous peripheral blood stem cell transplantation for multiple myeloma. *Oncol Nurs Forum* **35**, E53-61 (2008).
- 30. E. A. Coleman *et al.*, Feasibility of exercise during treatment for multiple myeloma. *Cancer Nurs* **26**, 410-419 (2003).
- 31. M. Jarden, K. Nelausen, D. Hovgaard, E. Boesen, L. Adamsen, The effect of a multimodal intervention on treatment-related symptoms in patients undergoing hematopoietic stem cell transplantation: a randomized controlled trial. *J Pain Symptom Manage* **38**, 174-190 (2009).
- 32. R. H. Knols *et al.*, Effects of an outpatient physical exercise program on hematopoietic stem-cell transplantation recipients: a randomized clinical trial. *Bone Marrow Transplant* **46**, 1245-1255 (2011).
- 33. J. Wiskemann *et al.*, Effects of a partly self-administered exercise program before, during, and after allogeneic stem cell transplantation. *Blood* **117**, 2604-2613 (2011).
- 34. E. D. Hacker *et al.*, Strength Training to Enhance Early Recovery after Hematopoietic Stem Cell Transplantation. *Biol Blood Marrow Transplant* **23**, 659-669 (2017).
- 35. S. Persoon *et al.*, Randomized controlled trial on the effects of a supervised high intensity exercise program in patients with a hematologic malignancy treated with autologous stem cell transplantation: Results from the EXIST study. *PLoS One* **12**, e0181313 (2017).
- 36. C. Chamorro-Vina *et al.*, Exercise during hematopoietic stem cell transplant hospitalization in children. *Med Sci Sports Exerc* **42**, 1045-1053 (2010).
- 37. S. Hayes, P. S. Davies, T. Parker, J. Bashford, B. Newman, Quality of life changes following peripheral blood stem cell transplantation and participation in a mixed-type, moderate-intensity, exercise program. *Bone Marrow Transplant* **33**, 553-558 (2004).
- 38. E. D. Hacker, J. L. Larson, D. Peace, Exercise in patients receiving hematopoietic stem cell transplantation: lessons learned and results from a feasibility study. *Oncol Nurs Forum* **38**, 216-223 (2011).

39. M. K. Schuler *et al.*, Feasibility of an exercise programme in elderly patients undergoing allogeneic stem cell transplantation - a pilot study. *Eur J Cancer Care (Engl)* **25**, 839-848 (2016).

Workout Session #1							Date:				
Stretching	Perform Before Every Exercise Session										
Foam Roll Legs (Front, Side)					. ,						
Foam Roll Calves											
Kneeling Hip Flexor Stretch											
Banded Lateral Walking											
Warm up	Get your body and mind ready for exercise by using a piece of cardio equipment that feels good to you for 5 minutes: stationary bike, treadmill, or elliptical										
	Seat Height and	Set	#1	Set #2		Set #3		Average			
Lower Body	Machine Settings	Weight	Reps	Weight	Reps	Weight	Reps	Weight	Reps		
Free Motion Squat Machine											
Quad Extension Machine											
Seated Hamstring Curl											
	Seat Height and	Set	#1	Set #2		Set #3		Average			
Upper Body	Machine Settings	Weight	Reps	Weight	Reps	Weight	Reps	Weight	Reps		
Free Motion Chest Press											
Free Motion Overhead Press											
Free Motion Row											
Free Motion Lat Pull-Down											
		Cot #1 Cot #3				Sot #3		Δνοτοσο			
Clutes and Core	Notos	Set #1		Set	#Z	Set	Set #5		Average		
Clute Bridge	Hold At Tan		Reps		ĸeps		Reps		керѕ		
Glute Bridge	Hold At Top										
Dead bug	Reep Abs Flexed										
	Record in Seconds										
1 Minute Challenge	Max Meters in 1 Minute on Rope Climb Machine. Distance I Completed Today:					Last Week:					
This program is personalized for you.	The goal is to complete betwee repetitions with excellent form become MORE capable.	n 10 - 15 repetit at your assigned	tions on each d weight. Eve	exercise with the	xcellent form. trying to do a	Let Jason know LITTLE bit more	/ if you can ea to encourage	sily do more than our body to ada	n 15 Ipt, grow, and		





Self-Efficacy

