1 Title

- 2 Differential asthma risk following respiratory infection in children from three minority
- 3 populations

4 Short Title

5 Asthma risk following respiratory infection in minority children

6 Authors

- 7 Eric M. Wohlford¹, Luisa N. Borrell^{2*}, Jennifer R. Elhawary¹, Brian Plotkin¹, Sam S. Oh¹,
- 8 Thomas J. Nuckton¹, Celeste Eng¹, Sandra Salazar¹, Michael A. LeNoir³, Kelley
- 9 Meade⁴, Harold J. Farber⁵, Denise Serebrisky⁶, Emerita Brigino-Buenaventura⁷, William
- 10 Rodriguez-Cintron⁸, Rajesh Kumar⁹, Shannon Thyne¹⁰, Max A. Seibold¹¹, José R.
- 11 Rodriguez-Santana¹², Esteban G. Burchard^{1,13}

12 Author affiliations

- 13 1 Department of Medicine, University of California, San Francisco, CA, USA.
- 14 2 Department of Epidemiology & Biostatistics, Graduate School of Public Health &
- 15 Health Policy, City University of New York, New York, NY, USA.
- 16 3 Bay Area Pediatrics, Oakland, CA, USA.
- 17 4 Children's Hospital and Research Center Oakland, Oakland, CA, USA.
- 18 5 Department of Pediatrics, Section of Pulmonology, Baylor College of Medicine and
- 19 Texas Children's Hospital, Houston, TX, USA.
- 20 6 Pediatric Pulmonary Division, Jacobi Medical Center, Bronx, NY, USA.

- 21 7 Department of Allergy and Immunology, Kaiser Permanente-Vallejo Medical Center,
- 22 Vallejo, CA, USA.
- 23 8 Veterans Caribbean Health Care System, San Juan, Puerto Rico, USA.
- 9 The Ann and Robert H. Lurie Children's Hospital of Chicago, Chicago, IL, USA.
- 10 Department of Pediatrics, University of California, Los Angeles, Los Angeles, CA,
- 26 USA.
- 11 National Jewish Health, Denver, CO, USA.
- 12 Centro de Neumologia Pediatrica, Caguas, Puerto Rico, USA.
- 13 Department of Bioengineering and Therapeutic Sciences, University of California,
- 30 San Francisco, CA, USA.

*** Corresponding author**

32 Email: Luisa.Borrell@sph.cuny.edu (LNB)

33 Author Contributions

EGB conceived of and designed the SAGE and GALA II studies. SSO and JRE carried
out formal analyses. SSO and EGB provided supervision and mentorship. SSO, JRE,
BP, LNB, TJN, and EMW wrote the manuscript. LNB carried out epidemiological
analysis. CE and SS oversaw all data curation. MAL, KM, HJF, DS, EB-B, WR-C, RK,
ST, MAS, and JRR-S were all responsible in the provision of patients, samples, and
instrumentation for the SAGE and GALA II studies. All authors reviewed, edited, and
approved of the final manuscript.

41 Conflict of interest statement

42 The authors declare that they have no relevant conflicts of interest.

43 Funding Sources

- 44 This work was supported in part by the Sandler Family Foundation, the American
- 45 Asthma Foundation, the RWJF Amos Medical Faculty Development Program, Harry
- 46 Wm. and Diana V. Hind Distinguished Professor in Pharmaceutical Sciences II, National
- 47 Institutes of Health R01HL117004, R01HL128439, R01HL135156, 1X01HL134589,
- 48 R01HL141992, 5T32GM007546, National Institute of Health and Environmental Health
- 49 Sciences R01ES015794, R21ES24844, the National Institute on Minority Health and
- 50 Health Disparities P60MD006902 RL5GM118984, R01MD010443 and the Tobacco-
- 51 Related Disease Research Program under Award Number 24RT-0025, 27IR-0030

52 Word count

53 2,621

54 Abstract

55 Rationale

- 56 Severe early-life respiratory illnesses, particularly those caused by respiratory syncytial
- virus (RSV) and human rhinovirus (HRV), are strongly associated with the development
- of asthma in children. Puerto Rican children in particular have a strikingly high asthma
- 59 burden, but prior studies of the potential associations between early-life respiratory
- 60 illnesses and asthma in Puerto Rican and other minority populations have been limited.

61 **Objectives**

- 62 We sought to determine whether early-life respiratory illness was associated with
- asthma in Puerto Rican children relative to other minority children.

64 Methods

Using a logistic regression analysis, we examined the association between early-life respiratory illnesses (report of upper respiratory infection (URI), pneumonia, bronchitis, and bronchiolitis/RSV) within the first two years of life and physician-diagnosed asthma after the age of two in a large cohort of minority children.

69 Measurements and Main Results

70 Early-life respiratory illnesses were associated with greater asthma risk in Puerto

- 71 Ricans relative to other racial/ethnic minority populations. Specifically, in Puerto Ricans,
- the odds was 6.15 (95% CI: 4.21-9.05) if the child reported at least one of the following

- respiratory illness: URI, pneumonia, bronchitis or bronchiolitis. The odds were also
- ⁷⁴ higher in Puerto Ricans when considering these conditions separately.

75 **Conclusions**

- 76 We observe population-specific associations between early-life respiratory illnesses and
- asthma, which was especially significant in Puerto Ricans. Taken together with the
- 78 known high burden of RSV in Puerto Rico, our results may help explain the high burden
- 79 of asthma in Puerto Ricans.

80 Word Count

81 249

82 Key Words

- 83 Puerto Rico, asthma, minority children, health disparity, human respiratory viruses, viral
- 84 infection, RSV, African American, Latino.

Abbreviations

- 86 RSV, respiratory syncytial virus; HRV, human rhinovirus; GALA II, Genes-Environments
- and Admixture in Latino Americans; SAGE, The Study of African Americans, Asthma,
- 88 Genes and Environments; URI, upper respiratory infection; SES, socioeconomic status

89 Introduction

90 Asthma is the most common chronic disease in children[1, 2], with genetic,

91 environmental, and infectious risk factors.[3-5] Though the global burden of asthma is

92 increasing, certain racial/ethnic and geographic populations are at especially high risk.

93 Puerto Ricans are among the most severely affected populations in the world.[4]

Approximately 36.5% of Puerto Ricans report they currently or previously had asthma,

compared to only 13.0% of African Americans, 12.1% of European Americans, and

96 7.5% of Mexican Americans.[6] These striking differences extend to asthma morbidity

and mortality, which are 2.4- and 4-fold higher in Puerto Ricans compared to whites,

98 respectively.[6, 7]

99 Several epidemiological studies have established a strong association between the

100 development of childhood asthma or recurrent wheeze with exposure to severe, early-

life respiratory illnesses across populations.[8-18] Associations with asthma were

strongest for infections caused by respiratory syncytial virus (RSV)[8, 11, 14-18] and

human rhinovirus (HRV).[12-14] Both RSV and HRV are known to cause bronchiolitis,

104 which can be a severe respiratory infection in children and is linked to later asthma

development.[16, 19] These viruses have a complex interaction between genetics and

106 environmental exposures in determining risks for asthma and related outcomes.[20, 21]

107 Additionally, Puerto Rico has an RSV season that is year-round whereas the mainland

108 United States only reports a 20-week season (**Fig 1**).[22, 23]

Fig 1. Respiratory syncytial virus (RSV) season in Puerto Rico (red) and the mainland United States (blue). RSV season begins when 10% or more of RSV tests are positive.

It is unclear at this time if differential responses to early-life respiratory illnesses 112 contribute to the striking asthma disparities seen across minority populations. It is 113 possible that genetic predisposition, environmental influences, and early-life respiratory 114 illness work together to increase asthma susceptibility in high-risk populations. Our aim 115 was to investigate the association of early-life respiratory illnesses with asthma 116 117 susceptibility seen in our large and well-phenotyped cohort of minority children. In light of disparate disease prevalence, [6, 22, 23] we examined this association in each 118 119 population (Puerto Ricans, African Americans and Mexican Americans), separately and

120 combined.

121 Methods

122 Study population

The analysis examined participants recruited through two parent studies: the Genes-123 Environments and Admixture in Latino Americans (GALA II) study and the Study of 124 African Americans, Asthma, Genes and Environments (SAGE), described in detail 125 elsewhere.[24-26] Briefly, GALA II and SAGE are parallel case-control studies of 126 asthma conducted between 2006 to 2014 in Latino (Mexican American and Puerto 127 Rican) and African American children aged 8-21, respectively. SAGE participants were 128 recruited from the San Francisco Bay Area and GALA II participants were recruited from 129 across the continental United States (Chicago, Houston, New York City, and the San 130

Francisco Bay Area) and Puerto Rico. Questionnaires were administered to adult participants and parents of minors. All participants provided consent to being in the study. Consent was obtained from all adult participants and parent/legal guardians of minor participants. All participating institutions obtained the appropriate approvals from their Institutional Review Boards.

Out of a total of 6,023, participants with asthma onset prior to the age of two (n=1,450)

and those individuals without complete illness and covariate data (n=2,301) were

excluded from the study. Those diagnosed with asthma before age two were initially

excluded from analysis to better delineate any causal association between early-life

respiratory illness before age two and subsequent diagnosis of asthma. The exclusion

141 criteria yielded an analytical sample of 2,824 subjects, including both cases (n=1,091)

142 and controls (n=1,733).

143 Outcomes

The outcome of our study was asthma diagnosed after the age of two by a
physician.[24] Eligible control subjects had no reported history of asthma, lung disease,
or chronic illness, and no reported symptoms of coughing, wheezing, or shortness of
breath in the two years before enrollment.

148 **Exposure**

149 The exposure was self-reported early-life respiratory illnesses within the first two years

of life. Early-life respiratory illnesses included upper respiratory infection (URI),

151 pneumonia, bronchitis, and bronchiolitis/RSV, which were analyzed separately. In

addition, the report of at least one of the four previous illnesses was used to categorizechildren with "any respiratory illness" and none otherwise.

154 **Covariates**

Consistent with previous studies, [27, 28] we considered factors known to be associated 155 with early-life respiratory illnesses and with asthma. These included sex, underweight at 156 birth (yes/no), maternal smoking during pregnancy (yes/no), whether the subject was 157 breastfed (yes/no), number of older siblings (none, one, two or more), socioeconomic 158 status (SES; low, medium, high; described further in Text A in S1 File), and recruitment 159 region; the latter of which was not used in the African American models because all the 160 participants were recruited from the same region. Global African and European ancestry 161 estimates (described further in Text B in S1 File) were additional covariates used in our 162 combined model to adjust for the underlying substructure of populations in the dataset. 163

164 Statistical Analysis

Descriptive statistics were calculated for the overall study population and for each minority population separately. Logistic regression was used to study the association between physician-diagnosed asthma and self-reported respiratory illnesses in the first two years of life. To account for the difference in asthma prevalence[6] and the fact that ethnicity acts as an effect modifier on the association, we also performed association tests stratified by population. The statistical programming language R version 3.5.1 was used to perform all analyses.

172 Data availability

- 173 Due to the sensitive nature of the data collected surrounding issues of race,
- socioeconomic status, and ancestry, data are available upon request. Access to the
- 175 limited dataset used in this study can be arranged by contacting Dr. Esteban Burchard
- 176 [email: esteban.burchard@ucsf.edu]. Of note, any data provided will be stripped of
- 177 specific subject identifiers that could be used to identify a specific child and/or his/her
- residence. All source code for this analysis can be accessed on our github page
- 179 <u>https://github.com/asthmacollaboratory/asthma-after-RI</u>.

180 **Results**

- 181 Descriptive characteristics of our final study population are presented in **Table 1**. When
- 182 compared with children without asthma, those with asthma were more likely to be male,
- to have a mother who smoked during pregnancy, and have lower SES. These
- distributions were observed in our sample with a few exceptions. For instance, while
- 185 Puerto Rican and Mexican American children without asthma were more likely to be
- breastfed than their counterparts with asthma, the opposite was true for African
- 187 Americans. Additionally, we found that Puerto Ricans were 4- to 9-fold more likely to
- report an RSV infection or bronchiolitis in the first two years of life than Mexican
- 189 Americans and African Americans, respectively (**Fig 2**).
- Fig 2. Population-specific proportions of selected respiratory illnesses during the
 first 2 years of life for our study population.

Table 1. Descriptive statistics of the final study population (N = 2,824).

	Puerto Rican				Mexican American				African American				Combined			
	Case		Control		Case		Control		Case		Control		Case		Control	
	Ν	%	Ν	%	Ν	%	Ν	%	Ν	%	Ν	%	Ν	%	Ν	%
Number of subjects	284	100	783	100	375	100	504	100	432	100	446	100	1091	100	1733	100
Males	150	52.8	361	46.1	212	56.5	210	41.7	218	50.5	187	41.9	580	53.2	758	43.7
Underweight at birth	32	11.3	61	7.8	24	6.4	35	6.9	36	8.3	45	10.1	92	8.4	141	8.1
In-utero smoke exposure	14	4.9	40	5.1	14	3.7	9	1.8	78	18.1	54	12.1	106	9.7	103	5.9
Breastfed	141	49.6	432	55.2	285	76	401	79.6	262	60.6	244	54.7	688	63.1	1077	62.1
Number of older siblings																
0	74	26.1	200	25.5	132	35.2	195	38.7	247	57.2	208	46.6	453	41.5	603	34.8
1	92	32.4	274	35	138	36.8	176	34.9	101	23.4	132	29.6	331	30.3	582	33.6
2 or more	118	41.5	309	39.5	105	28	133	26.4	84	19.4	106	23.8	307	28.1	548	31.6
Socioeconomic status*																
high	89	31.3	271	34.6	114	30.4	116	23	163	37.7	160	35.9	366	33.5	547	31.6
medium	51	18	136	17.4	79	21.1	84	16.7	113	26.2	137	30.7	243	22.3	357	20.6
low	144	50.7	376	48	182	48.5	304	60.3	156	36.1	149	33.4	482	44.2	829	47.8
Recruitment region																
Chicago	18	6.3	23	2.9	137	36.5	180	35.7	-	-	-	-	155	14.2	203	11.7
Houston	1	0.4	-	-	96	25.6	91	18.1	-	-	-	-	97	8.9	91	5.3
New York	27	9.5	34	4.3	20	5.3	64	12.7	-	-	-	-	47	4.3	98	5.7
San Francisco Bay Area	1	0.4	1	0.1	122	32.5	169	33.5	432	100	446	100	555	50.9	616	35.5
Puerto Rico	237	83.5	725	92.6	-	-	-	-	-	-	-	-	237	21.7	725	41.8
URI	57	20.1	35	4.5	24	6.4	27	5.4	67	15.5	16	3.6	148	13.6	78	4.5
Pneumonia	13	4.6	5	0.6	20	5.3	11	2.2	24	5.6	14	3.1	57	5.2	30	1.7
Bronchitis	36	12.7	10	1.3	37	9.9	10	2	16	3.7	6	1.3	89	8.2	26	1.5
Bronchiolitis/RSV	31	10.9	14	1.8	6	1.6	4	0.8	6	1.4	2	0.4	43	3.9	20	1.2
Any Listed	89	31.3	54	6.9	75	20	46	9.1	98	22.7	31	7	262	24	131	7.6

¹⁹³ *Socioeconomic status was derived from a combination of mother's education level, health insurance status, and household income

194 weighted by region, see **Text A in S1 File** for more information.

210	Fig 3. Odds ratios plotted on a log scale of the association between early-life
209	the participant reported having at least one early-life respiratory illness (Table 2, Fig 3).
208	bronchitis, 5.82 (95% CI: 3.26-10.8) for bronchiolitis/RSV, and 4.5 (95% CI: 3.52-5.78) if
207	5.96) for URI, 2.66 (95% CI: 1.67-4.3) for pneumonia, 7.04 (95% CI: 4.44-11.6) for
206	our combined analysis, we observed that the odds of asthma was 4.31 (95% CI: 3.15-
205	participant reported having at least one early-life respiratory illness (Table 2, Fig 3). In
204	bronchiolitis/RSV, and 3.07 (95% CI: 1.99-4.79) and 3.91 (95% CI: 2.55-6.12) if the
203	7.71) for bronchitis, 2.01 (95% CI: 0.552-8.15) and 2.9 (95% CI: 0.635-20.4) for
202	(95% CI: 0.967-3.86) for pneumonia, 4.78 (95% CI: 2.39-10.4) and 2.7 (95% CI: 1.08-
201	1.12-4.24) and 4.77 (95% CI: 2.76-8.71) for URI, 2.05 (95% CI: 0.956-4.61) and 1.9
200	asthma in Mexican Americans and African Americans respectively was 2.17 (95% CI:
199	least one early-life respiratory illness (Table 2, Fig 3). Comparatively, the odds of
198	bronchiolitis/RSV, and 6.15 (95% CI: 4.21-9.05) if the participant reported having at
197	pneumonia, 13 (95% CI: 6.51-28.2) for bronchitis, 7.27 (95% CI: 3.83-14.5) for
196	in Puerto Ricans was 5.25 (95% CI: 3.34-8.37) for URI, 7.23 (95% CI: 2.66-23) for
195	In the population-specific stratified analyses, we also observed that the odds of asthma

211 respiratory illnesses before the age of two and asthma diagnosis after the age of

two across our three main populations and combined.

Table 2. Odds ratios and confidence intervals from the population-specific and combined association analysis between respiratory

illnesses in the first two years of life and physician-diagnosed asthma after the age of two in the three main populations in GALA II and SAGE: 2006-2014.

	Puerto Rican (n=1,067)	Mexican American (n=879)	African American (n=878)	Combined (n=2,824)
URI	5.25 (3.34-8.37)	2.17 (1.12-4.24)	4.77 (2.76-8.71)	4.31 (3.15-5.96)
Pneumonia	7.23 (2.66-23)	2.05 (0.956-4.61)	1.9 (0.967-3.86)	2.66 (1.67-4.3)
Bronchitis	13 (6.51-28.2)	4.78 (2.39-10.4)	2.7 (1.08-7.71)	7.04 (4.44-11.6)
Bronchiolitis/RSV	7.27 (3.83-14.5)	2.01 (0.552-8.15)	2.9 (0.635-20.4)	5.82 (3.26-10.8)
Any Listed	6.15 (4.21-9.05)	3.07 (1.99-4.79)	3.91 (2.55-6.12)	4.5 (3.52-5.78)

216 **BOLD** values are statistically significant at p < 0.05.

To test statistically significant differences in odds ratios between racial/ethnic groups,
we performed a Z-test.[29] We found that there were significant differences (p < 0.05)
between Puerto Ricans and Mexican Americans for URI and Any Listed, and between
Puerto Ricans and African Americans for bronchitis and pneumonia (Table A in S2
File).

In a separate analysis, we performed these associations including those individuals diagnosed with asthma before the age of two (**Table B in S2 File**). Our results showed that all early-life respiratory illnesses were significantly associated (p < 0.05) with asthma diagnosis across our three populations and combined, with the exception of RSV in Mexican Americans (**Table C in S2 File**). Additionally, Puerto Ricans had a significantly higher risk of asthma after respiratory infection than Mexican Americans for every respiratory infection tested (**Table D in S2 File**).

We additionally assessed the differential risk of asthma after an early-life respiratory illness in Puerto Ricans living in Puerto Rico (Islanders) versus those living in the mainland United States (Mainlanders); descriptive characteristics for these populations can be found in **Table E in S2 File**. Our data suggest a high asthma risk following earlylife respiratory illness in Islanders relative to Mainlanders (**Table F in S2 File**) though due to the small number of Mainlanders we were underpowered to detect statistically significant differences between these groups.

236 **Discussion**

237 Overall, we found that early-life respiratory illnesses such as URI, pneumonia,

bronchitis, and bronchiolitis/RSV are significantly associated with asthma diagnosis

after the age of two regardless of race/ethnicity. However, Puerto Ricans displayed thestrongest associations.

241 Early-life respiratory illnesses increase risk for physician-

242 diagnosed childhood asthma

The association between early-life respiratory illnesses and asthma has been well documented.[8-18] Yet, little is known about racial/ethnic differences that underlie the associations between early-life respiratory illnesses and the development of asthma later on in childhood.

Previous research on the association between early-life respiratory infection and the 247 development of asthma has yielded conflicting results. Having a wheezing illness due to 248 249 RSV or HRV infection in early life has previously shown to be associated with a 2.6- and 9.8-fold increase, respectively, in asthma risk by age six in a mostly white 250 population.[12] Additionally, a recent study found that the severity of the RSV infection 251 was found to be strongly associated with childhood wheezing at age five in a mostly 252 white population.[11] However, the vast majority of children infected by respiratory 253 viruses like RSV do not go on to develop respiratory illnesses like recurrent wheezing 254 and asthma.[28] Currently, it is unclear why only a minority of children develop asthma 255 after exposure to an early-life respiratory illness. One plausible explanation is that these 256 257 respiratory illnesses alter the airway in early life, which leads to asthma later on in childhood; previous studies have shown that in adults with asthma, the airway 258 epithelium is altered in a heterogeneous manner.[30] Another possibility is that children 259 260 who are already genetically or environmentally prone to asthma present with early-life

respiratory illness as an early manifestation of asthma. Indeed, previous studies have 261 shown asymptomatic carriage of RSV and rhinovirus in children,[31-33] suggesting a 262 spectrum of disease which may be affected by underlying asthma predisposition.[34, 263 35] Our results show that asthma risk increases with early-life respiratory illnesses in all 264 populations studied, but that the increase is most dramatic in the Puerto Rican 265 266 population. These results support previous studies[35-37] that early-life respiratory illnesses are significantly associated with the development of asthma and asthma-267 related outcomes while additionally adding that these associations may vary by region 268 or population. 269

270 **Puerto Ricans are uniquely burdened compared to other**

271 groups

Most children who develop respiratory viral infections in early life experience minor 272 illness, but some develop much more severe illnesses that involve lower respiratory 273 symptoms like wheezing [9, 10, 31] These more severe wheezing illnesses at an early 274 stage in life are associated with a high risk for recurrent wheezing and asthma later in 275 childhood,[12, 13] particularly if these wheezing illnesses were caused by RSV or 276 277 HRV.[8, 11, 15-18] Puerto Rican children are especially at risk due to the year-round seasonality of RSV infections.[22, 23] In fact, 1,406 cases of bronchiolitis/RSV were 278 reported in children with a mean age of seven months from six Puerto Rican hospitals 279 280 over a period of nine months.[34] Our results showed that not only is the prevalence of bronchiolitis/RSV infection highest in Puerto Rican children in our study population, but 281 that the risk for asthma among children who have experienced a bronchiolitis/RSV 282

infection in early life is the highest for Puerto Rican Islanders. These observations
suggest that the burden of childhood asthma after bronchiolitis/RSV is particularly high
Puerto Rican children.

In our analysis of children diagnosed with asthma after two years of age, some 286 confidence intervals were too wide to make definitive conclusions regarding the relative 287 risk of asthma after infection in different racial/ethnic groups. However, when we 288 included those children diagnosed with asthma before two years of age, the increased 289 sample size allowed us to better delineate the association between different respiratory 290 291 infections and asthma development as stratified by racial/ethnic group. Interestingly, although Puerto Ricans and Mexican Americans are both considered Hispanic/Latino 292 293 for pulmonary function testing and other health metrics, these populations had widely divergent risks of asthma after early-life respiratory infections. Puerto Ricans 294 295 consistently had higher risk of asthma after all respiratory infections studied. Whether 296 the association between early-life respiratory infection and increased asthma risk is causal of the alarmingly high rate of asthma in Puerto Ricans requires further study. 297

While we are not able to draw definitive conclusions about the relative risk of asthma after respiratory infection in Puerto Rican Islanders versus Mainlanders due to the limited number of mainland US Puerto Rican patients studied, we find it interesting that Islander Puerto Ricans tended to have a lower risk of asthma after respiratory infections relative to Islanders. This may suggest an environmental effect about living in Puerto Rico that affects asthma risk. Further studies on Puerto Ricans living on the island and in the mainland US can help clarify this observation.

305 Strengths and limitations

Our study may have been limited because the majority (71%) of Puerto Ricans in the 306 GALA study were diagnosed with asthma before the age of two. In contrast, only 35% of 307 Mexican Americans, and 50% of African Americans were diagnosed with asthma before 308 309 two years old. Our secondary analysis including children diagnosed with asthma before two years of age addressed these differences. Additionally, our exposure 310 measurements were determined retrospectively, which may introduce biases in the 311 data. This study was underpowered to detect any significant associations of early-life 312 respiratory illnesses on asthma in Puerto Rican Mainlanders versus Puerto Rican 313 Islanders, and we were not able to determine whether the year-round RSV prevalence 314 in Puerto Rico modified the effect of the association between respiratory illnesses and 315 asthma. However, our study outlines that there are clear population-specific differences 316 in asthma susceptibility across a variety of respiratory illnesses using a large cohort of 317 minority children from three distinct populations. Our study also benefits from the wide 318 range of clinical, social, and genetic data available on these patients. 319

320 Conclusion

Early-life respiratory illnesses such as RSV, which is highly prevalent in Puerto Ricans, have previously been associated with asthma. Our results indicate that early-life respiratory infections are particularly associated with the later development of asthma among Puerto Rican children compared to other populations.

325

326 Acknowledgements

- 327 The authors acknowledge the families and patients for their participation and thank the
- numerous health care providers and community clinics for their support and
- 329 participation in GALA II / SAGE. In particular, the authors thank study coordinator
- 330 Sandra Salazar; the recruiters who obtained the data: Duanny Alva, MD, Gaby Ayala-
- Rodríguez, Lisa Caine, Elizabeth Castellanos, Jaime Colón, Denise DeJesus, Blanca
- López, Brenda López, MD, Louis Martos, Vivian Medina, Juana Olivo, Mario Peralta,
- 333 Esther Pomares, MD, Jihan Quraishi, Johanna Rodríguez, Shahdad Saeedi, Dean
- 334 Soto, Ana Taveras.

335 **References**

1. Asthma. World Health Organization (WHO). [Available from:

337 <u>http://www.who.int/respiratory/asthma/en/</u>.

338 2. Bloom B, Cohen RA, Freeman G. Summary health statistics for U.S. children:

National Health Interview Survey, 2009. Vital Health Stat 10. 2010(247):1-82.

340 3. Aligne CA, Auinger P, Byrd RS, Weitzman M. Risk factors for pediatric asthma.

341 Contributions of poverty, race, and urban residence. Am J Respir Crit Care Med.

342 2000;162(3 Pt 1):873-7.

4. Oh SS, White MJ, Gignoux CR, Burchard EG. Making Precision Medicine

344 Socially Precise. Take a Deep Breath. Am J Respir Crit Care Med. 2016;193(4):348-50.

5. Subbarao P, Mandhane PJ, Sears MR. Asthma: epidemiology, etiology and risk
factors. CMAJ. 2009;181(9):E181-90.

Barr RG, Aviles-Santa L, Davis SM, Aldrich TK, Gonzalez F, 2nd, Henderson
 AG, et al. Pulmonary Disease and Age at Immigration among Hispanics. Results from
 the Hispanic Community Health Study/Study of Latinos. Am J Respir Crit Care Med.
 2016;193(4):386-95.

Akinbami LJ. Asthma Prevalence, Health Care Use and Mortality: United States,
 2003-05: Center for Disease Control and Prevention: National Center for Health
 Statistics; 2010 [Available from: <u>https://www.cdc.gov/nchs/data/hestat/asthma03-</u>
 05/asthma03-05.htm.

Bacharier LB, Cohen R, Schweiger T, Yin-Declue H, Christie C, Zheng J, et al.
 Determinants of asthma after severe respiratory syncytial virus bronchiolitis. J Allergy
 Clin Immunol. 2012;130(1):91-100 e3.

Castro-Rodriguez JA, Forno E, Rodriguez-Martinez CE, Celedon JC. Risk and
 Protective Factors for Childhood Asthma: What Is the Evidence? J Allergy Clin Immunol
 Pract. 2016;4(6):1111-22.

10. Castro-Rodriguez JA, Holberg CJ, Wright AL, Martinez FD. A clinical index to
define risk of asthma in young children with recurrent wheezing. Am J Respir Crit Care
Med. 2000;162(4 Pt 1):1403-6.

11. Escobar GJ, Masaquel AS, Li SX, Walsh EM, Kipnis P. Persistent recurring
wheezing in the fifth year of life after laboratory-confirmed, medically attended
respiratory syncytial virus infection in infancy. BMC Pediatr. 2013;13:97.

Jackson DJ, Gangnon RE, Evans MD, Roberg KA, Anderson EL, Pappas TE, et
al. Wheezing rhinovirus illnesses in early life predict asthma development in high-risk
children. Am J Respir Crit Care Med. 2008;178(7):667-72.

13. Kusel MM, de Klerk NH, Kebadze T, Vohma V, Holt PG, Johnston SL, et al.
Early-life respiratory viral infections, atopic sensitization, and risk of subsequent
development of persistent asthma. J Allergy Clin Immunol. 2007;119(5):1105-10.

14. Lemanske RF, Jr., Jackson DJ, Gangnon RE, Evans MD, Li Z, Shult PA, et al.
Rhinovirus illnesses during infancy predict subsequent childhood wheezing. J Allergy
Clin Immunol. 2005;116(3):571-7.

376 15. Sigurs N. Epidemiologic and clinical evidence of a respiratory syncytial virusreactive airway disease link. Am J Respir Crit Care Med. 2001;163(3 Pt 2):S2-6. 377 378 16. Sigurs N, Biarnason R, Sigurbergsson F, Kjellman B, Respiratory syncytial virus bronchiolitis in infancy is an important risk factor for asthma and allergy at age 7. Am J 379 Respir Crit Care Med. 2000;161(5):1501-7. 380 381 17. Sigurs N, Bjarnason R, Sigurbergsson F, Kjellman B, Bjorksten B. Asthma and immunoglobulin E antibodies after respiratory syncytial virus bronchiolitis: a prospective 382 383 cohort study with matched controls. Pediatrics. 1995;95(4):500-5. 18. Sigurs N, Gustafsson PM, Bjarnason R, Lundberg F, Schmidt S, Sigurbergsson 384 F, et al. Severe respiratory syncytial virus bronchiolitis in infancy and asthma and 385 allergy at age 13. Am J Respir Crit Care Med. 2005;171(2):137-41. 386 19. Jartti T, Korppi M. Rhinovirus-induced bronchiolitis and asthma development. 387 Pediatr Allergy Immunol. 2011;22(4):350-5. 388

20. Caliskan M, Bochkov YA, Kreiner-Moller E, Bonnelykke K, Stein MM, Du G, et al.
Rhinovirus wheezing illness and genetic risk of childhood-onset asthma. N Engl J Med.
2013;368(15):1398-407.

Voraphani N, Stern DA, Wright AL, Guerra S, Morgan WJ, Martinez FD. Risk of
current asthma among adult smokers with respiratory syncytial virus illnesses in early
life. Am J Respir Crit Care Med. 2014;190(4):392-8.

McGuiness CB, Boron ML, Saunders B, Edelman L, Kumar VR, Rabon-Stith KM.
Respiratory syncytial virus surveillance in the United States, 2007-2012: results from a
national surveillance system. Pediatr Infect Dis J. 2014;33(6):589-94.

398 23. Molinari Such M, Garcia I, Garcia L, Puig G, Pedraza L, Marin J, et al.

Respiratory syncytial virus-related bronchiolitis in Puerto Rico. P R Health Sci J.

400 2005;24(2):137-40.

401 24. Nishimura KK, Galanter JM, Roth LA, Oh SS, Thakur N, Nguyen EA, et al. Early-

life air pollution and asthma risk in minority children. The GALA II and SAGE II studies.

403 Am J Respir Crit Care Med. 2013;188(3):309-18.

404 25. Oh SS, Tcheurekdjian H, Roth LA, Nguyen EA, Sen S, Galanter JM, et al. Effect
405 of secondhand smoke on asthma control among black and Latino children. J Allergy
406 Clin Immunol. 2012;129(6):1478-83 e7.

407 26. Thakur N, Oh SS, Nguyen EA, Martin M, Roth LA, Galanter J, et al.

408 Socioeconomic status and childhood asthma in urban minority youths. The GALA II and

409 SAGE II studies. Am J Respir Crit Care Med. 2013;188(10):1202-9.

Gillman MW, Barker D, Bier D, Cagampang F, Challis J, Fall C, et al. Meeting
report on the 3rd International Congress on Developmental Origins of Health and
Disease (DOHaD). Pediatr Res. 2007;61(5 Pt 1):625-9.

28. Pullan CR, Hey EN. Wheezing, asthma, and pulmonary dysfunction 10 years
after infection with respiratory syncytial virus in infancy. Br Med J (Clin Res Ed).
1982;284(6330):1665-9.

416	29.	Genome Toolbox. Test for a Difference in Two Odds Ratios 2014 [Available from:				
417	417 <u>http://genometoolbox.blogspot.com/2014/06/test-for-difference-in-two-odds-ratios.html</u> .					
	~~					
418	30.	Bai J, Smock SL, Jackson GR, Jr., MacIsaac KD, Huang Y, Mankus C, et al.				
419	Phenotypic responses of differentiated asthmatic human airway epithelial cultures to					
420	20 rhinovirus. PLoS One. 2015;10(2):e0118286.					
	0.4					
421	31.	Global Initiative for Asthma. Global Strategy for Asthma Management and				
422	Preve	ention 2017 [Available from: <u>www.ginasthma.org</u> .				
423	32.	Jansen RR, Wieringa J, Koekkoek SM, Visser CE, Pajkrt D, Molenkamp R, et al.				
	-					
424	Frequent detection of respiratory viruses without symptoms: toward defining clinically					
425	releva	ant cutoff values. J Clin Microbiol. 2011;49(7):2631-6.				
426	33.	Munywoki PK. Koech DC. Agoti CN. Bett A. Cane PA. Medley GF. et al.				
	_					
427	Frequent Asymptomatic Respiratory Syncytial Virus Infections During an Epidemic in a					
428	Rural	Kenyan Household Cohort. J Infect Dis. 2015;212(11):1711-8.				
429	34.	Matias I. Garcia I. Garcia-Fragoso L. Puig G. Pedraza L. Rodriguez L. et al.				
	_					
430	Trends of Respiratory Syncytial Virus Infections in Children under 2 Years of Age in					
431	Puert	o Rico. P R Health Sci J. 2015;34(2):98-101.				
132	35	Wennergren G. Kristiansson S. Relationshin between respiratory syncytial virus				
732	00.					
433	bronc	hiolitis and future obstructive airway diseases. Eur Respir J. 2001;18(6):1044-58.				

- 434 36. Burchard EG, Ziv E, Coyle N, Gomez SL, Tang H, Karter AJ, et al. The
- importance of race and ethnic background in biomedical research and clinical practice.
- 436 N Engl J Med. 2003;348(12):1170-5.
- 437 37. Szentpetery SE, Forno E, Canino G, Celedon JC. Asthma in Puerto Ricans:
- Lessons from a high-risk population. J Allergy Clin Immunol. 2016;138(6):1556-8.

439 Supplemental Information

- 440 **S1 File. Derivation of variables.** Text A-B.
- 441 **S2 File. Supplemental tables.** Tables A-F.







Puerto Rican
 Mexican
 African American
 Combined