The closed eye harbors a unique microbiome in dry eye disease

Kent A. Willis¹, Cameron K. Postnikoff^{2,3}, Amelia B. Freeman⁴, Gabriel Rezonzew⁴, Kelly K. Nichols², Amit Gaggar⁵, Charitharth V. Lal^{4,5}

¹ Division of Neonatology, Department of Pediatrics, College of Medicine, The University of Tennessee Health Science Center, Memphis, TN, USA

² School of Optometry, University of Alabama at Birmingham, Birmingham, AL, USA

³ CooperVision, Inc., Pleasanton, CA, USA

⁴ Division of Neonatology, Department of Pediatrics, College of Medicine, University of Alabama at Birmingham, Birmingham, AL, USA

⁵ Program in Protease and Matrix Biology, Department of Pediatrics, College of Medicine, University of Alabama at Birmingham, Birmingham, AL, USA

Dry eye affects millions of individuals. In experimental models, dry eye disease is associated with T helper cell 17-mediated inflammation of the ocular surface that may cause persistent damage to the corneal epithelium. However, the initiating and perpetuating factors associated with chronic inflammation of the ocular surface remain unclear. The ocular microbiota alters ocular surface inflammation and may influence dry eye disease development and progression. Here, we collected serial samples of closed eye tears during a randomized clinical trial of a non-pharmaceutical dry eye therapy and used 16S rRNA metabarcoding to characterize the microbiome. We show the closed dry eye microbiome is distinct from the healthy closed eye microbiome. The ocular microbiome was described only recently, and this report implicates a distinct microbiome in ocular disease development. Our findings suggest an interplay between microbial commensals and inflammation on the ocular surface. This information may inform future studies of the pathophysiological mechanisms of dry eye disease.

tears, microbial ecology, ophthalmology, randomized clinical trial

Introduction

Dry eye disease is a common condition that affects millions of individuals worldwide^{1,2}. Ocular surface inflammation has recently been recognized as a hallmark of dry eye disease^{3,4}. T helper type 17 (Th17) cells on the inflamed ocular surface mediate the long-term progression of dry eye disease in experimental models³, however, this same pathophysiology has yet to be demonstrated conclusively in humans. In chronic dry eye, persistent inflammation may eventually produce lasting corneal epithelial damage⁴. However, the factors that incite and perpetuate inflammation in dry eye disease remain unknown⁵. Microbial commensal organisms can alter Th17 populations in the host organism, both in homeostasis and when perturbed^{6,7}. In contrast to previous beliefs, the human eye hosts a resident microbiome⁸⁻¹¹. Culture and molecular-based swabs of the ocular surface have revealed an intrinsic

microbiome distinct from the surrounding skin, but information about the tear microbiome is scarce¹². Interestingly, the ocular microbiome impacts ocular surface inflammation¹³. Together, these findings suggest the ocular microbiome may cause or perpetuate the development of chronic dry eye. Here, we collected longitudinal samples of the bilateral closed eye tear microbiome from a randomized clinical trial for the treatment of dry eye disease. We aimed to identify a distinct closed eye microbiome in dry eye disease. We hypothesized that the chronic dry eye microbiome, if present, would be distinct from the normal closed eye microbiome.

Results

Cohort

Correspondence regarding preprint: Kent A. Willis, MD at kwill164@uthsc.edu or C. V. Lal, MD at clal@peds.uab.edu. Statement of Financial Support: This study was supported by AHA 17SDG32720009 (CVL), NIH K08HL141652 (CVL) and NIH R01HL102371 (AG).

Based on the power and pairwise sample-size estimator for permutational multivariate analysis of variance (PERMANOVA) application Micropower¹⁴, a low abundance 16S rRNA dataset similar to previous studies of the ocular micribome¹² would require a minimal sample size of 30 to generate a discriminatory power of 0.8 with a significance level of 0.05. Therefore, we aimed for at least 35 subjects per study arm. **Table** 1 shows the demographics and clinical characteristics of the enrolled participants. Based on a post-enrollment exam and two clinical surveys, after informed consent, participants were classified as dry eye or controls (**Supplemental S1**).

The resulting 36 control and 36 dry eye subjects were then stratified by clinical severity of eye disease (**Figure 1** and **Supplemental Table 1**). The study arms were randomized to daily saline eye wash upon awakening or non-intervention. A baseline closed eye tear sample was collected at randomization and the final sample after one month, yielding 144 samples. We observed no differences within the normal and dry eye cohorts (normal vs mild, or moderate vs severe). Therefore, all subsequent analyses were performed on normal or dry eye only (**Methods**).

Table 1. Participant demographics and clinical characteristics

	Normal Eye	Dry Eye	P value
Subjects, n	36	36	
Sex			1.00
Female, n (%)	23 (64)	23 (64)	
Male, n (%)	13 (36)	13 (36)	
Age, y ± SD	33 ± 11	52 ± 19	<0.0001
DEQ Score	1.2 ± 1.3	12 ± 3.3	<0.0001
Phenol Red Thread	22 ± 5.3	19 ± 8.6	0.08
Wetting length, mm ± SD			
NIKBUT, s ± SD	15 ± 5.2	10.7 ± 5.2	0.0007
Corneal Staining	0.47 ± 0.8	3.6 ± 3	<0.0001
InflammaDry Positive, n (%)	13 (36)	20 (55)	0.10
Randomization			
Control, n (%)	16 (44)	20 (56)	
Treatment, n (%)	20 (56)	16 (44)	



Fig. 1. Subjects were allocated, stratified and randomized to treatment or control. Schematic of collection of closed eye tears in sterile saline, followed by 16S rRNA metabarcoding of the bacterial microbiome. Diagram of subject allocation and randomization. Figure generated with BioRender.

Sequencing data

To investigate the bacterial microbiome of the closed dry eye, we used metabarcoding of the 16S rRNA gene. Sequence reads were generated using the MiSea system (57,022 Illumina ± 47.011 counts/sample, **Methods**). We assigned taxonomy using the Greengenes database¹⁵, producing 1,593 data points/sample. After excluding any reads aligned to chloroplasts or cyanobacteria, we examined the bacterial community composition using the remaining 1,195 data points/sample that were collectively assigned to 185 genera¹⁶. All tear samples had the standard > 1000 aligned reads/sample, so we retained all samples for further We used a log₂-transformation analvsis. of cumulative sum scaling¹⁷ (log₂ CSS) to normalize our dataset, consistently producing 8,700-10,000 reads per sample (Supplemental Fig. S2).

The closed eye microbiome is distinct in dry eye disease

In the taxonomic analysis of the closed eye microbiome, the microbiomes of individuals with chronic dry eye disease and those without form distinct communities. Dry eye microbial communities are more diverse as quantified by richness (ANOVA, f = 4.8, $P = 7.5 \times 10^{-5}$), evenness (ANOVA, f = 13, $P = 2.1 \times 10^{-12}$) and Shannon diversity (ANOVA, f = 12, P

=5.9 x10⁻¹², **Fig. 2a**). As demonstrated by principal coordinate analysis (PCoA) **Bray-Curtis** of dissimilarity, these communities form unique clusters as quantified by PERMANOVA (R^2 =0.21, P =0.00033. **2b**). Similarly, multivariate Fig. redundancy analysis (RDA, variance =95.25, f = 3.71, P = 0.001) and canonical correspondence analysis $(CCA, chi^2 = 0.16 f = 3.64, P = 0.001)$ showed these communities are distinct, and this factor accounts for the majority of variance in the data. As quantified by two-way ANOVA adjusted for false discovery rate (FDR), univariate analysis of the relative abundance of bacterial genera showed individuals with dry eye disease have differences in the relative abundance of 113 genera, with the most significant differences in Methylobacteriaceae, OPB56. Bacteroidetes. Pseudomonas, and Meiothermus (all with FDRadjusted $P < 2 \times 10^{-22}$, Supplemental Table 2). Similarly, mixed effects regression detected the most significant differences in the abundance of OPB56, Bacteroidetes, Pseudomonas, Meiothermus, and Methylobacteriaceae (all with FDR-adjusted P < 3.1x10⁻²⁰, Supplemental Table 3) among 49 genera with significant differences in relative abundance. Fig. 2c shows relative abundance at the order level. To further investigate the importance of particular bacterial operational taxonomic units (OTUs), we used Spearman network analysis (Fig. 2d).



Fig. 2. Patients with dry eye have a different closed eye microbiome.

(a) Individuals with chronic dry eye have different alpha diversity of the tear microbiome as quantified by the richness (ANOVA, f = 4.8, $P = 7.5 \times 10^{-5}$), evenness (ANOVA, f = 13, $P = 2.1 \times 10^{-12}$) and Shannon diversity (ANOVA, f = 12, $P = 5.9 \times 10^{-12}$) indices. (b) The beta diversity of the closed eye microbiome in individuals with dry eye disease is distinct by principal coordinate analysis (PCoA) of Bray-Curtis dissimilarity ($R^2 = 0.21 P = 0.00033$), redundancy analysis (RDA, variance =95.25, f = 3.71, P = 0.001) and canonical correspondence analysis (CCA, chi² = 0.16 f = 3.64 P = 0.001).

(c) Relative abundance of bacterial orders. $Log_2(CSS)$, log_2 transformation of cumulative-sum scaling. Two-way ANOVA, * P = 0.01, ** P = 0.001, *** P = 0.0001.

(d) Spearman network analysis at the OTU level. Positive correlations with a *P*-value < 0.05 are shown as an edge with the relative size determined by the importance of the taxa to the network.

The closed dry eye microbiome remains distinct despite daily eye wash

Daily eye wash on awakening with sterile saline is a proposed therapy for dry eye disease¹⁸. We tested if the microbial community composition could be normalized by the prescription of daily saline eye wash. However, the microbial communities of dry (**Supplemental Fig. S3**) and normal eyes (**Supplemental Fig. S4**) were relatively unaffected by daily eye wash. The diversity of subjects with dry eye remained higher than that of normal individuals (richness, ANOVA, f = 9.2, $P = 7.5 \times 10^{-5}$, evenness, ANOVA, f = 28, $P = 2.8 \times 10^{-14}$ and Shannon diversity,

ANOVA, f = 28, $P = 6.1 \times 10^{-14}$, **Fig. 3a**). Multivariate analysis of these communities also remained distinct (PCoA, PERMANOVA, $R^2 = 0.15$, P = 0.00033, RDA, variance =64.75, f = 5.73, P = 0.001 and CCA, chi² =0.11, f = 5.62, P = 0.001, **Fig. 3b**). Univariate analysis detected the most significant differences in the abundance of *MLE112*, *Lactobacillaceae*, *Streptococcus*, *Sphingobium*, *Caldicoprobacter* and *Anaerococcus* (ANOVA, P < 0.01, **Fig. 3c**). Core microbiome analysis highlighted key differences in the distribution of key genera, and network analysis demonstrated the importance of key taxa (**Fig. 3d**, **3e**).



Fig 3. Daily eye rinse does not alter the closed eye microbiome.

(a) Despite daily eye washes, the alpha diversity of the closed eye microbiome remains similar to baseline in individuals with and without dry eye disease as quantified by the richness (ANOVA, f = 9.2, $P = 7.5 \times 10^{-5}$), evenness (ANOVA, f = 28, $P = 2.8 \times 10^{-14}$) and Shannon diversity (ANOVA, f = 28, $P = 6.1 \times 10^{-14}$) indices.

(b) The beta diversity of the closed eye microbiome remains distinct in individuals with and without dry eye disease by principal coordinate analysis (PCoA) of Bray-Curtis dissimilarity (PERMANOVA, $R^2 = 0.15 P = 0.00033$), redundancy analysis (RDA, variance =64.75, *f* =5.73, *P* =0.001) and canonical correspondence analysis (CCA, chi² =0.11 *f* =5.62 *P* =0.001).

(c) Relative abundance of bacterial genera. $Log_2(CSS)$, log_2 transformation of cumulative-sum scaling. Two-way ANOVA, * P =0.01, ** P =0.001, ** P = 0.0001.

(d) Core microbiome analysis showing differences in microbial colonization at the genus level.

(e) Spearman network analysis at the OTU level. Positive correlations with a *P*-value < 0.05 are shown as an edge with the relative size determined by the importance of the taxa to the network.

The closed eye microbiome in dry eye disease is distinct at baseline

To gain further insight into the microbial community composition of the dry eye, we narrowed our focus to

examine the closed eye microbial composition at the time of randomization. As indicated by our more general analysis, both diversity (richness, ANOVA, f =6.1, P =0.016, evenness, ANOVA, f =24, P =5.8 x10⁻

⁶ and Shannon diversity. ANOVA. *f* =23. *P* =8.1 x10⁻ ⁸, Fig. 4a) and multivariate clustering of these communities are distinct at baseline (PCoA, PERMANOVA, R^2 =1.34 P =0.00033, RDA, variance =53.95, f =7.12, P =0.001 and CCA, chi² =0.09, f =7.12, P =0.001, Fig. 4b). Univariate analysis using negative binomial regression identified Methylobacteriaceae, Pseudomonas. Bradyrhizobium and Allobaculum as the five most differently abundant genera (all with FDR-adjusted P < 1.1 x10⁻¹¹, Supplemental Fig. S5a and Supplemental Table 4). At the phylum level, Firmicutes and Bacteroidetes remained unchanged,

while Verrucomicrobia (FDR $P = 4.6 \times 10^{-11}$) and Proteobacteria (FDR $P = 6.0 \times 10^{-10}$) showed the most significant differences. Core microbiome analysis identified 45 genera unique to dry eye with only 14 genera unique to the normal eye (**Fig. 4d**). Discriminant analysis of principal components at the order level revealed the abundances of orders OPB56 and Rhizobiales are important to discriminate the normal eye while the orders Halanaerobiales, Erysipelotrichales and Anaeroplasmatales are important to discriminate dry eye (**Fig. 4e**). Network analysis provided further evidence of these distinct communities (**Fig. 4f**).



Fig. 4. The closed eye microbiome of the in dry eye disease is distinct at baseline.

(a) The alpha diversity of the closed eye microbiome remains distinct as quantified by the richness (ANOVA, f = 6.1, P = 0.016), evenness (ANOVA, f = 24, $P = 5.8 \times 10^{-6}$) and Shannon diversity (ANOVA, f = 23, $P = 8.1 \times 10^{-8}$) indices.

(b) The beta diversity of the closed eye microbiome remains distinct as quantified by principal coordinate analysis (PCoA) of Bray-Curtis dissimilarity (PERMANOVA, $R^2 = 1.34 P = 0.00033$), redundancy analysis (RDA, variance =53.95, *f* =7.12, *P* =0.001) and canonical correspondence analysis (CCA, chi² =0.09, *f* =7.12, *P* =0.001).

(c) Relative abundance of bacterial orders. $Log_2(CSS)$, log_2 transformation of cumulative-sum scaling. Two-way ANOVA (***displaying only, P < 0.001).

(d) Core microbiome analysis showing differences in microbial colonization at the genus level.

(e) Discriminant analysis of principal components at the order level.

(f) Spearman network analysis at the OTU level. Positive correlations with a *P*-value < 0.05 are shown as an edge with the relative size determined by the importance of the taxa to the network.

Machine learning accurately classifies dry eye samples at baseline

To more precisely identify unique features with the potential to function as biomarkers for patients with dry eye, we used linear discriminant analysis of effect size. We identified 5 genera that reliably identified individuals with dry eye (*Pseudomonas, Methylobacteriaceae, Helicobacter, Acetobacter* and

Stenotrophomonas) with 3 genera that reliably identified normal patients (*Leuconostocaceae*, *Streptococcus* and *Calothrix*, **Supplemental Fig. S5b**). To further support the uniqueness of microbial communities in dry eye, we built a support vector machine using leave-one-out cross-validation that could identify samples with 94% accuracy. Similarly, when we developed a random forest classifier,

variable analysis identified the importance prevalence of the genera Methylobacterium. Megasphaera, Parabacteroides, S247. Bifidobacterium, Streptococcus. Desulfovibrio, Acetobacter, Dialister and Bacillus as particularly useful to identify samples from individuals with dry eye (Importance > 20, Supplemental Fig. S5c).

The closed eye microbiome in dry eye disease diverges after one month

We examined the stability of the dry eye microbiome by focusing on samples collected one month later in the same individuals. The dry eye microbiome remained distinct from the normal microbiome and remained consistent with the community composition noted at baseline (**Supplemental Fig. S3**, **S2**). Only slight divergence was noted over the course of a month (**Fig. 5**). The diversity of dry eye remained higher than that of the normal eye (richness, ANOVA, f=19, $P=5.2 \times 10^{-5}$, evenness, ANOVA, f=65, $P=1.9 \times 10^{-11}$ and Shannon diversity, ANOVA, f=60, $P=8.6 \times 10^{-11}$, **Fig. 5a**). The distinct clustering of these communities on multivariate analysis also persisted (PCoA of Bray-Curtis dissimilarity, PERMANOVA, R^2) =1.52, P =0.00033, RDA, variance =66.53, f =8.52, P =0.001 and CCA, chi^2 =0.10, f =8.14, P =0.001, Fig. **5b**). Similarly, negative binomial regression noted the greatest differences in the abundance of the genera Methylobacteriaceae, Pseudomonas, Azospirillum, Bradyrhizobium and Coriobacteriaceae (all with FDR-adjusted $P < 7.0 \times 10^{-18}$, Supplemental Fig. S6a, Supplemental Table 4). However, core microbiome analysis detected 76 unique genera in dry eye, 69 shared genera and 24 genera unique to the normal eye (Fig 5d). Similar to baseline, discriminant analysis of principal components at the order level identified OPB58 and Bacteroidetes as important discriminators of the normal eye. In individuals with dry eye, additional orders enabled further discrimination. with Flavobacteriales. Alteromonadales and Actinomycetes, Anaeroplasmatales and Desulfuromonadales functioning as the primary discriminators instead of Halanaeobiales and Erysipelotrichales (Fig. 5d).





(a) The alpha diversity of the closed eye microbiome remains distinct as quantified by the richness (ANOVA, f = 19, $P = 5.2 \times 10^{-5}$), evenness (ANOVA, f = 65, $P = 1.9 \times 10^{-11}$) and Shannon diversity (ANOVA, f = 60, $P = 8.6 \times 10^{-11}$) indices.

(b) The beta diversity of the closed eye microbiome remains distinct as quantified by principal coordinate analysis (PCoA) of Bray-Curtis dissimilarity (PERMANOVA, R^2 =1.52 P =0.00033), redundancy analysis (RDA, variance =66.53, *f* =8.52, P =0.001) and canonical correspondence analysis (CCA, chi² =0.10, *f* =8.14, P =0.001).

(c) Relative abundance of bacterial orders. $Log_2(CSS)$, log_2 transformation of cumulative-sum scaling. Two-way ANOVA, * P = 0.05, ** P = 0.01, *** P = 0.001.

(d) Core microbiome analysis showing differences in microbial colonization at the order level.

(e) Discriminant analysis of principal components at the order level.

(f) Spearman network analysis at the OTU level. Positive correlations with a *P*-value < 0.05 are shown as an edge with the relative size determined by the importance of the taxa to the network.

Machine learning more accurately identifies dry eye after a month

We used linear discriminant analysis of effect size to identify bacterial genera with the potential function as biomarkers and noted 10 genera unique to individuals with dry eye and 9 unique to individuals with a normal eye, supporting the limited divergence of these communities (**Supplemental Fig. S6b**). We also developed a support vector machine with leaveone-out cross-validation that identified dry eye samples with 97% accuracy. Variable importance analysis of a random forest classifier noted the prevalence of *Methylobacterium*, *Megasphaera*, *Parabacteroides*, S247, Bifidobacterium, *Streptococcus, Desulfovibrio* and *Acetobacter* as important identifiers (Importance > 20, **Supplemental Fig. S6c**).

The closed eye microbiome is unaltered by the cellular concentration of the tear fraction

To ensure our microbial ecological analysis was unaltered by the cellular content of the tear fraction, we compared the microbial composition of each cellular fraction collected. Three samples were excluded from this analysis due to incomplete data (n = 141). Additionally, community composition was similar when we collated high and low cellular fractions to increase discriminatory power (**Fig. 6**).



Fig. 6. The closed eye microbiome is unaltered by the cellular fraction.

a) The alpha diversity was not different as quantified by the richness (ANOVA, f = 1.3, P = 0.26), evenness (ANOVA, f = 1.8, P = 0.19) or Shannon diversity (ANOVA, f = 1.9, P = 0.17) indices.

(b) The beta diversity of the closed eye microbiome is unaltered by the cellular fraction as quantified by principal coordinate analysis (PCoA) of Bray-Curtis dissimilarity (PERMANOVA, R^2 =0.00894 *P* =0.124), redundancy analysis (RDA, variance =4.82, *f* =1.29, *P* =0.152) and non-metric multidimensional scaling (NMDS, stress = 0.228).

(c) Relative abundance of bacterial genera. $Log_2(CSS)$, log_2 -transformation of cumulative-sum scaling. Two-way ANOVA, * P = 0.05, ** P = 0.01, *** P = 0.001.

(d) Discriminant analysis of principal components.

(e) Spearman network analysis at the OTU level. Positive correlations with a *P*-value < 0.05 are shown as an edge with the relative size determined by the importance of the taxa to the network.

Discussion

Direct crosstalk between resident microbes and host immune cells at the mucosal surface is a critical determinant of inflammatory diseases¹⁹. Furthermore, Th17 and Treg cells are implicated in the development of chronic dry eye disease³ and particularly attuned to the ecology of the resident microbiota⁶. Therefore, the ocular surface microbiome may contribute to the pathogenesis of dry eye disease¹³. Understanding how the ocular microbiome influences ocular surface disease will be important for therapeutic development. In this report, we present initial evidence of a distinct closed eye microbiome in dry eye. Moreover, we provide the first

evidence linking microbiome changes with ocular surface disease.

In this study, human subjects collected samples of sterile saline rinse applied to the ocular surface upon awakening in the morning. Using saline eye wash to collect microbiome samples is novel and less invasive than swabs of the ocular surface, which has been routinely used for the collection of tear leukocytes^{18,20}. In a series of pooled samples from both eyes, the closed eye tear microbiome in individuals with dry eye disease was distinct from that in healthy controls. Compared to swabs, where the fiber content and limited sample area artificially reduce the number of detectable microbes²¹, saline wash may allow for a less biased sampling of the entire ocular surface, especially when pooled from both eyes. Furthermore, a general weakness of previous ocular microbiome studies is focusing on shifts in percent relative abundance to identify differences in bacterial taxa. A strength of our analysis is the rigorous log₂ normalization of the cumulative sum scaling transformation performed prior to analysis. With this normalization, our data more accurately reflect true shifts in relative abundance¹⁷. Using untransformed data, the most significant difference in our dataset was a nearly complete substitution of Staphylococcus spp. for Pseudomonas spp. in patients with drv eve. Doan et al. reported a similar shift in the unnormalized relative abundance of Staphylococcus spp. and Pseudomonas²². A more rigorous normalization performed on our dataset confirmed the reduction in Pseudomonas spp. but not in Staphylococcus spp. In addition, 48 other genera may warrant further investigation.

Previous research has focused on the open eve microbiome¹², while we focused on the closed eye microbiome. Every night during sleep, inflammatory species move into the closed eye tear film^{23,24}. perhaps in response to entrapped microbiota^{12,25}. If microbial abundance increases in this situation, differences in bacterial community composition could become more pronounced in the closed eye microbiome than in the open eye. This difference could lead to the closed eye producing more reliable disease states for markers of diagnostic development. Alternatively, this difference could help determine whether anti-inflammatory mechanisms in

the closed eye are sufficient to maintain a healthy ocular microbiome.

Graham and colleagues explored the microbiome of dry eye using culture and broad 16S rRNA-based PCR²³. Due to considerable differences in next sequencing and bioinformatics generation techniques in the last decade as well as our different sampling methods, directly comparing our 16S rRNAbased characterizations of the ocular microbiome is difficult. Nevertheless, our analysis does support their findings of increased bacterial diversity in patients with dry eye disease²³. Although we identified a broader number of genera than was feasible when Graham and colleagues²³ performed their analysis, we confirmed their identification of several genera in dry eye, including Streptococcus, Staphylococcus. Corvnebacterium and Propionibacterium. They nevertheless detected more colony forming units in patients with dry eye than in controls using culture-based techniques. These techniques are biased in low biomass samples because of the limited growth of some species in culture¹². Coagulase negative *Staphylococcus* spp. were the most commonly identified taxa via culture, and their PCR-based analysis showed the microbiome remained stable over three months. Another culture-based pilot study identified similar increased abundances of Staphylococcus aureus, coagulase negative Staphylococcus, Corvnebacterium and Propionibacterium in chronic dry eye²⁴. We also provided in-depth analysis of population dynamics and demonstrated possible divergence of microbial diversity and community structure of normal and dry eyes over time.

Increased bacterial diversity is considered favorable, as more diverse communities are often more resistant to perturbation²⁵. However, mounting evidence suggests the ocular microbiome deviates from this trend, likely due to lysozyme and antimicrobial compounds in tears²⁶. In dry eye disease, our results suggest increased microbial diversity is a hallmark of disease. Consistent with the behavior of the ocular microbiome in other pathological conditions¹², this finding may represent breakdown of the microbe-oriented homeostatic mechanisms of the host. This finding may also suggest increased diversity results from the incorporation of pathobionts into this community. Alternatively, the altered ecology may result from

changes within the ocular surface of the dry eye that support establishment of a wider diversity of microbes in this environment. These additional species may not be harmful but instead highlight the dynamic biome within the human eye. The evidence may also suggest the immune system of the closed eye cannot regulate the ocular surface microbiome during sleep. In the closed eye, preliminary evidence suggests higher numbers of neutrophils with an enhanced degranulation response in dry eye disease^{18,Postnikoff:2018fv; 20}, indicating a dysregulated immune response. This dysregulation may permit increased microbial diversity. The reverse could also be true: enhanced microbial diversity may induce a greater inflammatory response.

This dataset consistently demonstrates the microbial communities in dry eye disease are distinct and remain distinct even with daily saline eye wash. We asked subjects in the treatment arm to selfadminister saline eye wash immediately upon awakening. We hypothesized this time may represent the point of maximum difference since microbes likely accumulate on the ocular surface overnight. However, we did not observe reduced bacterial diversity with saline eyewash, with dry eye communities becoming more similar to control eye communities with treatment. Instead. the communities slightly diveraed: the microbial communities in dry eye remained different from those in healthy eyes. Thus, the mechanisms of microbial accumulation on the ocular surface may be more complex than moisture content or tear clearance. Intrinsic immune or mechanical alterations in dry eye disease may foster differences in commensal abundance, potentially explaining how altered bacterial communities on the ocular surface contribute to dry eye disease.

Methods

Participants

This study was performed in accordance to the Declaration of Helsinki under the supervision of the Institutional Review Board of the University of Alabama at Birmingham (UAB), Birmingham, AL, USA. We performed a secondary analysis of the closed eye microbiome from a prospective, randomized controlled trial of the efficacy of daily eye wash to ameliorate inflammation in individuals with dry eye disease (ClinicalTrials.gov NCT03332342).

Future studies may more accurately localize the innate microbes of the eye by examining ocular swabs and the tear microbiome simultaneously. Serial sampling throughout a day may demonstrate diurnal variation in these bacterial communities. However. aside from certain antibacterial medications²⁰, most human microbiome studies are observational by necessity. Gnotobiotic animals are often useful in unraveling causality in host-microbe interactions and could be the next step^{27,28}, especially explore interactions between the ocular to microbiome and host Th17 and Treg cells in dry eye disease. Building on these experiments may include identifying changes in microbial metabolites or exploring how the gut microbial communities influence ocular development and disease. Finally, aside from one study that examined the virome²⁹⁻³¹, most studies of the ocular microbiome have focused exclusively on bacteria. The fungal microbiome of the normal eye was described only recently²² and may warrant further investigation in dry eye disease. As performed recently for the gut³² and lung³³, modeling the initial succession of multi-kingdom microbial communities in newborns may identify ecological factors in the ocular microbiome.

The ocular microbiome potentially hosts a distinct core microbiome that may be perturbed in eye disease³⁴. Our study provides novel insight into the closed eye microbiome in dry eye disease. Th17 cells are sensitive to changes in the microbiome^{6,7} in experimental dry eye disease³. Therefore, increased microbial abundance in dry eye may promote the development of dry eye disease by altering T cell subpopulations on the ocular surface. More research is needed to reveal whether an altered microbiome drives the development of dry eye disease.

Informed consent was obtained from all participants prior to enrollment. Participants were selected based on the results of a physical exam and two clinical surveys, the Dry Eye Questionnaire 5 (DEQ5)³⁵ and the Ocular Surface Disease Index (OSDI)³⁶. Exclusion criteria included pregnancy, contact lens use within the past three months, glaucoma treatment, current tobacco use, systemic dry-eyeassociated inflammatory disease, any antiinflammatory therapy for dry eye treatment in the last

three months or other antimicrobial or antiinflammatory medications in the last month (**Supplemental Fig. S1**). Stratification into normal and dry eye was driven primarily by symptoms, as measured by the DEQ5 score, with further stratification among both groups determined by phenol red thread wetting length³⁷, non-invasive tear break-up time (NIKBUT) and the InflammaDry test (Quidel; San Diego, CA, USA. **Supplemental Table 1**). Dry eye subjects also had to endorse the sensation of dry eye or have a previous diagnosis of dry eye disease. Equal-weight randomization to either daily eye wash or control was performed using REDCap (Research Electronic Data Capture; Nashville, TN, USA)³⁸.

Eyewash treatment

Subjects were trained on the first visit to perform a daily eye wash. Briefly, as described previously³⁹⁻⁴¹, subjects were provided one 50 mL sterile centrifuge tube, and two 10 mL sterile polypropylene syringes (BD Biosciences; San Jose, CA, USA) with 5 mL of sterile saline per syringe. Immediately after awakening, subjects were instructed to wash their eyes by dispensing the saline gently across the ocular surface, with the runoff collected in a sterile Eppendorf tube. Runoff from both eyes was collected to produce a pooled sample of both eyes collected into the same tube. Subjects in the treatment group completed daily therapy for approximately 28 days.

Sample collection

We utilized the power and pairwise sample-size estimator for PERMANOVA application Micropower¹⁴, implemented as at https://fedematt.shinyapps.io/shinyMB/, to estimate the minimum number of samples to produce a power of 0.8 to detect a difference with a significance level of 0.05 via PERMANOVA. We processed the collected eyewash samples immediately after they were dropped off by study subjects. Pooled tear collections were centrifuged at 270x g, and the supernatant was aliquoted into 1.5 mL microcentrifuge tubes and stored at -80°C. If more than one million cells were recovered in the pooled tear collection, supernatant aliquots were spiked with cells from the precipitate to increase the potential vield of microbiome analysis by normalizing the cellular fraction.

Illumina MiSeq metabarcoding

In the Microbiome Core at UAB, microbial genomic DNA was isolated and PCR was used with unique bar-coded primers to amplify the V4 region of the 16S rRNA gene to create a 16S amplicon library of the samples. The PCR products were sequenced using the NextGen Illumina MiSeq System (Illumina Inc.; San Diego, CA, USA). The sequence data covered the 16S rRNA V4 region with a PCR product length of ~255 bases and 250 base paired-end reads. The raw dataset is available at the NCBI Sequence Read Archive:

http://www.ncbi.nlm.nih.gov/bioproject/597168.

Bioinformatics

Sequences were grouped into OTUs using QIIME v1.9⁴². We assigned taxonomy using the Ribosomal Database Project classifier (threshold 0.8) against the Greengenes database^{14,15}. We then imported these processed sequence reads into Calypso v8.84⁴³ for further processing and data analysis. After excluding any OTUs assigned to chloroplasts or cyanobacteria, we *a priori* determined to exclude any samples with < 1000 reads/sample from downstream analysis. We then utilized cumulative sum scaling¹⁷ to normalize the relative abundance and performed a log₂-transformation of the resulting data, centering the taxonomic counts to 0 and scaling to a range of -2 to 2 with a variance of 1. For alpha diversity analysis, samples were rarified to a depth of 7804 reads/sample. Alpha diversity was then quantified using the richness, evenness and Shannon diversity indices⁴⁴, testing for significant differences with ANOVA adjusted for FDR. For beta diversity, we visualized the data using PCoA, RDA, CCA or NMDS⁴⁵. We used PERMANOVA to determine significant differences in beta diversity⁴⁶. Relative abundance was guantified using ANOVA adjusted for FDR for multiple groups or negative binomial regression (DESeq2 function) for binary comparisons. For Spearman network analysis, we displayed positive correlations with an FDR-adjusted P < 0.05 as an edge. Relative size of the point represents the importance of the taxa to the resulting network. In addition, we used core microbiome analysis (of the 300 most abundant taxa), linear discriminant analysis of effect size, and discriminant analysis of principal components to select significant features. For analyses with two groups, we developed machine learning classifiers using support vector machine and random forest algorithms.

References

1. Farrand, K. F., Fridman, M., Stillman, I. Ö. & Schaumberg, D. A. Prevalence of Diagnosed Dry Eye Disease in the United States Among Adults Aged 18 Years and Older. *Am. J. Ophthalmol.* **182**, 90–98 (2017).

 Stapleton, F. *et al.* TFOS DEWS II Epidemiology Report. *The Ocular Surface* **15**, 334–365 (2017).

3. Chen, Y., Chauhan, S. K., Lee, H. S., Saban, D. R. & Dana, R. Chronic dry eye disease is principally mediated by effector memory Th17 cells. *Mucosal Immunol* **7**, 38–45 (2014).

4. Heidari, M., Noorizadeh, F., Wu, K., Inomata, T. & Mashaghi, A. Dry Eye Disease: Emerging Approaches to Disease Analysis and Therapy. *J Clin Med* **8**, 1439 (2019).

5. Gayton, J. L. Etiology, prevalence, and treatment of dry eye disease. *Clin Ophthalmol* **3**, 405–412 (2009).

6. Cheng, H., Guan, X., Chen, D. & Ma, W. The Th17/Treg Cell Balance: A Gut Microbiota-Modulated Story. *Microorganisms* **7**, 583 (2019).

7. Lee, Y. K., Menezes, J. S., Umesaki, Y. & Mazmanian, S. K. Proinflammatory T-cell responses to gut microbiota promote experimental autoimmune encephalomyelitis. *Proc Natl Acad Sci U S A* **108 Suppl 1**, 4615–4622 (2011).

8. Willcox, M. D. P. Characterization of the normal microbiota of the ocular surface. *Experimental Eye Research* **117**, 99–105 (2013).

9. Ozkan, J. *et al.* Temporal Stability and Composition of the Ocular Surface Microbiome. *Sci Rep* **7**, 9880–11 (2017).

10. de Paiva, C. S. *et al.* Altered Mucosal Microbiome Diversity and Disease Severity in Sjögren Syndrome. *Sci Rep* **6**, 23561–11 (2016).

11. Cavuoto, K. M., Banerjee, S. & Galor, A. Relationship between the microbiome and ocular health. *The Ocular Surface* **17**, 384–392 (2019).

12. Baim, A. D., Movahedan, A., Farooq, A. V. & Skondra, D. The microbiome and ophthalmic disease. *Exp. Biol. Med. (Maywood)* **244,** 419–429 (2019).

13. Narayanan, S., Redfern, R. L., Miller, W. L., Nichols, K. K. & McDermott, A. M. Dry Eye Disease and Microbial Keratitis: Is There a Connection? *The Ocular Surface* **11**, 75–92 (2013).

14. Kelly, B. J. *et al.* Power and sample-size estimation for microbiome studies using pairwise distances and PERMANOVA. *Bioinformatics* **31**, 2461–2468 (2015).

15. DeŚantis, T. Z. *et al.* Greengenes, a chimerachecked 16S rRNA gene database and workbench compatible with ARB. *Appl. Environ. Microbiol.* **72**, 5069– 5072 (2006).

16. Glassman, S. I. & Martiny, J. B. H. Broadscale Ecological Patterns Are Robust to Use of Exact Sequence Variants versus Operational Taxonomic Units. *mSphere* **3**, 5409 (2018). 17. Paulson, J. N., Stine, O. C., Bravo, H. C. & Pop, M. Robust methods for differential abundance analysis in marker gene surveys. *Nat. Methods* **10**, 1200–1202 (2013).

18. Postnikoff, C. K., Huisingh, C., McGwin, G. & Nichols, K. K. Leukocyte Distribution in the Open Eye Tears of Normal and Dry Eye Subjects. *Curr. Eye Res.* **43**, 1253–1259 (2018).

19. Ahern, P. P. & Maloy, K. J. Understanding immune-microbiota interactions in the intestine. *Immunology* **22**, 283 (2019).

20. Postnikoff, C. K. & Nichols, K. K. Neutrophil and T-Cell Homeostasis in the Closed Eye. *Invest. Ophthalmol. Vis. Sci.* **58**, 6212–6220 (2017).

21. Shin, H. *et al.* Changes in the Eye Microbiota Associated with Contact Lens Wearing. *mBio* **7**, e00198 (2016).

22. Doan, T. *et al.* Paucibacterial Microbiome and Resident DNA Virome of the Healthy Conjunctiva. *Invest. Ophthalmol. Vis. Sci.* **57**, 5116–5126 (2016).

23. Graham, J. E. *et al.* Ocular pathogen or commensal: a PCR-based study of surface bacterial flora in normal and dry eyes. *Invest. Ophthalmol. Vis. Sci.* **48**, 5616–5623 (2007).

24. Albietz, J. M. & Lenton, L. M. Effect of antibacterial honey on the ocular flora in tear deficiency and meibomian gland disease. *Cornea* **25**, 1012–1019 (2006).

25. Costello, E. K., Stagaman, K., Dethlefsen, Les, Bohannan, B. J. M. & Relman, D. A. The Application of Ecological Theory Toward an Understanding of the Human Microbiome. *Science* **336**, 1255–1262 (2012). 26. McDermott, A. M. Antimicrobial compounds in

tears. *Experimental Eye Research* **117**, 53–61 (2013). 27. Yin, V. T. *et al.* Antibiotic Resistance of Ocular Surface Flora With Repeated Use of a Topical Antibiotic After Intravitreal Injection. *JAMA Ophthalmol* **131**, 456– 461 (2013).

28. lovieno, A. *et al.* Preliminary evidence of the efficacy of probiotic eye-drop treatment in patients with vernal keratoconjunctivitis. *Graefes Arch. Clin. Exp. Ophthalmol.* **246**, 435–441 (2008).

29. Round, J. L. & Palm, N. W. Causal effects of the microbiota on immune-mediated diseases. *Sci Immunol* **3**, eaao1603 (2018).

30. Willis, K. A. *et al.* Perinatal maternal antibiotic exposure augments lung injury in offspring in experimental bronchopulmonary dysplasia. *American Journal of Physiology - Lung Cellular and Molecular Physiology* **3**, 21 (2019).

31. Dolma, K. *et al.* Effects of Hyperoxia on Alveolar and Pulmonary Vascular Development in Germ Free Mice. *American Journal of Physiology - Lung Cellular and Molecular Physiology* **295**, L86 (2019).

32. Wang, Y., Chen, H., Xia, T. & Huang, Y. Characterization of fungal microbiota on normal ocular surface of humans. *Clin. Microbiol. Infect.* (2019). doi:10.1016/j.cmi.2019.05.011

33. Willis, K. A. *et al.* Fungi form interkingdom microbial communities in the primordial human gut that develop with gestational age. *FASEB J.* **33**, 12825–12837 (2019).

34. Lal, C. V. *et al.* The airway microbiome at birth. *nature.com*

35. Chalmers, R. L., Begley, C. G. & Caffery, B. Validation of the 5-Item Dry Eye Questionnaire (DEQ-5): Discrimination across self-assessed severity and aqueous tear deficient dry eye diagnoses. *Contact Lens and Anterior Eye* **33**, 55–60 (2010).

36. Schiffman, R. M., Christianson, M. D., Jacobsen, G., Hirsch, J. D. & Reis, B. L. Reliability and Validity of the Ocular Surface Disease Index. *Arch Ophthalmol* **118**, 615–621 (2000).

37. Miller, W. L. *et al.* A Comparison of Tear Volume (by Tear Meniscus Height and Phenol Red Thread Test) and Tear Fluid Osmolality Measures in Non-Lens Wearers and in Contact Lens Wearers. *Eye & Contact Lens* **30**, 132–137 (2004).

38. Harris, P. A. *et al.* The REDCap consortium: Building an international community of software platform partners. *J Biomed Inform* **95**, 103208 (2019).

39. Postnikoff, C. K., Huisingh, C., McGwin, G. & Nichols, K. K. Leukocyte Distribution in the Open Eye

Tears of Normal and Dry Eye Subjects. *Curr. Eye Res.* **43**, 1253–1259 (2018).

40. Postnikoff, C. K., ophthalmology, K. N. I.2017. Neutrophil and T-cell homeostasis in the closed eye. *iovs.arvojournals.org*

41. Gorbet, M., Postnikoff, C. & Williams, S. The Noninflammatory Phenotype of Neutrophils From the Closed-Eye Environment: A Flow Cytometry Analysis of Receptor Expression. *Invest. Ophthalmol. Vis. Sci.* **56**, 4582–4591 (2015).

42. Caporaso, J. G. *et al.* QIIME allows analysis of high-throughput community sequencing data. *Nat. Methods* **7**, 335–336 (2010).

43. Zakrzewski, M. *et al.* Calypso: a user-friendly web-server for mining and visualizing microbiomeenvironment interactions. *Bioinformatics* **33**, 782–783 (2017).

44. Reese, A. T. & Dunn, R. R. Drivers of Microbiome Biodiversity: A Review of General Rules, Feces, and Ignorance. *mBio* **9**, 734 (2018).

45. Paliy, O. & Shankar, V. Application of multivariate statistical techniques in microbial ecology. *Mol. Ecol.* **25**, 1032–1057 (2016).

46. Anderson, M. J. *Permutational Multivariate Analysis of Variance (PERMANOVA)*. **18**, 1–15 (American Cancer Society, 2014).

Acknowledgements

The authors thank the Microbiome Resource at UAB for the performance of Illumina MiSeq. The following are acknowledged for their support of the Microbiome Resource: UAB School of Medicine, Comprehensive Cancer Center (P30AR050948), Center for AIDS Research (5P30AI027767), Center for Clinical Translational Science (UL1TR000165) and the Heflin Center. We also thank the Children's Foundation Research Institute at Le Bonheur Children's Hospital for providing scientific editing.

Author Contributions

CKP, AG and CVL conceptualized the study. CKP, KN facilitated the performance of the randomized clinical trial and oversaw the sample collection. GR performed the DNA extraction. KAW designed and performed the biostatistical analyses, drafted the manuscript and prepared the figures. CVL directed the performance of the study and provided funding. KAW, CKP, AF, GR, KN, AG and CVL edited the manuscript and approved the final version.

Additional Information

Supplementary information accompanies this preprint.

Competing financial interests. The authors declare no competing financial interests. The original randomized clinical trial was supported in part by Allergan, plc. (Dublin, Ireland). However, this study was an independent work supported by additional funding sources acknowledged above and no member of Allergan, plc. was involved in performance of this study, analysis of the resulting data, preparation of the manuscript or the decision to publish. These endeavors were solely the work of the authors. CKP is currently an employee of CooperVision, Inc. (Pleasanton, CA, USA), which was not involved in this study.

Supplement

The closed eye harbors a unique microbiome in dry eye disease

Kent A. Willis, Cameron K. Postnikoff, Amelia B. Freeman, Gabriel Rezonzew, Kelly K. Nichols, Amit Gaggar, Charitharth V. Lal

Supplemental Fig S1.



Fig. S1. Consolidated standards of reporting trials subject allocation diagram. CONSORT subject allocation diagram. Figure generated by BioRender.

Supplemental Fig S2.





- (a) Raw sequencing data.
- (b) Log₂-transformed relative abundance.
- (c) Sequencing depth.
- (d) Relative abundance at the phylum level.
- (e) Sequence reads per sample.
- (f) Read depth.



Supplemental Fig. S3.



- (a) Alpha diversity.
- (b) Beta diversity.
- (c) Relative abundance of bacterial genera. Two-way ANOVA, * P =0.05, ** P =0.01, *** P = 0.001.
- (d) Core microbiome analysis.
- (e) Discriminant analysis of principal components.
- (f) Spearman network analysis at the genus level.

Supplemental Fig. S4



Fig. S4. Limited changes occur in the normal eye microbiome with eye rinse.

- (a) Alpha diversity.
- (b) Beta diversity.
- (c) Relative abundance of bacterial orders. Two-way ANOVA, * *P* =0.05, ** *P* =0.01, *** *P* = 0.001.
- (d) Core microbiome analysis.
- (e) Discriminant analysis of principal components.
- (f) Spearman network analysis at the genus level.



Fig. S5. The acquisition of unique taxa separates the dry from the normal eye at baseline.

- (a) Negative binomial distribution (DESeq2 function).
- (b) Linear discriminant analysis of effect size (LEfSe).
- (c) Variable importance analysis of random forest classifier.



Supplemental Fig. S6.

Fig. S6. Additional unique taxa separate the dry from the normal eye after a month.

- (a) Negative binomial distribution (DESeq2 function).
- (b) Linear discriminant analysis of effect size (LEfSe).
- (c) Variable importance analysis of random forest classifier.

Supplemental Table 1.

	Normal		Dry Eye	
	Normal	Mild	Moderate	Severe
DEQ5	≤ 5	≤ 5	≥6	≥ 12
	All of the below:		At least one of the	e following:
Phenol Red	> 10, bilateral	> 10, bilateral	< 10, one eye	< 10, one eye
Thread, mm				
NIKBUT, s	> 5, bilateral	> 5, bilateral	< 10, one eye	< 5, one eye
InflammaDry	Negative, both	Positive, one or	Positive, one or	Positive, one or both
	eyes	both eyes	both eyes	eyes

Supplemental Table 2.

	P (biological	Adjusted P	
Taxa (genus)	condition)	(Bonferroni)	FDR
Unclassified.OPB56	< 1E-22	< 1E-22	< 1E-22
Unclassified.Methylobacteriaceae	< 1E-22	1E-21	2.00E-22
Unclassified.Bacteroidetes	< 1E-22	< 1E-22	< 1E-22
Pseudomonas	< 1E-22	< 1E-22	< 1E-22
Meiothermus	< 1E-22	< 1E-22	< 1E-22
Hydrogenophilus	2.5E-19	4.6E-17	7.7E-18
Unclassified.Chitinophagaceae	7.6E-19	1.4E-16	2E-17
Bradyrhizobium	6E-18	1.1E-15	1.4E-16
Unclassified	3.6E-16	6.6E-14	7.4E-15
Unclassified.Aeromonadaceae	7.2E-16	1.3E-13	1.3E-14
Raphanus	2.7E-15	5E-13	4.5E-14
Unclassified.Coriobacteriaceae	1.4E-14	2.6E-12	2.1E-13
Unclassified.Desulfovibrionaceae	4.8E-13	8.8E-11	6.8E-12
Brevundimonas	5.8E-12	1.1E-09	7.6E-11
Lactobacillus	5.4E-11	9.9E-09	6.6E-10
Leuconostoc	1.7E-10	0.00000031	0.00000002
Allobaculum	5.9E-10	0.0000011	6.4E-09
Unclassified.Beijerinckiaceae	1.1E-09	0.000002	0.00000011
Unclassified.Aurantimonadaceae	1.4E-09	0.0000026	0.00000014
Shewanella	1.8E-09	0.0000033	0.00000015
Corynebacterium	1.8E-09	0.0000033	0.00000015
Acinetobacter	1.8E-09	0.0000033	0.00000015
Roseomonas	2.2E-09	0.0000004	0.00000018
Phenylobacterium	4.7E-09	0.0000086	0.00000036
Azospirillum	5.8E-09	0.0000011	0.00000043
Aerococcus	0.00000025	0.0000046	0.0000018
Lysobacter	0.00000043	0.0000079	0.0000029
Pleomorphomonas	0.00000063	0.000012	0.00000041
Brevibacterium	0.00000068	0.000013	0.00000043
Morganella	0.00000094	0.000017	0.00000058
Symbiobacterium	0.00000097	0.000018	0.0000058
Peredibacter	0.000001	0.000018	0.00000058
Bacillus	0.0000012	0.000022	0.00000067
Unclassified.Solibacterales	0.0000024	0.000044	0.0000013
Jeotgalicoccus	0.0000031	0.000057	0.0000016
Sedimentibacter	0.0000088	0.00016	0.0000045
Caloramator	0.000001	0.00018	0.000005

Wolbachia	0.0000012	0.00022	0.0000058
Unclassified.Caulobacteraceae	0.0000013	0.00024	0.0000061
Streptococcus	0.0000022	0.0004	0.00001
Unclassified.Veillonellaceae	0.0000024	0.00044	0.000011
Unclassified.Neisseriaceae	0.0000041	0.00075	0.000018
Mycoplana	0.0000043	0.00079	0.000018
Mucispirillum	0.0000046	0.00085	0.000019
Sporomusa	0.0000056	0.001	0.000023
Caldicoprobacter	0.000077	0.0014	0.000031
Caldinitratiruptor	0.000013	0.0024	0.000051
Akkermansia	0.000015	0.0028	0.000058
Anaerococcus	0.000018	0.0033	0.000068
Leptotrichia	0.000024	0.0044	0.000088
Devosia	0.000025	0.0046	0.00009
Unclassified.Halanaerobiales	0.000027	0.005	0.000096
Neisseria	0.000032	0.0059	0.00011
Balneimonas	0.000033	0.0061	0.00011
Unclassified.Lactobacillaceae	0.000038	0.007	0.00013
Methanosarcina	0.000045	0.0083	0.00015
Desulfosporosinus	0.000048	0.0088	0.00015
Dietzia	0.000087	0.016	0.00028
Lactococcus	0.000093	0.017	0.00029
Unclassified.MLE112	0.00012	0.022	0.00037
Ruminococcus	0.00015	0.028	0.00045
Gordonia	0.00015	0.028	0.00045
Mycoplasma	0.00018	0.033	0.00053
Sutterella	0.00019	0.035	0.00055
Paenibacillus	0.00025	0.046	0.0007
Finegoldia	0.00025	0.046	0.0007
Unclassified.Paenibacillaceae	0.00027	0.05	0.00074
Rubrobacter	0.00032	0.059	0.00087
Helicobacter	0.0004	0.074	0.0011
Unclassified.EtOH8	0.00046	0.085	0.0012
Actinobaculum	0.00056	0.1	0.0015
Micrococcus	0.00063	0.12	0.0016
Lutispora	0.00078	0.14	0.002
Brachybacterium	0.00088	0.16	0.0022
Sphinaobium	0.0011	0.2	0.0027
Voaesella	0.0012	0.2	0.0029
Inclassified Lachnosniraceae	0.0012	0.22	0.0023
onerassijiea.eaennospiraceae	0.0015	0.24	0.005

Unclassified.Clostridiales	0.0013	0.24	0.003
Staphylococcus	0.0013	0.24	0.003
Moraxella	0.0013	0.24	0.003
Propionibacterium	0.0015	0.28	0.0034
Unclassified.Bacillales	0.0017	0.31	0.0038
Agrobacterium	0.0018	0.33	0.004
Rothia	0.002	0.37	0.0044
Unclassified.Barnesiellaceae	0.0021	0.39	0.0045
Unclassified.Acetobacteraceae	0.0021	0.39	0.0045
Alloiococcus	0.0026	0.48	0.0055
Unclassified.Aerococcaceae	0.0029	0.53	0.0061
Peptostreptococcus	0.0034	0.63	0.007
Escherichia	0.0037	0.68	0.0076
Coprococcus	0.0048	0.88	0.0096
Adlercreutzia	0.0048	0.88	0.0096
Oscillospira	0.0052	0.96	0.01
Comamonas	0.0063	1	0.012
Clostridium	0.0067	1	0.013
Unclassified.ML615J28	0.0068	1	0.013
Unclassified.Rikenellaceae	0.0083	1	0.016
Syntrophomonas	0.009	1	0.017
Unclassified.Ruminococcaceae	0.01	1	0.019
SMB53	0.011	1	0.02
Vibrio	0.012	1	0.022
Paraprevotella	0.012	1	0.022
Bacteroides	0.014	1	0.025
Unclassified.S247	0.016	1	0.028
Pantoea	0.016	1	0.028
Paracoccus	0.018	1	0.031
Haemophilus	0.018	1	0.031
Unclassified.03196G20	0.02	1	0.034
Rubellimicrobium	0.021	1	0.035
Peptoniphilus	0.022	1	0.037
Blautia	0.023	1	0.038
Candidatus_Arthromitus	0.028	1	0.046
Citrobacter	0.034	1	0.055
Veillonella	0.035	1	0.056
Capnocytophaga	0.036	1	0.058
Parabacteroides	0.037	1	0.058
Lachnospira	0.037	1	0.058

Roseburia	0.04	1	0.062
Anaeroplasma	0.047	1	0.073
Unclassified.Streptophyta	0.049	1	0.074
Unclassified.Gemellaceae	0.049	1	0.074
Enterococcus	0.049	1	0.074
Turicibacter	0.05	1	0.074
AF12	0.05	1	0.074
Collinsella	0.059	1	0.086
Chryseobacterium	0.059	1	0.086
Eubacterium	0.063	1	0.091
Kocuria	0.067	1	0.096
Desulfovibrio	0.073	1	0.1
Pelosinus	0.079	1	0.11
Faecalibacterium	0.08	1	0.11
Eikenella	0.08	1	0.11
Phascolarctobacterium	0.087	1	0.12
WAL_1855D	0.096	1	0.13
Arthrobacter	0.11	1	0.15
Shuttleworthia	0.13	1	0.17
Selenomonas	0.13	1	0.17
Actinomyces	0.13	1	0.17
Bifidobacterium	0.14	1	0.19
Lautropia	0.15	1	0.2
Unclassified.RF32	0.16	1	0.21
Sneathia	0.16	1	0.21
Phyllobacterium	0.17	1	0.22
Granulicatella	0.17	1	0.22
Varibaculum	0.18	1	0.23
Abiotrophia	0.18	1	0.23
Unclassified.Erysipelotrichaceae	0.19	1	0.24
Dorea	0.19	1	0.24
Dermabacter	0.2	1	0.25
X168	0.24	1	0.29
Unclassified.Bacteroidales	0.24	1	0.29
Achromobacter	0.26	1	0.31
Geobacter	0.28	1	0.34
Calothrix	0.32	1	0.38
Megasphaera	0.33	1	0.39
Aggregatibacter	0.34	1	0.4
Rheinheimera	0.39	1	0.45

Facklamia	0.39	1	0.45
Acetobacter	0.39	1	0.45
Dermacoccus	0.43	1	0.49
Stenotrophomonas	0.46	1	0.53
Unclassified.YS2	0.47	1	0.53
Enhydrobacter	0.47	1	0.53
Odoribacter	0.49	1	0.55
Dehalobacterium	0.49	1	0.55
Anaerostipes	0.54	1	0.6
Carnobacterium	0.56	1	0.62
Porphyromonas	0.58	1	0.63
Bilophila	0.58	1	0.63
rc44	0.59	1	0.64
Salinispora	0.61	1	0.65
Chroococcidiopsis	0.61	1	0.65
Prevotella	0.62	1	0.66
Atopobium	0.64	1	0.68
Fusobacterium	0.68	1	0.71
Campylobacter	0.69	1	0.72
Acidaminococcus	0.75	1	0.78
Methylobacterium	0.79	1	0.82
Mycobacterium	0.89	1	0.91
Proteus	0.9	1	0.92
Sphingomonas	0.91	1	0.93
Janthinobacterium	0.92	1	0.93
Dialister	0.93	1	0.94
Tissierella_Soehngenia	0.95	1	0.95

Supplemental Table 3.

	Р	FDR	
Таха	Assignment	Assignment	P adjusted Assignment
Unclassified.OPB56	< 1E-22	< 1E-22	< 1E-22
Unclassified.Bacteroidetes	< 1E-22	< 1E-22	< 1E-22
Pseudomonas	< 1E-22	< 1E-22	1E-22
Meiothermus	< 1E-22	2E-22	9E-22
Unclassified.Methylobacteriaceae	2E-22	6.3E-21	3.1E-20
Hydrogenophilus	5.8E-18	1.8E-16	1E-15
Unclassified.Chitinophagaceae	7.1E-18	1.9E-16	1.3E-15
Bradyrhizobium	1.4E-16	3.2E-15	2.5E-14
Unclassified	7.9E-15	1.6E-13	1.4E-12

Unclassified.Aeromonadaceae	1.1E-14	2E-13	1.9E-12
Raphanus	6.4E-14	1.1E-12	1.1E-11
Unclassified.Coriobacteriaceae	2.4E-13	3.7E-12	4.2E-11
Unclassified.Desulfovibrionaceae	8.6E-12	1.2E-10	1.5E-09
Brevundimonas	1.2E-10	1.6E-09	0.00000021
Lactobacillus	4.7E-10	5.8E-09	0.0000008
Unclassified.Aurantimonadaceae	0.00000002	0.00000023	0.0000034
Allobaculum	2.9E-09	0.00000031	0.0000049
Leuconostoc	3.1E-09	0.00000032	0.0000052
Unclassified.Beijerinckiaceae	0.00000009	0.00000087	0.0000015
Corynebacterium	0.00000021	0.00000019	0.000035
Shewanella	0.00000023	0.0000002	0.000038
Acinetobacter	0.00000028	0.0000023	0.0000046
Azospirillum	0.00000057	0.00000046	0.000092
Phenylobacterium	0.00000068	0.00000052	0.000011
Roseomonas	0.00000012	0.0000088	0.000019
Bacillus	0.0000023	0.0000016	0.000037
Lysobacter	0.0000025	0.0000017	0.00004
Aerococcus	0.0000033	0.0000022	0.000052
Pleomorphomonas	0.000006	0.000038	0.000094
Morganella	0.00000094	0.0000058	0.00015
Peredibacter	0.00000099	0.0000059	0.00015
Brevibacterium	0.0000022	0.000013	0.00034
Symbiobacterium	0.0000024	0.000013	0.00036
Unclassified.Solibacterales	0.0000033	0.000018	0.0005
Unclassified.Caulobacteraceae	0.0000096	0.00005	0.0014
Wolbachia	0.000014	0.00007	0.0021
Streptococcus	0.000014	0.00007	0.0021
Akkermansia	0.000018	0.000087	0.0026
Mycoplana	0.000024	0.00011	0.0035
Jeotgalicoccus	0.000033	0.00015	0.0048
Mucispirillum	0.000051	0.00023	0.0073
Unclassified.Neisseriaceae	0.000069	0.0003	0.0099
Caldinitratiruptor	0.00012	0.00051	0.017
Leptotrichia	0.00018	0.00075	0.025
Unclassified.Lactobacillaceae	0.00019	0.00076	0.027
Balneimonas	0.00019	0.00076	0.027
Devosia	0.00024	0.00094	0.033
Neisseria	0.00029	0.0011	0.04
Ruminococcus	0.00056	0.0021	0.076

Lactococcus	0.0008	0.0029	0.11
Dietzia	0.0008	0.0029	0.11
Anaerococcus	0.00089	0.0031	0.12
Sutterella	0.0014	0.0049	0.18
Mycoplasma	0.0019	0.0064	0.25
Finegoldia	0.0019	0.0064	0.25
Unclassified.Veillonellaceae	0.002	0.0066	0.26
Paenibacillus	0.0023	0.0073	0.29
Caloramator	0.0023	0.0073	0.29
Gordonia	0.0024	0.0075	0.3
Rubrobacter	0.0026	0.008	0.32
Unclassified.Halanaerobiales	0.0038	0.011	0.47
Caldicoprobacter	0.0038	0.011	0.47
Helicobacter	0.004	0.012	0.49
Actinobaculum	0.0044	0.013	0.53
Micrococcus	0.0048	0.014	0.58
Sporomusa	0.0052	0.014	0.62
Brachybacterium	0.0076	0.021	0.9
Sedimentibacter	0.0082	0.022	0.96
Unclassified.Lachnospiraceae	0.0084	0.022	0.97
Unclassified.MLE112	0.0089	0.023	1
Staphylococcus	0.0096	0.025	1
Vogesella	0.0099	0.025	1
Desulfosporosinus	0.01	0.025	1
Unclassified.Clostridiales	0.011	0.027	1
Unclassified.Acetobacteraceae	0.012	0.029	1
Unclassified.Aerococcaceae	0.013	0.031	1
Propionibacterium	0.014	0.033	1
Agrobacterium	0.014	0.033	1
Unclassified.Paenibacillaceae	0.015	0.034	1
Rothia	0.015	0.034	1
Moraxella	0.015	0.034	1
Unclassified.Bacillales	0.018	0.04	1
Alloiococcus	0.018	0.04	1
Peptostreptococcus	0.025	0.055	1
Escherichia	0.028	0.061	1
Coprococcus	0.03	0.064	1
Adlercreutzia	0.031	0.066	1
Unclassified.Barnesiellaceae	0.032	0.067	1
Sphingobium	0.041	0.084	1

Contantionals 0.041 0.044 1 Unclassified.EtOH8 0.046 0.092 1 Bacteroides 0.049 0.092 1 Oscillospira 0.049 0.092 1 Vibrio 0.05 0.098 1 Unclassified.Rikenellaceae 0.055 0.11 1 Methanosarcina 0.066 0.12 1 Unclassified.Ruminococcaceae 0.065 0.12 1 Unclassified.S247 0.073 0.14 1 Blautia 0.083 0.15 1 Lutispora 0.098 0.18 1 Veillonella 0.1 0.18 1 SMB53 0.1 0.18 1 Haemophilus 0.11 0.19 1 Unclassified.ML615J28 0.12 0.2 1 Paracoccus 0.12 0.2 1 Anaeroplasma 0.12 0.2 1 Anaeroplasma 0.15 0.25 1
Onlossified.Etons 0.046 0.092 1 Bacteroides 0.046 0.092 1 Oscillospira 0.049 0.097 1 Vibrio 0.05 0.098 1 Unclassified.Rikenellaceae 0.055 0.11 1 Methanosarcina 0.06 0.12 1 Unclassified.Ruminococcaceae 0.065 0.12 1 Unclassified.S247 0.073 0.14 1 Blautia 0.083 0.15 1 Lutispora 0.098 0.18 1 Veillonella 0.1 0.18 1 SMB53 0.1 0.18 1 Rubellimicrobium 0.11 0.19 1 Paraprevotella 0.12 0.2 1 Unclassified.ML615J28 0.12 0.2 1 Anaeroplasma 0.12 0.2 1 Qandidatus_Arthromitus 0.14 0.24 1 Parabacteroides 0.18 0.29
Batterbildes 0.046 0.092 1 Oscillospira 0.049 0.097 1 Vibrio 0.05 0.098 1 Unclassified.Rikenellaceae 0.055 0.11 1 Methanosarcina 0.066 0.12 1 Unclassified.Suminococcaceae 0.065 0.12 1 Unclassified.S247 0.073 0.14 1 Blautia 0.083 0.15 1 Lutispora 0.098 0.18 1 Veillonella 0.1 0.18 1 SMB53 0.1 0.18 1 Rubellimicrobium 0.11 0.19 1 Paraprevotella 0.11 0.19 1 Unclassified.ML615J28 0.12 0.2 1 Anaeroplasma 0.15 0.25 1 Paratoccus 0.15 0.25 1 Paratoea 0.15 0.25 1 Paratoea 0.16 0.26 1
Oscinospira 0.049 0.097 1 Vibrio 0.05 0.098 1 Unclassified.Rikenellaceae 0.055 0.11 1 Methanosarcina 0.06 0.12 1 Unclassified.Ruminococcaceae 0.065 0.12 1 Unclassified.S247 0.073 0.14 1 Blautia 0.083 0.15 1 Lutispora 0.098 0.18 1 Veillonella 0.1 0.18 1 SMB53 0.1 0.18 1 Rubellimicrobium 0.11 0.19 1 Paraprevotella 0.11 0.19 1 Unclassified.ML615J28 0.12 0.2 1 Paracoccus 0.12 0.2 1 Anaeroplasma 0.12 0.2 1 Candidatus_Arthromitus 0.14 0.24 1 Partoea 0.15 0.25 1 Peptoniphilus 0.16 0.26 1
Vibrio 0.05 0.058 0.058 1 Unclassified.Rikenellaceae 0.055 0.11 1 Methanosarcina 0.06 0.12 1 Unclassified.Ruminococcaceae 0.065 0.12 1 Unclassified.S247 0.073 0.14 1 Blautia 0.083 0.15 1 Lutispora 0.098 0.18 1 Veillonella 0.1 0.18 1 SMB53 0.1 0.18 1 Haemophilus 0.11 0.19 1 Rubellimicrobium 0.11 0.19 1 Paraprevotella 0.12 0.2 1 Anaeroplasma 0.12 0.2 1 Anaeroplasma 0.15 0.25 1 Petoniphilus 0.16 0.26 1 Parabacteroides 0.18 0.29 1 Syntrophomonas 0.19 0.31 1
Unclassified.Rikenellacede 0.055 0.11 1 Methanosarcina 0.06 0.12 1 Unclassified.Ruminococcaceae 0.065 0.12 1 Unclassified.S247 0.073 0.14 1 Blautia 0.083 0.15 1 Lutispora 0.098 0.18 1 Veillonella 0.1 0.18 1 SMB53 0.1 0.18 1 Haemophilus 0.11 0.19 1 Rubellimicrobium 0.11 0.19 1 Paraprevotella 0.12 0.2 1 Unclassified.ML615J28 0.12 0.2 1 Paracoccus 0.12 0.2 1 Anaeroplasma 0.12 0.2 1 Pantoea 0.15 0.25 1 Paraboterium 0.15 0.25 1 Peptoniphilus 0.16 0.26 1 Parabacteroides 0.18 0.29 1
Methanosarcina 0.06 0.12 1 Unclassified.Ruminococcaceae 0.065 0.12 1 Unclassified.S247 0.073 0.14 1 Blautia 0.083 0.15 1 Lutispora 0.098 0.18 1 Veillonella 0.1 0.18 1 SMB53 0.1 0.18 1 Haemophilus 0.1 0.18 1 Rubellimicrobium 0.11 0.19 1 Paraprevotella 0.12 0.2 1 Paracoccus 0.12 0.2 1 Anaeroplasma 0.12 0.2 1 Pantoea 0.15 0.25 1 Paraboateroides 0.18 0.29 1 Parabacteroides 0.18 0.29 1
Unclassified.Ruminococcaceae 0.065 0.12 1 Unclassified.S247 0.073 0.14 1 Blautia 0.083 0.15 1 Lutispora 0.098 0.18 1 Veillonella 0.1 0.18 1 SMB53 0.1 0.18 1 Haemophilus 0.1 0.18 1 Rubellimicrobium 0.11 0.19 1 Paraprevotella 0.12 0.2 1 Unclassified.ML615J28 0.12 0.2 1 Paracoccus 0.12 0.2 1 Anaeroplasma 0.12 0.2 1 Candidatus_Arthromitus 0.14 0.24 1 Pantoea 0.15 0.25 1 Peptoniphilus 0.16 0.26 1 Parabacteroides 0.18 0.29 1 Junclassified.NL615J28 0.16 0.26 1 Pantoea 0.15 0.25 1
Unclassified.S247 0.073 0.14 1 Blautia 0.083 0.15 1 Lutispora 0.098 0.18 1 Veillonella 0.1 0.18 1 SMB53 0.1 0.18 1 Haemophilus 0.1 0.18 1 Rubellimicrobium 0.11 0.19 1 Paraprevotella 0.11 0.19 1 Unclassified.ML615J28 0.12 0.2 1 Paracoccus 0.12 0.2 1 Anaeroplasma 0.12 0.2 1 Candidatus_Arthromitus 0.14 0.24 1 Partoea 0.15 0.25 1 Peptoniphilus 0.16 0.26 1 Parabacteroides 0.18 0.29 1 Syntrophomonas 0.19 0.31 1 Lachnospira 0.2 0.32 1
Blautia 0.083 0.15 1 Lutispora 0.098 0.18 1 Veillonella 0.1 0.18 1 SMB53 0.1 0.18 1 Haemophilus 0.1 0.18 1 Rubellimicrobium 0.11 0.19 1 Paraprevotella 0.11 0.19 1 Unclassified.ML615J28 0.12 0.2 1 Paracoccus 0.12 0.2 1 Anaeroplasma 0.12 0.2 1 Candidatus_Arthromitus 0.14 0.24 1 Partoea 0.15 0.25 1 Faecalibacterium 0.16 0.26 1 Parabacteroides 0.18 0.29 1 Syntrophomonas 0.19 0.31 1 Lachnospira 0.2 0.32 1
Lutispora 0.098 0.18 1 Veillonella 0.1 0.18 1 SMB53 0.1 0.18 1 Haemophilus 0.1 0.18 1 Rubellimicrobium 0.11 0.19 1 Paraprevotella 0.11 0.19 1 Unclassified.ML615J28 0.12 0.2 1 Paracoccus 0.12 0.2 1 Anaeroplasma 0.12 0.2 1 Candidatus_Arthromitus 0.14 0.24 1 Partoea 0.15 0.25 1 Peptoniphilus 0.16 0.26 1 Parabacteroides 0.18 0.29 1 Syntrophomonas 0.19 0.31 1 Lachnospira 0.2 0.32 1
Veillonella 0.1 0.18 1 SMB53 0.1 0.18 1 Haemophilus 0.1 0.18 1 Rubellimicrobium 0.11 0.19 1 Paraprevotella 0.11 0.19 1 Unclassified.ML615J28 0.12 0.2 1 Paracoccus 0.12 0.2 1 Anaeroplasma 0.12 0.2 1 Candidatus_Arthromitus 0.14 0.24 1 Pantoea 0.15 0.25 1 Faecalibacterium 0.16 0.26 1 Parabacteroides 0.18 0.29 1 Syntrophomonas 0.19 0.31 1 Lachnospira 0.2 0.32 1
SMB53 0.1 0.18 1 Haemophilus 0.1 0.18 1 Rubellimicrobium 0.11 0.19 1 Paraprevotella 0.11 0.19 1 Unclassified.ML615J28 0.12 0.2 1 Paracoccus 0.12 0.2 1 Anaeroplasma 0.12 0.2 1 Candidatus_Arthromitus 0.14 0.24 1 Patoea 0.15 0.25 1 Faecalibacterium 0.16 0.26 1 Parabacteroides 0.18 0.29 1 Syntrophomonas 0.19 0.31 1 Lachnospira 0.2 0.32 1
Haemophilus 0.1 0.18 1 Rubellimicrobium 0.11 0.19 1 Paraprevotella 0.11 0.19 1 Unclassified.ML615J28 0.12 0.2 1 Paracoccus 0.12 0.2 1 Anaeroplasma 0.12 0.2 1 Candidatus_Arthromitus 0.14 0.24 1 Pantoea 0.15 0.25 1 Faecalibacterium 0.16 0.26 1 Parabacteroides 0.18 0.29 1 Syntrophomonas 0.19 0.31 1 Lachnospira 0.2 0.32 1
Rubellimicrobium 0.11 0.19 1 Paraprevotella 0.11 0.19 1 Unclassified.ML615J28 0.12 0.2 1 Paracoccus 0.12 0.2 1 Anaeroplasma 0.12 0.2 1 Candidatus_Arthromitus 0.14 0.24 1 Pantoea 0.15 0.25 1 Faecalibacterium 0.16 0.26 1 Parabacteroides 0.18 0.29 1 Syntrophomonas 0.19 0.31 1 Lachnospira 0.2 0.32 1
Paraprevotella 0.11 0.19 1 Unclassified.ML615J28 0.12 0.2 1 Paracoccus 0.12 0.2 1 Anaeroplasma 0.12 0.2 1 Candidatus_Arthromitus 0.14 0.24 1 Pantoea 0.15 0.25 1 Faecalibacterium 0.15 0.25 1 Peptoniphilus 0.16 0.26 1 Parabacteroides 0.18 0.29 1 Syntrophomonas 0.19 0.31 1 Lachnospira 0.2 0.32 1
Unclassified.ML615J28 0.12 0.2 1 Paracoccus 0.12 0.2 1 Anaeroplasma 0.12 0.2 1 Candidatus_Arthromitus 0.14 0.24 1 Pantoea 0.15 0.25 1 Faecalibacterium 0.15 0.25 1 Peptoniphilus 0.16 0.26 1 Parabacteroides 0.18 0.29 1 Syntrophomonas 0.19 0.31 1 Lachnospira 0.2 0.32 1
Paracoccus0.120.21Anaeroplasma0.120.21Candidatus_Arthromitus0.140.241Pantoea0.150.251Faecalibacterium0.150.251Peptoniphilus0.160.261Parabacteroides0.180.291Syntrophomonas0.190.311Lachnospira0.20.321
Anaeroplasma0.120.21Candidatus_Arthromitus0.140.241Pantoea0.150.251Faecalibacterium0.150.251Peptoniphilus0.160.261Parabacteroides0.180.291Syntrophomonas0.190.311Lachnospira0.20.321
Candidatus_Arthromitus 0.14 0.24 1 Pantoea 0.15 0.25 1 Faecalibacterium 0.15 0.25 1 Peptoniphilus 0.16 0.26 1 Parabacteroides 0.18 0.29 1 Syntrophomonas 0.19 0.31 1 Lachnospira 0.2 0.32 1
Pantoea 0.15 0.25 1 Faecalibacterium 0.15 0.25 1 Peptoniphilus 0.16 0.26 1 Parabacteroides 0.18 0.29 1 Syntrophomonas 0.19 0.31 1 Lachnospira 0.2 0.32 1
Faecalibacterium 0.15 0.25 1 Peptoniphilus 0.16 0.26 1 Parabacteroides 0.18 0.29 1 Syntrophomonas 0.19 0.31 1 Lachnospira 0.2 0.32 1
Peptoniphilus 0.16 0.26 1 Parabacteroides 0.18 0.29 1 Syntrophomonas 0.19 0.31 1 Lachnospira 0.2 0.32 1
Parabacteroides 0.18 0.29 1 Syntrophomonas 0.19 0.31 1 Lachnospira 0.2 0.32 1
Syntrophomonas 0.19 0.31 1 Lachnospira 0.2 0.32 1
<i>Lachnospira</i> 0.2 0.32 1
<i>Capnocytophaga</i> 0.2 0.32 1
<i>Turicibacter</i> 0.21 0.33 1
<i>Roseburia</i> 0.22 0.34 1
Unclassified.Bacteroidales 0.23 0.35 1
Phascolarctobacterium 0.23 0.35 1
Unclassified.Gemellaceae 0.24 0.36 1
<i>Collinsella</i> 0.25 0.38 1
<i>Eubacterium</i> 0.28 0.41 1
Chryseobacterium 0.28 0.41 1
AF12 0.28 0.41 1
Unclassified.Streptophyta 0.29 0.42 1
Unclassified.03196G20 0.3 0.43 1
Dorea 0.32 0.46 1
<i>Desulfovibrio</i> 0.37 0.53 1

WAL_1855D	0.4	0.57	1
Enterococcus	0.44	0.61	1
Eikenella	0.44	0.61	1
Bifidobacterium	0.44	0.61	1
Citrobacter	0.45	0.62	1
Kocuria	0.48	0.65	1
Shuttleworthia	0.49	0.65	1
Clostridium	0.49	0.65	1
Arthrobacter	0.49	0.65	1
Actinomyces	0.55	0.73	1
Selenomonas	0.57	0.74	1
Lautropia	0.57	0.74	1
Unclassified.Erysipelotrichaceae	0.58	0.75	1
Prevotella	0.58	0.75	1
Sneathia	0.59	0.75	1
Unclassified.RF32	0.6	0.76	1
Varibaculum	0.62	0.78	1
Granulicatella	0.62	0.78	1
Abiotrophia	0.64	0.8	1
Pelosinus	0.67	0.83	1
Dermabacter	0.76	0.93	1
Achromobacter	0.8	0.97	1
Calothrix	0.84	1	1
Geobacter	0.85	1	1
Rheinheimera	0.87	1	1
Megasphaera	0.87	1	1
Bilophila	0.9	1	1
X168	0.91	1	1
Acetobacter	0.92	1	1
Dermacoccus	0.93	1	1
Stenotrophomonas	0.94	1	1
rc44	0.95	1	1
Unclassified.YS2	0.97	1	1
Odoribacter	0.97	1	1
Facklamia	0.97	1	1
Enhydrobacter	0.97	1	1
Phyllobacterium	0.98	1	1
Dehalobacterium	0.98	1	1
Tissierella_Soehngenia	1	1	1
Sphingomonas	1	1	1

Salinispora	1	1	1
Proteus	1	1	1
Porphyromonas	1	1	1
Mycobacterium	1	1	1
Methylobacterium	1	1	1
Janthinobacterium	1	1	1
Fusobacterium	1	1	1
Dialister	1	1	1
Chroococcidiopsis	1	1	1
Carnobacterium	1	1	1
Campylobacter	1	1	1
Atopobium	1	1	1
Anaerostipes	1	1	1
Aggregatibacter	1	1	1
Acidaminococcus	1	1	1

Supplemental Table 4.

	Fold	Adjusted P		
Taxa (Genus)	Change	P (DESeq2)	(Bonferroni)	FDR
Unclassified.Methylobacteriaceae	25.81	< 1E-22	< 1E-22	< 1E-22
Pseudomonas	18.38	< 1E-22	< 1E-22	< 1E-22
Azospirillum	102.9	3.5E-21	6.3E-19	1.9E-19
Bradyrhizobium	3.2	4.3E-21	7.8E-19	1.9E-19
Unclassified.Coriobacteriaceae	-78.88	1.9E-19	3.5E-17	7E-18
Unclassified.Bacteroidetes	-8.057	7.7E-19	1.4E-16	2.3E-17
Unclassified	1.389	8.8E-16	1.6E-13	1.8E-14
Unclassified.Beijerinckiaceae	39600	9E-16	1.6E-13	1.8E-14
Unclassified.Aeromonadaceae	-20.23	2E-15	3.7E-13	3.7E-14
Agrobacterium	5.308	1.8E-14	3.3E-12	3E-13
Unclassified.Desulfovibrionaceae	-36.35	2.1E-14	3.8E-12	3.2E-13
Phenylobacterium	69	7E-14	1.3E-11	9.8E-13
Lysobacter	12.23	1.3E-11	2.4E-09	1.7E-10
Allobaculum	-22.21	2.2E-11	3.9E-09	2.6E-10
Peredibacter	25.57	2.5E-11	4.5E-09	2.8E-10
Unclassified.Aurantimonadaceae	16.52	2.9E-11	5.3E-09	3.1E-10
Roseomonas	-21.81	1.1E-10	0.00000021	1.2E-09
Mucispirillum	-21.21	1.7E-10	0.00000031	1.6E-09
Unclassified.Caulobacteraceae	15.83	2.3E-10	0.00000042	2.1E-09
Akkermansia	-7.868	3.9E-10	0.0000007	3.3E-09
Mycoplana	3.14	8.5E-10	0.0000015	0.00000007

Meiothermus	-9.829	1.3E-09	0.0000023	9.7E-09
Corynebacterium	-4.932	1.3E-09	0.0000023	9.7E-09
Brevundimonas	-33.23	2.2E-09	0.0000004	0.000000016
Brevibacterium	-8.647	0.00000014	0.0000026	0.0000001
Unclassified.Chitinophagaceae	-20.34	0.00000024	0.0000044	0.00000016
Escherichia	1.325	0.00000041	0.0000074	0.00000026
Hydrogenophilus	-6.369	0.0000022	0.00004	0.0000014
Unclassified.OPB56	-8.359	0.00000024	0.000043	0.0000014
Leuconostoc	-8.183	0.0000005	0.000091	0.0000029
Unclassified.Lachnospiraceae	-1.012	0.0000058	0.0001	0.0000032
Methanosarcina	4.233	0.00000059	0.00011	0.0000032
Shewanella	-20.13	0.00000079	0.00014	0.0000042
Clostridium	1.871	0.0000023	0.00042	0.000012
Unclassified.EtOH8	6.491	0.0000027	0.00049	0.000014
Morganella	-5.069	0.0000036	0.00065	0.000018
Pleomorphomonas	8.764	0.0000044	0.00079	0.000021
Ruminococcus	1.151	0.0000097	0.0018	0.000045
Balneimonas	6.021	0.000012	0.0022	0.000054
Vibrio	-2.046	0.000018	0.0033	0.000081
Lutispora	4.099	0.000033	0.0061	0.00014
Anaerococcus	-3.812	0.000045	0.0081	0.00019
Jeotgalicoccus	-6.116	0.0001	0.019	0.00043
Paraprevotella	2.8	0.00014	0.026	0.00058
Aerococcus	-8.985	0.00023	0.042	0.0009
Vogesella	4.328	0.00024	0.043	0.0009
Propionibacterium	-3.183	0.00024	0.043	0.0009
Rubrobacter	-6.067	0.00025	0.046	0.00093
Coprococcus	1.133	0.00026	0.046	0.00093
Unclassified.Halanaerobiales	4.103	0.00029	0.053	0.001
Unclassified.Acetobacteraceae	13.38	0.0003	0.054	0.001
Rubellimicrobium	4.332	0.0003	0.054	0.001
Unclassified.Ruminococcaceae	1.386	0.00031	0.056	0.001
Gordonia	-1.118	0.00034	0.061	0.0011
Rheinheimera	1.18	0.00036	0.065	0.0012
Lactobacillus	-4.775	0.00046	0.083	0.0015
Lactococcus	-3.43	0.00056	0.1	0.0018
Dorea	1.36	0.00062	0.11	0.0019
Sporomusa	1.93	0.00064	0.12	0.0019
Dietzia	-4.13	0.00068	0.12	0.002
Citrobacter	1.767	0.00094	0.17	0.0027

Streptococcus	-2.691	0.001	0.18	0.0029
Anaeroplasma	1.15	0.001	0.18	0.0029
Odoribacter	1.734	0.0013	0.24	0.0037
Syntrophomonas	2.622	0.0017	0.3	0.0046
Neisseria	-5.043	0.0017	0.32	0.0047
Unclassified.Bacteroidales	1.442	0.0023	0.41	0.0061
Caloramator	1.715	0.0026	0.47	0.0068
Alloiococcus	-1.518	0.0026	0.47	0.0068
Helicobacter	-3.416	0.0031	0.57	0.008
Actinobaculum	-7.919	0.0034	0.62	0.0086
Sedimentibacter	1.655	0.0035	0.64	0.0087
Ignatzschineria	-14.77	0.0036	0.64	0.0087
Pantoea	2.364	0.0039	0.71	0.0094
Unclassified.Solibacterales	-6.776	0.005	0.91	0.012
Collinsella	1.413	0.0051	0.92	0.012
Adlercreutzia	-4.107	0.0073	1	0.017
Unclassified.Veillonellaceae	1.702	0.008	1	0.018
Varibaculum	1.724	0.0083	1	0.019
Phascolarctobacterium	1.498	0.01	1	0.023
Faecalibacterium	1.302	0.012	1	0.026
Arthrobacter	-4.483	0.012	1	0.027
Mycoplasma	2.988	0.015	1	0.032
Sphingobium	-3.533	0.019	1	0.04
Kocuria	2.79	0.021	1	0.043
Unclassified.Rikenellaceae	-1.203	0.022	1	0.046
Roseburia	-1.083	0.025	1	0.052
Unclassified.Lactobacillaceae	-3.374	0.027	1	0.055
Pelosinus	-1.596	0.03	1	0.061
AF12	1.225	0.036	1	0.072
Acinetobacter	-3.148	0.038	1	0.075
Geobacter	1.062	0.045	1	0.087
Unclassified.Neisseriaceae	-1.398	0.047	1	0.09
Paracoccus	-1.053	0.054	1	0.1
Enterococcus	-1.179	0.066	1	0.13
Leptotrichia	-3.279	0.076	1	0.14
Parabacteroides	-1.922	0.081	1	0.15
Sutterella	-5.098	0.088	1	0.16
Unclassified.Paenibacillaceae	-1.196	0.11	1	0.19
Symbiobacterium	-4.816	0.11	1	0.2
Staphylococcus	-1.869	0.12	1	0.21

Enhydrobacter	-2.418	0.12	1	0.21
Chryseobacterium	-2.675	0.12	1	0.22
Sphingomonas	-1.643	0.13	1	0.23
Haemophilus	-2.023	0.13	1	0.22
Wolbachia	-2.859	0.14	1	0.23
Aggregatibacter	1.088	0.14	1	0.24
Oscillospira	-1.383	0.15	1	0.25
Phyllobacterium	-1.332	0.16	1	0.26
Mycobacterium	-1.512	0.16	1	0.26
Rothia	-2.088	0.17	1	0.27
Pelotomaculum	-1.032	0.17	1	0.27
Unclassified.Streptophyta	-3.172	0.18	1	0.28
Dialister	1.108	0.19	1	0.29
Desulfosporosinus	-1.338	0.19	1	0.29
Chroococcidiopsis	12.31	0.2	1	0.31
Unclassified.RF39	-1.258	0.21	1	0.32
Carnobacterium	-1.16	0.21	1	0.32
Tepidimicrobium	1.105	0.22	1	0.33
Lautropia	-9.06	0.23	1	0.34
Achromobacter	-1.664	0.23	1	0.34
Unclassified.YS2	-3.569	0.25	1	0.36
Unclassified.Gemellaceae	-2.135	0.25	1	0.36
Capnocytophaga	-1.13	0.25	1	0.36
Tissierella_Soehngenia	-1.76	0.28	1	0.41
Bilophila	-1.311	0.29	1	0.42
Unclassified.Erysipelotrichaceae	1.091	0.34	1	0.48
Methylobacterium	-1.533	0.34	1	0.48
Bifidobacterium	-1.596	0.34	1	0.48
Bacteroides	-1.707	0.35	1	0.49
Unclassified.OPB54	-5.676	0.36	1	0.49
Burkholderia	-2.341	0.37	1	0.51
Lachnospira	-1.483	0.38	1	0.51
Facklamia	-1.657	0.39	1	0.52
Desulfovibrio	-1.809	0.39	1	0.52
Unclassified.MLE112	-3.853	0.4	1	0.53
WAL_1855D	-2.038	0.41	1	0.54
Fusobacterium	-1.756	0.41	1	0.54
Megasphaera	-2.338	0.43	1	0.56
Blautia	-1.234	0.43	1	0.56
Campylobacter	-1.373	0.44	1	0.56

Dermabacter	-4.032	0.45	1	0.57
Proteus	-1.954	0.47	1	0.59
Acidaminococcus	-1.064	0.49	1	0.61
Micrococcus	-2.801	0.52	1	0.65
Sneathia	-2.198	0.56	1	0.69
rc44	1.815	0.57	1	0.7
Veillonella	-1.787	0.59	1	0.72
Turicibacter	-2	0.6	1	0.72
Peptostreptococcus	-2.727	0.6	1	0.72
Dermacoccus	-2.61	0.6	1	0.72
Stenotrophomonas	-2.086	0.63	1	0.74
Unclassified.S247	-1.464	0.65	1	0.76
Unclassified.Bacillales	-1.809	0.66	1	0.76
Finegoldia	-3.337	0.66	1	0.76
Eikenella	-1.672	0.66	1	0.76
Candidatus_Arthromitus	-2.061	0.68	1	0.78
Janthinobacterium	-1.921	0.69	1	0.79
Moraxella	-1.631	0.71	1	0.8
Unclassified.ML615J28	-1.871	0.72	1	0.8
SMB53	-2.048	0.72	1	0.8
Dehalobacterium	-2.649	0.73	1	0.81
Prevotella	-1.739	0.75	1	0.83
Brachybacterium	-2.333	0.76	1	0.84
Peptoniphilus	-2.596	0.78	1	0.85
Anaerostipes	1.011	0.81	1	0.88
Unclassified.03196G20	-4.599	0.82	1	0.88
Unclassified.RF32	-1.961	0.85	1	0.9
Caldicoprobacter	-2.043	0.85	1	0.9
Shuttleworthia	-2.217	0.86	1	0.9
Pseudoxanthomonas	-3.164	0.86	1	0.9
Actinomyces	-1.658	0.86	1	0.9
Selenomonas	-3.757	0.89	1	0.93
Comamonas	-2.271	0.9	1	0.93
Porphyromonas	-1.242	0.94	1	0.97
Acetobacter	-2.159	0.95	1	0.98
Brevibacillus	-2.066	0.96	1	0.98
Atopobium	-1.961	0.97	1	0.98