

When will the battle against novel coronavirus end in Wuhan: a SEIR modeling analysis

Kangkang WAN^{1✉}, Jing CHEN^{2✉}, Changming LU¹, Lanlan DONG¹, Zhicheng WU¹, Lianglu ZHANG^{1✉✉}

¹Wuhan Ammunition Life-Tech Co., Ltd., Wuhan, Hubei, China,

²Clinics of Oilcrops Research Institute, CAAS, Wuhan, Hubei, China

✉First authors who contributed equally to this manuscript

✉✉Correspondence author, whose E-mail is zhangll@whammunition.com

Address: Building B10, High-Tech Medical Equipment Park, No. 818, High-Tech Avenue, Donghu High-Tech Development Zone, Wuhan. Hubei, China, 430200

Abstract

Recent outbreak of 2019-nCoV in Wuhan raised serious public health concerns. By February 15, 2020 in Wuhan, the total number of confirmed infection cases has reached 37,914, and the number of deaths has reached 1123, accounting for 56.9% of the total confirmed cases and 73.7% of the total deaths in China. People are eager to know when the epidemic will be completely controlled and when people's work and life will be on the right track. In this study we analyzed the epidemic dynamics and trend of 2019-nCoV in Wuhan by using the data after the closure of Wuhan city till February 12, 2020 based on the SEIR modeling method. The optimal parameters were estimated as $R_0=1.44$ (interquartile range: 1.40-1.47), $TI=14$ (interquartile range: 14-14) and $TE=3.0$ (interquartile range: 2.8-3.1). Based on these parameters, the number of infected individuals in Wuhan city may reach the peak around February 19 at about 45,000 people. Once entering March, the epidemic would gradually decline, and end around the late March. It is worth noting that the above prediction is based on the

assumption that the number of susceptible population $N = 200,000$ will not increase.

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

If the epidemic situation is not properly controlled, the peak of infected number can be further increased and the peak time will be a little postponed. It was expected that the epidemic would subside in early March, and disappear gradually towards the late March.

Keywords: novel coronavirus; epidemic dynamics; SEIR modeling; Wuhan

Introduction

Wuhan is the largest city in central China with a total population of more than 11 million^[1]. The epidemic of 2019-nCoV pneumonia has been raging in the whole country, especially in Hubei province for nearly a month. In late December 2019, 67 cases of 2019-nCoV pneumonia were reported in Wuhan^[2]. In order to prevent the further spread of 2019-nCoV, Wuhan began to close the city from 10:00 on January 23, banning all vehicles from entering and leaving the city. Tens of thousands of medical staff, soldiers and people from all walks of life have been involved in the campaign.

The spread of the epidemic has caused a huge threat to people's health and life safety, at the same time, it has a serious impact on China's social life and national economy. By February 15, 2020, the total number of confirmed cases has reached 37914, and the number of deaths has reached 1123 in Wuhan, accounting for 56.9% of the total confirmed cases and 73.7% of the total deaths in China^[3]. With the increase of medical staff from all over the country, the opening of several large novel hospitals, and the adoption of anti epidemic measures, more patients can get efficient and timely treatment. The number of confirmed cases increased sharply on February 12 and 13, while the total number of suspected cases decreased gradually^[3].

People are eager to know when the epidemic will be completely controlled and when people's work and life will be on the right track. In order to help the public to understand the future trend of the epidemic, we analyzed the epidemic dynamic and trend of 2019-nCoV in Wuhan city by using the SEIR modeling method based on the actual data and published references.

Methods

1. Epidemic transmission model

The SEIR model is a classical epidemic model for the flows of people between four states: susceptible (S), exposed (E), infected (I), and recovery (R). Each of those variables represents the number of people in those groups. The relationship among the four groups is elucidated in Figure 1, where βI is the probability of S to E after I contacts S, γI is the probability of E to I, and γ_2 is the probability of I to R. Since 2019-nCoV is also infectious in the incubation period, we introduced parameter β_2

here to represent the probability of S to E after E contact S. We used the “susceptible – exposed – infected – recovered” model^[4] to describe the prevalent characteristics of 2019-nCoV in Wuhan .




Fig. 1. Relationship among four groups according to SEIR model

This is an ordinary differential equation model, described by the following equations:

$$\begin{aligned}
 dS(t)/d(t) &= -\beta_1 * I(t) * (S(t)/N) - \beta_2 * E(t) * (S(t)/N) \\
 dE(t)/d(t) &= \beta_1 * I(t) * (S(t)/N) + \beta_2 * E(t) * (S(t)/N) - \gamma_1 * E(t) \\
 dI(t)/d(t) &= \gamma_1 * E(t) - \gamma_2 * I(t) \\
 dR(t)/d(t) &= \gamma_2 * I(t)
 \end{aligned}$$

Among which, $S(t)$, $E(t)$, $I(t)$ and $R(t)$ represent the number of people in the group of the susceptible, the exposed, the infected and the recovered on the day t , respectively. N is the total number of possible contact people, which is assumed to be fixed and $N = S + E + I + R$.

The simulation used the fourth-order Runge-Kutta algorithm^[5] to solve it numerically, with a step size fixed at 0.01 for R_0 , and at 0.1 for TE and TI.

2、 Estimation of parameters for the model

Parameters β_1 , β_2 , γ_1 and γ_2 were estimated according to the reference^[6] using the formula below:

$$\begin{aligned}\beta_1 &= R_0/TI \\ \beta_2 &= R_0/TE \\ \gamma_1 &= 1/TE \\ \gamma_2 &= 1/TI\end{aligned}$$

Among which, R_0 is the basic reproduction number, TI is the time of infectious period, TE is the time of incubation period, and β_1 , β_2 , γ_1 and γ_2 share the same meanings as in Figure 1. We set the range of TE to 1-7 days, TI to 1-14 days, R_0 to 1-5 days according to the reference^[7].

The optimized parameters of TE , TI and R_0 were determined on the condition of RMSE being the smallest.

$$\begin{aligned}RMSE_{(I)} &= \sqrt{1/n \sum (\hat{I} - I)^2} \\ RMSE_{(R)} &= \sqrt{1/n \sum (\hat{R} - R)^2}\end{aligned}$$

Where I and $\hat{I}$ indicate the real and simulated number of the infected, respectively while the R and $\hat{R}$ indicate the real and simulated number of the recovery in a specific time, respectively.

Calculate the real number of people (I and R) and the smallest RMSE of $\hat{I}$ and $\hat{R}$ obtained by the model, and choose the TE , TI and R_0 with the smallest RMSE as the optimal model parameters.

In order to avoid over-fitting in parameter optimization, we took 80% data each time, and repeated this process 100 times, and finally took the average value of R_0 , TI and TE of 100 times as the optimal parameter values.

3、Data source

The data were collected from the official website of Hubei Provincial Health Committee (<http://wjw.hubei.gov.cn/>)^[3], and shown in Table 1. We used the data of 22 days from January 22 to February 12 when Wuhan city was shut down and all the public transportation was suspended.

For construction of the model, data of 22 days were divided into two stages. The first stage is from January 23 to February 7, and the second stage is from February 8 to February 12. During the second stage, Wuhan city took a number of measures, including timely diagnosis, timely treatment and effective isolation of the infected population, which will have an important impact on the parameters of the model.

Table 1. Number of the infected and the recovery at specific date in Wuhan

Date	The infected	The recovery
22-Jan	425	28
23-Jan	495	31
24-Jan	572	39
25-Jan	618	40
26-Jan	698	42
27-Jan	1,590	45
28-Jan	1,905	75
29-Jan	2261	82
30-Jan	2639	103
31-Jan	3215	139
1-Feb	4109	171
2-Feb	5142	224
3-Feb	6384	303
4-Feb	8351	368
5-Feb	10117	431
6-Feb	11618	534
7-Feb	13603	698
8-Feb	14982	877
9-Feb	16902	1044
10-Feb	18454	1206
11-Feb	19588	1377
12-Feb	32944	1915

Source: <http://wjw.hubei.gov.cn/>

4、Initial parameter settings

To establish the model, we firstly estimated the parameters of the susceptible (S), the exposed (E), the infected (I) and the recovery(R) based on the latest data issued on February 12:

$N = 200000$, which is the total number of potential close contacts in Wuhan on February 12.

$S=N-I$, in which S is the number of the susceptible and I is the number of the infected.

$I_0 = 425$, which is the number of susceptible individuals at the beginning of the model run.

$E_0 = 426$, which is the number of exposed individuals at the beginning of the model run

$R_0 = 28$, which is the number of recovered individuals at the beginning of the model run

Results and Analysis

Epidemic prediction based on SEIR model

The epidemic of the novel coronavirus pneumonia in Wuhan was studied by SEIR modeling. The results showed that, at the time when Wuhan was closed, the number of initially infected individuals was $I_0 = 425$, the number of initially exposed individuals was $E_0 = 426$, and the number of initially recovered patients was $R_0 = 28$.

Next, we separated the data into two stages: January 22-February 7 and February 8-February 12. In the first stage, $TI = 14$ (interquartile range : 14-14) , $TE = 3.0$ (interquartile range : 2.8-3.1) , $R_0 = 1.44$ (interquartile range : 1.40-1.47) (Figure 2). The data showed that the infectious time of the infected person (I) is 14 days, and the incubation period is about 3 days, which is close to the data (Interquartile range: 3.0-7.2) estimated in the reference^[8]. The propagation base R_0 of this study is 1.44, which is significantly lower than the R_0 estimated by other papers before the closure of Wuhan^[9-11].




Fig2. Distribution of the R_0 (A), TI (B) and TE (C) for 100 random samplings

Table2. Distribution of the R_0 (A), TI (B) and TE (C) for 100 random samplings

Repeats	R_0	TI	TE	RMSE _(I)	RMSE _(R)	RMSE _(I) + RMSE _(R)
1	1.54	14.00	3.30	232.06	1108.30	1340.36
2	1.29	14.00	2.30	328.27	1514.78	1843.05
3	1.29	14.00	2.30	278.94	1446.31	1725.26
4	1.41	14.00	2.90	325.89	1335.01	1660.90
5	1.51	14.00	3.20	259.15	1237.79	1496.94
6	1.57	14.00	3.40	235.95	1115.25	1351.20
7	1.44	14.00	3.00	336.74	1549.45	1886.20
8	1.54	14.00	3.30	276.54	1203.22	1479.76
9	1.36	14.00	2.50	230.20	1110.36	1340.56
10	1.44	14.00	3.00	328.24	1525.51	1853.75
11	1.44	14.00	3.00	333.88	1510.17	1844.05
12	1.26	14.00	2.20	239.48	1420.70	1660.18
13	1.47	14.00	3.10	344.07	1492.23	1836.30
14	1.41	14.00	2.90	324.17	1307.48	1631.65
15	1.41	14.00	2.90	265.93	1492.69	1758.62
16	1.29	14.00	2.30	329.13	1524.16	1853.29
17	1.46	14.00	3.10	307.72	1181.57	1489.29
18	1.41	14.00	2.90	273.56	1492.94	1766.51
19	1.51	14.00	3.20	257.96	1180.79	1438.75
20	1.26	14.00	2.20	258.77	1409.80	1668.56
21	1.46	14.00	3.10	312.57	1389.26	1701.83
22	1.46	14.00	3.10	319.22	1223.58	1542.81
23	1.43	14.00	3.00	294.43	1188.60	1483.03
24	1.44	14.00	3.00	330.24	1507.94	1838.18
25	1.69	14.00	3.70	199.45	955.89	1155.34
26	1.41	14.00	2.90	324.01	1293.32	1617.33
27	1.46	14.00	3.10	314.87	1391.49	1706.36
28	1.29	14.00	2.30	316.01	1523.79	1839.79
29	1.47	14.00	3.10	343.78	1523.28	1867.06
30	1.51	14.00	3.20	264.95	1224.86	1489.81
31	1.52	14.00	3.30	325.20	1370.43	1695.63
32	1.43	14.00	3.00	202.23	1271.15	1473.37
33	1.44	14.00	3.00	333.66	1541.24	1874.90
34	1.48	14.00	3.10	235.99	1250.52	1486.51
35	1.41	14.00	2.90	267.35	1492.76	1760.11
36	1.38	14.00	2.80	259.48	1443.04	1702.52
37	1.64	14.00	3.60	288.99	1200.59	1489.58
38	1.29	14.00	2.30	331.33	1278.37	1609.69
39	1.51	14.00	3.20	257.86	1201.83	1459.70
40	1.64	14.00	3.60	283.26	1191.24	1474.50
41	1.26	14.00	2.20	303.57	1533.43	1837.00
42	1.73	14.00	3.80	195.97	739.28	935.25
43	1.26	14.00	2.20	258.73	1421.11	1679.84
44	1.44	14.00	3.00	329.85	1548.18	1878.03
45	1.32	14.00	2.40	342.06	1462.74	1804.80

46	1.44	14.00	3.00	334.80	1508.12	1842.92
47	1.43	14.00	2.70	280.35	1183.74	1464.09
48	1.47	14.00	3.10	340.39	1494.90	1835.28
49	1.46	14.00	3.10	304.57	1312.61	1617.19
50	1.69	14.00	3.70	199.44	940.55	1139.99
51	1.64	14.00	3.60	280.98	1168.77	1449.75
52	1.52	14.00	3.30	341.47	1403.35	1744.83
53	1.41	14.00	2.90	254.30	1460.80	1715.11
54	1.25	14.00	2.00	352.72	1318.67	1671.39
55	1.51	14.00	3.20	238.67	1079.43	1318.10
56	1.44	14.00	3.00	329.99	1356.57	1686.57
57	1.29	14.00	2.30	333.46	1480.18	1813.64
58	1.47	14.00	3.10	336.68	1492.01	1828.69
59	1.41	14.00	2.90	309.87	1361.03	1670.90
60	1.38	14.00	2.80	298.04	1366.97	1665.01
61	1.41	14.00	2.90	267.13	1480.91	1748.03
62	1.40	14.00	2.90	291.41	1200.51	1491.92
63	1.41	14.00	2.90	260.20	1491.40	1751.60
64	1.26	14.00	2.20	312.96	1346.16	1659.12
65	1.29	14.00	2.30	330.69	1508.49	1839.17
66	1.47	14.00	3.10	342.46	1501.42	1843.88
67	1.43	14.00	3.00	301.38	1182.60	1483.98
68	1.44	14.00	3.00	327.74	1479.68	1807.42
69	1.41	14.00	2.90	325.82	1356.72	1682.54
70	1.41	14.00	2.90	303.12	1561.21	1864.33
71	1.44	14.00	3.00	336.34	1537.86	1874.20
72	1.44	14.00	3.00	325.27	1553.87	1879.14
73	1.47	14.00	3.10	346.19	1514.79	1860.98
74	1.49	14.00	3.20	324.28	1427.12	1751.40
75	1.47	14.00	3.10	339.75	1489.98	1829.73
76	1.44	14.00	3.00	329.81	1554.92	1884.73
77	1.46	14.00	3.10	312.38	1201.54	1513.92
78	1.49	14.00	3.20	327.08	1425.26	1752.34
79	1.41	14.00	2.90	308.17	1557.02	1865.19
80	1.44	14.00	3.00	333.98	1498.83	1832.81
81	1.41	14.00	2.90	319.86	1361.46	1681.32
82	1.46	14.00	3.10	313.13	1342.61	1655.74
83	1.26	14.00	2.20	298.31	1543.74	1842.04
84	1.52	14.00	3.30	343.84	1389.07	1732.91
85	1.51	14.00	3.20	204.54	1142.99	1347.54
86	1.29	14.00	2.30	334.83	1322.84	1657.67
87	1.47	14.00	3.10	342.76	1475.58	1818.34
88	1.41	14.00	2.90	307.41	1545.55	1852.96
89	1.46	14.00	3.10	315.48	1362.47	1677.95
90	1.41	14.00	2.90	313.66	1367.89	1681.55
91	1.32	14.00	2.40	349.91	1487.17	1837.08
92	1.32	14.00	2.40	345.09	1478.18	1823.28
93	1.29	14.00	2.30	329.34	1497.61	1826.94

94	1.32	14.00	2.40	345.91	1489.51	1835.43
95	1.41	14.00	2.90	306.57	1490.55	1797.12
96	1.44	14.00	3.00	334.94	1498.89	1833.83
97	1.32	14.00	2.40	343.14	1462.04	1805.18
98	1.41	14.00	2.90	325.28	1313.11	1638.39
99	1.66	14.00	3.60	172.49	967.26	1139.74
100	1.25	14.00	2.00	412.51	1417.39	1829.90

In the second stage (after February 8), we set the number of susceptible population N to be fixed at 200000, and the infection cycle of infected population decreased from 14 days to 4 days, i.e. $TI = 4$, which was estimated according to the data of 5 days from February 8 to February 12, so as to get the epidemic development trend of 90 days since January 22, including the number of infected people, the number of latent people and the number of recovered people (Figure 3, Table 3). The results showed that the number of infected people increased slowly in the early stage (January 22-January 31), but during February 1-February 12, the number of infected people increased rapidly, which is expected to peak around February 19, reaching about 45000 people. Subsequently, the number of infections will decrease. Once entering March, the epidemic would gradually decline, and the epidemic would end around the end of March. It is worth noting that the above prediction is based on the assumption that the number of susceptible population $N = 200000$ will not increase.




Fig3. Epidemic trend of 2019-nCoV within 90 days after the closure of Wuhan

In Figure 3 and Figure 4, red line indicates the trend of cumulative infection number over time, blue line is the trend of cumulative rehabilitation number over time, and

green line is the trend of cumulative latent number over time. Vertical dash line indicates the peak time of cumulative infection number.




Fig4. Epidemic trend of 2019-nCoV within 90 days after the closure of Wuhan city assuming the number of susceptible people $n = 300000$ (A) and 400000 (B)

If the epidemic situation is not properly controlled, the number of susceptible population will continue to increase on the basis of current $N = 200000$. If the number of susceptible population increases to $N = 300000$, and other parameters remain unchanged, the peak number can increase to 75000, and the epidemic peak time will also be postponed at around February 21 (Figure 4). If it is increased to $N = 400000$ and other parameters remain unchanged, the peak number can be increased to 100000, and the epidemic peak will be postponed to around February 22 (Figure 4). Even in both cases, the epidemic would subside in early March, and disappear gradually towards the late March.

Table 3. Expected number of cases for infected, recovery and exposed

Date	Infected	Recovery	Exposed	Date	Infected	Recovery	Exposed
22-Jan	425	28	426	7-Mar	3121	185309	605
23-Jan	537	58	532	8-Mar	2542	186089	484
24-Jan	676	97	664	9-Mar	2068	186725	388
25-Jan	849	145	829	10-Mar	1680	187242	311
26-Jan	1064	206	1034	11-Mar	1364	187662	250
27-Jan	1333	282	1290	12-Mar	1106	188003	201
28-Jan	1668	377	1607	13-Mar	896	188279	161
29-Jan	2084	496	1999	14-Mar	726	188503	130
30-Jan	2602	645	2482	15-Mar	588	188685	104
31-Jan	3243	831	3075	16-Mar	476	188832	84
1-Feb	4037	1062	3799	17-Mar	385	188951	68
2-Feb	5015	1351	4677	18-Mar	311	189047	55
3-Feb	6216	1709	5733	19-Mar	252	189125	44
4-Feb	7683	2153	6989	20-Mar	203	189188	36
5-Feb	9464	2702	8465	21-Mar	164	189238	29
6-Feb	11610	3377	10172	22-Mar	133	189280	23
7-Feb	14171	4207	12108	23-Mar	107	189313	19
8-Feb	14664	7750	17346	24-Mar	87	189340	15
9-Feb	16780	11416	22496	25-Mar	70	189361	12
10-Feb	20084	15611	27606	26-Mar	57	189379	10
11-Feb	24265	20632	32449	27-Mar	46	189393	8
12-Feb	29015	26698	36597	28-Mar	37	189404	6
13-Feb	33960	33952	39545	29-Mar	30	189414	5
14-Feb	38652	42442	40874	30-Mar	24	189421	4
15-Feb	42614	52105	40409	31-Mar	19	189427	3
16-Feb	45430	62758	38286	1-Apr	16	189432	3
17-Feb	46835	74116	34898	2-Apr	13	189436	2
18-Feb	46759	85824	30762	3-Apr	10	189439	2
19-Feb	45323	97514	26371	4-Apr	8	189442	1
20-Feb	42782	108845	22107	5-Apr	7	189444	1
21-Feb	39456	119540	18213	6-Apr	5	189445	1
22-Feb	35663	129404	14809	7-Apr	4	189447	1
23-Feb	31684	138320	11925	8-Apr	4	189448	1
24-Feb	27738	146241	9537	9-Apr	3	189449	0
25-Feb	23982	153175	7591	10-Apr	2	189449	0
26-Feb	20517	159171	6025	11-Apr	2	189450	0
27-Feb	17396	164300	4774	12-Apr	1	189451	0
28-Feb	14639	168649	3781	13-Apr	1	189451	0
29-Feb	12239	172309	2995	14-Apr	1	189451	0
1-Mar	10178	175369	2374	15-Apr	1	189451	0
2-Mar	8424	177913	1883	16-Apr	1	189452	0
3-Mar	6946	180019	1496	17-Apr	1	189452	0
4-Mar	5708	181756	1191	18-Apr	0	189452	0
5-Mar	4678	183183	949	19-Apr	0	189452	0
6-Mar	3825	184352	757	20-Apr	0	189452	0

Discussion

Although some modeling studies on the epidemiological characteristics of 2019-nCoV epidemic have been reported so far, they had some limitations, such as the data come from the early stage of the epidemic. Due to the rapid change of the epidemic situation and the closure of Wuhan on January 23, many parameters related to the model have also changed, which affect the applicability and reliability of the model. This study used the latest 2019 nCoV data in Wuhan area, analyzed the epidemiological characteristics of 2019 nCoV epidemic after in Wuhan city was shut down. Compared with other studies, the R_0 value produced in this study is smaller, indicating that the closure and subsequent measures have played an important role in the spread of the epidemic.

The infection time index (TI) obtained in this study was higher than that of SARS[12] and MERS[13], but lower than that of 2019-nCoV in literatures[14] reported earlier. This result may be related to the sudden outbreak of the epidemic, the lack of medical resources for early response, and the failure of timely diagnosis and treatment of infected patients. A large number of mild patients and asymptomatic virus carriers were not isolated in time. The incubation period (TE) is about 3 days, which is close to the data in the reference^[14].

According to the latest reported data, the cumulative number of people infected on February 13 and 14 was 35991 and 37914 respectively, which is close to the number predicted by our estimation (Table 2). According to this study, the number of infected people will reach the peak in February 19 at about 45000 infected individuals.

Conclusions

With the implementation of more follow-up measures, including strict restrictions on people going out, accelerating the treatment of infected individuals, and clinical trials of new drugs, the development of 2019-nCoV epidemic in Wuhan will be effectively controlled, and the number of infected individuals will gradually decrease. It was expected that the epidemic would subside in early March, and disappear gradually towards the late March. If the epidemic situation is not properly controlled, the peak of infected number can be further increased and the peak time will be a little postponed.

Declarations

Ethics approval and consent to participate

The ethical approval or individual consent was not applicable.

Availability of data and materials

All data and materials used in this work were publicly available.

Consent for publication

Not applicable.

Funding

This work was not funded.

Disclaimer

The funding agencies had no role in the design and conduct of the study; collection, management, analysis, and interpretation of the data; preparation, review, or approval of the manuscript; or decision to submit the manuscript for publication.

Conflict of Interests

The author declared no competing interests.

Reference

1. Wee, Rolando Y. . "The Largest Cities In China." World Atlas, Jul. 9, 2019, worldatlas.com/articles/20-biggest-cities-in-china.html.
2. W. H. Organization, Coronavirus. World Health Organization, Available: <https://www.who.int/health-topics/coronavirus> (cited January 19, 2020).
3. The official website of Hubei Provincial Health Committee (<http://wjw.hubei.gov.cn/>)(cited February 19, 2020).
4. Pastor-Satorras R, Castellano C, Mieghem PV, et al. Epidemic processes in complex networks. Review of Modern Physics 2015, 87(3): 925-979.
5. Wallinga J, Lipsitch M, How generation intervals shape the relationship between growth rates and reproductive numbers. Proceedings of the Royal Society B: Biological Sciences, 2007, 274(1609):599-604.
6. Y Yang, Q Lu, M Liu, et al. Epidemiological and clinical features of the 2019 novel coronavirus outbreak in China. medRxiv 2020.02.10.20021675; doi: <https://doi.org/10.1101/2020.02.10.200216>.
7. Qun Li, Xuhua Guan, Peng Wu et al., Early Transmission Dynamics in Wuhan, China, of Novel Coronavirus-Infected Pneumonia, New Engl J Med, January 29, 2020,DOI: 10.1056/NEJMoa2001316
8. Y Yang, Q Lu, M Liu, et al. Epidemiological and clinical features of the 2019 novel coronavirus outbreak in China, medRxiv, 2020, January 29, 2020 DOI: 10.1056/NEJ Moe2001316

9. T Zhou, Q Liu, Z Yang, et al. Preliminary prediction of the basic reproduction number of the Wuhan novel coronavirus 2019-nCoV, arXiv, 2020 - arxiv.org.
 10. JM Read, JRE Bridgen, DAT Cummings, Novel coronavirus 2019-nCoV: early estimation of epidemiological parameters and epidemic predictions , medRxiv 2020.01.23.20018549; doi: <https://doi.org/10.1101/2020.01.23.20018549>
 11. Z Cao, Q Zhang, X Lu, et al. Incorporating Human Movement Data to Improve Epidemiological Estimates for 2019-nCoV. medRxiv 2020.02.07.20021071; doi: <https://doi.org/10.1101/2020.02.07.20021071>
 12. Donnelly CA, Ghani AC, Leung GM, et al. Epidemiological determinants of spread of causal agent of severe acute respiratory syndrome in Hong Kong. Lancet 361, 2003: 1761-1766.
 13. Assiri A, McGeer A, Perl TM, et al., Hospital outbreak of Middle East respiratory syndrome coronavirus. New Engl J Med 2013;369(5):407–16
 14. JT Wu, K Leung, GM Leung, Nowcasting and forecasting the potential domestic and international spread of the 2019-nCoV outbreak originating in Wuhan, China: a modelling study , The Lancet, 2020, [https://doi.org/10.1016/S0140-6736\(20\)30260-9](https://doi.org/10.1016/S0140-6736(20)30260-9)
-