> Non surgical procedure related postoperative Complications independently predicts perioperative mortality, in gastrointestinal and Hpb surgeries.- A retrospective Analysis of prospectively maintained data.

Dr.Bhavin Vasavada

Consultant hepatobiliary and liver transplant surgeon,

Shalby hospitals

Email: drbhavin.liversurgeon@gmail.com

Dr.Hardik patel

Consultant hepatobiliary and liver transplant surgeon

Ethical approval: taken from hospital ethical committee. IRB no:2342/2019 shalby hospitals

Funding: No funding disclosure needed

Conflict of interest: no conflict of interest.

Aim:

The Aim of the study was to evaluate relationship between non surgical procedure related complication and 30 days mortality.

Material and Methods:

All gastrointestinal and hepatobiliary procedures performed in last 3 years have been evaluated retrospectively. Non surgical procedure related postoperative complications were defined as perioperative complications non related to surgical procedures or techniques and related to patients' physiological health or comorbidities (e.g acute kidney injury, ARDS, acute respiratory failure, pre existing sepsis, etc.), Surgical related complications were defined as perioperative complications related to surgical procedures or techniques (e.g. bleeding, leaks, sepsis due to leaks etc.). Factors affecting 30 days mortality and morbidity were analysed using univariate and multivariate analysis. Chi square test was used for categorical values, Mann Whitney U test was used for numerical values. Multivariate logistic regression analysis was used for multivariate analysis. Statistical analysis was used suing SPSS version 21.

Results:

Total 325 major hepatobiliary and pancreatic surgery was done in our institute in last 2 years. 30 days overall mortality rate was 6.4%. In univariate analysis mortality was significantly associated with nonsurgical procedure related complications. (p < 0.0001). Surgical complications were not associated with mortality. On univariate analysis other factors associated with mortality were emergency surgeries, high CDC grade of surgery, higher ASA grades, increase operative duration, increased blood product requirements. However on multivariate analysis only nonsurgical procedure related postoperative

complications independently predicted mortality. (p=0.001).

Conclusions:

Non surgical procedure related post operative complications (Physiological) is strongly associated with 30 days mortality, suggesting improved perioperative care can help to reduce post operative mortality.

Introduction:

Surgical Complications are major cause of mortality and morbidity. [1] Surgical complications can be as high as 30% in some group of patients. [2,3]. Surgical complications generally consist of two type of complications as per our knowledge. One is technique or surgical procedure related e.g. bleeding, anastomotic leaks. The other is non surgical technique related complications or complications due to change in physiology due to surgical stress, for example ARDS, Acute kidney Injury, Post operative Acute left ventricular failure or Post operative acute delirium. In non surgical technical related complication prehabilitation can work. [4]. We hypothesised that non surgical technique related complications are much more common than surgical technique related or surgical complications.

Aim:

The Aim of the study was to evaluate relationship between non surgical procedure related complication and 30 days mortality.

Material and Methods:

All gastrointestinal and hepatobiliary procedures performed in last 3 years have been evaluated retrospectively.

Nonsurgical procedure related postoperative complications were defined as perioperative complications non related to surgical procedures or techniques and related to patients' physiological health or comorbidities (e.g acute kidney injury, ARDS, acute respiratory failure, pre existing sepsis, etc.).

Surgical related complications were defined as perioperative complications related to surgical procedures or techniques (e.g., bleeding, leaks, sepsis due to leaks, etc.). Factors affecting 30 days mortality were analysed using univariate and multivariate analysis.

Statistics:

Fisher t test was used for categorical values, Mann Whitney U test was used for numerical values. Multivariate logistic regression analysis was used for multivariate analysis. Statistical analysis was used suing SPSS version 21.

RESULTS:

Total 325 major hepatobiliary and pancreatic surgery was done in our institute in last 2 years. 30 days overall mortality rate was 6.4%.

In univariate analysis mortality was significantly associated with nonsurgical procedure related complications. (p < 0.0001). Surgical complications were not associated with mortality. (p=0.235)

On univariate analysis other factors associated with mortality were emergency

surgeries, high CDC grade of surgery, higher ASA grades, increase operative duration, increased blood product requirements. [Table 1]

However on multivariate analysis only nonsurgical procedure related postoperative complications independently predicted mortality. (p < 0.001). [Table 2] Odds ratio was 0.007 with 95% confidence interval 0.001 to 0.038.

Kaplan Meier 30 days survival curve also showed mortality was significantly higher in patients with nonsurgical technique related complications. P value was less than 0.0001 with log rank test. [Figure 1]. However there was no difference in patients' with surgical technique related complications.

Factors	No mortality	Mortality. (n= 21)	P value
	(n=304)		
Nonsurgical	6	16	< 0.0001
(physiological)			
complication.			
(n=22)			
Surgical site	21	0	P= 0.232
infection (n=21)			
Emergency	48	10	P= 0.001
Surgery (n= 58)			
AGE [mean(SD)]	51.2 (12.6)	57.1(12.8)	P= 0.07
GRADE OF	3.04(0.59)	3.4 (0.58)	P<0.0001
SURGERY			
[mean(SD)]			
Blood products	1.59 (1.68)	2.0 (3.0)	P<0.001

[mean(SD)]				
ASA	stage	2.8 (0.75)	3.2 (0.81)	P< 0.001
[mean(SD)]				
Operative	time	155 (81)	185 (143)	P< 0.001
[mean(SD)]				
(Minutes)				

Table :1 Univariate analysis for 30 days mortality

		Table 2. multivariate logistic regression								
									95% C.I.for EXP(B)	
			В	S.E.	Wald	df	Sig.	Exp(B)	Lower	Upper
	Step 1 ^a	gradeofsurgery	.072	.770	.009	1	.926	1.074	.237	4.861
		bloodprodcuts	.329	.207	2.524	1	.112	1.389	.926	2.084
		ASA	.124	.508	.059	1	.808	1.132	.418	3.065
		EMMERGENCY(1)	-1.069	1.138	.882	1	.348	.343	.037	3.197
		operativetime	.001	.003	.237	1	.626	1.001	.995	1.008
		Physiological (1)	-4.893	.830	34.744	1	.000	.007	.001	.038
		Constant	.360	3.150	.013	1	.909	1.434		
ľ	a. Variable(s) entered on step 1: gradeofsurgery, bloodprodcuts, ASA, EMMERGENCY, operativetime, PHYSIOLOGICAL.							OGICAL.		

Sig.

.000



Overall Comparisons

228.483

The vector of trend weights is -1, 1. This is the default.

Log Rank (Mantel-Cox) df

1

Figure 1 30 day survival comparison with non surgical complication and surgical complication with log rank analysis. (p < 0.0001)

df

Overall Comparisons



Figure 2: There was no difference in 30 days survival in patient with surgical technique related complications.

Discussion:

Perioperative mortality is the one the most important problem surgical community is facing. Peri operative mortality range from 0.1% to as high as 27%-30%

according to various types of surgeries. [5]. Gastro intestinal and hepatobiliary surgeries are technically demanding procedures and has one of the highest perioperative mortality rates. [6,7,8]

Surgeons are always worried about surgical techniques and surgical techniques related mortalities, how ever a very few studies are done to look at the impact of such complications on perioperative mortality. There are various perioperative complications which are not actually related to surgical techniques, but depend on many factors like patients' pre operative conditions as well as perioperative anaesthetic course and complications due to that. These complications can include but not limited to acute kidney injury, adult respiratory distress syndrome, post operative delirium, non surgery related sepsis and septic shock. These complications can contribute to mortality significantly. [9.10]

Aim of our study to look for effect of nonsurgical technique related complications and surgical technique related complications on mortality. For gastrointestinal and hepatobiliary surgeries we defined anastomotic leaks, sepsis due to leaks, intraoperative bleeding ,iatrogenic injuries to surrounding structure as surgical technique related complications and other complications like acute kidney injury, ards (acute respiratory distress syndrome), sepsis not related to surgical technique etc we defined as non surgical technique related complications of physiological complications.

Surgery related complications were not associated with mortality. On univariate analysis, physiological complications, ASA score, blood product requirements, operative time, CDC grade of surgery [11], and emergency surgery were associated with 30 days mortality. [table 1] Suggesting patient related factors were associated with 30 days mortality rather than surgical technique related

> complications. Which is also supported by current literature available. [12,13,14,15]

> However on multivariate analysis only occurrence of nonsurgical technique related or what we call as physiological complications predicted, survival outcomes. [table 2]. Kaplan Meier survival curves also predicted lower 30 days survival for patients who had physiological complications. [figure 1] but there was no difference in 30 days survival in patient who has surgical technique related complications. [Figure 2]

We do not want to say that surgical technique related complications are not harmful but our purpose is to point out importance of nonsurgical technique related complications and their impact on surgical mortality. This study also shows importance of critical care management in reducing postoperative mortality. [16,17,18,19].

Being a retrospective study this study has inherent limitations associated with it. Another limitation is its includes wide variety of sub specialities. However we believe these findings will remain same across the specialities.

In conclusion Non surgical procedure related post operative complications (Physiological) is strongly associated with 30 days mortality, suggesting improved perioperative critical care can help to reduce post operative mortality.

References:

- Tevis SE, Kennedy GD. Postoperative complications and implications on patient-centered outcomes. J Surg Res. 2013;181(1):106–113. doi:10.1016/j.jss.2013.01.032
- Longo WE, Virgo KS, Johnson FE, Oprian CA, Vernava AM, Wade TP, Phelan MA, Henderson WG, Daley J, Khuri SF. Dis Colon Rectum. 2000 Jan; 43(1):83-91.
- Mayo NE, Feldman L, Scott S, et al. Impact of preoperative change in physical function on postoperative recovery: argument supporting prehabilitation for colorectal surgery. Surgery. 2011;150(3):505–514. doi:10.1016/j.surg.2011.07.045
- Hijazi Y, Gondal U, Aziz O. A systematic review of prehabilitation programs in abdominal cancer surgery. Int J Surg. 2017;39:156–162. doi:10.1016/j.ijsu.2017.01.111
- Ng-Kamstra JS, Arya S, Greenberg SLM, et al. Perioperative mortality rates in low-income and middle-income countries: a systematic review and metaanalysis. BMJ Glob Health. 2018;3(3):e000810. Published 2018 Jun 22. doi:10.1136/bmjgh-2018-000810
- Csikesz NG, Simons JP, Tseng JF, Shah SA. Surgical specialization and operative mortality in hepato-pancreatico-biliary (HPB) surgery. J Gastrointest Surg. 2008;12(9):1534–1539. doi:10.1007/s11605-008-0566-z
- Kneuertz PJ, Pitt HA, Bilimoria KY, et al. Risk of morbidity and mortality following hepato-pancreato-biliary surgery. J Gastrointest Surg. 2012;16(9):1727–1735. doi:10.1007/s11605-012-1938-y

- Sørensen LT, Malaki A, Wille-Jørgensen P, et al. Risk factors for mortality and postoperative complications after gastrointestinal surgery. J Gastrointest Surg. 2007;11(7):903–910. doi:10.1007/s11605-007-0165-4
- A. Miskovic, A. B. Lumb, Postoperative pulmonary complications, BJA: British Journal of Anaesthesia, Volume 118, Issue 3, March 2017, Pages 317– 334, https://doi.org/10.1093/bja/aex002
- Gameiro J, Fonseca JA, Neves M, Jorge S, Lopes JA. Acute kidney injury in major abdominal surgery: incidence, risk factors, pathogenesis and outcomes. Ann Intensive Care. 2018;8(1):22. Published 2018 Feb 9. doi:10.1186/s13613-018-0369-7
- Onyekwelu I, Yakkanti R, Protzer L, Pinkston CM, Tucker C, Seligson D. Surgical Wound Classification and Surgical Site Infections in the Orthopaedic Patient. J Am Acad Orthop Surg Glob Res Rev. 2017;1(3):e022. Published 2017 Jun 13. doi:10.5435/JAAOSGlobal-D-17-00022
- 12. Chu CL, Chiou HY, Chou WH, Chang PY, Huang YY, Yeh HM. Leading Comorbidity associated with 30-day post-anesthetic mortality in geriatric surgical patients in Taiwan: a retrospective study from the health insurance data. BMC Geriatr. 2017;17(1):245. Published 2017 Oct 24. doi:10.1186/s12877-017-0629-7
- Chou WC, Chang PH, Lu CH, et al. Effect of Comorbidity on Postoperative Survival Outcomes in Patients with Solid Cancers: A 6-Year Multicenter Study in Taiwan. J Cancer. 2016;7(7):854–861. Published 2016 Apr 28. doi:10.7150/jca.14777

- Clough RA, Leavitt BJ, Morton JR, et al. The Effect of Comorbid Illness on Mortality Outcomes in Cardiac Surgery. Arch Surg. 2002;137(4):428–433. doi:10.1001/archsurg.137.4.428
- Oliver K. Jawitz, Zuoheng Wang, Daniel J. Boffa, Frank C. Detterbeck, Justin D. Blasberg, Anthony W. Kim, The differential impact of preoperative comorbidity on perioperative outcomes following thoracoscopic and open lobectomies, European Journal of Cardio-Thoracic Surgery, Volume 51, Issue 1, January 2017, Pages 169–174
- 16. Frankel HL, Butler KL, Cuschieri J, et al. The role and value of surgical critical care, an essential component of Acute Care Surgery, in the Affordable Care Act: a report from the Critical Care Committee and Board of Managers of the American Association for the Surgery of Trauma. J Trauma Acute Care Surg. 2012;73(1):20–26.
- McMillen MA, Boucher N, Keith D, Gould DS, Gave A, Hoffman D. Maintaining quality of care 24/7 in a nontrauma surgical intensive care unit. J Trauma Acute Care Surg. 2012;73(1):202–208.
- Park CM, Chun HK, Lee DS, Jeon K, Suh GY, Jeong JC. Impact of a surgical intensivist on the clinical outcomes of patients admitted to a surgical intensive care unit. Ann Surg Treat Res. 2014;86(6):319–324. doi:10.4174/astr.2014.86.6.319
- Rohrig SAH, Lance MD, Faisal Malmstrom M. Surgical intensive care current and future challenges?. Qatar Med J. 2020;2019(2):3. Published 2020 Jan 13.