Chloroquine, hydroxychloroquine, and COVID-19: systematic review and narrative synthesis of efficacy and safety

Systematic review of (hydroxy)chloroquine efficacy and safety

Michael Takla^{1,2}, Kamalan Jeevaratnam^{1*}

¹ Faculty of Health and Medical Science, University of Surrey, Guildford, GU2 7AL, United Kingdom

² Christ's College, University of Cambridge, Cambridge, CB2 3EG, United Kingdom

Corresponding author

*Kamalan Jeevaratnam - Faculty of Health and Medical Sciences, University of Surrey, Guildford, GU2 7AL, United Kingdom Email: drkamalanjeeva@gmail.com Telephone: +441483 682395

<u>Abstract</u>

Background: The COVID-19 pandemic has required clinicians to urgently identify new treatment options or the repurposing of existing drugs. Several drugs are now being repurposed with the aim of identifying if these drugs provide some level of disease resolution. Of particular interest are chloroquine (CQ) and hydroxychloroquine (HCQ), first developed as an antimalarial therapy. There is increasing concern with regards to the efficacy and safety of these agents. The aims of this review are to systematically identify and collate studies describing the use of CQ and HCQ in human clinical trials and provide a detailed synthesis of evidence of its efficacy and safety.

Methods and Findings: Searches for ("COVID" AND "chloroquine"[title/abstract] AND "outcomes"[full text]) and two ("COVID" AND "hydroxychloroquine"[title/abstract] AND "outcomes"[full text]) yielded 272 unique articles. Unique articles were manually checked for inclusion and exclusion criteria and also subjected to a quality appraisal assessment. A total of 19 articles were included in the systematic review. Seventy-five percent of observational studies employing an endpoint specific to efficacy recorded no significant difference in the attainment of outcomes, between COVID-19 patients given a range of CQ and/or HCQ doses, and the control groups. All clinical trials and 82% of observational studies examining an indicator unique to drug safety discovered a higher probability of adverse events in those treated patients suspected of, and diagnosed with, COVID-19. Seventy-five percent of the total papers focusing on cardiac side-effects found a greater incidence among patients administered a wide range of CQ and/or HCQ doses, with QTc prolongation the most common finding, in addition to its consequences of VT and cardiac

arrest. Of the total studies using mortality rate as an end-point, 60% reported no significant change in the risk of death, while 30% showed an elevation, and 10% a depression, in treated relative to control patients.

Conclusion: The strongest available evidence suggests that, relative to standard in-hospital management of symptoms, the use of CQ and HCQ to treat hospitalised COVID-19 patients has likely been unsafe. At the very least, the poor quality of data failing to find any significant changes in the risk of VT should preclude definitive judgment on drug safety until the completion of high-quality randomised clinical trials.

1.0 Introduction

In the final week of December 2019, the Hubei Integrated Chinese and Western Medicine hospital in Wuhan, reported a clustered point-source outbreak of pneumonia [1], of unknown viral origin. Within 30 days, the rapid geographic expansion of the disease, which the International Committee on Taxonomy of Viruses later coined Coronavirus Disease 2019 (COVID-19) [2], implied propagation by human-to-human transmission. On March 11 2020, the World Health Organisation (WHO) designated COVID-19 a pandemic [3]. As of May 26 2020, COVID-19 has been confirmed as the cause of 5,508,904 cases and 346, 326 deaths [4] globally.

In the absence of specific antiviral pharmacotherapy, the repositioning of existing drugs represents an attractive clinical option. Selecting which drugs to repurpose, however, hinges on the compatibility of their mechanisms of action with the disease progression of severe acute respiratory syndrome (SARS), and with the biology of the recently emerged pathogenic agent that causes it.

With a likely evolutionary origin in bats [5], the novel (beta)-coronavirus, SARS-CoV-2, probably acquired the ability to zoonotically infect humans via natural selection of the receptor-binding domains of its spike (S) proteins in an intermediate mammalian host [6]. Indeed, compared to SARS-CoV-1, the highly homologous [7] coronavirus responsible for the SARS pandemic [8], the S protein has a 10-20-fold greater affinity for the ACE2 receptor [9] predominantly expressed by pulmonary and intestinal epithelia and vascular endothelia [10]. In fact, *in silico* analysis has demonstrated that the expression of sialic acid further

facilitates viral entry, whereby binding of human gangliosides impairs inhibitory interactions between the S protein and the plasma membrane [11]. The resulting receptor-mediated endocytosis precedes endosomal cathepsin and TMPRSS2-mediated [12] cleavage of the S protein, permitting fusion of the viral lipid envelope and human vesicular membrane, whereupon RNA entry into the cytosol enables viral replication, maturation [13], and budding. The initial innate immune response to the subsequent dissemination of SARS-CoV-2 throughout the patient's extracellular fluid elicits a wave of pro-inflammatory cytokines, including IL-1(beta), IL-6, and TNF-(alpha), that recedes upon lymphopenia, only to return at higher concentrations in a cytokine storm [14] that predisposes to a potentially lethal acute respiratory distress syndrome (ARDS). Additionally contributing to the mortality of critically ill COVID-19 patients is the significantly elevated incidence of often pulmonary thromboembolic events [15][16][17].

With antiviral [18][19], anti-inflammatory [20], and anti-thrombotic [21][22][23][24] effects, chloroquine (CQ), and its less oculotoxic [25][26] derivative, hydroxychloroquine, (HCQ) were among the first drugs selected for repurposing to treat COVID-19 patients. However, the ability of 4-aminoquinolones to prolong the QT interval [27][28] increases the risk of *de novo* ventricular tachyarrhythmias (VTs), calling into question their cardiac safety [29].

Here, we systematically review existing clinical trial data to provide a detailed synthesis of evidence for the efficacy and safety of CQ and HCQ. We also aim to clarify if the use of such drugs in COVID-19 patients in the absence of rigorous evidence may not only

have had little efficacy, but, owing to their lack of (especially cardiac) safety, may have been responsible for excess mortality.

2.0. Review Methodology

2.1 Objectives

This systematic review seeks to clarify the strength of evidence for the relative efficacy and safety of CQ and HCQ treatment in patients suspected of, and diagnosed with, COVID-19.

2.2 Methods

In line with the PICOT format [30] of framing subjects for clinical research, this study centres on answering the question: 'In patients suspected of, and diagnosed with, COVID-19, how efficacious and safe are CQ and HCQ, relative to standard symptomatic care?'

The subsequent elaboration of the systematic review and narrative synthesis adhered to the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines [31] for evidence-based assessment of published research.

2.3 Search Strategy

In light of the current global health emergency, and the requisite rapid turnover of publications to meet the consequently urgent need to obtain and analyse the results that they present, several authors have resorted to the use of preprint servers to disseminate their findings. Despite the evident shortfalls inherent in referring to data, whose quality has not been peer-reviewed, the present paucity of published original research on the efficacy and safety of CQ and HCQ in COVID-19 patients demands exceptional measures. As such, this

systematic review will take into account non-peer-reviewed work, provided that they have been submitted to a preprint server, where they are available in open-access form. Nevertheless, given its focus on data quality, this review will make unambiguous every instance in which data from such sources are used.

Therefore, on May 26 2020, MEDRXIV, along with PubMed, Web of Science and Embase, acted as the databases for the initial search of items relevant to the PICOT-formatted question. The preliminary use of the search terms "COVID", "chloroquine", and "hydroxychloroquine" yielded a large number of results that bore little relevance to the research topic. Combining such terms into phrases – "COVID" AND "chloroquine", and "COVID" AND "hydroxychloroquine" in the title or abstract – and requiring the term "outcomes" anywhere in the full text considerably focused the responses. The subsequent application of identical phase stenography to each database ensured internal consistency.

2.4 Search Attrition Criteria

The aim of this review being to establish the weight of evidence for the use of a therapy in patients, data able to answer such a question must derive from primary research. Moreover, owing to the international extent of the present health crisis, any imposition of an original language requirement would exclude useful and otherwise rare resources. As such, following the collation of items in EndNote and the removal of duplicates, application of these criteria excluded unique items for which there was either no English version or no original data.

Screening of the resulting papers against the criteria established by the PICOTformatted question – namely, the requirement that data be collected from COVID-19 patients treated with CQ and/or HCQ – included only controlled trials and observational studies.

2.5 Article Processing and Selection

Having applied the exclusion and inclusion criteria to all search results and removing duplicates at all stages where necessary, two investigators independently reviewed the final repertoire of studies.

2.6 Quality Appraisal

Rather than merely verifying the relevance and scope of the material in the final library, holistic analysis of each item of research in line with the framework set out in the Checklist of Review Criteria provided by the Task Force of Academic Medicine and GEA-RIME committee [32] ensured stringent appraisal of study quality.

Indeed, the identification of – among other facets of robust research – appropriate study design, statistical analysis, and quality control (Table 1) permitted only papers with sufficient scientific merit to pass onto the data extraction stage.

									<u>.</u>					9
ublication st author)	Problem Statement, Conceptual Framework, and Research Question	Reference to the Literature and Documentation	Relevance	Research Design	Instrumentation, Data Collection, and Quality Control	Population and Sample	Data Analysis and Statistics	Reporting of Statistical Analyses	Presentation of Results	Discussion and Conclusion: Interpretation	Title, Authors, and Abstract	Presentation and Documentation	Scientific Conduct	Total Criteria Met ⊽
ang [33]	✓	✓	✓	✓	✓	✓	✓	✓	✓	~	✓	✓	✓	13 ng
orba [34]	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	13 🗟 🗧
autret [35]	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	13 g v
uang [36]	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	13 🔬 🗐
ehra [37]	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	13 8 6
ingh [38]	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	13 🗟 🔤
lp [39]	✓	√	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	13 🚽 🤶
osenberg [40]	~	✓	~	~	✓	~	~	~	~	\checkmark	~	\checkmark	~	13 der o
eleris [41]	✓	✓	✓	✓	✓	✓	✓	√	✓	✓	✓	✓	✓	13 0
illion [42]	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	13 <mark>?</mark> s
Yu [43]	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	13 🝟 ट
lagagnoli [44]	~	~	~	~	~	~	~	~	~	~	~	√	~	
Kim [45]	✓	✓	✓	✓	✓	✓	✓	✓	✓	~	✓	✓	✓	13 4
aleh [46]	✓	√	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	13 🛓
Aahevas [47]	~	~	~	~	√	~	~	~	~	\checkmark	~	√	~	13 0 Int
amireddy [48]	~	✓	~	~	~	~	~	~	~	√	~	√	~	13 ernat
rcuro [49]	✓	✓	✓	~	✓	~	✓	✓	✓	✓	✓	√	~	13 🧧 🗧
horin [50]	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	13 😐 🚆
lallat [51]	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	13 ਨ ਕਿ

Table 1: Quality appraisal of the 19 papers passing through search attrition.

2.7 Data Extraction

Among the research items constituting the final library for analysis, there exists a wide variation in study design, results, and, crucially, the extent to which each distinct aspect of the PICOT-formatted question is answered. As such, a specialised data extraction table collates and summarises the most important information in every paper (Table 2). In particular, emphasis on the different sample sizes and structures, doses of drug used, primary and/or secondary outcomes and overall design limitations, facilitates both clarity and caution when making comparisons between the sets of results presented.

Study	Sample	Age	Male	Symptom	Relevant	Efficacy in meeting	Safety	Limitations
	size	(mean/	(%)	severity	treatment	outcome		
		median)						
Tang [33]	150	46	55	Mild to	Standard of	No significant difference	30% of HCQ patients reported	Absence of placebo.
(RCT)				moderate	care +	in probability of negative	adverse events, compared to	
					HCQ 1200	conversion of SARS-	only 9% given the standard of	Lack of masking.
					mg day ⁻¹	CoV-2 at end time-point,	care only.	
					loading dose	and at all specific time-	The most frequent adverse	Underpowered sample size.
					for 3 days,	points.	event in HCQ patients was	
					then 800 mg		diarrhoea (10%), with HCQ	Premature termination of trial censored data on
					day ⁻¹	No significant difference	discontinued in 1 patient with	primary outcome.
					maintenance	in median time to	blurred vision.	
					dose for	negative conversion.	2 patients in the HCQ group	No assessment of antiviral efficacy within 48h of
					remainder of	_	experienced serious events	onset due to enrolment of hospitalised patients.
					2 weeks if	No significant difference	related to COVID-19	
					mild/	in median time to	progression and upper	Short-term period of follow-up underestimates
					moderate, or	symptom recession	respiratory tract infection.	frequency of QT prolongation.
					3 weeks if			
					severe		No cardiac arrhythmic events,	Possible underestimation of retinal damage
							including QT prolongation, in	suggested by detection of early harm caused by 800-
							either group.	1200 mg day ⁻¹ HCQ in a sensitive screening test.
Borba [34]	81	51	75	Severe	CQ 600 mg	High dose associated	Immediate interruption of high-	Placebo not given alone, resulting in reliance on
(RCT)					orally, 2 day	with lethality, but not	dose group for all ages and	historical data for similar patients not receiving CQ.
					¹ for 10 days	when corrected for age.	reversion to low dose.	
								Recruitment of patients with suspected, rather than
							15.1% had QTcF > 500 ms,	confirmed, COVID-19.
							more frequent in the high	
							(19%) than low (11%) dose	Randomisation with small sample size resulted in
							group.	more older patients with heart disease receiving a
							2.7% in the high-dose group	high CQ dose.
							experienced VT before death.	
								Where influenza was suspected, 150 mg day ⁻¹
							Both CK and CKMB were	oseltamivir for 5 days in 93% and 87% of high and

Table 2: Data extraction from the 19 papers passing quality appraisal.

							elevated in almost 40% of	low dose groups, confounding QTc increase.
							patients, more frequent in the	
							high (50%) than low (32%)	All patients also received I.V. ceftriaxone (2 g day ⁻¹ for
							dose group.	7 days) and azithromycin (500 mg day ⁻¹ for 5 days).
							No differences in	
							haematological or renal toxicity	
							between high and low dose	
							groups	
Gautret	36	45	42	No	HCQ 200 mg	Higher probability of viral	N.A.	Lack of randomisation.
[35]				symptoms	3 day ⁻¹ for 10	clearance after 6 days in		
(NRCT)				(17%),	days	HCQ (70%) than control		Absence of a placebo.
				upper		(13%) patients.		
				(61%) and	In 6 patients,			Small sample size.
				lower	addition of	Among patients given		
				(22%)	azithromycin	HCQ, higher probability		HCQ patients were significantly older than the control
				respiratory	500 mg on	of viral clearance after 6		group (51 vs 37).
				tract	day 1, then	days with (100%)		
				infections	250 mg day	compared to without		Short-term period of follow-up.
					on days 2-5	(57%) azithromycin.		
								6 HCQ patients (23%), but 0 control patients, lost in
						Significantly larger effect		follow-up due to transfer to ICU (50%), cessation due
						of HCQ in upper and		to nausea, and disenrollment.
						lower respiratory tract		
						infection, compared to		Viral clearance defined as $C_t > 35$, contrary to
						asymptomatic patients.		standard definition of $C_t > 40$
Huang [36]	22	44	59	Moderate	CQ 500 mg	(i) Slightly higher	5 (50%) CQ patients had 9	Absence of a placebo.
(RCT)				to severe	orally, 2 day	proportion tested	adverse events: vomiting,	.
					' for 10 days	negative on days 7, 10,	abdominal pain, nausea,	Small sample size.
						and 14.	diarrhoea, rash, shortness of	A- - - - - - - - - -
						(II) More than double	breath	Q1c not measured as endpoint.
						incidence rate of lung		
						improvement on CT (rate	No significant decrease in	
						ratio 2.1, 95% CI 0.81-	CD3+1 cell count	
						6.62) by day 14.		
						(III) All patients		
						discharged by day 14,		

						compared to 50% in the		
						control group.		
Mehra [37] (ROS)	96, 032	54	54	Variable or unclear	Within 48h of diagnosis: CQ only, 765 mg for 6.6 days HCQ only, 596 mg for 4.2 days CQ + macrolide, 790 mg for 6.8 days HCQ + macrolide, 597 mg for 4.3 days	No observable benefit. Treatment groups are associated with higher mortality than control (p<0.0001). Independently, CQ only (16%), CQ + macrolide (22%), HCQ alone (18%), and HCQ + macrolide (24%), groups exhibited elevated in-hospital mortality rates with respect to the controls (9%).	Independently, CQ only (4%), CQ + macrolide (7%), HCQ alone (6%), and HCQ + macrolide (8%), groups exhibited augmented risk of de novo in-hospital ventricular arrhythmias with respect to the controls (0.3%).	Lack of randomisation. Possible influence of unmeasured confounder(s). However, tipping point analyses demonstrated that such a factor would have to exist with a prevalence of 37-50% and HR 1.5-2.0 to make observed differences non-significant. No direct analysis of cause-effect relationship between either drug(s) and survival, or cardiovascular risk and mortality.
Singh [38] (ROS)	3,372	62	52	Variable or unclear	Unspecified dose of HCQ In 71% of patients, combination with azithromycin	No significant difference in mortality or need for mechanical ventilation.	No significant difference in incidence of <i>de novo</i> VT, fibrillation or SCD in HCQ compared to control patients.	Lack of randomisation.
lp [39] (ROS)	2,512	64	62	Variable or unclear	Median 5 days after symptoms, variable doses, e.g.: HCQ 800 mg	No significant difference in mortality.	Significantly higher proportion of mortality attributable to cardiac causes in patients treated with HCQ (21%) compared to the control group (16%).	Lack of randomisation. Significantly lower age, but later presentation in clinical course and greater symptomatic disease in the treatment group. Possible misclassification due to manual abstraction

					on day 1 + 400 mg on		Similar incidence of arrhythmias and	of HER data.
					days 2-5		cardiomyopathy in treated (5%	Possible sampling bias due to use of a convenience
					(80%) then		and 1%) and control (4% and	sample for data collection
					200 mg TID		1%) patients	
					(4%)or other		i voj padonio.	
Rosenberg	1,438	61-66	60	Variable or	Unspecified	No significant differences	Significantly higher incidence	Lack of randomisation.
[40] (ROS)				unclear	dose of HCQ	in mortality rate between	of cardiac arrest in the HCQ +	
,					±	all, following adjustment	azithromycin, compared to the	No measurement of common inflammatory marker
					Unspecified	for demography, hospital,	standard of care only, group,	confounders.
					duse of	symptom seventy and	and in the nod only,	Higher provalence of checity and dispetes in the HCO
					aziunomycin	pre-existing conditions.	only, group.	+ azithromycin, and of chronic lung and cardiovascular diseases in HCO only groups
							No significant differences in	compared to no treatment
							incidence of abnormal ECG	Both treatment groups had more clinically severe
							findings between either the	disease than the no treatment group.
							HCQ + azithromycin groups	
							and the patients receiving	Adverse events were defined as occurring at any time
							standard of care only.	point during each stay, confounding association to drug administration.
								Possible detection bias due to heightened clinical
								vigilance for arrhythmias among the HCQ \pm
								azithromycin patients.
Geleris	1,376	Not given	57	Variable or	HCQ loading	No significant difference	N.A.	Lack of randomisation.
[41] (ROS)		in main		unclear	dose 600 mg	in probability of mortality		
		text			2 day ^{⁻1} on	or intubation between		Some HCQ patients were also administered
					day 1, 400	HCQ and control		sarilumab.
					mg day ⁻¹ on	patients, under primary		Single-centre design reduces representativeness of
					days 2-5	multivariable analysis.		the sample.
					±	However, the confidence		
					Azithromycin	interval was relatively		Lower baseline PaO2:FiO2 in HCQ compared to
					500 mg on	wide (0.82 to 1.32).		control patients.
					day 1, 250			
			1		mg day on			Missing data for some variables.

		days 2-5			Possible inaccuracios in electronic health records
		~~,~ = ~			
46	Mild	HCQ 200 mg 3 day ⁻¹ for 10 days + Azithromycin 500 mg on day 1 and 250 mg on days 2-5	Significantly lower mean HCQ dosage on day 2 in the group of patients with good clinical outcome. However, multivariate analysis demonstrated no significant association between HCQ and poor clinical outcome.	9 patients had QTc prolongation > 60 ms, but final QTc did not exceed 500 ms. No observation of arrhythmic events or SCDs. All deaths in the group of patients receiving HCQ + azithromycin for 3 or more days arose from respiratory failure; no ECG recordings showed <i>torsades de pointe</i> .	Lack of randomisation. Data incomplete for some patients. Co-administration of ceftriaxone and ertapenem in patients with NEWS score of 5 or higher, and pneumonia. 32% of patients with poor clinical outcome had a lower plasma HCQ concentration than the therapeutic target (including 2 patients in whom HCQ was absent).
					Possible lack of adherence to prescribed treatment due to lack of control of therapy intake.
63	Severe	HCQ 200 mg 2 day ⁻¹ for 7- 10 days	Significantly lower mortality rate in HCQ (19%) compared to control (47%) patients. No significant difference in average hospital stay. Among patients who died, significantly longer hospital stay for the HCQ (15 days) compared to the control (8 days) group. Significant reduction in plasma IL-6 in HCQ, but not control patients. Among patients with IL-6 > 60 pg ml ⁻¹ , HCQ treatment, but not control	N.A.	Lack of randomisation. Considerable imbalance in the sample size of the treatment and control groups. Interferon application reached 11% in the control, but 0% in the treatment group.
	63	46 Mild 63 Severe	 46 Mild HCQ 200 mg 3 day⁻¹ for 10 days + Azithromycin 500 mg on day 1 and 250 mg on days 2-5 63 Severe HCQ 200 mg 2 day⁻¹ for 7- 10 days 	46 Mild HCQ 200 mg 3 day ⁻¹ for 10 days Significantly lower mean HCQ dosage on day 2 in the group of patients with good clinical outcome. 47 Azithromycin 500 mg on day 1 and 250 mg on days 2-5 However, multivariate analysis demonstrated no significant association between HCQ and poor clinical outcome. 63 Severe HCQ 200 mg 2 day ⁻¹ for 7- 10 days Significantly lower mortality rate in HCQ (19%) compared to control (47%) patients. No significant difference in average hospital stay. Among patients who died, significantly longer hospital stay for the HCQ (15 days) compared to the control (8 days) group. Significant reduction in plasma IL-6 in HCQ, but not control patients. Among patients with IL-6 > 60 pg ml ⁻¹ , HCQ treatment, but not control treatment reversed the trend after 10 days. and	46 Mild HCQ 200 mg 3 day' ¹ for 10 days + Azithromycin 500 mg on day 1 and 250 mg on days 2-5 Significantly lower mean HCQ dosage on day 2 in the group of patients with good clinical outcome. However, multivariate analysis demonstrated no significant association between HCQ and poor clinical outcome. 9 patients had QTc prolongation > 60 ms, but final QTc did not exceed 500 ms. No observation of arrhythmic events or SCDs. 63 Severe HCQ 200 mg 2 day' ¹ for 7- 10 days Significantly lower mortality rate in HCQ (19%) compared to control (47%) patients. N.A. 63 Severe HCQ 200 mg 2 day' ¹ for 7- 10 days Significantly lower mortality rate in HCQ (19%) compared to control (47%) patients. N.A. 63 Severe HCQ 200 mg 2 day' ¹ for 7- 10 days Significant difference in average hospital stay. Among patients who died, significant ly longer hospital stay for the HCQ (15 days) compared to the control (8 days) group. N.A. Significant reduction in plasma IL-6 in HCQ, but not control patients. Among patients with IL-6 > 60 pg m ¹ , HCQ Significant reduction in plasma IL-6 in HCQ, but not control patients. Among patients with IL-6 > 60 pg m ¹ , HCQ

						significantly reduced fatality.		
						Early start of HCQ within 5 days of admission		
						reduced fatality		
						compared to a late start.		
						However, this difference		
						was not statistically		
						significant.		
Magagnoli	368	68-70	100	Variable or	Unspecified	No significant difference	N.A.	Lack of randomisation.
[44] (ROS)				unclear	dose of HCQ	in ventilation risk, or		
					±	mortality following		Absence of a control.
					doso of	trootmont group		Higher probability of proscribing $HCO + azithromycin$
					azithromycin	compared to the control		to patients with more severe metabolic
					azitinomyoni	compared to the control.		haematological and ventilatory symptoms of COVID-
						Higher mortality risk in		
						HCQ only, but not HCQ +		
						azithromycin, groups		Significant differences in demography, vital signs,
						compared to the control.		prescription drug use, comorbidities, and disease
								severity.
								However, all adjusted by propensity score.
								The vast majority of the sample were African
								American and > 65 yrs, with both groups exhibiting
								disproportionately high rates of hospitalisation.
Kim [45]	270	38	36	Mild to	HQ 200 mg	No significant difference	Conservative management	Lack of randomisation.
(ROS)				moderate	2 day +	In risk of mortality or ICU	only experienced significantly	The concernative core only group subilited forwar
					unspecified	transier.	newer adverse events,	The conservative care only group exhibited lewer
					antibiotics	Significantly shorter time	and nauses and vomiting than	
						to complete or probable	patients in either active drug	Use of antivirals in the control group only
						viral clearance and to	aroup	However, HQ + antibiotic outcomes were superior to
						resolution of fever and	However, adverse effects did	those of healthier baseline patients given
						cough symptoms in the	not significantly differ between	conservative management.
						HQ + antibiotics group.	the HQ + antibiotic (12) and	

						Significantly shorter hospital stay with HQ + antibiotics compared to either of the control groups.	Lop/R + antibiotic (7) groups. No observed serious cardiac toxicity. However, this was not actively searched for.	No measurement of either QT prolongation or retinopathy.
Saleh [46] (ROS)	201	59	58	Moderate to severe	CQ 500 mg 2 day ¹ on day 1, and 500 mg day ¹ on days 2-5 OR HCQ 400 mg 2 day ¹ on day 1, and 200 mg 2 day ¹ on days 2-5 \pm Azithromycin 500 mg on days 1-5	N.A.	Significant increase in QTc to peak and post-treatment. Significantly shorter maximum QTc in CQ/HCQ only compared to CQ/HCQ ± azithromycin patients. 8% of patients experienced de novo atrial fibrillation, and 3% had monomorphic ventricular tachycardia, which was non- sustained for all but one patient. No observation of <i>torsades de</i> <i>pointe</i> . However, 4% of patients had to prematurely discontinue HCQ owing to QT prolongation	Lack of randomisation. Absence of a control. Small sample size relative to total cohort population treated. Lack of reporting of instances of torsades de pointe may be influenced by reporting bias. QT intervals in MCOT patches while on therapy were not correlated to baseline ECGs.
Mahevas [47] (ROS)	173	60	72	Moderate	Within 48h of admission: HCQ 600 mg day ⁻¹	No significant differences in overall survival rate, survival rate without transfer to ICU, survival rate without ARDS, time to weaning from O2 therapy, or time to discharge.	10% of HCQ patients experienced averse ECG modifications requiring cessation of treatment after a median of 4 days. Among them, 88% had a QTc prolongation > 60 ms (including > 500 ms in one	Lack of randomisation. Small sample size. Lower probability of co-administration of azithromycin in the HCQ (18%) compared to the control (29%) group. Higher probability of co-administration of amoxicillin

						Likewise for patients with better prognoses upon admission and less severe COVID-19 symptoms.	patient). One of these patients presented with a 1 st degree AV block after 2 days despite a lack of concomitant proarrhythmic medication.	and clavulanic acid in the HCQ (52%) compared to the control (28%) group. HCQ patients had lower prevalence of comorbidities, except hepatic cirrhosis.
								The 4 covariates exceeding the standardised difference threshold were excluded from the final propensity score model. Imbalance in the number of HCQ patients between centres not taken into account by the propensity score model.
Ramireddy [47] (ROS)	98	62	61	Variable or unclear	Unspecified dose of HCQ ±	N.A.	Significantly longer QTc, even when corrected with the Fridericia formula, with drug	Lack of randomisation. Absence of control.
					Unspecified dose of azithromycin		administration. Mean QT prolongation with HCQ ± azithromycin was	Small sample size.
							several-fold higher (17.2 ms vs. 0.5 ms) than with azithromycin only.	Variation in dosing patterns and duration for each drug.
							12% of patients reached QTc of 500 ms (if QRS < 120 ms).	Possible underestimation of critical QTc prolongation due to inconsistency in ECG measurement.
							of 550 ms (if QRS > 120 ms), or prolongation > 60 ms.	Higher probability of coadministration of HCQ + azithromycin in patients with shorter baseline QTc, and of monotherapy in those with longer baseline
							No observation of <i>torsades de pointe</i> , syncope, or lethal arrhythmias.	QTc.
Mercuro [49] (POS)	90	60	51	Moderate to severe	HCQ 400 mg 2 day ⁻¹ on day 1, 400	N.A.	11% had QTc prolongation > 60 ms, including 3% of HCQ only and 13% of concomitant	Lack of randomisation.
					mg day ⁻¹ on days 2-5		HCQ + azithromycin patients.	Possible underestimation of QTc due to short follow-
					± Azithromycin		Final QTc exceeded 500 ms in 20% of patients, including 19%	up period. Possible role of COVID-19-associated myocarditis

						of HCQ only and 21% of concomitant HCQ + azithromycin patients. 11% of patients ceased HCQ before day 5 due to arrhythmic and GI adverse events, as well as a case of hypoglycaemia. One such patient developed <i>torsades de pointe</i> 3 days after cessation.	 and/or stress cardiomyopathy in observed adverse events. Baseline QTc was shorter in HCQ + azithromycin compared to HCQ only patients. Most patients had at least 1 cardiac comorbidity, and were taking 2 or more drugs prolonging QTc.
Chorin [50] 84 (ROS)	63	74	Variable or unclear	HCQ 200 mg 3 day ⁻¹ for 10 days + azithromycin 500 mg on day 1, then 250 mg day ⁻¹ on days 2-5	N.A.	Significant prolongation of mean QTc from 435 ± 24 ms at baseline, to 463 ± 32 ms at maximum, arising 3.6 ± 1.6 days after therapy initiation. 11% of patients developed severe prolongation to QT > 500 ms.	Lack of randomisation. Absence of control. Small sample size.
Mallat [51] 34 (ROS)	37	74	Mild to moderate	HCQ 400 mg 2 day ⁻¹ on day 1, HCQ 400 mg day ⁻¹ for 10 days	Significantly longer time to viral clearance in HCQ (17 days) compared to control (10 days) patients, even after adjustment for confounders, such as symptoms and pneumonia or oxygen therapy. By day 14, significantly lower proportion of HCQ (48%) compared to control (91%) patients tested negative for SARS-CoV-2.	No observed side-effects of HCQ. No significant changes in plasma counts of leukocytes, lymphocytes, or concentrations of CRP and ferritin in either HCQ or control groups.	Lack of randomisation. Small sample size. No direct measurement of QTc prolongation. HCQ group had both significantly higher comorbidities and D-dimer levels

Abbreviations: RCT = randomised controlled trial; NRCT = non-randomised controlled trial; ROS = retrospective observational study; POS =

prospective observational study; SCD = sudden cardiac death; MCOT = mobile cardiac outpatient telemetry

3.0 Results

3.1 Search Breakdown

The results of each search, and the number failing and passing exclusion and inclusion criteria, respectively, have been summarised in a flowchart (Figure 1).

On May 26 2020, searches for phrases one ("COVID" AND "chloroquine"[title/abstract] AND "outcomes"[full text]) and two ("COVID" AND "hydroxychloroquine"[title/abstract] AND "outcomes"[full text]) yielded results distributed as follows: 53 on PubMed; one on Web of Science; and 40 on Embase. The subsequent removal of 49 duplicates left 45 unique items from these three databases. In parallel, use of the same stenography in searching MEDRXIV yielded 227 unique results.

Of the unique papers discovered in the first three databases, 21 failed the exclusion criteria, distributed categorically as follows: eight reviews; five case reports; four systematic reviews; three commentaries; and one editorial. Likewise, of those unique items found on the preprint servers, 82 were excluded by the same criteria, distributed categorically as follows: 39 systematic reviews; 18 models of COVID-19 spread and/or symptoms; 12 study proposals and/or protocols; 10 reviews; and three commentaries.

As part of the search attrition methodology, parallel application of the inclusion criteria to each set of remaining unique results left eight and 13 items from the first three, and preprint, databases, respectively. Of these 21 studies, two were duplicates, such that 19 papers answering the research question passed onto quality appraisal.

3.2 Quality Appraisal

The use of 13 criteria created in accordance with the aforementioned Checklist of Review Criteria elaborated by the GERA-RIME committee, meeting 12 of which was judged to be indicative of scientific rigour, did not exclude any of the remaining 19 items (Table 2).

3.3 Study Design

As a result of the relatively broad scope of the research question, the authors of the 19 papers passing quality appraisal employed a variety of study types, therapeutic doses, and primary and/or secondary outcomes (Table 2). It is thus essential to distinguish results by study design in order to prevent invalid inferences drawn from comparison of data sets.

3.3.1 Study type

Although the research question requires the use of CQ and/or HCQ in patients suspected of, and diagnosed with, COVID-19, there exists a range of possible approaches to the collection of data obtained from such patients.

The gold standard of primary clinical research into the efficacy and safety of drugs administered to humans is the randomised controlled trial, evidence from which may be further buttressed by masking of subjects, experimenters, or both, as well as the use of a placebo in the control group. However, only 21% of items passing quality appraisal were clinical trials, of which, though ³/₄ were randomised, none employed either masking or an exclusively placebo control. While unacceptable under normal circumstances, the absence of both masking and a placebo is admissible in light of the ethical violation that would otherwise result from the use of either in the context of patient consent being unlikely.

The remaining 79% of papers were observational studies, of which 93% retrospectively searched hospital databases to, and 7% prospectively, collect clinical data obtained by following up on cohorts of patients from the time they received CQ and/or HCQ, or the standard of care only, until a defined end-point. Despite the vast majority (73%) using a case-control structure, a minority (27%) constituted case series focusing on the cardiac safety of drugs administered to hospitalised COVID-19 patients for a given duration of time.

3.3.2 Therapeutic doses

The deliberate absence of a specified dose in the research question accounts for the diversity of administration regimens among the 19 papers. Indeed, the doses used largely reflect the studies being conducted on different dates, which, in turn, influences the relative sway of either federal healthcare guidelines or the results of prior clinical research on regimen selection.

Every clinical trial tested a distinct dosing scheme: 500 mg CQ, twice a day for 10 days [36]; 600 mg CQ, twice a day for 10 days [34]; a loading dose of 1200 mg HCQ per day for 3 days, followed by a maintenance dose of 800 mg per day for 2 or 3 weeks if symptoms are mild/moderate or severe, respectively [33]; and 200 mg CQ three times a day for 10 days, with or without 500 mg azithromycin for 1 day, followed by 250 mg per day for 4 days [35]. Likewise, each clinical trial treated its control group differently, from: 400/100 mg lopinavir/ritonavir twice a day for 10 days [36]; to 450 mg CQ and one placebo tablet twice a day for 1 day, followed by 450 mg CQ and one placebo tablet first and 4 placebo tablets second for 4 days, followed by 4 placebo tablets twice a day for 5 days [34]; and standard of care only [33][35].

By contrast, the retrospective and often multi-centre nature of many observational studies has resulted in 40% using variable or undeclared doses, all collecting data from some patients taking azithromycin in combination with the HCQ. The remaining 60% of items relied on highly divergent dosing regimens (Table 2). One popular iteration administered 200 mg HCQ twice a day, along with: undeclared antibiotics [45]; 500 mg azithromycin per day for 1 day, followed by 250 mg per day for 4 days [50][42]; no antibiotics, and no time declaration [43]. The most common higher dose of choice involved giving 400 mg HCQ, twice a day for 1 day, followed either by: 500 mg per day for 10 days [51]; 400 mg per day for 4 days, with or without an undeclared dose of azithromycin [49]; or 200 mg twice per day for 4 days, with or without 500 mg azithromycin per day for 5 days [46]. In contrast, two studies relied on a much higher loading dose of 600 mg HCQ, either once per day [47] throughout, or twice a day for 1 day, followed by 400 mg per day for 4 days, and with or without 500 mg azithromycin per day for 1 day, followed by 250 mg per day for 4 days [41]. Only one set of authors also analysed data for patients taking CQ, at a dose of 500 mg twice a day for 1 day, followed by 500 mg per day for 4 days, with or without 500 mg azithromycin per day for 5 days [46]. Similarly, the 73% of studies with a control treated the group differently, giving standard of care without (55%), or with declaration of additional antivirals (18%), antibiotics (18%), or both (9%).

3.3.3 Primary and/or secondary outcomes

Given the vast array of possible measures of CQ and HCQ efficacy and safety in COVID-19 patients, the different authors outlined distinct primary and/or secondary outcomes.

The primary outcome in 75% of clinical trials was viral clearance, the definition for which varied from $C_t > 40$ (67%) to $C_t > 35$ (33%) for PCR amplification of SARS-CoV-2 RNA. Likewise, 53% of observational studies directly measured a specific indicator of efficacy other than mortality rate, using similar outcomes to the clinical trials, as well as the duration of hospital stay, need for mechanical ventilation, and probability of transfer to an intensive care unit (ICU).

As regards direct measurement of safety, 75% of clinical trials and 73% of observational studies recorded specific adverse events as an indicator of CQ and/or HCQ safety in COVID-19 patients, with 63% of the total actively focusing on cardiac pathology.

Notably, 53% of studies used mortality rate as an end-point. In isolation, however, risk of death could be indicative of either safety or efficacy. As such, this review reports the findings on mortality rate separately from those pertaining to outcomes that are specific measures of one of efficacy or safety.

3.4 Results

Of the clinical trials providing data on a specific indicator of CQ and/or HCQ efficacy in patients suspected of, and diagnosed with, COVID-19, 67% showed a significant increase in the probability of viral clearance in the treatment, compared to the control, group [36][35]. Conversely, 33% failed to find any significant difference in the likelihood of negative SARS-CoV-2 conversion [33], despite using a larger dose of HCQ in patients, of whom 99% exhibited only mild to moderate symptoms.

In contrast, 75% of observational studies employing an endpoint specific to efficacy recorded no significant difference in the attainment of outcomes, such as duration of hospital

stay, need for mechanical ventilation, and probability of transfer to an intensive care unit (ICU), between COVID-19 patients given a range of CQ and/or HCQ doses, and the control groups. One such study, however, discovered a significantly lower mean HCQ dose in patients with better clinical outcomes [42], while the two remaining sets of authors found either a significant deceleration [51] or acceleration [45] in viral clearance, the latter conflicting with its own data on an unchanged probability of ICU transfer.

All clinical trials and 82% of observational studies examining an indicator unique to drug safety discovered a higher probability of adverse events in those treated patients suspected of, and diagnosed with, COVID-19. Seventy-five percent of the total papers focusing on cardiac side-effects found a greater incidence among patients administered a wide range of CQ and/or HCQ doses, with QTc prolongation the most common finding, in addition to its consequences of VT and cardiac arrest.

Of the total studies using mortality rate as an end-point, 60% reported no significant change in the risk of death, while 30% showed an elevation, and 10% a depression, in treated relative to control patients.

4.0 Discussion

The absence of a pharmacological treatment tailored to COVID-19 has rendered urgent the search to find alternative therapies by repositioning drugs with the theoretical potential to alleviate symptoms. However, that a solution is hypothetically plausible is insufficient grounds for translation into clinical practice. Indeed, any therapeutic repurposing must only proceed in light of strong evidence for the pre-clinical basis, and clinical efficacy and safety, of the drug in question. This review finds that, while such evidence certainly

exists for the former, it does not for the latter, calling into question any clinical use of CQ and/or HCQ in COVID-19 patients in the absence of high-quality randomised clinical trial data.

4.1 Pre-clinical indications of the potential of CQ and HCQ to treat COVID-19

Pre-clinical studies performed *in vitro* provide strong evidence for the theoretical utility of CQ and HCQ in inhibiting all stages of viral entry, maturation, and spread.

In vitro, CQ blocks infection both at, and after, entry of SARS-CoV-2 into Vero E6 cells, with an EC₅₀ of 1.13 μ M [19]. Indeed, although therapeutic doses of CQ do not seem to alter S protein glycosylation [52], whose pattern is distinct from that of SARS-CoV-1 [53], they may inhibit biosynthesis of sialic acid [54], *N*-glycosylation of ACE2, as well as downregulating the expression of PICALM [55] in the clathrin-dependent endocytosis machinery.

Furthermore, immunofluorescence analysis of the amount of NP in distinct vesicular compartments of the host cell has demonstrated that treatment of infected cells with CQ and HCQ stalls transfer of viruses from early to late endosomes [56]. In fact, by increasing the pH of the early endosome, CQ has the potential to reduce acid-dependent proteolytic cleavage of the S protein by cathepsin and TMPRSS2, thereby inhibiting viral uncoating, genomic replication and particle maturation [57]. Despite its similar effect on viral distribution, as well as its comparable cytotoxicity [58], to CQ, HCQ appeared to amplify and enlarge the late endosomes, implying a slightly distinct mechanism of action. Furthermore, there exists conflicting evidence for the relative *in vitro* efficacy of the two drugs [59].

It is nonetheless clear that the initial basis for investigating the translatability of CQ

and/or HCQ to the treatment of hospitalised COVID-19 patients was predicated on highquality evidence for its pre-clinical antiviral efficacy.

4.2 Clinical evidence of the efficacy of CQ and HCQ in the treatment of COVID-19

That CQ and HCQ can reduce viral entry, trafficking, and budding *in vitro* constitutes evidence of translational potential relies on the underlying assumption that symptom severity is a function of viral replication. Yet, while viral load may influence severity in the very early stages of COVID-19 [60] – as in SARS [61] – , subsequent symptoms result firstly from the initial cytokine wave of the innate immune response [62], then a state of immunodeficiency and lymphopenia [63], and, finally, a potentially lethal cytokine storm [64]. The causal distinction between these symptomatic phases highlights not only the difficulty in repurposing a single drug for use at all time-points, but also the need to approach with caution the comparison of trial data collected from patients given drugs at different times post-infection.

The sample of one of the first clinical trials performed on patients testing positive upon PCR amplification of SARS-CoV-2 RNA comprised asymptomatic patients (17%), as well as those with upper (61%) and lower (22%) respiratory tract infections, thereby capturing the range of symptom severity. After 6 days of treatment, patients given HCQ alone had a higher probability of viral clearance compared to those given the standard of care only (57% vs. 13%), rising to 100% in patients also given azithromycin [35]. That the authors additionally discovered a greater drug effect on patients with upper and lower respiratory tract infections than on asymptomatic individuals raises the possibility that the potential therapeutic benefit of HCQ in COVID-19 patients lies in its capacity for immunomodulation.

On a theoretical level, the anti-inflammatory effects of HCQ render such an effect possible. Indeed, through alkalinisation of early endosomes, CQ and HCQ could impair: PAMPinduced activation of TLR7 and TLR9 [65], and, by extension, MMP-9 expression [66]; antigen presentation by major histocompatibility complexes (MHCs) [67][68]; prostaglandin and thromboxane production [69]; and T and B cell activation [70], differentiation, and proliferation [20]. Importantly, both SARS-CoV-2 [71] [72] and related coronaviruses, such as SARS-CoV-1 [73] and MERS-CoV [74], may incur pulmonary damage through TNF-(alpha) [75], which CQ and HCQ can down-regulate through p38 MAPK inhibition [76].

However, this study had several considerable limitations. In addition to the lack of randomisation and the use of a C_t of 35 rather than 40 as the threshold for viral clearance, the sample size of 36 was very small. Moreover, 23% of patients in the treatment, but none of those in the control, group were lost in the follow-up due to transfer to the ICU, disenrollment, or premature cessation, leaving the sample even further underpowered. Indeed, Bayesian reanalysis of the data demonstrates that the statistical evidence for efficacy weakens to anecdotally positive upon the exclusion of untested patients, and even to anecdotally negative with the assumption that untested patients were infected with SARS-CoV-2 [77].

Other research coming to the same conclusion regarding the efficacy of CQ and/or HCQ in COVID-19 patients similarly exhibited numerous shortcomings. For instance, although a randomised controlled trial found that, compared to the control group administered lopinavir/ritonavir, patients with moderate and severe symptoms given CQ exhibited more than double the rate of improvement in CT scan indicators of pulmonary

health, the sample consisted of only 22 individuals [36]. Likewise, half of the observational studies showing a lower mortality rate or higher probability of viral clearance with low dose CQ and/or HCQ treatment in patients with severe symptoms were severely underpowered, with a significant imbalance in size of the treatment (48) and control (502) groups [43].

By contrast, the majority of observational studies failing to find a significant difference in the mortality rate between COVID-19 patients treated with CQ and/or HCQ and those given the standard of care (with or without additional antibiotics or antivirals) were sufficiently powered. Furthermore, the homogeneity of, and correction for, baseline characteristics in the case and control cohorts further buttresses the reliability of the evidence presented by studies with sample sizes of 1,061 [42], 1,376 [41], 1,438 [40], 3,372 [38], and 96, 032 [37]. Although one study with a sample size of 2,512 used a treatment cohort with a significantly lower age but greater symptomatic disease compared to the control [39], the vast majority of papers coming to this conclusion despite a significant difference in sample structure and comorbidities were underpowered and thus unlikely to influence the conclusion of this review [47]. Likewise, the only retrospective cohort study suggesting that HCQ delayed viral clearance in COVID-19 patients with mild to moderate symptoms suffered from both a very small sample size of 34, and considerably more frequent comorbidities in the treatment group.

As such, it is reasonable to conclude that the weight of present evidence does not come down in favour of either CQ or HCQ being efficacious in the treatment of COVID-19 patients, relative to the standard in-hospital management of symptoms. Indeed, even if the evidence demonstrating a lack of efficacy were absent, the poor quality of data suggesting

any significant benefit should preclude any definitive judgment until the completion of highquality randomised controlled trials.

4.3 Clinical evidence of the safety of CQ and HCQ in the treatment of COVID-19

Robust evidence for the safety of an otherwise efficacious drug is a prerequisite for its widespread application in any clinical setting. In light of the inefficacy – or, at least, wholly unsubstantiated benefit – of CQ and/or HCQ administration in patients with COVID-19, there exists an even more compelling imperative to ensure that any compassionate use did, and does, not contribute to excess mortality.

As an aminoquinolone, CQ, and its derivative, HCQ, are proarrhythmic [78]. Arrythmias arise from an imbalance of the normal physiological variables influencing the activation and inactivation kinetics of the cardiac ion channels that permit the transmembrane currents forming the foundation of the cardiac action potential. In a healthy *milieu intérieur*, the waveform of this cardiac action potential is quadriphasic [79]. Electrical diastole (IV) precedes a rapid depolarisation (0), after which a refractory period is followed by a slow hyperpolarisation (I), a 300 ms plateau (II), and then a period of repolarisation (III) to resting membrane potential (E_m). By blocking – in order of increasing potency – the delayed (I_{Kr}) and inwardly-rectifying (I_{K1}) K⁺ currents [80], and the latter preferentially at depolarised E_m , CQ and HCQ significantly prolong the QT interval and slow ventricular conduction, thereby predisposing to early-after-depolarisation and, by extension, *torsades de pointe*. Combined with their tonic block of voltage-gated Na⁺ and L-type Ca²⁺ currents at low channel opening

frequencies, QT interval prolongation thus significantly increases the risk of potentially fatal VTs.

Indeed, there is considerable evidence that CQ and/or HCQ treatment predisposes COVID-19 patients to tachyarrhythmia. In fact, a significant association of high doses of CQ with lethality in patients with severe symptoms forced the premature termination of a randomised controlled trial [34]. Despite this relationship with mortality risk disappearing upon correction for age, there remained a significantly higher proportion of patients in the high (19%) compared to the low (11%) dose group with QTcF > 500 ms, including 2.7% patients who experienced VT before death. However, given the abortion of the study, as well as the co-administration of QT-prolonging oseltamivir [81] confounding the causal link to cardiac side-effects, these data, alone, are insufficient to conclude that CQ is unsafe in COVID-19 patients.

However, several highly powered retrospective observational studies have found significant excess mortality in patients treated with CQ and/or HCQ relative to controls, providing further evidence of a lack of drug safety. For instance, in 96, 032 patients with no significant differences in comorbidities between groups, patients given CQ (4%) or HCQ (6%) alone had a significantly augmented risk of *de novo* in-hospital ventricular arrhythmias, compared to controls (0.3%) given standard therapy, including remdesivir [37]. Notably, the addition of a macrolide antibiotic further increased VT risk, owing to synergistic prolongation of QTc. Furthermore, ³/₄ of the remaining studies with smaller cohorts – but still over 1,000 – demonstrated either a significant increase in the risk of VT or of cardiac-related mortality incurred by the administration of CQ. Moreover, despite being individually

underpowered, several smaller retrospective database searches focusing specifically on the QTc duration have consistently and independently supported a significant prolongation in hospitalised COVID-19 patients treated with a range of CQ and/or HCQ doses [46][49][50][48].

From a theoretical standpoint, however, the cardiac safety risk of CQ and HCQ use is unlikely uniform among COVID-19 patients. Indeed, there is a significant positive correlation between baseline QTc and age [82], QTc prolongation and antidysrhythmic, antipsychotic or macrolide antibiotic coadministration [83], and QT interval dispersion and mortality risk in type II diabetes mellitus [84]. Given the relative risk conferred by both older age [85] and type II diabetes mellitus [86] on symptom severity and consequent probability of hospitalisation of COVID-19 patients, the data for in-hospital CQ/HCQ safety may not be extrapolable to many infected individuals in the population due to selection bias [87]. Regardless, drawing causal inferences from such observational studies is inadvisable given the lack of randomisation and absence of a placebo in the control groups, leaving the data susceptible to confounders. Moreover, the wide variation in average sample patient age, symptom severity and drug dosing regimen (Table 2) further complicates inferences of reliable agreement between the papers. Yet, for the most highly powered of the studies, tipping point analysis suggested that any confounding factor not taken into account would have to exist with a prevalence of 37-50% to render the observed differences insignificant [34]. Furthermore, the strength of the correlative relationship alone between CO/HCO use and cardiac side-effects in hospitalised patients is sufficient to warrant extreme caution when administering them to patients in such a clinical setting.

Therefore, as of 26 May 2020, the strongest available evidence suggests that, relative to standard in-hospital management of symptoms, the use of CQ and HCQ to treat hospitalised COVID-19 patients has likely been unsafe. At the very least, the poor quality of data failing to find any significant changes in the risk of VT should preclude definitive judgment on drug safety until the completion of high-quality randomised clinical trials.

5.0 Limitations

Crucial to the understanding of the conclusions drawn in this systematic review is an appreciation of its many limitations, which relate to both the search methodology and data analysis.

Insofar as peer-reviewed publications are concerned, this review searched three databases to yield only 45 unique results, leaving the authors to also seek the findings of 227 papers on a preprint server. Despite facilitating the collection of a more representative sample of current research on the subject in question, the absence of documented expert scrutiny ought to prevent their data from influencing clinical decisions. Nevertheless, to compensate for the lack of peer review, rigorous application of the quality appraisal criteria established by the GEA-RIME committee and the Task Force of Academic Medicine ensured that only data from adequately designed studies were taken into account. Importantly, however, that peer-reviewed journal material was no longer a prerequisite for inclusion may have slightly reduced the effects of positive publication bias [88] on the results of this systematic review.

However, the predominant shortcoming of the review is its inability to completely disentangle the large differences in study design when making comparisons between different data sets from the included papers. Indeed, despite stressing the obvious invalidity of cutting across distinct sample sizes, baseline characteristics, drug doses, and individual limitations, a systematic review, by nature, does exactly that. The categorisation of the results and data analysis by randomisation, COVID-19 symptom severity, and HCQ/CQ dosage constitutes an attempt to reduce this problem of comparative inferences as greatly as possible.

6.0 Summary and Conclusions

On March 18 2020, the WHO announced the launch of an international phase III-IV randomised controlled trial, with four arms, measuring the efficacy and safety of: (1) remdesivir; (2) lopinavir and ritonavir; (3) lopinavir, ritonavir, and IFN-(beta); and (4) CQ or HCQ [89]. In the meantime, amidst a dearth of high-quality evidence from randomised clinical trials, the U.S. Food and Drug Administration (FDA) issued an emergency use authorisation of CQ and HCQ in COVID-19 patients [90].

Since then, data from the most robust case-control studies have failed to find any significant efficacy, and, indeed, a notable lack of (particularly cardiac) safety for the use of CQ and/or HCQ to treat hospitalised COVID-19 patients. On May 25 2020, the WHO suspended the fourth arm of the Solidarity Trial, citing these safety concerns.

The urge to begin the Solidarity Trial arose from the understandable, and – as aforementioned – necessary desire to rapidly compensate for the prior and present scarcity of randomised data on the efficacy and safety of CQ and/or HCQ in patients infected with SARS-CoV-2. Yet the clinical knowledge of, and subsequent evidence for, the cardiac sideeffects of CQ and HCQ, call into question the scientific prudence of the FDA's initial decision to authorise, and only belatedly caution against, their use in hospitalised patients already at risk owing to comorbidities. In any case, any resumption in administration of either CQ or HCQ in a hospital setting would require strong substantiation of both their safety and efficacy from high-quality randomised controlled data.

Acknowledgements

N.A.

Conflict of interest statement

None to declare.

List of abbreviations

- CoV-coronavirus
- COVID-19 Coronavirus Disease 2019
- CQ chloroquine
- FDA Food and Drug Administration
- HCQ hydroxychloroquine
- ICU intensive care unit
- MERS Middle East Respiratory Syndrome
- PICOT Population, intervention, comparison, outcome, time
- PRISMA Preferred Reporting Items for Systematic Reviews and Meta-Analyses
- QTcF The corrected QT interval by Fredericia
- SARS Severe Acute Respiratory Syndrome
- WHO World Health Organisation
- VT ventricular tachyarrythmia

References

- Wu Z, McGoogan JM. Characteristics of and Important Lessons From the Coronavirus Disease 2019 (COVID-19) Outbreak in China. JAMA. 2020. doi:10.1001/jama.2020.2648
- Gorbalenya AE, Baker SC, Baric RS, de Groot RJ, Drosten C, Gulyaeva AA, et al. The species Severe acute respiratory syndrome-related coronavirus: classifying 2019-nCoV and naming it SARS-CoV-2. Nature Microbiology. 2020. doi:10.1038/s41564-020-0695-z
- 3. Cucinotta D, Vanelli M. WHO declares COVID-19 a pandemic. Acta Biomedica. 2020. doi:10.23750/abm.v91i1.9397
- Dong E, Du H, Gardner L. An interactive web-based dashboard to track COVID-19 in real time. The Lancet Infectious Diseases. 2020. doi:10.1016/S1473-3099(20)30120-1
- Zhou P, Yang X Lou, Wang XG, Hu B, Zhang L, Zhang W, et al. A pneumonia outbreak associated with a new coronavirus of probable bat origin. Nature. 2020. doi:10.1038/s41586-020-2012-7
- Tang X, Wu C, Li X, Song Y, Yao X, Wu X, et al. On the origin and continuing evolution of SARS-CoV-2. Natl Sci Rev. 2020. doi:10.1093/nsr/nwaa036
- Jaimes JA, André NM, Chappie JS, Millet JK, Whittaker GR. Phylogenetic Analysis and Structural Modeling of SARS-CoV-2 Spike Protein Reveals an Evolutionary Distinct and Proteolytically Sensitive Activation Loop. J Mol Biol. 2020. doi:10.1016/j.jmb.2020.04.009
- Marra MA, Jones SJM, Astell CR, Holt RA, Brooks-Wilson A, Butterfield YSN, et al. The genome sequence of the SARS-associated coronavirus. Science (80-). 2003. doi:10.1126/science.1085953
- Wrapp D, Wang N, Corbett KS, Goldsmith JA, Hsieh CL, Abiona O, et al. Cryo-EM structure of the 2019-nCoV spike in the prefusion conformation. Science (80-). 2020. doi:10.1126/science.aax0902
- Hamming I, Timens W, Bulthuis MLC, Lely AT, Navis GJ, van Goor H. Tissue distribution of ACE2 protein, the functional receptor for SARS coronavirus. A first step in understanding SARS pathogenesis. J Pathol. 2004. doi:10.1002/path.1570

- Fantini J, Di Scala C, Chahinian H, Yahi N. Structural and molecular modelling studies reveal a new mechanism of action of chloroquine and hydroxychloroquine against SARS-CoV-2 infection. Int J Antimicrob Agents. 2020. doi:10.1016/j.ijantimicag.2020.105960
- Hoffmann M, Kleine-Weber H, Schroeder S, Krüger N, Herrler T, Erichsen S, et al. SARS-CoV-2 Cell Entry Depends on ACE2 and TMPRSS2 and Is Blocked by a Clinically Proven Protease Inhibitor. Cell. 2020. doi:10.1016/j.cell.2020.02.052
- Coutard B, Valle C, de Lamballerie X, Canard B, Seidah NG, Decroly E. The spike glycoprotein of the new coronavirus 2019-nCoV contains a furin-like cleavage site absent in CoV of the same clade. Antiviral Res. 2020. doi:10.1016/j.antiviral.2020.104742
- McGonagle D, Sharif K, O'Regan A, Bridgewood C. The Role of Cytokines including Interleukin-6 in COVID-19 induced Pneumonia and Macrophage Activation Syndrome-Like Disease. Autoimmunity Reviews. 2020. doi:10.1016/j.autrev.2020.102537
- Poissy J, Goutay J, Caplan M, Parmentier E, Duburcq T, Lassalle F, et al. Pulmonary Embolism in COVID-19 Patients: Awareness of an Increased Prevalence. Circulation. 2020. doi:10.1161/circulationaha.120.047430
- 16. Klok FA, Kruip MJHA, van der Meer NJM, Arbous MS, Gommers DAMPJ, Kant KM, et al. Incidence of thrombotic complications in critically ill ICU patients with COVID-19. Thromb Res. 2020. doi:10.1016/j.thromres.2020.04.013
- 17. Zhang L, Yan X, Fan Q, Liu H, Liu X, Liu Z, et al. D-dimer levels on admission to predict in-hospital mortality in patients with Covid-19. J Thromb Haemost. 2020. doi:10.1111/jth.14859
- Ozden S, Lucas-Hourani M, Ceccaldi PE, Basak A, Valentine M, Benjannet S, et al. Inhibition of Chikungunya virus infection in cultured human muscle cells by furin inhibitors: Impairment of the maturation of the E2 surface glycoprotein. J Biol Chem. 2008. doi:10.1074/jbc.M802444200
- Wang M, Cao R, Zhang L, Yang X, Liu J, Xu M, et al. Remdesivir and chloroquine effectively inhibit the recently emerged novel coronavirus (2019-nCoV) in vitro. Cell Research. 2020. doi:10.1038/s41422-020-0282-0
- 20. Yang J, Yang X, Yang J, Li M. Hydroxychloroquine inhibits the differentiation of Th17 cells in systemic lupus erythematosus. J Rheumatol. 2018. doi:10.3899/jrheum.170737

- Carter AE, Eban R. Prevention Of Postoperative Deep Venous Thrombosis In Legs By Orally Administered Hydroxychloroquine Sulphate. Br Med J. 1974. doi:10.1136/bmj.3.5923.94
- Pilcher DB. Hydroxychloroquine sulfate in prevention of thromboembolic phenomena in surgical patients. Am Surg. 1975.
- Rand J, Wu X-X, Ashton A. Hydroxychloroquine Protects the Annexin A5 Anticoagulant Shield from Disruption by Antiphospholipid Antibodies. Blood. 2008. doi:10.1182/blood.v112.11.404.404
- 24. Bertrand E, Cloitre B, Ticolat R, Kouad Bile R, Gautier C, Obou Abiyou G, et al. Antiaggregation of chloroquine. Med Trop. 1990.
- Raines MF, Bhargava SK, Rosen ES. The blood-retinal barrier in chloroquine retinopathy. Investig Ophthalmol Vis Sci. 1989.
- 26. Yam JCS, Kwok AKH. Ocular toxicity of hydroxychloroquine. Hong Kong Medical Journal. 2006.
- Chen CY, Wang FL, Lin CC. Chronic hydroxychloroquine use associated with QT prolongation and refractory ventricular arrhythmia. Clin Toxicol. 2006. doi:10.1080/15563650500514558
- 28. A. N, A. T, E. W. Long QT and hydroxychloroquine; a poorly recognised problem in rheumatology patients. Arthritis and Rheumatism. 2013.
- Jeevaratnam K. Chloroquine and hydroxychloroquine for COVID-19: implications for cardiac safety. Eur Hear journal Cardiovasc Pharmacother. 2020. doi:10.1093/ehjcvp/pvaa041
- Riva JJ, Malik KMP, Burnie SJ, Endicott AR, Busse JW. What is your research question? An introduction to the PICOT format for clinicians. J Can Chiropr Assoc. 2012.
- L. S, D. M, M. C, D. G, A. L, M. P, et al. Preferred reporting items for systematic review and meta-analysis protocols (prisma-p) 2015: Elaboration and explanation. BMJ (Online). 2015.
- Bordage G, Caelleigh AS, Steinecke A, Bland CJ, Crandall SJ, McGaghie WC, et al. Joint task force of academic medicine and the GEA-RIME committee. Academic Medicine. 2001.

- 33. Tang W, Cao Z, Han M, Wang Z, Chen J, Sun W, et al. Hydroxychloroquine in patients with COVID-19: an open-label, randomized, controlled trial. medRxiv. 2020. doi:10.1101/2020.04.10.20060558
- 34. Borba MGS, Val FFA, Sampaio VS, Alexandre MAA, Melo GC, Brito M, et al. Effect of High vs Low Doses of Chloroquine Diphosphate as Adjunctive Therapy for Patients Hospitalized With Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) Infection. JAMA Netw Open. 2020. doi:10.1001/jamanetworkopen.2020.8857
- Gautret P, Lagier J-C, Parola P, Hoang VT, Meddeb L, Mailhe M, et al. Hydroxychloroquine and azithromycin as a treatment of COVID-19: results of an open-label non-randomized clinical trial. Int J Antimicrob Agents. 2020. doi:10.1016/j.ijantimicag.2020.105949
- Huang M, Tang T, Pang P, Li M, Ma R, Lu J, et al. Treating COVID-19 with Chloroquine. J Mol Cell Biol. 2020. doi:10.1093/jmcb/mjaa014
- Mehra MR, Desai SS, Ruschitzka F, Patel AN. Hydroxychloroquine or chloroquine with or without a macrolide for treatment of COVID-19: a multinational registry analysis. Lancet. 2020.
- Singh S, Khan A, Chowdhry M, Chatterjee A. Outcomes of Hydroxychloroquine Treatment Among Hospitalized COVID-19 Patients in the United States-Real-World Evidence From a Federated Electronic Medical Record Network. medRxiv. 2020.
- Ip A, Berry DA, Hansen E, Goy AH, Pecora AL, Sinclaire BA, et al. Hydroxychloroquine and Tocilizumab Therapy in COVID-19 Patients - An Observational Study. medRxiv. 2020; 2020.05.21.20109207. doi:10.1101/2020.05.21.20109207
- 40. Rosenberg ES, Dufort EM, Udo T, Wilberschied LA, Kumar J, Tesoriero J, et al. Association of treatment with hydroxychloroquine or azithromycin with in-hospital mortality in patients with COVID-19 in New York state. JAMA. 2020.
- 41. Geleris J, Sun Y, Platt J, Zucker J, Baldwin M, Hripcsak G, et al. Observational study of hydroxychloroquine in hospitalized patients with Covid-19. N Engl J Med. 2020.
- 42. Million M, Lagier J-C, Gautret P, Colson P, Fournier P-E, Amrane S, et al. Full-length title: Early treatment of COVID-19 patients with hydroxychloroquine and azithromycin: A retrospective analysis of 1061 cases in Marseille, France.

Travel Med Infect Dis. 2020; 101738.

- 43. Yu B, Li C, Chen P, Zhou N, Wang L, Li J, et al. Low dose of hydroxychloroquine reduces fatality of critically ill patients with COVID-19. Sci China Life Sci. 2020; 1.
- 44. Magagnoli J, Narendran S, Pereira F, Cummings T, Hardin JW, Sutton SS, et al. Outcomes of hydroxychloroquine usage in United States veterans hospitalized with Covid-19. medRxiv. 2020. doi:10.1101/2020.04.16.20065920
- 45. Kim MS, Jang S-W, Park Y-K, Kim B, Hwang T-H, Kang SH, et al. Treatment Response to Hydroxychloroquine, Lopinavir/Ritonavir, and Antibiotics for Moderate COVID 19: A First Report on the Pharmacological Outcomes from South Korea. medRxiv. 2020.
- Saleh M, Gabriels J, Chang D, Kim BS, Mansoor A, Mahmood E, et al. The Effect of Chloroquine, Hydroxychloroquine and Azithromycin on the Corrected QT Interval in Patients with SARS-CoV-2 Infection. Circ Arrhythmia Electrophysiol. 2020. doi:10.1161/CIRCEP.120.008662
- 47. Mahévas M, Tran V-T, Roumier M, Chabrol A, Paule R, Guillaud C, et al. Clinical efficacy of hydroxychloroquine in patients with covid-19 pneumonia who require oxygen: observational comparative study using routine care data. bmj. 2020;369.
- Ramireddy A, Chugh HS, Reinier K, Ebinger J, Park E, Thompson M, et al. Experience with Hydroxychloroquine and Azithromycin in the COVID-19 Pandemic: Implications for QT Interval Monitoring. medRxiv. 2020. doi:10.1101/2020.04.22.20075671
- 49. Mercuro NJ, Yen CF, Shim DJ, Maher TR, McCoy CM, Zimetbaum PJ, et al. Risk of QT interval prolongation associated with use of hydroxychloroquine with or without concomitant azithromycin among hospitalized patients testing positive for coronavirus disease 2019 (COVID-19). JAMA Cardiol. 2020.
- Chorin E, Dai M, Shulman E, Wadhwani L, Cohen RB, Barbhaiya C, et al. The QT Interval in Patients with SARS-CoV-2 Infection Treated with Hydroxychloroquine/Azithromycin. medRxiv. 2020. doi:10.1101/2020.04.02.20047050
- 51. Mallat J, Hamed F, Balkis M, Mohamed MA, Mooty M, Malik A, et al. Hydroxychloroquine is associated with slower viral clearance in clinical COVID-19 patients with mild to moderate disease: A retrospective study. medRxiv. 2020.

- Vincent MJ, Bergeron E, Benjannet S, Erickson BR, Rollin PE, Ksiazek TG, et al. Chloroquine is a potent inhibitor of SARS coronavirus infection and spread. Virol J. 2005. doi:10.1186/1743-422X-2-69
- Kumar S, Maurya VK, Prasad AK, Bhatt MLB, Saxena SK. Structural, glycosylation and antigenic variation between 2019 novel coronavirus (2019-nCoV) and SARS coronavirus (SARS-CoV). VirusDisease. 2020. doi:10.1007/s13337-020-00571-5
- 54. Kwiek JJ, Haystead TAJ, Rudolph J. Kinetic Mechanism of Quinone Oxidoreductase 2 and Its Inhibition by the Antimalarial Quinolines. Biochemistry. 2004. doi:10.1021/bi035923w
- 55. Wolfram J, Nizzero S, Liu H, Li F, Zhang G, Li Z, et al. A chloroquine-induced macrophage-preconditioning strategy for improved nanodelivery. Sci Rep. 2017. doi:10.1038/s41598-017-14221-2
- 56. Liu J, Cao R, Xu M, Wang X, Zhang H, Hu H, et al. Hydroxychloroquine, a less toxic derivative of chloroquine, is effective in inhibiting SARS-CoV-2 infection in vitro. Cell Discovery. 2020. doi:10.1038/s41421-020-0156-0
- 57. Wang H, Yang P, Liu K, Guo F, Zhang Y, Zhang G, et al. SARS coronavirus entry into host cells through a novel clathrin- and caveolae-independent endocytic pathway. Cell Res. 2008. doi:10.1038/cr.2008.15
- Liu W, Li H. COVID-19: Attacks the 1-Beta Chain of Hemoglobin and Captures the Porphyrin to Inhibit Human Heme Metabolism. ChemRxiv. 2020. doi:10.26434/chemrxiv.11938173.v6
- Yao X, Ye F, Zhang M, Cui C, Huang B, Niu P, et al. In Vitro Antiviral Activity and Projection of Optimized Dosing Design of Hydroxychloroquine for the Treatment of Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2). Clin Infect Dis. 2020. doi:10.1093/cid/ciaa237
- He X, Lau EHY, Wu P, Deng X, Wang J, Hao X, et al. Temporal dynamics in viral shedding and transmissibility of COVID-19. Nat Med. 2020. doi:10.1038/s41591-020-0869-5
- Cheng VCC, Hung IFN, Tang BSF, Chu CM, Wong MML, Chan KH, et al. Viral Replication in the Nasopharynx Is Associated with Diarrhea in Patients with Severe Acute Respiratory Syndrome. Clin Infect Dis. 2004. doi:10.1086/382681
- 62. Yang Y, Shen C, Li J, Yuan J, Yang M, Wang F, et al. Exuberant elevation of IP-10, MCP-3 and IL-1ra during SARS-

CoV-2 infection is associated with disease severity and fatal outcome. medRxiv. 2020. doi:10.1101/2020.03.02.20029975

- 63. Hadjadj J, Yatim N, Barnabei L, Corneau A, Boussier J, Pere H, et al. Impaired type I interferon activity and exacerbated inflammatory responses in severe Covid-19 patients. medRxiv. 2020. doi:10.1101/2020.04.19.20068015
- 64. Zhou F, Yu T, Du R, Fan G, Liu Y, Liu Z, et al. Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. Lancet. 2020. doi:10.1016/S0140-6736(20)30566-3
- Saitoh SI, Miyake K. Regulatory molecules required for nucleotide-sensing Toll-like receptors. Immunological Reviews. 2009. doi:10.1111/j.1600-065X.2008.00729.x
- 66. Lim EJ, Lee SH, Lee JG, Chin BR, Bae YS, Kim JR, et al. Activation of toll-like receptor-9 induces matrix metalloproteinase-9 expression through Akt and tumor necrosis factor-α signaling. FEBS Lett. 2006. doi:10.1016/j.febslet.2006.06.100
- 67. Roche PA, Furuta K. The ins and outs of MHC class II-mediated antigen processing and presentation. Nature Reviews Immunology. 2015. doi:10.1038/nri3818
- Guerriero JL. Macrophages: Their Untold Story in T Cell Activation and Function. Int Rev Cell Mol Biol. 2019. doi:10.1016/bs.ircmb.2018.07.001
- Nosál' R, Jančinová V. Cationic amphiphilic drugs and platelet phospholipase A2 (cPLA2). Thromb Res. 2002. doi:10.1016/S0049-3848(02)00036-1
- Goldman FD, Gilman AL, Hollenback C, Kato RM, Premack BA, Rawlings DJ. Hydroxychloroquine inhibits calcium signals in T cells: A new mechanism to explain its immunomodulatory properties. Blood. 2000. doi:10.1182/blood.v95.11.3460.011k26_3460_3466
- 71. Chen G, Wu D, Guo W, Cao Y, Huang D, Wang H, et al. Clinical and immunologic features in severe and moderate Coronavirus Disease 2019. J Clin Invest. 2020. doi:10.1172/JCI137244
- 72. Conti P, Gallenga CE, Tetè G, Caraffa A, Ronconi G, Younes A, et al. How to reduce the likelihood of coronavirus-19 (CoV-19 or SARS-CoV-2) infection and lung inflammation mediated by IL-1. Journal of biological regulators and

homeostatic agents. 2020. doi:10.23812/Editorial-Conti-2

- 73. Wang SF, Tseng SP, Yen CH, Yang JY, Tsao CH, Shen CW, et al. Antibody-dependent SARS coronavirus infection is mediated by antibodies against spike proteins. Biochem Biophys Res Commun. 2014. doi:10.1016/j.bbrc.2014.07.090
- 74. Mahallawi WH, Khabour OF, Zhang Q, Makhdoum HM, Suliman BA. MERS-CoV infection in humans is associated with a pro-inflammatory Th1 and Th17 cytokine profile. Cytokine. 2018. doi:10.1016/j.cyto.2018.01.025
- Malaviya R, Laskin JD, Laskin DL. Anti-TNFα therapy in inflammatory lung diseases. Pharmacology and Therapeutics.
 2017. doi:10.1016/j.pharmthera.2017.06.008
- 76. Kono M, Tatsumi K, Imai AM, Saito K, Kuriyama T, Shirasawa H. Inhibition of human coronavirus 229E infection in human epithelial lung cells (L132) by chloroquine: Involvement of p38 MAPK and ERK. Antiviral Res. 2008. doi:10.1016/j.antiviral.2007.10.011
- Hulme OJ, Wagenmakers E-J, Damkier P, Madelung CF, Siebner HR, Helweg-Larsen J, et al. Reply to Gautret et al.
 2020: A Bayesian reanalysis of the effects of hydroxychloroquine and azithromycin on viral carriage in patients with
 COVID-19. medRxiv. 2020. doi:10.1101/2020.03.31.20048777
- 78. Khobragade SB, Gupta P, Gurav P, Chaudhari G, Gatne MM, Shingatgeri VM. Assessment of proarrhythmic activity of chloroquine in in vivo and ex vivo rabbit models. J Pharmacol Pharmacother. 2013. doi:10.4103/0976-500X.110892
- 79. Noble D. A modification of the Hodgkin—Huxley equations applicable to Purkinje fibre action and pacemaker potentials. J Physiol. 1962. doi:10.1113/jphysiol.1962.sp006849
- Sánchez-Chapula JA, Salinas-Stefanon E, Torres-Jácome J, Benavides-Haro DE, Navarro-Polanco RA. Blockade of currents by the antimalarial drug chloroquine in feline ventricular myocytes. J Pharmacol Exp Ther. 2001.
- Hama R, Bennett CL. The mechanisms of sudden-onset type adverse reactions to oseltamivir. Acta Neurologica Scandinavica. 2017. doi:10.1111/ane.12629
- Reardon M, Malik M. QT interval change with age in an overtly healthy older population. Clin Cardiol. 1996. doi:10.1002/clc.4960191209

- Al-Khatib SM, Allen LaPointe NM, Kramer JM, Califf RM. What Clinicians Should Know about the QT Interval. Journal of the American Medical Association. 2003. doi:10.1001/jama.289.16.2120
- 84. Giunti S, Amione C, Gruden G, Ghezzo G, Fornengo P, Cavallo-Perin P, et al. Increased QT nterval dispersion predicts 15-year cardiovascular mortality in type 2 diabetic subjects: The population-based Casale Monferrato Study. Diabetes Care. 2012. doi:10.2337/dc11-1397
- The epidemiological characteristics of an outbreak of 2019 novel coronavirus diseases (COVID-19) in China.
 Zhonghua Liu Xing Bing Xue Za Zhi. 2020. doi:10.3760/cma.j.issn.0254-6450.2020.02.003
- Richardson S, Hirsch JS, Narasimhan M, Crawford JM, McGinn T, Davidson KW, et al. Presenting Characteristics, Comorbidities, and Outcomes among 5700 Patients Hospitalized with COVID-19 in the New York City Area. JAMA - J Am Med Assoc. 2020. doi:10.1001/jama.2020.6775
- Henderson M, Page L. Appraising the evidence: What is selection bias? Evidence-Based Mental Health. 2007. doi:10.1136/ebmh.10.3.67
- Mlinarić A, Horvat M, Smolčić VŠ. Dealing with the positive publication bias: Why you should really publish your negative results. Biochemia Medica. 2017. doi:10.11613/BM.2017.030201
- Kupferschmidt K. WHO launches global megatrial of the four most promising coronavirus treatments. Science (80-).
 2020. doi:10.1126/science.abb8497
- Piller C. Former FDA leaders decry emergency authorization of malaria drugs for coronavirus. Science (80-). 2020. doi:10.1126/science.abc1337

Figure Legend

Fig 1: The attrition processes for publication. This comprised: (1) searching of three databases for peer-reviewed papers; (2) acquisition of submitted, but not yet peer-reviewed, items from a preprint server; and (3) application of exclusion and inclusion criteria. The removal of duplicates at each step, where necessary, left 19 unique items to pass onto the quality appraisal stage.

