#### 1 ORIGINAL ARTICLE

2

# High incidence of glucocorticoid-induced hyperglycaemia in inflammatory bowel disease; metabolic and clinical predictors identified by machine learning

#### 5 6 TITLE PAGE

## 78 Author List:

- 9 Martin McDonnell<sup>1</sup><sup>\$</sup>, Richard J Harris<sup>1</sup><sup>\$</sup>, Florina Borca<sup>1,2</sup>, Tilly Mills<sup>1</sup>, Louise Downey<sup>1</sup>, Suranga Dharmasiri<sup>1</sup>,
- 10 Mayank Patel<sup>3</sup>, Benjamin Zaré<sup>1</sup>, Matt Stammers<sup>1,2</sup>, Trevor R Smith<sup>1</sup>, Richard Felwick<sup>1</sup>, JR Fraser Cummings<sup>1,4</sup>,
- 11 Hang T.T. Phan<sup>2</sup>, Markus Gwiggner<sup>1</sup>
- <sup>1</sup>Department of Gastroenterology, University Hospital Southampton, Southampton, United Kingdom
- 13 <sup>2</sup> University of Southampton NIHR Biomedical Research Facility, Southampton, United Kingdom
- 14 <sup>3</sup> Department of Diabetes and Endocrinology, University Hospital Southampton, Southampton, United Kingdom
- <sup>4</sup>Faculty of Medicine, University of Southampton, Southampton, United Kingdom
- 16 <sup>\$</sup> Joint first authors17

#### 18 Corresponding Author:

- 19
- 20 Dr Martin McDonnell
- 21 Department of Gastroenterology
- 22 Mail point 190
- 23 University Hospital Southampton
- 24 Tremona Road
- 25 Southampton
- 26 SO16 6YD
- 27
- 28 <u>m.j.mcdonnell@soton.ac.uk</u>
- 29
- 30

#### 31 ABSTRACT

32

#### 33 Background

- 34 Glucocorticosteroids (GC) are long-established, widely used agents for induction of remission in inflammatory
- 35 bowel disease (IBD). Hyperglycaemia is a known complication of GC treatment with implications for morbidity and
- 36 mortality. Published data on prevalence and risk factors for GC-induced hyperglycaemia in the IBD population are
- 37 limited. We prospectively characterise this complication in our cohort, employing machine-learning methods to
- 38 identify key predictors of risk.

## 39

## 40 Methods

- 41 We conducted a prospective observational study of IBD patients receiving intravenous hydrocortisone (IVH).
- 42 Electronically triggered three times daily capillary blood glucose (CBG) monitoring was recorded alongside diabetes
- 43 mellitus (DM) history, IBD biomarkers, nutritional and IBD clinical activity scores. Hyperglycaemia was defined as
- $\label{eq:cbg_eq} 44 \qquad CBG \geq 11 \cdot 1 \text{mmol/L and undiagnosed DM as HbA1c} \geq 48 \text{ mmol/mol. Random Forest regression models were used to}$
- 45 extract predictor-patterns present within the dataset.
- 46

### 47 Findings

- 48 94 consecutive IBD patients treated with IVH were included. 60% (56/94) of the cohort recorded an episode of
- 49 hyperglycaemia, including 57% (50/88) of those with no prior history of DM, of which 19% (17/88) and 5% (4/88)
- 50 recorded a CBG  $\geq$  14mmol/L and  $\geq$  20mmol/L, respectively. The Random Forest models identified increased CRP
- 51 followed by a longer IBD duration as leading risk predictors for significant hyperglycaemia.
- 52

#### 53 Interpretation

- 54 Hyperglycaemia is common in IBD patients treated with intravenous GC, therefore CBG monitoring should be
- 55 included in routine clinical practice. Machine learning methods can identify key risk factors for clinical
- 56 complications. Physicians should consider steroid-sparing strategies in high-risk patients such as those with high
- 57 admission CRP or a longer IBD duration. There is an emergent case for research to explore steroid-free treatment
- **58** regimens for hospitalised patients with severe IBD flares.
- 59

#### 60 Funding

61 No specific funding was received for this study.

#### 62 Evidence before this study

63 Glucocorticosteroids (GC) are long-established induction agents in the management of inflammatory bowel disease

- 64 (IBD). They are recommended first-line therapy in consensus guidelines and prescribing remains widespread, with
- an estimated 30% of IBD patients exposed annually. Hyperglycaemia is a known complication of GC and has been
- 66 linked to increased length of hospital stay, morbidity and mortality. Small case series of GC treated medical patients
- 67 suggest a higher risk of hyperglycaemia in the hospitalised population but have suffered from a lack of systematic
- 68 blood glucose monitoring.
- 69

#### 70 Added value of this study

- 71 This is the first study utilising prospective, systematic monitoring of capillary blood glucose (CBG) to determine the
- 72 frequency of hyperglycaemia in a GC-treated hospitalised IBD population. We report that more than half of IBD
- 73patients without prior diabetes mellitus treated with intravenous hydrocortisone (IVH), will develop hyperglycaemia74(CBG  $\geq$ 11·1mmol/L). Random Forest regressors pinpointed CRP and IBD duration as the strongest predictor of this75adverse outcome.
- 76

## 77 Implications of all the available evidence

- 78 Hyperglycaemia is a common complication of IVH therapy in hospitalised IBD patients, particularly in those with
- 79 high inflammatory burden. The monitoring and management of this complication, which has potential implications
- 80 for the morbidity, mortality and subsequent risk of diabetes diagnosis should become part of routine clinical
- 81 practice.
- 82 83

## 84 Contribution statement:

85 MM, RJH, SD and MG were responsible for the original concept and planning of the study. MM, RJH, TM, SD, BZ

and LD were responsible for clinical data collection and analysis. FB and HP were responsible for data extraction,

analysis and modelling. RJH and MM contributed equally to this work and drafted the manuscript, which RF, TS,

- 88 MP, JRFC, HP, MS and MG critically reviewed and revised.
- 89
- 90

- 92
- 93

#### 94 INTRODUCTION

95

96 In the 1950's Truelove and Witts performed the first randomised controlled trial of glucocorticosteroids (GC) in

- active ulcerative colitis (UC), demonstrating a reduction in mortality from 10.9% to 4.6% after two months of
- 98 treatment.<sup>1</sup> 70 years later, GC remain a common treatment for inflammatory bowel disease (IBD); a recent UK
- 99 multi-centre study identified 30% of patients were exposed to GC in the preceding 12 months.<sup>2</sup> GC are first-line
- 100 induction agents for moderate to severe Crohn's Disease (CD) and UC in consensus guidelines and intravenous GC
- are widely used as the initial medical therapy for IBD patients admitted to hospital with severe disease flares.<sup>3,4</sup>
   102
- 103 The mechanism of action of GC is mediated via the cytosolic glucocorticoid receptor (cGR). After GC binding,
- 104 cGR-associated proteins dissociate, and the GC-cGR complex translocates to the nucleus. The nuclear GC-cGR
- 105 complex binds to DNA, upregulating the transcription of anti-inflammatory cytokines (e.g. IL-10, annexin 1) and
- downregulating transcription of pro-inflammatory mediators (e.g. IL-1 $\alpha$ , IL-1 $\beta$ , IL-8, TNF $\alpha$ ). More rapid, non-
- 107 genomic, effects occur via both the cGR dissociated proteins and likely via other novel transmembrane receptors.<sup>5</sup>
- 108 Steroid-induced hyperglycaemia is mediated by decreased  $\beta$ -cell insulin production, increased insulin resistance and 109 increased gluconeogenesis.<sup>6</sup>
- 110

111 Studies in other GC-treated populations demonstrate both short and long-term severe adverse effects including

112 ocular, musculoskeletal, cardiovascular, metabolic and infectious complications.<sup>7</sup> In addition, de-novo

113 hyperglycaemia and deterioration of glycaemic control in those with pre-existing diabetes mellitus (DM) have been

recognised adverse effects of GC since the 1950s.<sup>8</sup> The specific risk in IBD hospitalised patients with severe IBD

- 115 flares, however, is not known.
- 116

117 Hyperglycaemia has been associated with increased length of stay, morbidity and mortality in non-critical care

118 medical and surgical patients.<sup>9</sup> At least 67 cases of GC induced metabolic emergencies including ketoacidosis and

119 hyperosmolar hyperglycaemic states have been reported, 11 of which were fatal.<sup>10</sup> Following 2 cases of GC-related

- 120 diabetic crisis at our institution in IBD patients with no prior diabetes, electronically triggered systematic capillary
- 121 blood glucose (CBG) monitoring was introduced for all prescribed intravenous hydrocortisone (IVH). This allowed
- the systematic study of hyperglycaemia in hospitalised IBD patients receiving IVH.
- 124 MATERIALS AND METHODS
- 125

We conducted a prospective observational study of consecutive admissions of IBD patients treated with IVH
 between October 2017 and December 2018. Included subjects had a confirmed diagnosis of IBD (either historic or

- 128 during admission episode) and received at least four doses of 100mg IVH over 24 hours.
- 129

130 Subjects had three times daily CBG monitoring automatically triggered by electronic prescription of IVH. CBG

131 results were automatically digitally captured in the electronic patient record. Predictor variables included admission

132 IBD severity score (Harvey Bradshaw Index or partial Mayo Score), serum biochemistry, HbA1c, faecal

133 calprotectin, C-reactive protein, potential DM risk factors (BMI, family history, concomitant medications), self-

134 reported weight loss and MUST (Malnutrition Universal Screening Tool) score.<sup>11</sup>

135

**136** Patients developing a CBG  $\geq$  14mmol/l were referred to the diabetes team for consideration of treatment.

137 Undiagnosed DM was excluded by baseline HbA1c <48 mmol/mol or a pre-IVH CBG <11.1 mmol/L.<sup>12</sup> Biochemical

- follow up was performed at one year to assess repeat HbA1c measurements and serum and faecal biomarker
- assessments.
- 140
- 141 Laboratory values outside the quantification limit were substituted with the upper/lower limit value. Categorical and
- 142 continuous predictors were handled and formatted appropriately. Data were analysed using Python v3.7.4 and R

- 143 v3.6.1. Descriptive statistics were applied to cohorts, and Pearson correlations between independent and dependent
- variables were ascertained with matplotlib/seaborn heat-maps. Random Forest (RF) machine learning models using
- 145 the scikit-learn package v0.22.1 were constructed to regress the phenotypic and admission clinical predictors.<sup>13,14</sup>
- Pairwise correlations between input features (see supplementary data) were identified to eliminate similar features.
- 147 Then multivariable RF models were constructed and optimised using random grid search. Subsequently 5-fold cross-
- validation with a train-test split was used to test effectiveness at predicting glucose levels in the internal holdout set.
- 149 Mean squared error was the reported metric.
- 150

152

151 The protocol was reviewed and approved by the UK Health Research Authority - reference 19/HRA/0033.

#### 153 **RESULTS**

154

156

## 155 Demographics

157 94 IBD inpatient episodes met the inclusion criteria; 54 for CD, 36 for UC and 4 for IBD-as yet unclassified

(IBDU). The median length of IVH treatment was four days (range 1-11).

159

160 5/94 episodes were for patients with a prior diagnosis of Type 2 DM. One further subject had a pre-GC treatment

161 CBG of 13.9mmol/L and was thus reclassified as undiagnosed Type 2 DM.

162

		CD	IBDU	UC
	n	54	4	36
Age		42.2 (37.9 - 46.5)	49.8 (28.4 - 71.1)	43.1 (37.0 - 49.2)
Gender	Female	25 (46.3)	2 (50.0)	19 (52.8)
BMI	<18.5	6 (11.1)	1 (25.0)	3 (8.3)
	18.5 - 25	28 (51.9)	2 (50)	16 (44.4)
	25-30	13 (24.1)	1 (25)	14 (38.9)
	>30	7 (13.0)	0 (0.0)	3 (8.3)
MUST Score	0	17 (31.5)	1 (25.0)	17 (48.6)
	1	13 (24.1)	1 (25.0)	10 (28.6)
	2	13 (24.1)	2 (50.0)	4 (11.4)
	≥3	11 (20.4)	0 (0.0)	4 (11.4)
Length of diagnosis (years)		7.3 (4.6 - 10.0)	3.3 (0 - 8.4)	5.0 (3.2 - 6.8)
History of Diabetes	No	49 (90.7)	4 (100.0)	35 (97.2)
	Type 2 DM	5 (9.3)	0 (0.0)	1 (2.8)
Admission HBA1C		38.5 (35.5 - 41.5)	37.0 (33.1 - 40.8)	37.7 (35.8 - 39.6)
Admission Calprotectin		2854 (1980 - 2298)	2491 (NA)	3159 (2302 - 4071)
Admission CRP		75.9 (50.5 - 101.3)	138.0 (106.7 - 169.3)	88.9 (53.6 - 124.1)
PMS			5.50 (4.5 - 6.5)	7.78 (7.2 - 8.3)
HBI		14.7 (13.0-16.5)		
Current Treatment	Oral steroids	8 (14.8)	1 (25)	6 (16.7)
	Oral 5ASA	2 (3.7)	2 (50)	20 (55.6)
	Immunomodulator	18 (33.3)	0 (0)	6 (16.7)
	Anti-TNF	10 (19.2)	0 (0)	10 (29.4)
	Vedolizumab	2 (3.8)	0 (0)	0 (0.0)
	Ustekinumab	7 (13.5)	0 (0)	0 (0.0)
Smoking	Current	12 (23.5)	0 (0)	0 (0.0)
	Ex-smoker	9 (17.7)	1 (25.0)	13 (36.1)
	Never smoked	30 (58.8)	3 (75.0)	23 (63.9)
Table 1: Demographic	cs			

<sup>163</sup> 164

165 For categorical variables n (%) shown, for continuous variables mean (95% Confidence Interval) shown.

166 Abbreviations: '..' = not applicable, MUST (Malnutrition Universal Screening Tool), BMI (Body Mass Index), PMS

167 (Partial Mayo Score), HBI (Harvey Bradshaw Index), 5-ASA (5-aminosalicylic acid), Anti-TNF (anti-Tumour

168 Necrosis Factor), Immunomodulator (azathioprine, mercaptopurine or methotrexate).

171	Incidence and management of hyperglycaemia
170	
169	

173 The overall global incidence of hyperglycaemia in our cohort of hospitalised IBD patients receiving IVH was 60%
174 (56/94), with intervention for diabetes initiated in 20% (11/56) of these patients. For those without a prior diagnosis

of DM, 57% (50/88) developed diabetic-range CBG ( $\geq$ 11·1mmol/L) while 19% (17/88) and 5% (4/88) developed a CBG >14mmol/L and >20mmol/L, respectively. 25/50 of those with de-novo hyperglycaemia had CD, 22/50 UC,

and 3/50 IBDU with a combined mean age of 44.8 years (Standard Deviation 17.4). In patients without a prior

diagnosis of DM 14% (7/50) had diabetic interventions; 6/50 started an oral hypoglycaemic and 1/50 commenced
 insulin therapy.

180

172

All six patients with prior DM (5 CD and 1 UC) had an episode of hyperglycaemia and 4/6 recorded a CBG
 ≥20mmol/L. 50% (3/6) were started on insulin, and 17% (1/6) commenced oral hypoglycaemic therapy.

183

184 Of the 50 admissions in patients without prior DM recording a CBG  $\geq 11.1$  mmol/L, 22% (11/50) were preceded

directly by a course of oral steroids compared to 8% (3/38) without hyperglycaemia (p = 0.09, Fisher's exact test).

186 In 60% (30/50) peak CBG occurred on day two of IVH treatment, but in 14% (7/50) of cases peak CBG was not

187 reached until seven days after the first dose.

188

189 During index admission, 54% (30/56) of hyperglycaemic patients and 50% (19/38) of normoglycaemic patients were

190 started on biologic therapy. Four patients underwent emergency inpatient colectomy, all of whom had recorded a

191 hyperglycaemic episode. Mean ( $\pm$  95% Confidence Interval) length of stay was 12.6 ( $\pm$  3.0) days for

hyperglycaemic patients and  $9.1 (\pm 2.0)$  days for normoglycaemic patients (p = 0.053, t-test).

193

194 Physicians' decision making was influenced by steroid-induced hyperglycaemia, particularly in those with a

persistent CBG  $\geq$ 14mmol/L (trigger for diabetes specialist review). Conventional post-IVH eight-week tapering

course of oral prednisolone prescribing differed. Only 30% of patients with CBG  $\geq$ 14mmol/L received oral

197 prednisolone without intervention (e.g. diabetic medication or rapid steroid taper) versus 58% of those with

198 normoglycemia.199



200201 Figure 1: Maximum recorded CBG for each admission plotted in ascending order

#### 202 Predictors of hyperglycaemia

203 Maximum CBG was positively correlated with CRP, platelet count, and HbA1c and negatively correlated with

haemoglobin (see correlation heat-map in supplementary data). For non-diabetic patients, the RF model was fit 500

times with a maximum depth of 90 trees using a random grid search method. CRP was the most critical predictor

identified, followed by disease duration, platelet count, admission haemoglobin and HbA1C. The final mean squarederror of the optimised non-diabetic patient model was 1.876.

208

209 The variables and their respective importance of contribution to the RF modelling are shown in figure 2.



Relative importance
 Figure 2: Relative variable importance for Random Forest modelling of maximum CBG (subjects with history of DM excluded)

214 RF models optimised by grid search will rank the relative importance of each variable against the most critical

variable (in this case, CRP). This provides a sense of how much each variable contributes to the predictive power ofthe overall model. In this instance, the length of IVH usage and metabolic variables are not as crucial to predicting

217 hyperglycaemia as blood markers of inflammation.

Once the optimised RF model had been developed, blood glucose predictions were made on the holdout set using 5 fold cross validation to reduce overfitting.

221 222





Figure 3: Predicted highest CBG vs measured highest CBG using model (subjects with history of DM excluded)

Figure 3 reveals that the test-set predictions were comparable to the training set predictions. This pattern was present across all folds of the cross-validation; this graph is given only as an example of the model's performance.

228

229 When the RFR model included the six patients with DM, the leading predictor of CBG elevation became HbA1c.

230 Due to high CBG elevation ( $\geq$ 20mmol/L) in subjects with known diabetes, particularly those with evidence of

inadequate control by admission HbA1c, the RFR modelling shown above excluded these patients. (See

- supplementary data) for the variable importance, correlation matrix and model with the patients with previous
- diabetes included. Their inclusion increased the mean squared error to 2.158, which is not surprising given thediffering pre-admission characteristics in this sub-population.
- 235

## Additional reported side effects and metabolic instability237

32% (30/94) of episodes included a patient-reported symptom of anxiety or mood change, 11% (10/94) of poor sleep
and 10% (9/94) a report of appetite change. Electrolyte depletion, like hyperglycaemia, is a marker of metabolic and
nutritional dysregulation and was common in the whole cohort with no statistically significant differences between
hyperglycaemic and normoglycemic patients. 62% (58/94) recorded a serum potassium below the lower limit of our
laboratory reference range (<3.5mmol/L), 71% (67/94) low serum magnesium (<0.7mmol/l) and 84% (79/94) low</li>
serum phosphate (<0.8mmol/L). See supplementary data.</li>

244

66% (37/56) of hyperglycaemic patients and 57% (21/38) of normoglycaemic patients reported weight loss in the six
months before admission. All patients were routinely assessed by a specialist dietitian who prescribed nutritional
support as indicated; 53/94 had oral nutritional supplements, 5/94 nasogastric feeding and 5/94 parenteral nutrition.
There was no significant difference in the proportion of hyperglycaemic versus normoglycaemic patients receiving
nutritional therapy.

## 251 One-year follow-up

252

250

There was a significant decrease in faecal calprotectin over the 12-month follow-up. For those without
hyperglycaemia during the acute episode, there was a trend to lower mean calprotectin at 12 months, 210 (± 707)
versus 738 (± 640 p=0·18 t-test). In non-diabetic patients with raised calprotectin at admission and paired results
35% (6/17) normalised by 12 months in those who had hyperglycaemia and 75% (9/12) normalised in those without
hyperglycaemia p=0·06 (Fisher's exact test). There was no significant change in HbA1c over the one-year follow-up
for any groups.

## 260 **DISCUSSION**

261

262 Our data demonstrate that hyperglycaemia is common in IBD patients treated with IVH. By instituting electronically 263 triggered regular CBG monitoring of IBD patients treated with IVH, we have uncovered the high prevalence of this 264 relevant complication. Our monitoring regime identified patients who required diabetic intervention or modification 265 of their treatment in response to hyperglycaemia. The void of data regarding steroid induced diabetes in IBD 266 suggests that in current practice it may often be under-recognised, and thus under-treated, potentially exposing 267 patients to the risk of complications such as increased length of stay and infection. Hyperglycaemia has also been 268 mechanistically linked to intestinal barrier dysfunction in both murine models and non-IBD human subjects.<sup>15</sup> 269 Through this mechanism, it is at least theoretically possible to link hyperglycaemia directly to intestinal 270 inflammation and IBD outcomes.<sup>16</sup>

- 271
- 272 Until now there has been no accurate prospective evidence of the extent and severity of IVH-induced
- 273 hyperglycaemia in IBD. A previous retrospective study of steroid side effects in elderly Crohn's disease patients
- estimated an incidence of just 17%, but did not have systematic CBG monitoring.<sup>17</sup> Likewise, studies of GC-treated

- 275 rheumatology patients report the occurrence of hyperglycaemia at 9 42%.<sup>18-20</sup> A potential explanation for the
- higher frequency seen here is that our cohort only included hospitalised patients, who were both monitored
- 277 frequently and likely to have had more significant systemic inflammation. The incidence we report is similar to that
- 278 in a smaller study of a more elderly, heterogeneous group of hospitalised patients, prescribed GC.<sup>21</sup>
- 279

Machine learning techniques can be applied to clinical problems for risk prediction. We chose RF as it is a powerful
 ensemble machine learning method for predictive modelling, known to perform better than other methods on smaller
 datasets. Random Forests work by constructing a large number of smaller decision trees and averaging the output of
 individual trees and are capable of capturing complex dependency patterns among multi-variate input features.

284

The RF model accurately predicted the degree of hyperglycaemia for those without a history of DM. Principal independent determinants were admission CRP, length of disease followed by thrombocytosis and anaemia, suggesting that systemic inflammatory burden and duration of IBD determines the risk of clinically significant hyperglycaemia in IVH-treated IBD patients. No data of historic GC prescriptions were available so we cannot speculate further on whether the relationship with disease duration is a consequence of past GC exposure or a consequence of the disease process itself.

291

By using RF for early identification of those most at risk of significant GC-induced hyperglycaemia, alternative
 treatment strategies can be considered, including a rapid steroid taper, early use of biologics and avoidance of oral
 steroids in those demonstrating adequate response to biologic induction. Awareness of significant steroid-induced
 diabetes in this cohort appeared to influence physicians' decisions with an observed reduction in follow-on
 prescriptions for conventional steroid taper.

297

Although hyperglycaemia appears to resolve on stopping GC, this is not always the case. A large population-based study associated previous oral GC with an increased incidence of future DM and in rheumatic diseases cumulative GC dose has been shown to be a risk factor for DM.<sup>22,23</sup> The IBD population, regardless of GC exposure, may be at higher risk of DM. A recent Danish population-wide cohort study suggests a higher incidence of type 1 diabetes amongst subjects with IBD and a Korean cohort study found IBD patients under 40 to have an increased rate of incident diabetes diagnosis over five years, compared to age, sex and BMI-matched controls.<sup>24,25</sup>

304

305 This study has limitations. While we note that hyperglycaemia occurred during the admission of subjects treated 306 with IVH, we do not know whether another aspect of the disease or its treatments was the principal cause. It is 307 known that de-novo hyperglycaemia may also occur in hospitalised patients independently of GC as a result of the neurohormonal milieu of acute illness.<sup>26</sup> As current guidelines recommend the use of high dose GC in moderate to 308 309 severe acute IBD, there was no comparable cohort of patients managed without IVH treatment.<sup>3</sup> Despite an attempt 310 to measure baseline HbA1c in all patients starting steroids, 29% did not have this checked, and 3% had neither 311 HbA1c nor CBG pre-IVH, so some cases of undiagnosed diabetes mellitus may have been missed. Future research 312 should aim to validate the model in a wider population and determine any effect of hyperglycaemia on clinical

- 313 outcomes, which our study was not powered to achieve.
  - 314

Intravenous GC have undoubtedly saved the lives of many IBD patients. However, there is significant associated
morbidity, as acknowledged seven decades earlier by Truelove and Witts in their seminal work.<sup>1</sup> It is questionable if
GCs would be licensed in the regulatory framework currently applied to newer agents, given their side effect profile.
In the ever-evolving IBD therapeutic landscape, it may be necessary to re-evaluate the position of GCs in the
management of IBD. In UC, ciclosporin and infliximab are of equivalent efficacy as 'rescue' therapy for those
demonstrating inadequate response to IVH; whether and for how long the preceding and concomitant GC course
should be given for is not known.<sup>27</sup> The JAK inhibitor tofacitinib has demonstrated onset within three days in

- 322 moderate to severe UC and has been used effectively alongside shorter IVH courses and budesonide as induction
- 323 therapy in an acute severe UC case series.<sup>28,29</sup>

#### 324

- In acute CD the evidence for the benefit of GC to induce remission is less secure.<sup>4</sup> Nutritional therapies can if given
   as exclusive enteral nutrition, be of similar efficacy as CD induction agents to oral GC. They may offer an
   alternative treatment strategy for those in whom steroids are contraindicated.<sup>30</sup> When co-prescribed for a primary
- 328 nutritional indication in our cohort, nutritional therapies did not increase the risk of hyperglycaemia.
- 329
- 330 Our data supports the inclusion of CBG monitoring into standard clinical practice for IBD patients receiving IVH.
- 331 Machine learning identified patients with a high inflammatory disease burden and a longer disease duration as being
- 332 at the greatest risk of significant hyperglycaemia. Given the common complications of GCs and an enlarging
- 333 landscape of biological, small molecule and nutritional therapies, there is an emergent case for researching steroid-
- **334** free treatment strategies for acute IBD.
- 335
- 336
- 337

#### 338 ACKNOWLEDGEMENTS

339 We would like to thank the diabetes multidisciplinary specialist team at UHS for helping to look after these patients.

## 340 **References**

341

Truelove SC, Witts L. Cortisone in ulcerative colitis. *British medical journal* 1955;
 2(4947): 1041.

Selinger CP, Parkes GC, Bassi A, et al. A multi-centre audit of excess steroid use in
 1176 patients with inflammatory bowel disease. *Alimentary pharmacology & therapeutics* 2017;
 46(10): 964-73.

347 3. Gomollon F, Dignass A, Annese V, et al. 3rd European Evidence-based Consensus on
348 the Diagnosis and Management of Crohn's Disease 2016: Part 1: Diagnosis and Medical
349 Management. *Journal of Crohn's & colitis* 2017; **11**(1): 3-25.

4. Ford AC, Bernstein CN, Khan KJ, et al. Glucocorticosteroid therapy in inflammatory bowel disease: systematic review and meta-analysis. *Am J Gastroenterol* 2011; **106**(4): 590-9; quiz 600.

Hayashi R, Wada H, Ito K, Adcock IM. Effects of glucocorticoids on gene transcription.
 *European journal of pharmacology* 2004; **500**(1-3): 51-62.

355 6. Schäcke H, Döcke W-D, Asadullah K. Mechanisms involved in the side effects of glucocorticoids. *Pharmacology & therapeutics* 2002; **96**(1): 23-43.

Waljee AK, Rogers MA, Lin P, et al. Short term use of oral corticosteroids and related
 harms among adults in the United States: population based cohort study. *bmj* 2017; **357**: j1415.

8. Conn J, Fajans S. Influence of adrenal cortical steroids on carbohydrate metabolism in
 man. *Metabolism: clinical and experimental* 1956; 5(2): 114-27.

Umpierrez GE, Hellman R, Korytkowski MT, et al. Management of hyperglycemia in
 hospitalized patients in non-critical care setting: an endocrine society clinical practice guideline.
 *The Journal of clinical endocrinology and metabolism* 2012; **97**(1): 16-38.

Braithwaite SS. Detection and Management of Diabetes Mellitus During Glucocorticoid
 Therapy of Nonendocrine Disease. Endocrine Replacement therapy in Clinical Practice:
 Humana Press; 2003: 254-5.

367 11. Weekes CE, Elia M, Emery PW. The development, validation and reliability of a nutrition
368 screening tool based on the recommendations of the British Association for Parenteral and
369 Enteral Nutrition (BAPEN). *Clinical nutrition (Edinburgh, Scotland)* 2004; **23**(5): 1104-12.

12. International Expert Committee Report on the Role of the A1C Assay in the Diagnosis of Diabetes. *Diabetes care* 2009; **32**(7): 1327-34.

- 372 13. Breiman L. Random forests. *Machine learning* 2001; **45**(1): 5-32.
- 14. Pedregosa F, Varoquaux G, Gramfort A, et al. Scikit-learn: Machine learning in Python.
  Journal of machine learning research 2011; **12**(Oct): 2825-30.

Thaiss CA, Levy M, Grosheva I, et al. Hyperglycemia drives intestinal barrier dysfunction
and risk for enteric infection. *Science* 2018; **359**(6382): 1376-83.

16. McGuckin MA, Eri R, Simms LA, Florin TH, Radford-Smith G. Intestinal barrier

- 378 dysfunction in inflammatory bowel diseases. *Inflammatory bowel diseases* 2009; **15**(1): 100-13.
- 379 17. Akerkar GA, Peppercorn MA, Hamel MB, Parker RA. Corticosteroid-associated
- complications in elderly Crohn's disease patients. *American Journal of Gastroenterology* 1997;
   92(3).
- Ha YJ, Lee K, Jung S, Lee S, Park Y. Glucocorticoid-induced diabetes mellitus in
   patients with systemic lupus erythematosus treated with high-dose glucocorticoid therapy.
   *Lupus* 2011; **20**(10): 1027-34.
- 385 19. Panthakalam S, Bhatnagar D, Klimiuk P. The prevalence and management of

386 hyperglycaemia in patients with rheumatoid arthritis on corticosteroid therapy. Scottish medical

387 *journal* 2004; **49**(4): 139-41.

388 20. Ito S, Ogishima H, Kondo Y, et al. Early diagnosis and treatment of steroid-induced
 389 diabetes mellitus in patients with rheumatoid arthritis and other connective tissue diseases.
 390 *Modern rheumatology* 2014; **24**(1): 52-9.

391 21. Donihi AC, Raval D, Saul M, Korytkowski MT, DeVita MA. Prevalence and predictors of
392 corticosteroid-related hyperglycemia in hospitalized patients. *Endocr Pract* 2006; **12**(4): 358-62.
393 22. Gulliford MC, Charlton J, Latinovic R. Risk of diabetes associated with prescribed

394 glucocorticoids in a large population. *Diabetes care* 2006; **29**(12): 2728-9.

- 395 23. Movahedi M, Beauchamp M-E, Abrahamowicz M, et al. Risk of Incident Diabetes
  396 Mellitus Associated With the Dosage and Duration of Oral Glucocorticoid Therapy in Patients
  397 With Rheumatoid Arthritis. *Arthritis & Rheumatology* 2016; **68**(5): 1089-98.
- Halling ML, Kjeldsen J, Knudsen T, Nielsen J, Hansen LK. Patients with inflammatory
   bowel disease have increased risk of autoimmune and inflammatory diseases. *World journal of gastroenterology* 2017; 23(33): 6137-46.
- 401 25. Kang EA, Han K, Chun J, et al. Increased Risk of Diabetes in Inflammatory Bowel
- 402 Disease Patients: A Nationwide Population-based Study in Korea. *J Clin Med* 2019; **8**(3): 343.
- 403 26. Dungan KM, Braithwaite SS, Preiser J-C. Stress hyperglycaemia. *The Lancet* 2009;
  404 **373**(9677): 1798-807.
- Williams JG, Alam MF, Alrubaiy L, et al. Infliximab versus ciclosporin for steroid-resistant
  acute severe ulcerative colitis (CONSTRUCT): a mixed methods, open-label, pragmatic
  randomised trial. *The Lancet Gastroenterology & Hepatology* 2016; 1(1): 15-24.
- 408
   408
   408
   408
   409
   409
   409
   409
   409
   409
   409
   409
   409
   409
   409
   400
   400
   400
   400
   400
   400
   400
   400
   400
   400
   400
   400
   400
   400
   400
   400
   400
   400
   400
   400
   400
   400
   400
   400
   400
   400
   400
   400
   400
   400
   400
   400
   400
   400
   400
   400
   400
   400
   400
   400
   400
   400
   400
   400
   400
   400
   400
   400
   400
   400
   400
   400
   400
   400
   400
   400
   400
   400
   400
   400
   400
   400
   400
   400
   400
   400
   400
   400
   400
   400
   400
   400
   400
   400
   400
   400
   400
   400
   400
   400
   400
   400
   400
   400
   400
   400
   400
   400
   400
   400
   400
   400
   400
   400
   400
   400
   400
   400
   400
   400
   400
   400
   400
   400
   400
- 410 *Hepatology* 2019; **17**(1): 139-47.
- 411 29. Berinstein JA, Steiner CA, Regal RE, et al. Efficacy of Induction Therapy With High412 Intensity Tofacitinib in 4 Patients With Acute Severe Ulcerative Colitis. *Clinical Gastroenterology*413 and Hepatology 2019; **17**(5): 988-90.e1.
- 414 30. Narula N, Dhillon A, Zhang D, Sherlock ME, Tondeur M, Zachos M. Enteral nutritional
- 415 therapy for induction of remission in Crohn's disease. *Cochrane Database of Systematic* 416 *Reviews* 2018; (4).
- 417

## 418 SUPPLEMENTARY DATA

419

#### 420 Input features for RF model:

421 MUST score, weight loss during the past six months (<5%, 5-10%, >10%), diagnosis (CD, UC, IBDU), time since

422 diagnosis, admission faecal calprotectin and admission measurements of serum haemoglobin, platelets, magnesium,

423 phosphate, potassium and CRP.

424

427

## 425 GridSearch optimised parameters of the RF model: 426 • For patients with no history of DM: 'n estimat

- For patients with no history of DM: 'n\_estimators': 733, 'min\_samples\_split': 10, 'min\_samples\_leaf': 4, 'max\_features': 'sqrt', 'max\_depth': 90, 'bootstrap': True
- For all patients regardless of DM history: 'n\_estimators': 3400, 'min\_samples\_split': 2, 'min\_samples\_leaf':
  429
  4, 'max\_features': 'sqrt', 'max\_depth': 80, 'bootstrap': False
- 430
- 431

432

433



#### 435 Supplementary figure 1: Correlation heat map of Random Forest parameters (subjects with history of DM excluded)

#### 438 Supplementary figure 2: Predicted highest CBG vs measured highest CBG (all patients)



439 440

Supplementary figure 3: Variable importance for Random Forest modelling (all patients)



441

442



#### 444 Supplementary figure 4: Correlation heat map of Random Forest parameters (all patients)

445 446 447

448

-<del>--</del>-0









454 Supplementary figure 6: Highest recorded blood glucose stratified by nutritional supplements given (all patients)

455 NG = Nasogastric tube feeding, Oral = Oral nutritional supplements, TPN = total parenteral nutrition

