

1 **SARS-CoV-2 spread across the Colombian-Venezuelan border**

2 Alberto Paniz-Mondolfi^{1*}, Marina Muñoz², Carolina Florez³, Sergio Gomez³, Angelica
3 Rico³, Lisseth Pardo³, Esther C. Barros³, Carolina Hernández², Lourdes Delgado², Jesús E.
4 Jaimes², Luis Pérez², Aníbal A. Teherán⁴, Hala Alejel Alshammary¹, Ajay Obla⁵, Zenab
5 Khan⁵, Ana S. Gonzalez-Reiche⁵, Matthew M. Hernandez¹, Emilia Mia Sordillo¹, Viviana
6 Simon^{6,7,8}, Harm van Bakel^{5,9}, Juan David Ramírez²

7 ¹ Department of Pathology, Molecular and Cell Based Medicine, Icahn School of Medicine
8 at Mount Sinai, New York, NY 10029, USA

9 ²Grupo de Investigaciones Microbiológicas-UR (GIMUR), Departamento de Biología,
10 Facultad de Ciencias Naturales, Universidad del Rosario, Bogotá, Colombia

11 ³ Instituto Nacional de Salud, Bogotá, Colombia

12 ⁴ Fundación Juan N Corpas, Bogotá, Colombia

13 ⁵ Department of Genetics and Genomic Sciences, Icahn School of Medicine at Mount Sinai,
14 New York, NY 10029, USA

15 ⁶Department of Microbiology, Icahn School of Medicine at Mount Sinai, New York, NY
16 10029, USA

17 ⁷ The Global Health and Emerging Pathogens Institute, Icahn School of Medicine at Mount
18 Sinai, New York, NY 10029, USA

19 ⁸ Division of Infectious Diseases, Department of Medicine, Icahn School of Medicine at
20 Mount Sinai, New York, NY 10029, USA

2

21 ⁹ Icahn Institute for Data Science and Genomic Technology, Icahn School of Medicine at
22 Mount Sinai, New York, NY 10029, USA

23 *Correspondence: Alberto.Paniz-mondolfi@mountsinai.org

24

25 **ABSTRACT**

26 **Introduction:** Venezuela and Colombia both adopted measures of containment early in
27 response to the COVID-19 pandemic. However, Venezuela's ongoing humanitarian crisis
28 has decimated its health care system, and forced millions of Venezuelans to flee through its
29 porous border with Colombia. The extensive shared border, and illegal cross-border transit
30 through improvised trails between the two countries are major challenges for public health
31 authorities. We report the first SARS-CoV-2 genomes from Venezuela, and present a
32 snapshot of the SARS-CoV-2 epidemiologic landscape in the Colombian-Venezuelan
33 border region.

34 **Methods:** We sequenced and assembled viral genomes from total RNA extracted from
35 nasopharyngeal (NP) clinical specimens using a custom reference-based analysis pipeline.
36 Three assemblies obtained were subjected to typing using the Phylogenetic Assignment of
37 Named Global Outbreak LINEages 'Pangolin' tool. A total of 376 publicly available SARS-
38 CoV-2 genomes from South America were obtained from the GISAID database to perform
39 comparative genomic analyses. Additionally, the Wuhan-1 strain was used as reference.

40 **Results:** We found that two of the SARS-CoV-2 genomes from Venezuela belonged to the
41 B1 lineage, and the third to the B.1.13 lineage. We observed a point mutation in the Spike
42 protein gene (D614G substitution), previously reported to be associated with increased

43 infectivity, in all three Venezuelan genomes. An additional three mutations (R203K/G204R
44 substitution) were present in the nucleocapsid (N) gene of one Venezuelan genome.

45 **Conclusions:** Genomic sequencing demonstrates similarity between SARS-CoV-2 lineages
46 from Venezuela and viruses collected from patients in bordering areas in Colombia and
47 from Brazil, consistent with cross-border transit despite administrative measures including
48 lockdowns. The presence of mutations associated with increased infectivity in the 3
49 Venezuelan genomes we report and Colombian SARS-CoV-2 genomes from neighboring
50 borders areas may pose additional challenges for control of SARS-CoV-2 spread in the
51 complex epidemiological landscape in Latin American countries. Public health authorities
52 should carefully follow the progress of the pandemic and its impact on displaced
53 populations within the region.

54

55 **Keywords:** SARS-CoV-2; COVID-19; Colombia; Venezuela; Border; Novel Coronavirus.

56

57 **1. INTRODUCTION**

58 As Severe Acute Respiratory Syndrome Coronavirus -2 (SARS-CoV-2) spreads throughout
59 the Western hemisphere, Latin America has become an epicenter for the Coronavirus
60 Disease 2019 (COVID-19) pandemic. Since the report on March 13, 2020 of the first two
61 cases diagnosed with COVID-19 in Venezuela, SARS-CoV-2 has spread rapidly across the
62 country. The areas of highest transmission flank the Colombian-Venezuelan border, where
63 the reported incidence reaches 47,9% with a 34,2% case fatality rate (“Coronavirus
64 Venezuela,” n.d.)

65 Venezuela's ongoing humanitarian crisis has had a severe impact on its health care system:
66 the country has experienced a massive exodus of medical personnel, shortages of treatment
67 and supplies, as well as a systematic dismantling of public health infrastructure (Daniels,
68 2020). In addition, the increased poverty and violence, plus the abandonment of all
69 epidemiological surveillance programs, has set the stage for the re-emergence of vaccine-
70 preventable and vector-borne diseases (Grillet et al., 2019; Paniz-Mondolfi et al., 2019).
71 The Venezuelan public health system cannot withstand current autochthonous threats, let
72 alone emerging infectious agents such as SARS-CoV-2. Furthermore, the political and
73 economic turmoil in Venezuela has precipitated one of the largest refugee crises witnessed
74 in the hemisphere. This has resulted in the establishment of disease corridors to neighboring
75 countries---particularly Colombia, which houses 1.4 million Venezuelan migrants (Daniels,
76 2020; Torres and Castro, 2019). Despite early implementation of lockdown and border
77 restrictions starting in March 2020, control of cross-border migration has remained a
78 challenge due to the length (approximately 2,219 km) of the Venezuelan-Colombian
79 border, and the unregulated transit of individuals through illegal trails known as “trochas”.

80 As of July 8th 2020, the majority of COVID-19 cases in Venezuela have been reported from
81 three border states, Apure, Táchira and Zulia. (“Estadísticas Venezuela | COVID-19 en
82 Venezuela,” n.d.).

83 Here we report sequences for three of five SARS-CoV-2-positive samples, representing the
84 first viral genomes from Venezuela, and providing a snapshot of the epidemiological
85 landscape across the Colombian-Venezuelan border.

86

87 **2. METHODS**

88 ***2.1. Patients, sampling and demographic data***

89 Newly arrived migrants from Venezuela meeting case-definition criteria established by the
90 Colombian Ministry of Health and Social Protection were screened for SARS-CoV-2
91 infection at different hospitals and healthcare centers in Norte de Santander and Bolivar
92 Departments of Colombia between March 31st and May 1st, 2020. Molecular detection of
93 SARS-CoV-2 in nasopharyngeal swab specimens in viral transport media (NP-VTM) was
94 performed using the Berlin Charité protocol. SARS-CoV-2 -positive specimens from five
95 individuals who were tested within 24 hours of their arrival in Colombia were referred for
96 further characterization. Four of the five individuals already were symptomatic at the time
97 of arrival from Venezuela, and one was a close contact of a confirmed COVID-19 patient.
98 Complete viral genomes were generated from specimens from three of these five cases. The
99 patients, aged 30 (♂), 42 (♂) and 56 (♀) years, came from different regions, Cucuta, Norte
100 de Santander, and Bolivar, (Fig 1A). Sequencing was unsuccessful for the other 2
101 specimens, most likely due to insufficient target material.

102 ***2.2.Phylogenetic analysis***

103 We sequenced and assembled viral genomes from total RNA extracted from NP viral
104 transport medium (VTM) clinical specimens, as described previously (Gonzalez-Reiche et
105 al., 2020; Ramirez et al., 2020). The data generated by high throughput sequencing was
106 assembled using a custom reference-based analysis pipeline (Gonzalez-Reiche et al., 2020).
107 The complete genomes were typed using the Phylogenetic Assignment of Named Global
108 Outbreak LINEages ‘Pangolin’ tool (Rambaut et al., 2020).

109 A total of 376 publicly available SARS-CoV-2 genomes encompassing the lineage
110 diversity from South America were downloaded from the GISAID database for
111 comparative genomic analyses. These sequences were aligned in MAFFT (Katoh et al.,
112 2018), using the Wuhan-1 strain (NC_045512.2) as reference. The complete data set was
113 subjected to the same typing scheme using Pangolin tool. Thus far, 28 pangolin lineages
114 have been reported for SARS-CoV2 strains circulating in South American countries (Fig.
115 1B), with B1 as the predominant lineage representing 62.9% of the total reported genomes.
116 Four other lineages include B.1.5 (12.5%), A.5 (4.5%), A.2 and B (3.4% each) account for
117 an additional 23.8%. Each of the remaining 15 lineages accounted for fewer than 10
118 genomes, with several having only a single representative. A time-scaled maximum
119 likelihood (ML) phylogeny based on TreeTime built in IQTREE (Rambaut et al., 2016;
120 Sagulenko et al., 2018) revealed that there was no clustering by originating country in the
121 reconstruction, although a general clustering by pangolin lineages was observed (Fig. 1C).
122 These analyses are in agreement with the simultaneous circulation of SARS-CoV-2
123 lineages from different geographical origins.

124 A detailed screening of single-nucleotide polymorphisms (SNPs) in important open reading
125 frames (ORFs) of SARS-CoV-2 was then conducted, and substitutions in Spike and
126 Nucleocapsid sequences were evaluated. The alignment was inspected using Ugene
127 (<http://ugene.net/>). The regions of interest were exported considering the ORFs described
128 for the reference strain Wuhan-1 (NC_045512.2), as previously described in NCBI:
129 [https://www.ncbi.nlm.nih.gov/nucore/?term = Severe + acute + respiratory + syndrome +](https://www.ncbi.nlm.nih.gov/nucore/?term=Severe+acute+respiratory+syndrome+coronavirus+2+isolate+Wuhan-Hu-1)
130 [coronavirus + 2 + isolate + Wuhan-Hu-1.](https://www.ncbi.nlm.nih.gov/nucore/?term=Severe+acute+respiratory+syndrome+coronavirus+2+isolate+Wuhan-Hu-1)

131 3. RESULTS

132 We found that two of the SARS-CoV-2 genomes obtained (VEN-89312 and VEN-95072)
133 were identical and belonged to the B1 lineage, while the third (VEN-95070) belonged to
134 the B.1.13 lineage, suggesting two separate introductions (Candido et al., 2020; Ramirez et
135 al., 2020). We analyzed the individual genomic information for the 3 Venezuelan viruses
136 reported in this study within the context of the larger global viral diversity to determine
137 potential dates for the introduction of SARS-CoV2 into Venezuela, and similarly for 3
138 viruses collected from patients in bordering areas in Colombia (Ramirez et al., 2020) (Table
139 1).

140

141 **Table 1.** Potential introduction dates of SARS-CoV-2 from Venezuelan patients and
142 patients in related border areas in Colombia.

Origin	ID	Node date	[95%CI]
Venezuela	VEN-89312	19/3/20	[18/3/20 to 19/3/20]

	VEN-95070	29/3/20	[28/3/20 to 29/3/20]
	VEN-95072	29/3/20	[28/3/20 to 29/3/20]
Colombian	GVI-97588		[16/3/20 to
related		17/3/20	17/3/20]
areas	GVI-97769	20/3/20	[19/3/20 to 20/3/20]
	GVI-97760	30/3/20	[29/3/20 to 30/3/20]

143

144 Phylogenetic analysis revealed that two of the Venezuelan SARS-CoV-2 genomes (VEN-
145 95070 and VEN-95072) closely resembled genomes from neighboring Colombia (Cesar
146 and Norte de Santander Departments) (Ramirez et al., 2020) while the third (VEN-89312)
147 was related to genomes from Brazil (Fig. 1C) (Candido et al., 2020).

148 The three Venezuelan genomes carried a G-to-A point mutation at position 23,403 resulting
149 in a D614G substitution in the spike (S) protein. This mutation characterizes the B.1
150 lineages and has been associated with enhanced viral entry into host cells, and potentially
151 with increasing infectivity and transmissibility (Bhattacharyya et al., 2020; Korber et al.,
152 2020).

153 Additionally, we identified three substitutions in the nucleocapsid (N) gene of VEN-89312
154 changing GGG-to-AAC at positions 28,616-28,618 resulting R203K/G204R substitutions
155 according to the whole-genome position after removing the 5'UTR. These R203K/G204R
156 substitutions, which have been reported previously in other South American genomes, were
157 absent from the two other Venezuelan genomes we sequenced. The alignment is shown
158 highlighting the SNPs found in the N gene (Fig. 1D) (Crooks et al., 2004).

159 **4. DISCUSSION**

160 The presence of mutations previously reported to be associated with increased infectivity
161 and transmissibility in the genomic sequences we determined for SARS-CoV-2 from both
162 Venezuela and neighboring border regions of Colombia may represent an additional
163 challenge for control of the COVID-19 pandemic in Latin America. The spectre of a
164 SARS-CoV-2 variant with increased infectivity further complicates the already-complex
165 epidemiological landscape with its chronic limitations of biomedical understaffing, poor
166 healthcare infrastructure, limited or inadequate diagnostic capacities, and poor compliance
167 of the general population with disease containment measures (Miller et al., 2020).

168 Following the first report of SARS-CoV-2 in Brazil in late February 2020 (Andrus et al.,
169 2020; Rodriguez-Morales et al., 2020), SARS-CoV-2 has spread rapidly across the region,
170 resulting in nearly two million cases as of June 25th, 2020, with Brazil and Peru reporting
171 the highest burden of disease (“Epidemic Diseases - Cumulative suspected and confirmed
172 COVID-19 cases reported by countries and territories in the Americas,” n.d.).

173 Although Colombia and Venezuela both implemented early lockdown and containment
174 strategies since March 10th and March 17th, respectively, (“CORONAVIRUS (COVID-
175 19),” n.d., “Presidente Maduro anuncia que este martes todo el país entra en cuarentena
176 social • Ministerio del Poder Popular para Relaciones Exteriores,” n.d.), according to
177 current official records there have been over 124,500 COVID-19 cases in Colombia and
178 over 7,700 cases in Venezuela (“CORONAVIRUS (COVID-19),” n.d., “Estadísticas
179 Venezuela | COVID-19 en Venezuela,” n.d.). However, these numbers have to be
180 contextualized because: (i) Colombia has exclusively performed state-of-the-art RT-PCR-
181 based SARS-CoV-2 diagnosis whereas Venezuelan authorities have relied mostly on

182 serological testing that are considerably less specific (Chia et al., 2020) and less useful than
183 molecular methods for acute case detection; (ii) Venezuela has precarious epidemiological
184 surveillance and contact tracing systems; and (iii) Venezuela has reduced international
185 traffic through airports secondary to extensive departure of multiple airlines.

186 The concurrent humanitarian crisis has forced millions of Venezuelans to flee to
187 neighboring countries ---mainly Colombia--- seeking economic and social stability (Torres
188 and Castro, 2019; Tuite et al., 2018). Venezuelans with ongoing health issues also travel to
189 Colombia to obtain high-quality healthcare such as surgical procedures and hemodialysis
190 (Daniels, 2020). However, massive Venezuelan migration has resulted in an unprecedented
191 infectious disease exodus, representing one of the most concerning public health threats in
192 the region (Grillet et al., 2019; Torres and Castro, 2019). COVID-19 has deepened the
193 situation and has prompted xenophobia and further marginalization of Venezuelan migrants
194 and refugees at the Venezuelan-Colombian border (Daniels, 2020).

195 The shared presence of B lineages in SARS-CoV-2 from Venezuela and Colombia
196 reinforces the close interactions of persons living in border regions and the difficulty of
197 containment across a porous border. Similarly, the detection of the B.1.13 lineage, only
198 previously described in cases in Spain, England and Australia (Batty et al., 2020) further
199 underlines the rapid global spread of SARS-CoV-2 through interconnected populations.
200 Additionally, the presence of substitution D614G in the spike protein of the three viruses
201 from patients residing in the current hotspots of COVID-19 in Venezuela may correlate
202 with the reported increased infectivity (Korber et al., 2020) observed in SARS-CoV-2-
203 infected patients in the state of Zulia (“Zulia suma 15 muertes confirmadas por coronavirus
204 y seis no reportadas - Efecto Cocuyo,” n.d.).

205 An important limitation to our study is that the limited number of genomes currently
206 available from Venezuela could potentially result in sampling bias. Given the difficulty in
207 obtaining samples, the extent to which our findings truly reflect Venezuela's overall
208 phylogenetic landscape remains to be determined. Future studies are needed to expand the
209 SARS-CoV-2 genome repertoire in Venezuela and related areas, and to enable better
210 understanding of the interplay between genotype and phenotype, and their relevance for
211 disease surveillance and containment.

212 **5. CONCLUSION**

213 The Venezuelan humanitarian and refugee crisis, coupled with uncontrolled migration
214 across the Colombian-Venezuelan border, is a devastating reminder of the potential effects
215 of infectious disease spillover on the already vulnerable public health systems of
216 neighboring countries. As SARS-CoV-2 continues to spread across Latin America, public
217 health authorities and the international community should carefully follow the impact of the
218 pandemic on displaced populations. Intensive efforts are urgently needed to help minimize
219 the impact of Venezuela's crisis on the COVID-19 pandemic crisis.

220 **6. ACKNOWLEDGEMENTS**

221 Funding was provided by the University of Glasgow, Scottish Funding Council and the
222 Global Challenges Research Fund (GCRF) and GCRF Research Network EP/T003782/1.

223 **REFERENCES**

224 Andrus, J.K., Evans-Gilbert, T., Santos, J.I., Guzman, M.G., Rosenthal, P.J., Toscano, C.,
225 Valenzuela, M.T., Siqueira, M., Etienne, C., Breman, J.G., 2020. Perspectives on
226 Battling COVID-19 in Countries of Latin America and the Caribbean. *Am. J. Trop.*

- 227 Med. Hyg. 1–4. <https://doi.org/10.4269/ajtmh.20-0571>
- 228 Batty, E.M., Kochakarn, T., Panthan, B., Kumpornsin, K., Jiaranai, P., Wangwiwatsin, A.,
229 Kotanan, N., Jaruampornpan, P., Watthanachockchai, T., Rakmanee, K., Sensorn, I.,
230 Sungkanuparph, S., Pasomsub, E., Chookajorn, T., Chantratita, W., 2020. Genomic
231 surveillance of SARS-CoV-2 in Thailand reveals mixed imported populations, a local
232 lineage expansion and a virus with truncated ORF7a. medRxiv 2020.05.22.20108498.
233 <https://doi.org/10.1101/2020.05.22.20108498>
- 234 Bhattacharyya, C., Das, C., Ghosh, A., Singh, A.K., Mukherjee, S., Majumder, P.P., Basu,
235 A., Biswas, N.K., 2020. Global Spread of SARS-CoV-2 Subtype with Spike Protein
236 Mutation D614G is Shaped by Human Genomic Variations that Regulate Expression
237 of TMPRSS2 and MX1 Genes. bioRxiv 2020.05.04.075911.
238 <https://doi.org/10.1101/2020.05.04.075911>
- 239 Candido, D. da S., Claro, I.M., de Jesus, J.G., Souza, W.M. de, Moreira, F.R.R., Dellicour,
240 S., Mellan, T.A., du Plessis, L., Pereira, R.H.M., Sales, F.C. da S., Manuli, E.R.,
241 Theze, J., Almeida, L., de Menezes, M.T., Voloch, C.M., Fumagalli, M.J., Coletti, T.
242 de M., Silva, C.A.M., Ramundo, M.S., Amorim, M.R., Hoeltgebaum, H., Mishra, S.,
243 Gill, M., Carvalho, L.M., Buss, L.F., Prete, C.A., Ashworth, J., Nakaya, H., Peixoto,
244 P. da S., Brady, O.J., Nicholls, S.M., Tanuri, A., Rossi, A.D., Braga, C.K.V., Gerber,
245 A.L., Guimaraes, A.P., Gaburo, N., Alencar, C.S., Ferreira, A.C. de S., Lima, C.X.,
246 Levi, J.E., Granato, C., Ferreira, G.M., Francisco, R. da S., Granja, F., Garcia, M.T.,
247 Moretti, M.L., Perroud, M.W., Castineiras, T.M.P.P., Lazari, C.D.S., Hill, S.C.,
248 Santos, A.A. de S., Simeoni, C.L., Forato, J., Sposito, A.C., Schreiber, A.Z., Santos,
249 M.N.N., Sa, C.Z., Souza, R.P., Resende Moreira, L.C., Teixeira, M.M., Hubner, J.,
250 Leme, P.A.F., Moreira, R.G., Nogueira, M.L., Ferguson, N., Costa, S.F., Proenca-

- 251 Modena, J.L., Vasconcelos, A.T., Bhatt, S., Lemey, P., Wu, C.-H., Rambaut, A.,
252 Loman, N.J., Aguiar, R.S., Pybus, O.G., Sabino, E.C., Faria, N.R., 2020. Evolution
253 and epidemic spread of SARS-CoV-2 in Brazil. medRxiv 2020.06.11.20128249.
254 <https://doi.org/10.1101/2020.06.11.20128249>
- 255 Chia, W.N., Tan, C.W., Foo, R., Kang, A.E.Z., Peng, Y., Sivalingam, V., Tiu, C., Ong,
256 X.M., Zhu, F., Young, B.E., Chen, M.I.C., Tan, Y.-J., Lye, D.C., Anderson, D.E.,
257 Wang, L.-F., 2020. Serological differentiation between COVID-19 and SARS
258 infections. *Emerg. Microbes Infect.* 1–23.
259 <https://doi.org/10.1080/22221751.2020.1780951>
- 260 CORONAVIRUS (COVID-19) [WWW Document], n.d. URL
261 https://www.minsalud.gov.co/salud/publica/PET/Paginas/Covid-19_copia.aspx
262 (accessed 7.8.20).
- 263 Coronavirus Venezuela [WWW Document], n.d. URL [https://coronavirus-venezuela-](https://coronavirus-venezuela-esriven.hub.arcgis.com/)
264 [esriven.hub.arcgis.com/](https://coronavirus-venezuela-esriven.hub.arcgis.com/) (accessed 7.3.20).
- 265 Crooks, G.E., Hon, G., Chandonia, J.-M., Brenner, S.E., 2004. WebLogo: a sequence logo
266 generator. *Genome Res.* 14, 1188–1190. <https://doi.org/10.1101/gr.849004>
- 267 Daniels, J.P., 2020. Venezuelan migrants “struggling to survive” amid COVID-19. *Lancet*
268 (London, England) 395, 1023. [https://doi.org/10.1016/S0140-6736\(20\)30718-2](https://doi.org/10.1016/S0140-6736(20)30718-2)
- 269 Epidemic Diseases - Cumulative suspected and confirmed COVID-19 cases reported by
270 countries and territories in the Americas [WWW Document], n.d. URL
271 <https://ais.paho.org/phil/viz/COVID19Table.asp> (accessed 6.26.20).
- 272 Estadísticas Venezuela | COVID-19 en Venezuela [WWW Document], n.d. URL
273 <https://covid19.patria.org.ve/estadisticas-venezuela/> (accessed 7.8.20).
- 274 Gonzalez-Reiche, A.S., Hernandez, M.M., Sullivan, M.J., Ciferri, B., Alshammary, H.,

275 Obla, A., Fabre, S., Kleiner, G., Polanco, J., Khan, Z., Albuquerque, B., van de
276 Guchte, A., Dutta, J., Francoeur, N., Melo, B.S., Oussenko, I., Deikus, G., Soto, J.,
277 Sridhar, S.H., Wang, Y.-C., Twyman, K., Kasarskis, A., Altman, D.R., Smith, M.,
278 Sebra, R., Aberg, J., Krammer, F., García-Sastre, A., Luksza, M., Patel, G., Paniz-
279 Mondolfi, A., Gitman, M., Sordillo, E.M., Simon, V., van Bakel, H., 2020.
280 Introductions and early spread of SARS-CoV-2 in the New York City area. *Science*
281 (80-). eabc1917. <https://doi.org/10.1126/science.abc1917>

282 Grillet, M.E., Hernández-Villena, J. V., Llewellyn, M.S., Paniz-Mondolfi, A.E., Tami, A.,
283 Vincenti-Gonzalez, M.F., Marquez, M., Mogollon-Mendoza, A.C., Hernandez-Pereira,
284 C.E., Plaza-Morr, J.D., Blohm, G., Grijalva, M.J., Costales, J.A., Ferguson, H.M.,
285 Schwabl, P., Hernandez-Castro, L.E., Lamberton, P.H.L., Streicker, D.G., Haydon,
286 D.T., Miles, M.A., Acosta-Serrano, A., Acquattella, H., Basañez, M.G., Benaim, G.,
287 Colmenares, L.A., Conn, J.E., Espinoza, R., Freilij, H., Graterol-Gil, M.C., Hotez,
288 P.J., Kato, H., Lednicky, J.A., Martinez, C.E., Mas-Coma, S., Morris, J.G., Navarro,
289 J.C., Ramirez, J.L., Rodriguez, M., Urbina, J.A., Villegas, L., Segovia, M.J., Carrasco,
290 H.J., Crainey, J.L., Luz, S.L.B., Moreno, J.D., Noya Gonzalez, O.O., Ramírez, J.D.,
291 Alarcón-de Noya, B., 2019. Venezuela’s humanitarian crisis, resurgence of vector-
292 borne diseases, and implications for spillover in the region. *Lancet Infect. Dis.* 19,
293 e149–e161. [https://doi.org/10.1016/S1473-3099\(18\)30757-6](https://doi.org/10.1016/S1473-3099(18)30757-6)

294 Katoh, K., Rozewicki, J., Yamada, K.D., 2018. MAFFT online service: Multiple sequence
295 alignment, interactive sequence choice and visualization. *Brief. Bioinform.* 20, 1160–
296 1166. <https://doi.org/10.1093/bib/bbx108>

297 Korber, B., Fischer, W.M., Gnanakaran, S., Yoon, H., Theiler, J., Abfalterer, W.,
298 Hengartner, N., Giorgi, E.E., Bhattacharya, T., Foley, B., Hastie, K.M., Parker, M.D.,

- 299 Partridge, D.G., Evans, C.M., Freeman, T.M., de Silva, T.I., McDanal, C., Perez, L.G.,
300 Tang, H., Moon-Walker, A., Whelan, S.P., LaBranche, C.C., Sapphire, E.O.,
301 Montefiori, D.C., Group, S.C.-19 G., 2020. Tracking changes in SARS-CoV-2 Spike:
302 evidence that D614G increases infectivity of the COVID-19 virus. *Cell*.
303 <https://doi.org/10.1016/j.cell.2020.06.043>
- 304 Miller, M.J., Loaiza, J.R., Takyar, A., Gilman, R.H., 2020. Covid-19 in latin america:
305 Novel transmission dynamics for a global pandemic? *PLoS Negl. Trop. Dis.* 14, 3–7.
306 <https://doi.org/10.1371/JOURNAL.PNTD.0008265>
- 307 Paniz-Mondolfi, A.E., Tami, A., Grillet, M.E., Márquez, M., Hernández-Villena, J.,
308 Escalona-Rodríguez, M.A., Blohm, G.M., Mejías, I., Urbina-Medina, H., Rísquez, A.,
309 Castro, J., Carvajal, A., Walter, C., López, M.G., Schwabl, P., Hernández-Castro, L.,
310 Miles, M.A., Hotez, P.J., Lednicky, J., Morris, J.G., Crainey, J., Luz, S., Ramírez,
311 J.D., Sordillo, E., Llewellyn, M., Canache, M., Araque, M., Oletta, J., 2019.
312 Resurgence of Vaccine-Preventable Diseases in Venezuela as a Regional Public
313 Health Threat in the Americas. *Emerg. Infect. Dis.* 25, 625–632.
314 <https://doi.org/10.3201/eid2504.181305>
- 315 Presidente Maduro anuncia que este martes todo el país entra en cuarentena social •
316 Ministerio del Poder Popular para Relaciones Exteriores [WWW Document], n.d.
317 URL <http://mppre.gob.ve/2020/03/16/Venezuela-cuarentena-social/> (accessed
318 6.26.20).
- 319 Rambaut, A., Holmes, E.C., Hill, V., O’Toole, Á., McCrone, J.T., Ruis, C., du Plessis, L.,
320 Pybus, O.G., 2020. A dynamic nomenclature proposal for SARS-CoV-2 to assist
321 genomic epidemiology. *bioRxiv* 2020.04.17.046086.
322 <https://doi.org/10.1101/2020.04.17.046086>

- 323 Rambaut, A., Lam, T.T., Carvalho, L.M., Pybus, O.G., 2016. Exploring the temporal
324 structure of heterochronous sequences using TempEst (formerly Path-O-Gen). *Virus*
325 *Evol.* 2, 1–7. <https://doi.org/10.1093/ve/vew007>
- 326 Ramirez, J.D., Florez, C., Munoz, M., Hernandez, C., Castillo, A., Gomez, S., Rico, A.,
327 Pardo, L., Barros, E.C., Castaneda, S., Ballesteros, N., Martinez, D., Vega, L., Jaimes,
328 J., Cruz-Saavedra, L., Patino, L.H., Teheran, A., Gonzalez-Reiche, A., Hernandez, M.,
329 Sordillo, E., Simon, V., van Bakel, H., Paniz-Mondolfi, A., 2020. The arrival and
330 spread of SARS-CoV2 in Colombia. *medRxiv* 2020.06.11.20125799.
331 <https://doi.org/10.1101/2020.06.11.20125799>
- 332 Rodriguez-Morales, A.J., Gallego, V., Escalera-Antezana, J.P., Méndez, C.A., Zambrano,
333 L.I., Franco-Paredes, C., Suárez, J.A., Rodriguez-Enciso, H.D., Balbin-Ramon, G.J.,
334 Savio-Larriera, E., Riquez, A., Cimerman, S., 2020. COVID-19 in Latin America:
335 The implications of the first confirmed case in Brazil. *Travel Med. Infect. Dis.* 35,
336 101613. <https://doi.org/10.1016/j.tmaid.2020.101613>
- 337 Sagulenko, P., Puller, V., Neher, R.A., 2018. TreeTime: Maximum-likelihood
338 phylodynamic analysis. *Virus Evol.* 4, 1–9. <https://doi.org/10.1093/ve/vex042>
- 339 Torres, J.R., Castro, J.S., 2019. Venezuela’s migration crisis: a growing health threat to the
340 region requiring immediate attention. *J. Travel Med.* 26.
341 <https://doi.org/10.1093/jtm/tay141>
- 342 Tuite, A.R., Thomas-Bachli, A., Acosta, H., Bhatia, D., Huber, C., Petrasek, K., Watts, A.,
343 Yong, J.H.E., Bogoch, I.I., Khan, K., 2018. Infectious disease implications of large-
344 scale migration of Venezuelan nationals. *J. Travel Med.* 25, 1–8.
345 <https://doi.org/10.1093/jtm/tay077>
- 346 Zulia suma 15 muertes confirmadas por coronavirus y seis no reportadas - Efecto Cocuyo

347 [WWW Document], n.d. URL <https://efectococuyo.com/coronavirus/zulia-suma-15->
348 [muertes-confirmadas-por-coronavirus-y-seis-no-reportadas/](https://efectococuyo.com/coronavirus/zulia-suma-15-muertes-confirmadas-por-coronavirus-y-seis-no-reportadas/) (accessed 6.27.20).

349

350

351 **Figure Legends**

352 **Figure 1. Regional comparative genomic analysis of SARS-CoV-2. A.** Geographical
353 distribution of the sequences from Venezuelan genomes and the available Colombian
354 genomes analyzed in this study. The pink symbols indicate the Venezuelan patients
355 identified in Colombia with their respective Venezuelan state origin. **B.** Stacked bar plot of
356 the number of genomes per lineage determined using Phylogenetic Assignment of Named
357 Global Outbreak LINEages ‘Pangolin’ tool. Three SARS-CoV-2 genomes from Venezuela
358 were compared with 376 assemblies from other 7 South American countries (Argentina,
359 Brazil, Chile, Colombia, Ecuador, Peru and Uruguay) using the publicly available GISAID
360 EpiCoVTM database (<https://www.gisaid.org/>). Frequencies are discriminated by country of
361 origin. The Venezuelan lineages are identified by black arrows. **C.** Maximum likelihood
362 tree built in IQtree shows the phylogenetic relationships between genomes from Venezuela
363 (pink dots) and the closest Colombian regions (yellow dots) with other South American
364 genomes. The branches were colored according to the country of origin, using the color
365 code of A panel. The clustering of the most frequent pangolin lineages ($n > 10$) is
366 represented on the right side. The black dots represent highly supported nodes. **D.** Multiple
367 alignment of the Nucleocapsid gene showing the substitutions found using the Wuhan-1
368 sequence as reference.

369

18

370

371

