

Sample Size Calculation for Phylogenetic Case Linkage

Shirlee Wohl, John R. Giles, and Justin Lessler

Affiliations

Johns Hopkins Bloomberg School of Public Health, Department of Epidemiology, Baltimore, MD, USA

Correspondence to: justin@jhu.edu

Abstract

Sample size calculations are an essential component of the design and evaluation of scientific studies. However, there is a lack of clear guidance for determining the sample size needed for phylogenetic studies, which are becoming an essential part of studying pathogen transmission. We introduce a statistical framework for determining the number of true infector-infectee transmission pairs identified by a phylogenetic study, given the size and population coverage of that study. We then show how characteristics of the criteria used to determine linkage and aspects of the study design can influence our ability to correctly identify transmission links, in sometimes counterintuitive ways. We test the overall approach using outbreak simulations and provide guidance for calculating the sensitivity and specificity of the linkage criteria, the key inputs to our approach. The framework is freely available as the R package *phylosamp*, and is broadly applicable to designing and evaluating a wide array of pathogen phylogenetic studies.

22 Introduction

23 As the cost of pathogen sequencing has declined, the number and size of studies based on pathogen sequence
 24 analysis has increased dramatically (Neher and Bedford 2018). Traditionally, researchers have sequenced
 25 convenience samples collected as part of routine clinical or public health activities (e.g., diagnostic specimens
 26 collected as part of an outbreak response), or as part of studies where specimens are collected for other purposes.
 27 However, the analysis of pathogen genomic sequences is increasingly becoming a primary goal of both research
 28 studies and public health surveillance efforts (Gardy et al. 2011; Jackson et al. 2016; Quick et al. 2016; Snider et al.
 29 2016). This shift has been driven by the apparent utility of pathogen sequence data for understanding aspects of
 30 pathogen spread ranging from the frequency and source of introductions into a region (Nelson et al. 2007; Lei and
 31 Shi 2011; Thézé et al. 2018; Weill et al. 2019; Gonzalez-Reiche et al. 2020), to identifying endogenous spread of
 32 emerging diseases (Carroll et al. 2015; Park et al. 2015), to understanding the role of “hotspots” in maintaining
 33 broader community epidemics (Ratmann et al. 2020), to understanding transmission patterns at an individual or
 34 “microscale” level (Gardy et al. 2011; Salje et al. 2012).

35 Despite these many examples, there is a lack of clear and accessible guidance for appropriately designing and
 36 sizing studies aimed at understanding pathogen transmission, or for evaluating the design and conclusions of past
 37 studies. Without such guidance, it is difficult for researchers to design studies in a way that maximizes the chances
 38 of success, and difficult for reviewers to appropriately evaluate papers and grant applications centered around
 39 molecular or phylogenetic outcomes (Volz and Frost 2013; Frost et al. 2015). In particular, undersampling or biased
 40 sampling can lead to poorly supported inferences about patterns of disease spread (Grabowski and Lessler 2017;
 41 Mavian et al. 2020). While there are examples of researchers conducting careful *a priori* analyses of sampling
 42 strategies (Network and Others 2013; Farhat et al. 2014; Kelly et al. 2015), these have largely relied on
 43 sophisticated techniques that are not broadly generalizable. Hence, there is a need for broadly accepted and
 44 accessible guidance for the selection of specimens for sequencing and phylogenetic analyses.

45 As noted above, pathogen sequences have been used to understand multiple aspects of infectious disease
 46 transmission at scales ranging from the global (e.g., movement of pathogens between countries) to the individual
 47 (e.g., reconstruction of individual transmission chains). Arguably, all such analyses can be reduced to the basic
 48 question of whether pairs of infected individuals are related within a particular number of generations of
 49 transmission. Therefore, developing tools for assessing the number of sequences needed to confidently identify
 50 linked pairs (infections separated by no more than a specific number of generations of transmission) is a good place
 51 to start building a theory for power calculations for phylogenetic inference. In this paper, we present a framework
 52 for making critical decisions about study design when the goal is to identify infector-infectee pairs, and we
 53 illustrate this approach with simulation studies.

54 Approach

55 General Principles

56 In this paper we will deal with studies that aim to identify infector-infectee pairs from phylogenetic analysis of
 57 pathogen sequence data collected from infected individuals. We assume the study aims to achieve some level of
 58 certainty that identified infector-infectee pairs are correct, and may also require identification of some minimum
 59 number of pairs. Below we lay out a precise terminology (**Table 1**) and general principles.

60 **Table 1: Parameters used in calculations and simulations.**

Parameter	Description
M	Number of infections sampled
N	Total number of (relevant) infected individuals in an outbreak
ρ	Proportion of outbreak infections sampled (M/N)
η	Sensitivity of the linkage criteria
χ	Specificity of the linkage criteria
ϕ	Probability that an identified link represents a true transmission event (1-False Discovery Rate)
R	Reproductive number of a pathogen
R_{pop}	Average reproductive number of a pathogen in a finite population (always <1)
μ	Mutation rate of the pathogen (in mutations per genome per transmission event)

61
 62 To start, we define the term *linkage criteria* to represent all the criteria used to infer whether a set of infected
 63 individuals are linked to one another by direct transmission. The *linkage criteria* can be derived from a combination
 64 of genetic distance between pathogens isolated from different individuals, tree structure (e.g., clade support), and
 65 epidemiologic information (e.g., relative dates of symptom onset). We refer to infections inferred to be connected
 66 by transmission using this criteria as *linked pairs*. Some of these linked pairs will represent actual transmission
 67 events (*true transmission pairs*) and some will be false positives. We want to determine the sample size (M) and
 68 proportion of the population (ρ) required to recover a predetermined number of linked pairs, while keeping the *false*
 69 *discovery rate* (the proportion of these linked pairs that are false positives) below a predetermined threshold. When
 70 applied to a study where design was dictated by other factors (e.g., specimen availability), the same methods can be
 71 used to determine the *false discovery rate*, which will inform the confidence we have in any conclusions about
 72 disease transmission in that study.


73 To capture *true transmission pairs*, the infector and their partner infectee must both be in the sample. Therefore,
 74 correctly identifying direct transmission links (and, conversely, calculating the false discovery rate) depends on the
 75 sampling fraction (ρ), which is equal to the sample size (M) divided by the total number of infected individuals in
 76 the relevant population (N). Identification of these links will further depend on the *sensitivity* (η) and *specificity* (χ)
 77 of the criteria used to define linkage. We define sensitivity as the probability that the linkage criteria will identify a
 78 true transmission pair as a linked pair given that both the infector and infectee are in the sample. Similarly, the
 79 specificity is the probability that two infections not linked by transmission are not linked by the linkage criteria.

Here we show that, if we have reasonable estimates of the sampling fraction, sensitivity, and specificity, we can, for a sample of size M , estimate the false discovery rate. The relationship between these parameters can then be used to design studies with a sample size and sampling fraction that minimizes the false discovery rate and therefore maximizes our ability to draw inferences from identified infections.

Calculating sample size and false discovery rate

Single link and single true transmission

We start with the simple example of identifying the correct infector of a particular infection (Volz and Frost 2013). In this scenario, we make assumptions about transmission that simplify the relationship between sample size and false discovery rate. Namely, we assume that each infected individual is connected by transmission to exactly one other individual, and that the linkage criteria similarly identifies exactly one probable link for each infection. Under these assumptions, we can calculate the probability of correctly identifying a true transmission pair, ϕ (equal to one minus the false discovery rate), as a function of the sensitivity and specificity of the linkage criteria, the proportion sampled, and the sample size. **Figure 1** provides some intuition as to the form of this probability expression under the stated assumptions of single linkage and single transmission (see **Text S1** for full derivation).



94

Figure 1. Visual derivation of the probability of correctly identifying a true transmission pair. Blue shaded regions represent correct identification of the true transmission partner of a random infection i . Red shaded regions represent linkage of i to an infection that is not its true transmission partner. White shaded regions represent the probability of no linkage occurring.

98

99 The probability of correctly identifying a true transmission pair (ϕ) under the assumptions of single transmission
100 and single linkage is:

$$\phi = \frac{\eta\rho}{\eta\rho + (1 - \chi^{M-2})(1 - \eta)\rho + (1 - \chi^{M-1})(1 - \rho)} \quad (1)$$


101 Under the same assumptions, we can also calculate the expected total number of true transmission pairs that will be
102 identified in our sample, $\mathbb{E}[\text{number of true pairs}]$, as:

$$\mathbb{E}[\text{number of true pairs}] = \frac{M}{2}\eta\rho$$

103 Through algebraic rearrangement of these equations, we can determine the expected number of linked pairs
104 (identified with the linkage criteria) observed in this sample ($\mathbb{E}[\text{number of pairs observed}]$):

$$\mathbb{E}[\text{number of pairs observed}] = \frac{M}{2}[\eta\rho + (1 - \chi^{M-2})(1 - \eta)\rho + (1 - \chi^{M-1})]$$

105 These equations can be used to determine the false discovery rate ($1 - \phi$) and the expected number of linked pairs
106 given a particular criteria, sample size, and sampling proportion. Additionally, we can use these equations to
107 observe how the expected number of links and the true discovery rate vary with the proportion sampled and the
108 sample size (**Fig 2A**). For a given sensitivity and specificity of the linkage criteria, we observe that the false
109 discovery rate *increases* with sample size if the proportion sampled remains constant, suggesting that studies aimed
110 at correctly identifying the highest proportion of transmission links should prioritize sampling proportion over
111 number of samples. Additionally, the relationship between false discovery rate and sampling proportion is
112 dependent on the sample size needed to obtain that sampling proportion such that the impact of sampling proportion
113 increases with sample size. We also observe the effects of changing sensitivity and specificity on the false
114 discovery rate and find that the specificity of the linkage criteria is of key importance when attempting to minimize
115 the false discovery rate of transmission pairs (**Fig 2B**).



116

117 **Figure 2. Sample size and false discovery rate given single linkage and single transmission.** (A) Effect of sample size
118 (red lines) or proportion sampled (blue lines) on the expected number of linked pairs (upper plots) or the false discovery rate of
119 linked pairs (lower plots). The specificity and sensitivity are held constant. (B) Effect of varying the sensitivity and specificity of
120 the linkage criteria on the false discovery rate (FDR).

121 *Multiple links and multiple true transmissions*

122 In many cases, we will be interested in linking an infected individual to both their infector and anyone they infect.
 123 Therefore, we must account for the fact that each infection in an outbreak may be linked by transmission to
 124 multiple other infections, only some of which may have been sampled. If the goal is to identify all such true
 125 transmission pairs in the sample, the linkage criteria used must similarly allow for multiple linkages. Here, we
 126 calculate the false discovery rate for transmission pairs under these assumptions.

127 The average number of transmission links per infection is determined by the epidemiological parameter R , the
 128 expected number of other individuals each infected individual infects. However, sampled infections come from a
 129 bounded source population. In this finite sampling frame, the average number of infectees per infector, denoted
 130 R_{pop} , may differ from R (in fact, R_{pop} must be less than 1, see below). Because each infection is linked to, on
 131 average, R_{pop} infectees as well as its infector, each infection has $R_{\text{pop}} + 1$ true transmission partners. If we assume
 132 that the distribution of the number of transmission partners per infection is Poisson distributed, we get the following
 133 equation for the true discovery rate, ϕ (see **Text S1** for full derivation):

$$\phi = \frac{\eta\rho(R_{\text{pop}} + 1)}{\eta\rho(R_{\text{pop}} + 1) + (1 - \chi)(M - \rho(R_{\text{pop}} + 1) - 1)} \quad (2)$$

134 Under the same assumptions, we can calculate the total number of sampled true pairs, $\mathbb{E}[\text{number of true pairs}]$, as:

$$\mathbb{E}[\text{number of true pairs}] = \frac{M\rho(R_{\text{pop}} + 1)\eta}{2}$$

135 Through algebraic rearrangement of these equations we can determine the expected number of pairs observed in
 136 this sample, $\mathbb{E}[\text{number of pairs observed}]$:

$$\mathbb{E}[\text{number of pairs observed}] = \frac{M}{2}[\eta\rho(R_{\text{pop}} + 1) + (1 - \chi)(M - \rho(R_{\text{pop}} + 1) - 1)]$$

137 Again, we observe that the false discovery rate increases with the sample size, but decreases with the proportion
 138 sampled, and we see the important effect of the specificity of the linkage criteria on the false discovery rate (**Fig 3**).

139 **Estimating the average reproductive number**

140 In the previous section, we distinguished R , the basic reproductive number of a pathogen, from R_{pop} , the *average*
 141 reproductive number in a bounded population. This is an important distinction because we can show that the
 142 average reproductive number (R_{pop}) is at most one. This is because any sampling frame contains a finite number of
 143 infected individuals. Therefore, there will always be more infections than infection events (at minimum, all
 144 infectees in a transmission chain plus a single index case, see **Fig S1**). Hence, R_{pop} , which is equal to the number of
 145 actual transmission events divided by the number of infections, will be at most one.

146 In epidemic situations where there is a single introduction, R_{pop} will be close to one, as the number of infections will
 147 exceed the number of infection events by precisely one. In situations where there are multiple introductions (e.g.,

transmission chains that are persistently seeded from sources outside the sampling frame) then R_{pop} may be substantially less than one. Specifically:

$$R_{pop} = \frac{\text{cases} - \text{introductions}}{\text{cases}}$$

The examples shown in this paper focus on epidemics seeded by a single introduction, where R_{pop} is approximately equal to one.




Figure 3. Sample size and false discovery rate given multiple linkage and multiple transmissions. (A) Effect of sample size (red lines) or proportion sampled (blue lines) on the expected number of linked pairs (upper plots) or the false discovery rate of linked pairs (lower plots). The specificity and sensitivity are held constant. (B) Effect of varying the sensitivity and specificity of the linkage criteria on the false discovery rate (FDR). White dots: theoretical sensitivity and specificity values at different genetic distance thresholds for a hypothetical pathogen with mutation rate = 1 mutation/genome/transmission and $R=2$ (see 'Determining sensitivity and specificity' below for details). In both panels, $R_{pop} = 1$.

Determining sensitivity and specificity

In the framework presented here, the sensitivity and specificity of the linkage criteria are needed to estimate the false discovery rate from sample size and vice versa. This criteria can be based on a number of phylogenetic and epidemiological metrics, and may depend on the data available for a particular study. In this section, we outline two methods for approximating the sensitivity and specificity of a simple genomic metric: genetic distance.

Both methods involve determining these parameters from the discrete distributions of genetic distances between linked and unlinked infections, but they differ in how these distributions are obtained. Given the distributions, we can consider a number of different genetic distance thresholds (e.g., 2 mutations between sequences) that could be used as the criteria for differentiating between linked and unlinked pairs, and we can calculate the sensitivity and specificity at each. The optimal threshold and its associated sensitivity and specificity can be selected in a variety of ways (Youden 1950; Perkins and Schisterman 2006; Liu 2012; Zou et al. 2013) based on the specific study goals.


Below, we describe two ways to obtain the genetic distance distributions of linked and unlinked infection pairs for a hypothetical pathogen with $R=2$ and a mutation rate (μ) of 1 mutation per genome per generation. Here and

172 henceforth, “generation” refers to a generation of transmission (i.e., the mutation rate provides the number of
173 mutations expected per transmission event, not per viral replication).

174 *Empirical method*

175 One way to estimate the relevant genetic distance distributions is to use existing data. Specifically, we need a
176 subsample of infections for which sequencing data is available and we have a high degree of confidence—based on
177 epidemiological data—of the true transmission relationships between included infections. For example, infected
178 individuals who share a household versus community members with no known relationship. We can compute the
179 genetic distance between every pair of pathogen sequences from this subsample and use the results to approximate
180 the underlying genetic distance distributions between linked and unlinked infections in the population.

181 We illustrate this method on a simulated outbreak of approximately 1500 infections (data available at
182 <https://github.com/HopkinsIDD/phylosamplesize>), created using the *outbreaker* R package (R Core Team 2013;
183 Jombart et al. 2014) (see ‘Outbreak simulations’ below). To create our known subsample, we selected a small
184 number of infections from early in the outbreak and extracted their true transmission links and simulated genomes.
185 We then calculated the genetic distance matrix of sequences in this subsample and determined the genetic distance
186 distributions (**Fig 4A**). Next, we estimated the sensitivity and specificity at every mutation threshold (0 mutations, 1
187 mutation, etc.) and used the point closest to the (0,1) corner to determine the optimal threshold for differentiating
188 between linked and unlinked infections. In this case, the optimal threshold was 3 mutations, which had a sensitivity
189 of 0.95 and a specificity of 0.88.



190

191 **Figure 4. Determining the sensitivity and specificity of a genetic distance threshold.** (A) Empirical distribution of genetic
192 distances for linked (purple) and unlinked (yellow) infections for 50 infections selected from early in a simulated outbreak ($\mu = 1$
193 mutation/genome/generation, $R=2$). Inset: receiver operating characteristic (ROC) for all possible genetic distance thresholds.
194 Optimal threshold shown as green dot (ROC) and dashed vertical line (distribution). (B) Estimated distribution of genetic
195 distances for linked and unlinked infections generated by the mutation rate method. Parameters and plots are as in (A).

196

197 *Mutation rate method*

198 Pathogen mutation rates can also be used to estimate the genetic distance distributions, especially when a
199 subsample of infections with known transmission histories is not available. If we assume that the number of
200 mutations between two linked infections is Poisson distributed around the mutation rate and that we know the
201 distribution of the number of generations between infections in the population, the probability of observing a
202 specific genetic distance (d) between the sequences from any two infected individuals linked by transmission is:

$$\frac{1}{\sum_{i=1}^{g_{link}} g(i)} \sum_{i=1}^{g_{link}} g(i) \cdot f(d; i \cdot \mu) \quad (3)$$

203 where $g(i)$ is the probability of observing i generations between infections, g_{link} is the maximum number of
204 generations between infections considered linked, $f(d; i \cdot \mu)$ is the probability of observing d mutations between
205 two infections separated by i generations, and μ is the mutation rate per genome per generation (see **Text S2**).

206 Similarly, the probability of observing a genetic distance d between two infections not linked by transmission is:

$$\frac{1}{\sum_{i=g_{link}+1}^{g_{max}} g(i)} \sum_{i=g_{link}+1}^{g_{max}} g(i) \cdot f(d; i \cdot \mu) \quad (4)$$

207 Where g_{max} is the maximum number of generations considered.

208 Determining the distribution of generations between infections is a non-trivial task (Dobrow 1996; Mahmoud and
209 Neining 2003; Salje et al. 2016), and depends on several factors, including the shape of the epidemic and the
210 period of time from which infections are sampled (**Fig S2**). In the examples included herein, we use simulations to
211 empirically approximate this distribution (see **Text S2**), but it is likely that adequate approximations can be
212 obtained by other means—or that more sophisticated approaches can be employed to directly estimate the necessary
213 genetic distance distributions (Worby et al. 2014).

214 Given the approximate generation distribution between infections, we calculated the genetic distance distributions
215 for linked and unlinked infections for the pathogen described above. The optimal genetic distance threshold for
216 distinguishing between linked and unlinked infections was 4 mutations (sensitivity=0.98, specificity=0.99) (**Fig**
217 **4B**). The empirical and mutation rate methods result in a similar, but not identical, optimal threshold for the
218 pathogen in this example, likely due to sparse sampling in the empirical case.

219 Additionally, we note that the clear threshold (and high sensitivity and specificity) observed here only occurs when
220 the mutation rate is high enough (and the reproductive number low enough) that a significant number of mutations
221 occur between infections considered linked (Campbell et al. 2018). For pathogens that do not meet these criteria, it
222 may not be possible to use genetic distance alone to distinguish between linked and unlinked infections (**Fig S3**).

223 **Methods**

224 **Outbreak simulations**

225 We used outbreak simulations to validate our approach. We simulated outbreaks using the ‘simOutbreak’ function
 226 implemented in the *outbreaker* R package (Jombart et al. 2014). For all simulations we assumed a large number of
 227 susceptible individuals in the population ($n_{\text{hosts}}=100,000$), a genome length of 1,000 nucleotides, and no
 228 importation events (single source outbreak). We also assumed every infected individual transmitted their infection
 229 exactly one time step after infection, and ran the simulation for the number of generations needed to achieve a final
 230 outbreak size of approximately 1,000 infections ($\ln(1,000)/\ln(R)$). After simulating the source population, we
 231 randomly selected a predetermined proportion of infections from that population.

232 For each sampling proportion, we simulated outbreaks over a variety of mutation rates and reproductive numbers.
 233 We allowed the mutation rate to vary between 0.0001–4 mutations per genome per generation, and allowed the
 234 reproductive number to vary between 1.3–18. We chose these ranges to encompass mutation rates and reproductive
 235 numbers observed in actual human pathogens. We divided each parameter range into 100 discrete values and ran
 236 simulations with all combinations of mutation rate and reproductive number, for a total of 10,000 simulations for
 237 each sampling proportion. We required simulated outbreaks to contain at least 100 and no more than 2000
 238 infections.

239 **Implementation**

240 Functions for calculating the necessary sample size based on a desired false discovery rate are implemented in the
 241 R package *phylosamp*, freely available at: <https://github.com/HopkinsIDD/phylosamp>. This package also includes
 242 functions for calculating the false discovery rate for a specific sample size or proportion, and functions to estimate
 243 the number of transmission pairs that will be observed given a sample size and a set of assumptions (e.g., multiple
 244 links and multiple transmissions, single link and single transmission, etc.). We also provide generation distributions
 245 for values of R between 1.3–18, derived from the simulations described in **Text S2**.

246 **Results**

247 **Method performance with known sensitivity and specificity**

248 We used simulated outbreaks to validate the relationship between sample size and false discovery rate using genetic
 249 distance as our linkage criteria. We subsampled each outbreak and, using the known transmission relationships and
 250 genetic distances between simulated infections, calculated the false discovery rate at each possible genetic distance
 251 threshold in the subsample (“simulated FDR”). For each simulation (before subsampling), we also calculated the
 252 actual specificity and sensitivity at every relevant genetic distance threshold. We used these values and the
 253 observed R_{pop} (roughly equal to one in most simulations) to then calculate the theoretical false discovery rate at a
 254 particular sampling proportion using **Equation 2**. We find that the theoretical false discovery rate is consistent with
 255 the simulated value for a wide array of pathogen mutation rates and reproductive numbers (**Fig 5**).




Figure 5: Predicted versus observed false discovery rate in outbreak simulations. Theoretical versus simulated false discovery rate (FDR) for each genetic distance threshold in 10,000 simulations of varying mutation rate and reproductive number. White line: smoothed conditional mean; grey dashed line: $y=x$ line. Increasing values of the sample size (M) are plotted in darker color; because the maximum outbreak size is fixed at 2000, the maximum sample size differs for each sampling proportion. Increasing both the sample size and proportion reduces bias and error, see **Table 2**.

Overall, the bias of our estimate of the false discovery rate approached zero for all sampling proportions. The average error was less than 4% in each case, decreasing significantly with increased sample size or proportion sampled (**Table 2**, **Table S1**). We note that special care should be taken with low sample sizes and low theoretical false discovery rates, as error rates can be particularly high in this range. Additionally, while our method is an unbiased estimator and overall correct in expectation, it is always possible for performance in a particular set of individuals sampled from a population to deviate substantially from expectation (for example, when a subsample happens to contain no true transmission pairs), particularly when sample sizes are low.

Table 2: Bias and error of calculated false discovery rate for simulations with fixed sampling proportion.


Bias	$\rho=0.10$	$\rho=0.25$	$\rho=0.50$	$\rho=0.75$	All ρ values	N
FDR=0.00-0.25	-0.0006	0.0045	0.0001	0.0036	0.0022	17,900
FDR=0.25-0.50	0.0044	0.0045	0.0009	0.0032	0.0032	31,633
FDR=0.50-0.75	0.0064	0.0039	0.0006	0.001	0.0029	51,069
FDR=0.75-1.00	0.0001	0.0001	<0.0001	<0.0001	0.0001	965,125
All FDR Values	0.0005	0.0005	0.0001	0.0002	0.0003	1,065,727
N	261,360	267,239	268,900	268,228	1,065,727	

Error	$\rho=0.10$	$\rho=0.25$	$\rho=0.50$	$\rho=0.75$	All ρ values	N
FDR=0.00-0.25	0.2135	0.1359	0.0799	0.0401	0.098	17,900
FDR=0.25-0.50	0.2751	0.1583	0.079	0.0416	0.1275	31,633
FDR=0.50-0.75	0.2057	0.0979	0.0478	0.0259	0.092	51,069
FDR=0.75-1.00	0.0155	0.0069	0.0035	0.002	0.007	965,125
All FDR Values	0.032	0.0181	0.0097	0.0052	0.0161	1,065,727
N	261,360	267,239	268,900	268,228	1,065,727	

To better understand why the error rate of our estimator increases as the false discovery rate decreases, we stratified the simulation data by the sensitivity and specificity given a particular genetic distance threshold. We found that the error is highest when sensitivity is low and specificity is high (**Fig S4A-B**), which occurs when a high genetic distance threshold is used. This combination often produces low false discovery rates, but is highly dependent on sampling (namely, if any true positives or false positives are sampled). This leads to highly variable simulated false discovery rates and consequently higher error rates. Unsurprisingly, this analysis also highlights that a discrete threshold like genetic distance produces a limited number of possible sensitivity and specificity combinations (**Fig S4C-D**). Therefore, obtaining reasonable estimates for these values in tandem is of key importance when using our method to estimate the false discovery rate of a phylogenetic study.

Method performance with estimated sensitivity and specificity

We repeated the false discovery rate comparison described above, but instead of using the actual sensitivity and specificity observed in each simulation, we calculated these parameters from the mutation rate used to generate that simulated outbreak (**Fig 6**). To reduce reliance on simulation data to calculate necessary parameters, we used $R_{pop} = I$, rather than the empirical value.




287

288 **Figure 6: Validation of mutation rate method to calculate sensitivity and specificity.** Theoretical versus simulated false
289 discovery rate (FDR) for each genetic distance threshold in 10,000 simulations of varying mutation rate and reproductive
290 number. White line: smoothed conditional mean; grey dashed line: $y=x$ line. Increasing values of the sample size (M) are plotted
291 in darker color; increasing both the sample size and proportion reduces bias and error, see **Tables S2** and **S3**.

292 Under this more realistic set of assumptions, we observe a slight bias, though overall values remain less than one
293 percent (**Table S2**, **Table S3**). However, while mean bias is very low on average, it is greater when the theoretical
294 false discovery rate is low, reaching nearly 8% for predicted false discovery rates less than 25%. Average error
295 rates were similarly slightly increased, but remained less than 4% overall.

296 Given that correct sensitivity and specificity values are an important component of calculating the theoretical false
297 discovery rate, we looked at the specific estimates for these parameters generated by our mutation rate method.
298 When considering only direct transmissions as linked (as we do throughout these simulations), **Equation 3**
299 simplifies to simply a poisson distribution around the mutation rate, resulting in highly accurate and precise
300 sensitivity estimates (**Fig S5**). However, we find that our estimates for specificity (**Fig S6**) have some positive bias
301 (and large error, particularly for low sample sizes). We hypothesized that inaccuracies in the estimated specificity
302 were due to the distribution of generations between infections used in our calculation; as discussed in **Approach**,
303 this is a non-trivial distribution that we estimated by averaging over many simulations (see **Text S2** for details).

To test this hypothesis, we used the actual distribution of generations between infections from each simulation in our calculation of specificity (sensitivity estimates are unaffected by this distribution when considering only direct transmissions, as described above). We find that this does in fact reduce bias in our specificity estimates (**Fig 7**) and leads to largely unbiased ($<2\%$) estimates of the false discovery rate, even at low theoretical false discovery rate values (**Fig S7, Table S4**).



309

Figure 7: Effect of the generation distribution on specificity of the linkage criteria. Theoretical versus simulated specificity for each genetic distance threshold in 10,000 simulations of varying mutation rate and reproductive number (proportion sampled = 0.75). White line: smoothed conditional mean; grey dashed line: $y=x$ line. Increasing values of the sample size (M) are plotted in darker color. (A) Theoretical sensitivity and specificity calculated using average distribution of generations between infections from simulations (see **Text S2**). (B) Theoretical sensitivity and specificity calculated using the actual distribution of generations between infections from that simulated outbreak.

316 Discussion

We have developed a mathematical framework for making informed sampling decisions in pathogen genome sequencing studies. Specifically, this framework allows for easy calculation of the relationship between the number or proportion of infections sampled during an outbreak and the ability of some phylogenetic or epidemiological criteria to correctly identify infections within this sample that are linked by direct transmission. Understanding this relationship is crucial to making correct inferences about pathogen transmission patterns, especially as genomic studies are becoming more feasible and widely used to answer both scientific and public health questions.

This framework is broadly applicable to a variety of phylogenetic or epidemiological approaches, as long as the sensitivity and specificity of the criteria can be approximated. With a basic understanding of the pathogen and the criteria being used, researchers can more effectively design studies that correctly identify transmission pairs with a known level of confidence. Additionally, this generalizable method (available as a free software, the R package *phylosamp*) provides a metric by which reviewers of these studies can evaluate their conclusions. We apply our method to simulated outbreaks using genetic distance as the linkage criteria and find that we can effectively estimate the false discovery rate for a variety of pathogen mutation rates, reproductive numbers, and relevant

330 genetic distance thresholds. It is important to note, however, that for a given sensitivity and specificity, there may
331 not always be a study design that achieves the desired false discovery rate.

332 Performance of the method presented depends on our ability to estimate the sensitivity and specificity of a
333 particular linkage criteria. While we present two methods for doing this—empirically and theoretically using the
334 mutation rate of the pathogen—implementing either in practice is not without challenges, and improved estimation
335 of these values may be a fruitful area for future research. For instance, the mutation rate based approach also
336 depends on the distribution of the number of generations of transmission between infections in the underlying
337 population. Although distributions derived from simulations (provided as part of the *phylosamp* package) provide a
338 reasonable proxy, estimates of sensitivity and specificity are much improved when using the exact generation
339 distribution, which currently can only be determined from complete knowledge of all transmission events. Further
340 research into all the factors affecting this distribution will be necessary to improve its estimation. Likewise, there
341 are challenges to the empirical approach, particularly for novel pathogens.

342 Better performance can likely be obtained by not restricting ourselves to genetic distance alone when determining a
343 linkage criteria. Genetic distance is easy to determine from sequence data, but this simple metric does not take into
344 account ancestral relationships or uncertainty around these relationships, and is limited to discrete mutational
345 changes. Applying more complex phylogenetic criteria may allow us to learn more about transmission
346 relationships, though there is a limit to the extent to which genetic data can be used to distinguish infections in fast-
347 spreading (or slow-mutating) pathogen outbreaks. There are several examples of outbreaks in which multiple
348 infected individuals have the same consensus viral genome (Campbell et al. 2018). In this case, incorporating
349 epidemiological data (e.g., location, time of symptom onset) may be important in determining which infections are
350 unlikely to be linked. Doing so is part of a larger effort to better integrate epidemiological and genomic data into
351 pathogen transmission studies (Morelli et al. 2012; Ypma et al. 2012; Jombart et al. 2014; Klinkenberg et al. 2017).

352 While in this manuscript we have focused on direct transmission pairs, our framework is designed to be extensible
353 to alternative definitions of linkage; for example, infections connected within a specified number of transmission
354 events. Expanding the definition of linkage to include such indirect transmissions has a number of useful
355 applications in outbreak research, such as identifying and connecting transmission clusters. This method could also
356 be extended to more complex direct transmission relationships, for example when within-host evolution results in
357 the existence of viral quasispecies within infected individuals, each of which has some potential of being
358 transmitted. In all of these scenarios, it is equally important to understand the sample size needed to make the
359 desired inferences.

360 We hope that this work represents a step towards developing a larger theory of study design for making inferences
361 from pathogen sequence data, but recognize it is only a step. The focus of this paper is sample size, but which
362 infections are sampled may be equally important (Stack et al. 2010; de Silva et al. 2012; Hall et al. 2016). For
363 example, understanding routes of direct transmission may require dense sampling of a small group of highly-
364 connected individuals, while understanding general transmission trends over the course of a geographically-

365 dispersed outbreak may require us to sample broadly over space and time. Additionally, the goal of linking
 366 infections is seldom the linkages themselves, but the larger inferences about risk and transmission derived from
 367 those linkages. Adapting the techniques here to more directly link sample size calculations to these outcomes is an
 368 important next step.

369

370 **Acknowledgements**

371 We thank Stuart Ray for his insightful comments on the manuscript. Funding was provided by Bill and Melinda
 372 Gates Foundation OPP1195157 (S.W. and J.L.).

References

- Campbell F, Strang C, Ferguson N, Cori A, Jombart T. 2018. When are pathogen genome sequences informative of transmission events? *PLoS Pathog.* 14:e1006885.
- Carroll MW, Matthews DA, Hiscox JA, Elmore MJ, Pollakis G, Rambaut A, Hewson R, García-Dorival I, Bore JA, Koundouno R, et al. 2015. Temporal and spatial analysis of the 2014-2015 Ebola virus outbreak in West Africa. *Nature* 524:97–101.
- Dobrow RP. 1996. On the distribution of distances in recursive trees. *J. Appl. Probab.* 33:749–757.
- Farhat MR, Shapiro BJ, Sheppard SK, Colijn C, Murray M. 2014. A phylogeny-based sampling strategy and power calculator informs genome-wide associations study design for microbial pathogens. *Genome Med.* 6:101.
- Frost SDW, Pybus OG, Gog JR, Viboud C, Bonhoeffer S, Bedford T. 2015. Eight challenges in phylodynamic inference. *Epidemics* 10:88–92.
- Gardy JL, Johnston JC, Ho Sui SJ, Cook VJ, Shah L, Brodtkin E, Rempel S, Moore R, Zhao Y, Holt R, et al. 2011. Whole-genome sequencing and social-network analysis of a tuberculosis outbreak. *N. Engl. J. Med.* 364:730–739.
- Gonzalez-Reiche AS, Hernandez MM, Sullivan MJ, Ciferri B, Alshammery H, Obla A, Fabre S, Kleiner G, Polanco J, Khan Z, et al. 2020. Introductions and early spread of SARS-CoV-2 in the New York City area. *Science* [Internet]. Available from: <http://dx.doi.org/10.1126/science.abc1917>
- Grabowski MK, Lessler J. 2017. Phylogenetic insights into age-disparate partnerships and HIV. *Lancet HIV* 4:e8–e9.
- Hall MD, Woolhouse MEJ, Rambaut A. 2016. The effects of sampling strategy on the quality of reconstruction of viral population dynamics using Bayesian skyline family coalescent methods: A simulation study. *Virus Evol* 2:vew003.
- Jackson BR, Tarr C, Strain E, Jackson KA, Conrad A, Carleton H, Katz LS, Stroika S, Gould LH, Mody RK, et al. 2016. Implementation of Nationwide Real-time Whole-genome Sequencing to Enhance Listeriosis Outbreak Detection and Investigation. *Clin. Infect. Dis.* 63:380–386.
- Jombart T, Cori A, Didelot X, Cauchemez S, Fraser C, Ferguson N. 2014. Bayesian reconstruction of disease outbreaks by combining epidemiologic and genomic data. *PLoS Comput. Biol.* 10:e1003457.
- Kelly BJ, Gross R, Bittinger K, Sherrill-Mix S, Lewis JD, Collman RG, Bushman FD, Li H. 2015. Power and sample-size estimation for microbiome studies using pairwise distances and PERMANOVA. *Bioinformatics* 31:2461–2468.
- Klinkenberg D, Backer JA, Didelot X, Colijn C, Wallinga J. 2017. Simultaneous inference of phylogenetic and transmission trees in infectious disease outbreaks. *PLoS Comput. Biol.* 13:e1005495.
- Lei F, Shi W. 2011. Prospective of Genomics in Revealing Transmission, Reassortment and Evolution of Wildlife-Borne Avian Influenza A (H5N1) Viruses. *Curr. Genomics* 12:466–474.
- Liu X. 2012. Classification accuracy and cut point selection. *Stat. Med.* 31:2676–2686.
- Mahmoud HM, Neininger R. 2003. Distribution of distances in random binary search trees. *Ann. Appl. Probab.* 13:253–276.
- Mavian C, Marini S, Manes C, Capua I, Prosperi M, Salemi M. 2020. Regaining perspective on SARS-CoV-2 molecular tracing and its implications. *medRxiv:2020.03.16.20034470*.
- Morelli MJ, Thébaud G, Chadœuf J, King DP, Haydon DT, Soubeyrand S. 2012. A Bayesian inference framework to reconstruct transmission trees using epidemiological and genetic data. *PLoS Comput. Biol.* 8:e1002768.
- Neher RA, Bedford T. 2018. Real-Time Analysis and Visualization of Pathogen Sequence Data. *J. Clin. Microbiol.* [Internet] 56. Available from: <http://dx.doi.org/10.1128/JCM.00480-18>
- Nelson MI, Simonsen L, Viboud C, Miller MA, Holmes EC. 2007. Phylogenetic analysis reveals the global migration of seasonal influenza A viruses. *PLoS Pathog.* 3:1220–1228.

- Network HPT, Others. 2013. HPTN 071: population effects of antiretroviral therapy to reduce HIV transmission (PopART): a cluster-randomized trial of the impact of a combination prevention package on population-level HIV incidence in Zambia and South Africa.
- Park DJ, Dudas G, Wohl S, Goba A, Whitmer SLM, Andersen KG, Sealfon RS, Ladner JT, Kugelman JR, Matranga CB, et al. 2015. Ebola Virus Epidemiology, Transmission, and Evolution during Seven Months in Sierra Leone. *Cell* 161:1516–1526.
- Perkins NJ, Schisterman EF. 2006. The inconsistency of “optimal” cutpoints obtained using two criteria based on the receiver operating characteristic curve. *Am. J. Epidemiol.* 163:670–675.
- Quick J, Loman NJ, Duraffour S, Simpson JT, Severi E, Cowley L, Bore JA, Koundouno R, Dudas G, Mikhail A, et al. 2016. Real-time, portable genome sequencing for Ebola surveillance. *Nature* 530:228–232.
- Ratmann O, Kagaayi J, Hall M, Golubchick T, Kigozi G, Xi X, Wymant C, Nakigozi G, Abeler-Dörner L, Bonsall D, et al. 2020. Quantifying HIV transmission flow between high-prevalence hotspots and surrounding communities: a population-based study in Rakai, Uganda. *Lancet HIV* 7:e173–e183.
- Salje H, Cummings DAT, Lessler J. 2016. Estimating infectious disease transmission distances using the overall distribution of cases. *Epidemics* 17:10–18.
- Salje H, Lessler J, Endy TP, Curriero FC, Gibbons RV, Nisalak A, Nimmannitya S, Kalayanaroj S, Jarman RG, Thomas SJ, et al. 2012. Revealing the microscale spatial signature of dengue transmission and immunity in an urban population. *Proc. Natl. Acad. Sci. U. S. A.* 109:9535–9538.
- de Silva E, Ferguson NM, Fraser C. 2012. Inferring pandemic growth rates from sequence data. *J. R. Soc. Interface* 9:1797–1808.
- Snider CJ, Diop OM, Burns CC, Tangermann RH, Wassilak SGF. 2016. Surveillance Systems to Track Progress Toward Polio Eradication--Worldwide, 2014-2015. *MMWR Morb. Mortal. Wkly. Rep.* 65:346–351.
- Stack JC, Welch JD, Ferrari MJ, Shapiro BU, Grenfell BT. 2010. Protocols for sampling viral sequences to study epidemic dynamics. *J. R. Soc. Interface* 7:1119–1127.
- Team RC, Others. 2013. R: A language and environment for statistical computing. Available from: <http://finzi.psych.upenn.edu/R/library/dplR/doc/intro-dplR.pdf>
- Thézé J, Li T, du Plessis L, Bouquet J, Kraemer MUG, Somasekar S, Yu G, de Cesare M, Balmaseda A, Kuan G, et al. 2018. Genomic Epidemiology Reconstructs the Introduction and Spread of Zika Virus in Central America and Mexico. *Cell Host Microbe* 23:855–864.e7.
- Volz EM, Frost SDW. 2013. Inferring the source of transmission with phylogenetic data. *PLoS Comput. Biol.* 9:e1003397.
- Weill F-X, Domman D, Njamkepo E, Almesbahi AA, Naji M, Nasher SS, Rakesh A, Assiri AM, Sharma NC, Kariuki S, et al. 2019. Genomic insights into the 2016-2017 cholera epidemic in Yemen. *Nature* 565:230–233.
- Worby CJ, Chang H-H, Hanage WP, Lipsitch M. 2014. The distribution of pairwise genetic distances: a tool for investigating disease transmission. *Genetics* 198:1395–1404.
- Youden WJ. 1950. Index for rating diagnostic tests. *Cancer* 3:32–35.
- Ypma RJF, Bataille AMA, Stegeman A, Koch G, Wallinga J, van Ballegooijen WM. 2012. Unravelling transmission trees of infectious diseases by combining genetic and epidemiological data. *Proc. Biol. Sci.* 279:444–450.
- Zou KH, Yu C-R, Liu K, Carlsson MO, Cabrera J. 2013. Optimal thresholds by maximizing or minimizing various metrics via ROC-type analysis. *Acad. Radiol.* 20:807–815.