# 1 Full Title

2	The Mean Unfulfilled Lifespan (MUL):
3	A new indicator of the impact of mortality shocks on the individual lifespan,
4	with application to global 2020 quarterly mortality from COVID-19
5	
6	Short Title
7	The Mean Unfulfilled Lifespan:
8	A new indicator of the impact of mortality shocks on the individual lifespan
9	
10	Updated Version, 2/18/2020
11	Original Manuscript Submitted 8/9/2020, Published 8/13/2020 at:
12	https://www.medrxiv.org/content/10.1101/2020.08.09.20171264v1
13	
14	Patrick Heuveline, Ph.D.
15	California Center for Population Research (CCPR)
16	University of California, Los Angeles (UCLA)
17	
18	Corresponding author: Patrick Heuveline <heuveline@soc.ucla.edu></heuveline@soc.ucla.edu>
19 20	

# 21 Abstract

Declines in period life expectancy at birth (PLEB) provide intuitive indicators of the impact of a cause of death on the individual lifespan. Derived under the assumption that future mortality conditions will remain indefinitely those observed during a reference period, however, the intuitive interpretation of a PLEB becomes problematic when that period conditions reflect a temporary mortality "shock", resulting from a natural disaster or the diffusion of a new epidemic in the population for instance.

Rather than to make assumptions about future mortality, I propose measuring the difference between a period average age at death and the average expected age at death of the same individuals (death cohort): the Mean Unfulfilled Lifespan (MUL). For fine-grained tracking of the mortality impact of an epidemic, I also provide an empirical shortcut to MUL estimation for small areas or short periods.

33 For illustration, quarterly MUL values in 2020 are derived from estimates of COVID-19 34 deaths in 159 national populations and 122 sub-national populations in Italy, Mexico, Spain and 35 the US. The highest quarterly values in national populations are obtained for Ecuador (5.12) 36 years, second quarter) and Peru (4.56 years, third quarter) and, in sub-national populations, for 37 New York (5.52 years), New Jersey (5.56 years, second quarter) and Baja California (5.19 years, 38 fourth quarter). Using a seven-day rolling window, the empirical shortcut suggests the MUL 39 peaked at 9.12 years in Madrid, 9.20 years in New York, and 9.15 years in Baja California, and 40 in Guayas (Ecuador) it even reached 12.6 years for the entire month of April. 41 Based on reported COVID-19 deaths that might substantially underestimate overall

42 mortality change in affected populations, these results nonetheless illustrate how the MUL tracks

- 43 the mortality impact of the pandemic, or any mortality shock, retaining the intuitive metric of
- 44 differences in PLEB, without their problematic underlying assumptions.

45

### 47 Introduction

For months, the numbers of deaths from the novel coronavirus disease 2019 (COVID-19) have become part of the daily news cycle the world over. Even when related to the population size in deaths per capita ratio, however, these numbers do not really provide any intuition for the magnitude nor the dynamics of the pandemic. Quite useful for between-population comparisons, the age standardization of these ratios does not make them more easily interpretable.

The period life expectancy at birth (PLEB) is probably the most readily interpretable of the period indicators mortality. Translating a number of deaths from a given cause into its impact on PLEB involves multiple steps but is fairly straightforward.<sup>1, 2</sup> Unfortunately, the intuitive appeal of the PLEB, its interpretation as a measure of the individual lifespan, derives from the assumption that period mortality conditions will continue to prevail indefinitely. Declines in PLEB induced by a relatively rapid and likely temporary increase in mortality, such as currently experienced with the COVID-19 pandemic, can thus provide misleading indicators of changes in the individual lifespan.<sup>3</sup>

60 The assumption is even more problematic when PLEB or changes therein are estimated for 61 smaller populations and shorter periods of time. Tracking the pandemic at a finer-grained geographical 62 and temporal scale undoubtedly provides better insights on the pandemic than annual, national averages.<sup>4</sup>, <sup>5, 6</sup> But while the assumption underlying an annual PLEB estimate—that mortality conditions of a given 63 64 year will be repeated year after year in the future—may seem unlikely, the seasonality of mortality makes 65 the indefinite repetition of the mortality conditions in any fraction of a year not just unlikely but impossible.<sup>7</sup> When referring to mortality conditions not only in a short period but also in a small area, the 66 assumption on which PLEB estimates are build resort to a Groundhog-Day-like<sup>8</sup> time loop repeating itself 67 68 in a small area from which individuals are unable to leave.

For a both interpretable and scalable, over space and time, measure of the impact of changing
mortality conditions on the individual lifespan, I suggest an alternative to the reduction in PLEB, the
Mean Unfulfilled Lifespan (MUL). Making no assumption about future mortality, the MUL translates

72 past changes in mortality into average difference in the length of lived lives. That difference is obtained 73 by comparing the actual average age at death during a given period and the expected ages at death of the 74 same individuals in the absence of mortality changes, whether induced by a specific cause of death or by 75 an event affecting multiple causes of death. The MUL can thus be estimated for populations of any size 76 and for periods of any length, from data on excess deaths or cause-specific deaths, as illustrated here with 77 COVID-19 mortality data from 159 national populations and 122 sub-national populations in Italy, 78 Mexico, Spain and the USA, for each sex and each quarter in 2020. The MUL also equals the product of 79 (1) the proportion of deaths in the population during a reference period that are due to a specific cause or 80 event and (2) the average reduction in length of life among individuals who died from that specific cause 81 or due to that specific event in the population during the reference period. In the case of COVID-19, I 82 show that for a given population the value of that average reduction only changes very slowly over time, 83 providing an easy short-cut for fine-grained tracking of the pandemic. 84 *Conceptual Detour* 85 Assessing PLEB reductions induced by a specific cause or event requires two period life tables, one 86 representing the prevailing mortality conditions and another one representing the counterfactual mortality 87 conditions expected in the absence of that cause or event. The assessment involves a relatively copious 88 amount of life table manipulations, but decades ago Nathan Keyfitz provided most useful insights as to 89 what these manipulations boil down to. Considering the related issue of estimating the increase in PLEB 90 brought by the permanent elimination of a cause of death, he summarized that the increase "depends on the average time that elapses before the persons rescued will die of some other cause."9 Conversely, the 91 92 decrease induced by a new cause of death depends on the average time that would have elapsed before the 93 persons who died from the new cause would have died from other causes.

This average time can be derived from the synthetic cohort approach modelled in the period life table where each death at age *a* from a cause *C*,  $d^{C}(a)$ , reduces the number of person-years lived by the life expectancy at age *a* in the absence of that new cause,  $e^{o-C}(a)$ . This assumes that persons dying from the new cause would have had the same life expectancy in the absence of that cause as same-age persons

98 who survived that cause. This common assumption may appear unlikely, and interactions between causes 99 of death can be incorporated instead, but the data requirements are substantial. Under the common 100 assumption, the difference in PLEB is thus the average over all members of the synthetic cohort,  $l_0$  (the 101 radix of the life table), of the difference in person-years lived by cohort members:

$$\Delta PLEB = \frac{1}{l_0} \cdot \int_0^{\omega} (d^C(a) \cdot e^{o-C}(a)) da$$

Keyfitz' insight relates to the concept of "potential years of life lost"<sup>10</sup> developed a couple of decades earlier still. The initial approach, designed to measure premature mortality, compared ages at death to a fixed value (70 or 75 years).<sup>11</sup> This approach is not suited to study cause of deaths at older ages since deaths at ages above the fixed value are not considered.<sup>12</sup> In burden-of-disease assessments, it has become customary to estimate Years of Life Lost (YLL) as:

$$YLL = \int_0^{\omega} (D^C(a) \cdot e^{o*}(a)) da$$

107 where  $D^{C}(a)$  is the number of deaths from a certain cause *C* at age *a* observed in the population during a 108 reference period and  $e^{o^{*}}(a)$  is life expectancy at age *a* in a counterfactual life table. YLL to COVID-19 109 have been estimated using this approach.<sup>13, 14, 15</sup>

110 Three differences between the two above equations can be observed. First, YLL are estimated 111 from actual numbers of deaths by age rather than from numbers of life table decrements. This implies that 112 the YLL estimate is sensitive to the age distribution of the population, as in turn it affects the distribution of deaths by age. While the PLEB is not an age-standardized measure.<sup>16</sup> it equals the inverse of the 113 114 "stationary" death rate, that is, a weighted average of the period age-specific death rates with weights 115 derived from these death rates through life table construction. Using these internally-derived weights 116 rather than an external, standard age distribution, the stationary death rate and PLEB are independent of 117 the actual age composition of the population. This relative advantage of the difference in PLEB comes at 118 the cost of using a "stationary" age distribution of deaths, however, represented by the life table 119 decrements that result from indefinitely subjecting the population to the mortality conditions of the

period. As discussed in the introduction, this assumption is precisely what is problematic for studying the impact of a mortality shock or an emerging disease such as COVID-19. Moreover, the actual distribution of deaths can be used in a population of any size and periods of any length, allowing for the mortality impact to be tracked on short temporal and small spatial scales for which interpreting differences in PLEB hardly makes sense.

The second difference refers to the counterfactual life expectancies. In global burden-of-disease assessments, a universal life table representing optimal survival conditions is typically used. This has the advantage of making YLL for different populations additive allowing for the derivation of a global estimate of YLL due to a cause by simple summation. However, using a universal life table may misrepresent the actual gains from averting a death in a specific population.

130 The last difference concerns the denominator, or lack thereof in YLL. The use of a denominator 131 in the difference in PLEB allows for relating a total number of years in a population, measured by YLL, 132 to a number of persons, and thus for a more intuitive interpretation as an average difference in years lived 133 per person. Two ratios involving YLL can be found in the literature. First, the average YLL (AYLL) 134 relates YLL to all the deaths from that cause in the population during the reference period, *D<sup>C</sup>*:

$$AYLL = \frac{\int_{0}^{\omega} (D^{C}(a) \cdot e^{0^{*}}(a)) da}{D^{C}} = \int_{0}^{\omega} \left( \frac{D^{C}(a)}{\int_{0}^{\omega} D^{C}(a) da} \right) \cdot e^{0^{*}}(a) da$$

135 The AYLL thus represents the average (universal) life expectancy left to population members who died 136 from the specific cause during a given period. On the one hand, it is a coherent measure as its 137 denominator includes all the deaths that contribute to lost years in the numerator. On the other hand, it is 138 only a function of the distribution of deaths by age, irrespective of the prevalence of that cause of death. 139 The AYLL thus cannot provide a measure of the intensity of a mortality shock. The second measure is 140 YLL per capita. While it does depend on the prevalence of the different causes of death, it is a less 141 coherent measure as it includes in the denominator all the individuals in the population, including many 142 that survived the mortality shock and do not contribute to YLL in the numerator. In turn, this complicates 143 providing a precise interpretation for the value of YLL per capita.

144 Considering the advantages and limitations of the extant measures, I propose to add one measure 145 of the mortality impact of a cause of death or an event on the individual lifespan. This measure, the Mean 146 Unfulfilled Lifespan (MUL), is intended for situations where the underlying assumptions of PLEB might 147 be implausible, and thus based, as the estimate of YLL, on actual numbers of deaths in a population 148 during a period rather than on life table decrements. To retain its intuitive interpretation, however, the 149 MUL is structured like the difference in PLEB as summarized by Keyfitz and, using counterfactual life 150 expectancies representing the mortality conditions in the population of interest, similarly expressed as an 151 average difference in person-years lived per person. Since the life table radix,  $l_0$ , equals the sum of all 152 decrements at all ages, the structural equivalence is maintained by defining the MUL as:

$$MUL \stackrel{\text{def}}{=} \frac{1}{D} \cdot \int_0^{\omega} (D^C(a) \cdot e^{o-C}(a)) da$$

153 where *D* is the total number of deaths (from all causes at all ages) during the reference period.

154 This intuitive interpretation of the MUL can be derived by rewriting this defining equation as:

$$MUL = \frac{\int_{0}^{\omega} \left( \left( D^{c}(a) \cdot \left( a + e^{o^{-c}(a)} - a \right) \right) + \left( D^{-c}(a) \cdot \left( a - a \right) \right) \right) da}{\int_{0}^{\omega} \left( D^{c}(a) + D^{-c}(a) \right) da}$$
$$= \frac{\int_{0}^{\omega} \left( \left( D^{c}(a) \cdot \left( a + e^{o^{-c}(a)} \right) \right) + \left( D^{-c}(a) \cdot a \right) \right) da}{\int_{0}^{\omega} \left( D^{c}(a) + D^{-c}(a) \right) da} - \frac{\int_{0}^{\omega} \left( \left( D^{c}(a) + D^{-c}(a) \right) \cdot a \right) da}{\int_{0}^{\omega} \left( D^{c}(a) + D^{-c}(a) \right) da}$$

155 where  $D^{-C}(a)$  is the number of deaths from all causes but C at age a observed in the population during the 156 reference period. The second term represents the average age at death in a given period. If we assume that 157 individuals who die of other causes than C die at the same age as they would have in the absence of cause 158 C (no indirect effect of cause C on other causes of death), and that individuals who die of cause C at age a 159 would have otherwise lived to age a+e(a), the first term represents the average expected age at death of 160 the same individuals in the absence of cause C. The MUL is thus the difference between the average age 161 at death in the population during a given period and the average expected age at death of the same 162 individuals in the absence of cause C.

163 The assumption that cause C has no indirect effect on other causes of death is actually not 164 required. As will be shown below, one can also derive the MUL from data on all deaths, and 165 distinguishing between deaths that, based on counterfactual "benchmark" mortality conditions, were 166 expected to occur in that period and those that were not, use the latter number of "excess" deaths instead 167 of deaths from a specific cause. Finally, note that the MUL differs from changes in average ages at death 168 across periods, which can be readily measured but may actually be positive with the emergence of a new 169 cause of death if that cause affects people who are older on average that those dying from other causes. 170 To sum up this conceptual detour, the MUL thus complements existing indicators of the impact on the 171 individual lifespan of a cause of death by providing a measure of the average potential years of life lost to 172 a specific cause of death, or to any mortality shock, for all population members dying in a certain period, 173 regardless of their cause of death.

174 Empirical Shortcut

175 Calculating the MUL for small geographical areas and short periods is not conceptually problematic since 176 it captures the actual length of lives that ended there and then, unlike differences in life expectancies that 177 require assumptions about future conditions. The demand on data (including a separate counterfactual life 178 table for each population of interest) is substantial, however, and the life table manipulations are not 179 particularly straightforward.

#### 180 To simplify the estimation, the MUL can be rewritten as:

$$MUL = \frac{D^{C}}{D} \cdot \frac{\int_{0}^{\omega} D^{C}(a) \cdot e^{o-C}(a) da}{D^{C}} = \frac{D^{C}}{D} \cdot PAYLL$$

181 The second term is similar to the AYLL in burden-of-disease assessments, but again based on population-182 specific, counterfactual life expectancies instead of universal ones, and to underscore the difference is 183 termed here the Population AYLL (PAYLL). In a given population, the PAYLL is a weighted average of 184 counterfactual life expectancies that are estimated from prior conditions and do not change over time. 185 Meanwhile, the weights are the ratios:

$$\frac{D^{C}(a)}{D^{C}} = \frac{M^{C}(a) \cdot N(a)}{M^{C} \cdot N} = \frac{M^{C}(a)}{M^{C}} \cdot \frac{N(a)}{N}$$

186 where  $M^{C}$  and  $M^{C}(a)$  are the all-age death rate and the death rate at age *a* from a specific cause, and *N* and 187 *N*(*a*) are the total population size and number of individuals at age *a* in the population. These weights 188 should also be expected to vary little within short periods because their values depend on the population 189 composition, *N*(*a*)/*N*, and on the age pattern of cause-specific death rates,  $M^{C}(a)/M^{C}$ , both of which should 190 vary little within short periods. To confirm this, Figure 1 compares the distribution of provisional 191 COVID-19 death counts in the USA at four points in time: on May 13 (one of the earliest dates for which

this distribution is available) and on July 1<sup>st</sup>, September 30<sup>th</sup> and December 30<sup>th</sup>.







- 195 (54,860 deaths), 7/1 (112,223 deaths), 9/30 (194,087 deaths) and 12/30 (301,671 deaths)
- 196 Source: Centers for Disease Control and Prevention (CDC)

197 This suggests that the value of PAYLL can be expected to only change slowly over time and to be

198 relatively close across populations with similar life expectancies and population compositions. In

particular, MUL values for a sub-population or during a sub-period can be approximated as the product of the PAYLL for the whole population or the entire period and the variable all-age ratio of deaths from a specific cause to all deaths,  $D^{C}/D$ , in the sub-population or during that sub-period. As noted above, this can be readily extended to any mortality shock, with data on excess deaths and all-cause deaths.

203 Materials and Methods

The equations defining YLL and the MUL look deceivingly simple. To implement them, one has to apply estimates of life expectancies, which refer to exact ages, to numbers of deaths that are available or can be estimated only for age intervals. The varying value of life expectancy on a closed age interval is typically approximated by linear interpolation,<sup>17</sup> in which case the contribution of the closed age interval between ages *x* and *x*+*n* to the MUL equals:

$$\frac{\int_0^n D^C(x+a) \cdot e^{o-C}(x+a) da}{D} = \frac{n D_x^C}{D} \cdot e^{o-C} (x+n a_x^C)$$

where  ${}_{n}D_{x}{}^{C}$  is the number of individuals between ages *x* and *x*+*n* who died of cause *C* during the reference period and  ${}_{n}a_{x}{}^{C}$  is the average number of years lived after age *x* by these individuals. In turn, life expectancy at exact age  $x+{}_{n}a_{x}{}^{C}$  can be derived by linear interpolation between the values of life expectancy at ages *x* and *x*+*n*.

213 The linear interpolation is more problematic for wider the age intervals and for older age groups. In the 2018 US life table for males for instance.<sup>18</sup> life expectancy declines by 2.9 years between ages 75 214 215 and 80 and by 2.4 years between ages 80 and 85. In this case, the linear approximation over-estimate life 216 expectancy in the interval. The average number of years lived after age 75 by individuals dying between 217 ages 75 and 85 is 5.4 years, and linear interpolation would yield a life expectancy at age 80.4 years of 8.5 218 years, whereas life expectancy has already dropped to 8.4 years at age 80. This upward bias is particularly 219 undesirable in the case of COVID-19 deaths at those ages because COVID-19 victims are more likely to 220 suffer from other long-term conditions that more likely reduced their life expectancies compared to same-221 age individuals. Even more problematic with linear interpolation is the open-ended interval, as it requires 222 that an arbitrary upper age limit be set. Unfortunately, these couple of issues may concern a large share of

223 COVID-19 deaths: as shown in Figure 1, close to 60% of US deaths are above age 75 years and reported 224 in just one ten-year closed interval (75 to 84 years) and one open age interval (over 85 years). 225 In this respect, working with all deaths in a given period rather than with deaths from a specific 226 cause of interest presents an important empirical advantage in addition to the benefit of assessing both the 227 direct and the indirect effects of that cause of death. Using all deaths in a closed age interval between ages 228 x and x+n,  $_{D}D_{x}$ , one may first calculate the contribution to the MUL of the deaths that were expected to 229 occur under the counterfactual mortality conditions,  ${}_{n}D_{x}^{-C}$ . For these deaths, the difference in length of life 230 averages to the difference between the average number of years lived in the age interval under the 231 prevailing conditions,  ${}_{n}a_{x}$ , and under the counterfactual conditions,  ${}_{n}a_{x}^{-C}$ . The advantage of this approach 232 is that setting an arbitrary upper value for the open-ended age interval becomes unnecessary because, 233 having already reached age N, individuals who contributed to the number of deaths in the open age 234 interval would all have been expected to die in the same open-ended age interval under the counterfactual 235 mortality conditions, albeit possibly not in that period. For all the open-interval deaths over age  $N, D_{N+}$ , 236 the difference in length of life thus averages to the difference between life expectancy at age N under the 237 prevailing conditions,  $e^{o}(N)$ , and under the counterfactual conditions,  $e^{o-C}(N)$ .

This approach limits the issue of estimating the average reduction in length of lived lives on closed age intervals from values of life expectancies that vary on these age intervals to "excess" deaths,  ${}_{n}D_{x} - {}_{n}D_{x}^{-C}$ . An alternative to linear interpolation derives from the fact that an individual's expected length of life ( $e^{o}(x)+x$ ) gradually increases with age *x*. Life expectancy at any age is thus larger than the difference between life expectancy at an earlier age and the difference between the two ages:

$$\int_{0}^{n} D(x+a) \cdot e^{o}(x+a) da > \int_{0}^{n} D(x+a) \cdot (e^{o}(x)-a) da = {}_{n} D_{x} \cdot (e^{o}(x)-{}_{n}a_{x})$$

243 Applying this approximation to excess deaths on any closed age interval will induce some

underestimation of the average life expectancy of individuals dying on the interval, which appears

- 245 preferable in this case to the overestimation induced by linear interpolation.
- Adding the contributions of the different types of deaths yields:

$$MUL \cong \frac{1}{D} \cdot \left( \sum_{x=0,n}^{N-n} {}_{n} D_{x}^{-C} \cdot \left( {}_{n} a_{x}^{-C} - {}_{n} a_{x} \right) + \left( D_{N+\cdot} \left( e^{o-C}(N) - e^{o}(N) \right) \right) + \sum_{x=0,n}^{N-n} \left( {}_{n} D_{x} - {}_{n} D_{x}^{-C} \right) \cdot \left( e_{x}^{o-C} - {}_{n} a_{x} \right) \right)$$

Rearranging the sums corresponding to the closed age intervals and using the multiple-decrement life table relationship  $e^{o}(N)/e^{o-C}(N) = D^{-C}{}_{N+}/D_{N+}$ , this can be rewritten as:

$$MUL \cong \frac{1}{D} \left( \sum_{x=0,n}^{N-n} \left( {}_{n}D_{x} \cdot \left( {}_{n}a_{x}^{-C} - {}_{n}a_{x} \right) + \left( {}_{n}D_{x} - {}_{n}D_{x}^{-C} \right) \cdot \left( e_{x}^{o-C} - {}_{n}a_{x}^{-C} \right) \right) + \left( \left( D_{N+} - D_{N+}^{-C} \right) \cdot e_{N}^{o-C} \right) \right)$$

Using this approximation to estimate the impact of COVID-19 on the individual lifespan in each quarter of 2020 first requires a life table representing survival conditions in 2020 in the absence of COVID-19 whose values of  $e_x^{o-C}$  and  $_na_x^{-C}$  can be used. Combined with the number of individuals by sex and age-group, the life table values of age-specific death rates,  $_nm_x^{-C}$ , then provide the expected numbers of deaths  $_nD_x^{-C}$  in the absence of COVID-19. Population data and life table functions for countries were obtained from the UN Population Division.<sup>19</sup> Corresponding data for sub-populations in Italy, Spain, and the US were obtained from national statistical agencies.<sup>20, 21, 22, 23</sup>

256 New life tables representing actual mortality conditions (with COVID-19) in each quarter must 257 then be derived to calculate the corresponding values of  $na_x$ . The construction of these life tables requires 258 the quarterly numbers of deaths by sex and age-group. In countries where vital statistics are incomplete 259 not available yet, but estimates of COVID-19 deaths are available, the total number of deaths,  $_nD_x$ , can be 260 obtained by adding these estimates (through a multi-decrement life table to adjust for competing risks of 261 deaths, which typically assumies no indirect effects on other causes of death)<sup>24</sup> to the expected numbers of deaths in the absence of Covid-19 ( $_nD_x^{-C}$ ). When COVID-19 estimates are not broken down by sex and 262 263 age-group, an alternative is to use a reference set of age-and-sex death rates from COVID-19 from another population for which these rates are deemed reliable.<sup>25</sup> Centers for Disease Control and 264 265 Prevention (CDC) data provided the reference set of age-and-sex death rates from COVID-19.<sup>26</sup> Estimates

- of COVID-19 deaths by March 31, June 30, September 30 and December 31, 2020 were taken from the
- 267 IHME.<sup>27</sup> All of these data were downloaded from institutional websites.
- 268 Results
- 269 Figure 2 shows MUL values for selected national and sub-national populations in each of the four
- 270 quarters of 2020 (for both sexes combined). First-quarter MUL values illustrate substantial impacts of
- 271 COVID-19 on longevity already in parts of Spain (Madrid, 3.26 years) and Italy (Lombardy 2.76 years).
- In the second quarter, the impact continued to increase in these two nations, with the MUL reaching 4.32
- 273 years in Catalonia for instance, but also in other European nations such as Belgium (2.80 years).
- However, the second-quarter MUL values reached their highest values in parts of the USA (New Jersey,
- 5.56 years, and New York, 5.52 years). Very high second-quarter values are also estimated for parts of
- 276 South America (Ecuador, 5.12 years) and Mexico (Baja California, 5.10 years).



277

Fig 2: Quarterly Mean Unfulfilled Lifespan (MUL) for both sexes, in years, selected populations
(populations with 5,000 or more COVID-19 deaths by January 1<sup>st</sup>, 2021, and the highest annual MUL
values for 2020).

In the third quarter, MUL values continue to increase or remain high in parts of South America (Peru, 4.56 years) and Mexico (Baja California, 4.39 years, and Nuevo Leon, 4.03 years), whereas they were declining in Europe and the USA. In the last quarter of 2020, MUL values decline in South America, whereas the highest values remain for parts of Mexico (Baja California, 5.19 years, and Mexico City, 4.64 years). Fourth-quarter MUL values also rebound in parts of Europe (Belgium, 3.00 years) and the USA (Illinois, 3.07 years). MUL values for men are typically higher than for women, and the average age at death of men who died in New Jersey during the second quarter was nearly 6 years younger than

their average expected age at death without COVID-19 (5.97 years, compared to 5.14 years for women,



290 The estimated quarterly PAYLL values (see supplementary file, Table S1) can be used to apply

- the suggested empirical short-cut. These values range from a low of 9.7 years in Bulgaria to a high of 26.4
- 292 years in Qatar. These differences can be explained by different age compositions, with younger
- 293 compositions giving more weight to remaining life expectancies at younger ages, which are obviously
- higher. As expected, however, these values change relatively little from one quarter to the next for a given
- 295 population. Figure 3 shows peak MUL values for a rolling seven-day period, derived from quarterly
- 296 PAYLL values, were reached between mid-March and the end of June in Madrid (9.12 years on 3/29), in
- 297 New York (9.20 years on 4/13) and in Baja California (9.15 years on 6/9).





300	The same approximation but for sub-populations is illustrated by focusing on the province of
301	Guayas in Ecuador. Data on the monthly number of deaths by province show the marked increase in
302	March, April and May from a baseline of 1,700-2,000 per month in January, February and again in June. <sup>28</sup>
303	In April, the number of deaths reached 12,004, of which the January-February-June average suggests only
304	17.2% might be estimated to be from causes other than COVID-19 (without adjustment for competing
305	causes). No specific life table is available to estimate MUL values for the province directly, but based on
306	the second-quarter PAYLL derived for Ecuador, individuals appeared to have died 12.6 years younger, on
307	average, that their expected age at death in April in the province of Guayas.
308	Discussion

309 The MUL is proposed here as an alternative to induced changes in PLEB to assess the impact of a cause 310 of death of the individual lifespan in situations where the assumptions underlying life table construction 311 are implausible, invalidating the usual interpretation of the PLEB. Complementing existing measures 312 such as the AYLL and the YLL per capita, the MUL is similarly based on estimating the number of 313 potential years of life lost corresponding to deaths from a specific cause or excess deaths from a specific 314 event in a given period. As shown above, by averaging estimated potential years of life lost over the total 315 number of deaths in the period, the MUL is structured like a difference in PLEB. The MUL retains the 316 intuitive interpretation of a difference in PLEB as a change in average length of life, but for an actual 317 cohort of individuals (those dying in the period) subjected to prevailing mortality conditions rather than 318 for a synthetic cohort subjected to indefinitely constant conditions.

To illustrate the derivation and interpretation of the MUL in the context of COVID-19 mortality, MUL values were derived for a total of 281 populations for each sex and each quarter of 2020. As argued above, the MUL is best estimated from data on numbers of excess deaths by age and sex in a given period, estimated from data on all-cause mortality. At this writing, however, these data are not available from the vast majority of the national and subnational populations around the world. Instead, total numbers of quarterly COVID-19 deaths in each of these populations were used and their age-and-sex breakdown derived with the assumption of a shared age-and-sex COVID-19 mortality "pattern" (i.e., the

326 same ratio of age-and-sex-specific death rates to the all-age, both-sex death rate in all populations).

- 327 Excess deaths by age and sex were then derived through life table adjustments for competing causes of
- death (as COVID-19 deaths reduced exposure to other causes of deaths).

329 The adjustment for competing causes of death, as well as the estimate of potential years of life 330 lost corresponding to each excess death is based on the assumption that, in the absence of a certain cause 331 of death, individuals who died from that cause would have faced the same risk of death from other causes 332 as any individual of the same age and sex. As discussed above, this standard assumption of YLL-based 333 measures is problematic in the case of COVID-19 due to the higher proportion of several underlying long-334 term conditions (e.g., obesity) observed among COVID-19 victims. Adjusting for differences in long-335 term conditions prevalence is very data demanding, however, and the impact might not be as large as 336 expected, a study finding that performing this adjustment reduces the AYLL for COVID-19 in the United 337 Kingdom was from 13 to 12 years (average for both sex).<sup>29</sup> An alternative strategy was proposed here that 338 underestimates the value of potential years of life lost over an age interval by estimating its value at the 339 beginning of the age interval. As shown in the supplementary files, this yields quarterly PAYLL values 340 for the United Kingdom that vary between 11.9 and 12.0 years (Table S1), values thus quite close to what 341 the adjustment for underlying long-term conditions might have provided. While there is of course no 342 guarantee that this strategy would apply equally well to causes of death other than COVID-19, ignoring 343 comorbidities will likely lead to some overestimation of AYLL for most causes of death, and purposely 344 underestimating the number of potential years of life lost will likely be preferable in other situations as 345 well.

The MUL estimates derived here can also be improved by using actual age-and-sex breakdowns of COVID-19 deaths in populations of interest, which are becoming available for an increasing number of nations.<sup>30</sup> Here too, however, the actual impact on MUL estimates might not be as large as expected. The age patterns of CoViD-19 death rates available so far similarly exhibit remarkable regularities, with some modest variation in the slope of these age patterns at the oldest ages, probably due to the number of fatalities in nursing homes across Europe and the USA.<sup>31</sup> Considering the two countries with the largest

number of COVID-19 deaths in 2020, substituting the actual sex and age pattern in Brazil to the one
derived by assuming the same mortality pattern as in the USA only changes the estimated reduction in
2020 PLEB for Brazil from 1.72 to 1.67 years.<sup>32</sup> The structural similarity between the MUL and
differences in PLEB suggest than using population-specific data on COVID-19 mortality by age and sex
might only improve MUL estimates by a similarly modest order of magnitude.

357 To date, by far the strongest reason to caution against taking these MUL estimates at face value 358 originates in lacking or incomplete data on the total number of deaths in 2020 for many populations and 359 how this number might eventually relate to the number of reported COVID-19 deaths. Even in countries 360 with reliable vital statistics, annual deaths data might not be finalized until several months into the 361 following year. Moreover, the estimation of excess deaths is performed by comparison with benchmark 362 mortality rates to represent counterfactual mortality conditions, and the results can be quite sensitive to 363 the choice of a benchmark period (e.g., the last five years) and the estimation methodology (e.g., assumptions about mortality trends).<sup>33</sup> In a study of 21 high-income nations,<sup>34</sup> the official number of 364 365 COVID-19 deaths from mid-February to end of May was found to be lower than the lower bound of the 366 95% credible interval for excess deaths during that period in 5 countries. The discrepancy was highest in 367 Spain, with the credible interval from 47% to 91% more excess deaths than reported COVID-19 deaths. In the USA, between January 26<sup>th</sup> and October 3<sup>rd</sup>, excess deaths were also estimated to be 38% higher 368 than the COVID-19 death tally over the same period.<sup>35</sup> Unfortunately, these ratios exhibit insufficient 369 370 empirical regularities across populations (and possibly over time) to uniformly adjust numbers of reported 371 COVID-19 deaths across populations for which the number of excess deaths remains unknown at this 372 point.

373 More generally, another advantage of the MUL is that it remains interpretable regardless of its 374 temporal and geographical scale, pertaining to deaths in a given population in a given period, whereas we 375 argued the PLEB becomes difficult to interpret as a short-term, micro-level indicator. With the PAYLL 376 values provided in supplementary file (Table S1), MUL values can easily be approximated as the product 377 of the corresponding PAYLL value and the ratio of COVID-19 to total deaths in the population during a

378 given period. The approximation can be used for short-duration periods as long as PAYLL can be 379 assumed to be almost constant over time and quarterly PAYLL estimates are found indeed to change little 380 from one quarter to the next. This approximation can also be used for sub-populations for which the 381 necessary data are only provided of the entire population, but only when the age compositions of the sub-382 populations can be held as relatively similar. 383 This relates more generally to the fact that, like YLL-based measures, the MUL is not sex- or 384 age-standardized and MUL comparisons across populations that differ markedly in age composition will 385 be biased. On the one hand, all else equal, a younger population composition yields a younger distribution

386 of deaths and a higher PAYLL value. On the other hand, known variations of COVID-19 mortality with

387 age<sup>36</sup> suggest that an older population distribution contributes to a higher proportion of COVID-19 deaths

relative to all deaths. Only by accident would the two, opposite age-composition effects cancel each other

and in general the MUL is not a standardized measure. This represents one disadvantage compared to

390 differences in PLEB. However, this advantage over the MUL again comes at a cost, since the internal

derivation assumes the period mortality conditions will become permanent.

392 When this assumption is not tenable, the MUL provides an unstandardized alternative to 393 differences in PLEB, similarly structured as an average difference in length of lives lived per person. 394 Related to other unstandardized measures such as the AYLL or YLL per capita, the MUL's interpretation 395 pertains to an actual death cohort, that is, population members who died during a certain period rather 396 than to a synthetic cohort as represented in the life table. To reiterate, the MUL indicates the difference 397 between their average age at death and their average expected age at death had a temporary mortality 398 shock not occurred, and its measurement requires no assumption that these temporary conditions will 399 either pass entirely or extend indefinitely into the future.

400

389

#### 401 Acknowledgments

- 402 The author benefited from facilities and resources provided by the California Center for
- 403 Population Research at UCLA (CCPR), which receives core support (P2C-HD041022) from the
- 404 Eunice Kennedy Shriver National Institute of Child Health and Human Development (NICHD).
- 405 The author thanks Michel Guillot, Philippe Bocquier, Tim Riffe, Michel Garenne, Christophe
- 406 Guilmoto and Sam Preston for comments on an earlier draft of this manuscript.

## 408 **References**

<sup>1</sup> Beltrán-Sánchez H, Preston SH & Canudas-Romo V. An integrated approach to cause-of-death analysis:

Cause-deleted life tables and decompositions of life expectancy. Demographic Research 2008;19: 1323-

1350. doi: 10.4054/DemRes.2008.19.35

<sup>2</sup> Ho J. The contribution of drug overdose to educational gradients in life expectancy in the

United States, 1992–2011. Demography 2017;54: 1175-1202. doi: 10.1007/s13524-017-0565-3

<sup>3</sup> Goldstein JR & Lee RD. Demographic perspectives on the mortality of COVID-19 and other epidemics.

Proc. Natl Acad. Sci. 220; 117(36): 22035-22041. doi: 10.1073/pnas.2006392117

<sup>4</sup> Ghislandi S et al. 2020. News from the Front: Estimation of excess mortality and life expectancy in the major epicenters of the COVID-19 pandemic in Italy. MedRxiv [preprint] 2020. doi:

10.1101/2020.04.29.20084335

<sup>5</sup> Trias-Llimos S, Riffe T & Bilal U. Monitoring life expectancy levels during the COVID-19 pandemic:

Example of the unequal impact of the first wave in Spanish regions. PLOS One 2020. doi:

10.1371/journal.pone.0241952

<sup>6</sup> Aburto JM, Kashyap R, Schöley J et al. 2020. Estimating the burden of COVID-19 on mortality, life expectancy and lifespan inequality in England and Wales: A population-level study. MedRxiv [preprint] 2020. doi: 10.1101/2020.07.16.20155077

<sup>7</sup> Ho T & Noymer A. Summertime, and the livin' is easy: Winter and summer pseudoseasonal life expectancy in the United States. Demographic Research 2017; 37(45): 1445-1476. doi:

10.4054/DemRes.2017.37.45

<sup>8</sup> Ramis H (director and writer) & Rubin D (writer). Groundhog day. [Film] 1993. Washington, D.C., United States: National Film Registry.

<sup>9</sup> Keyfitz N. What difference would it make if cancer were eradicated? An examination of the Taeuber paradox. Demography 1977; 14: 411-418. https://link.springer.com/content/pdf/10.2307/2060587.pdf <sup>10</sup> Dempsey M. Decline in tuberculosis - The death rate fails to tell the entire story. Am Rev Tuberc. 1947; 56: 157-164.

<sup>11</sup> OECD. Potential years of life lost (indicator). [Cited February 18, 2021]. doi: 10.1787/193a2829-en

<sup>12</sup> Mitra AK, Payton M, Kabir N, et al. Potential years of life lost due to COVID-19 in the United States,

Italy, and Germany: An old formula with newer Ideas. Int. J. Environ. Res. Public Health 2020; 17: 4392.

doi: 10.3390/ijerph17124392

<sup>13</sup> Mohanty SK, Dubey M, Mishra US & Sahoo U. Impact of COVID-19 attributable deaths on longevity,

premature mortality and DALY: Estimates of USA, Italy, Sweden and Germany. MedRxiv [preprint] 2020.

doi: 10.1101/2020.07.06.20147009

<sup>14</sup> Wetzler HP, Wetzler EA & Cobb HW. COVID-19: How many years of life lost? MedRxiv [preprint] 2020.

doi: 10.1101/2020.06.08.20050559

<sup>15</sup> Pifarré i Arolas H, Acosta E, López Casasnovas, G et al. Global years of life lost to COVID-19. MedRxiv

[preprint] 2020. doi: 10.1101/2020.06.19.20136069

<sup>16</sup> Modig K, Rau R & Ahlbom A. Life Expectancy: What Does it Measure? BMJ Open 2020; 10.

https://bmjopen.bmj.com/content/10/7/e035932

<sup>17</sup> Aragon TJ, Lichtensztajn DY, Katcher BS et al. Calculating expected years of life lost for assessing local ethnic disparities in causes of premature death. BMC Public Health 2020; 8: 116. doi: 10.1186/1471-

2458-8-116

<sup>18</sup> Arias E & Xu JQ. United States Life Tables, 2018. National Vital Statistics Reports 2020; 69(12).

Hyattsville, MD: National Center for Health Statistics.

<sup>19</sup> United Nations, Department of Economic and Social Affairs, Population Division. World population prospects 2019, online edition. rev. [Cited June 28, 2020].

https://population.un.org/wpp/Download/Standard/CSV/.

<sup>20</sup> Istituto Nazionale di Statistica (Istat). Life tables. [Cited June 28, 2020].

http://dati.istat.it/Index.aspx?QueryId=19053&lang=en.

<sup>21</sup> Instituto Nacional Estadistica (INE). Mortality tables. [Cited June 28, 2020].

https://www.ine.es/dyngs/INEbase/en/categoria.htm?c=Estadistica\_P&cid=1254734710984.

<sup>22</sup> United States Census Bureau. Explore census data. [Cited May 11, 2020]. https://data.census.gov.

<sup>23</sup> Centers for Disease Control and Prevention (CDC). CDC wonder. [Cited June 28, 2020].

https://wonder.cdc.gov/cmf-icd10.html.

<sup>24</sup> Preston SH, Heuveline P & Guillot M. Demography: Measuring and modeling population processes.

Malden, MA: Blackwell; 2001.

<sup>25</sup> Guilmoto CZZ. COVID-19 death rates by age and sex and the resulting mortality vulnerability of

countries and regions in the world. MedRxiv [preprint] 2020. doi: 10.1101/2020.05.17.20097410

<sup>26</sup> Centers for Disease Control and Prevention. Provisional COVID-19 death counts by sex, age and state.

[Cited December 31, 2020]. https://data.cdc.gov/NCHS/Provisional-COVID-19-Death-Counts-by-Sex-Ageand-S/9bhg-hcku.

<sup>27</sup> University of Washington, Institute for Health Metrics and Evaluation. COVID-19 projections. [Cited January 2, 2021]. https://covid19.healthdata.org/.

<sup>28</sup> Dirección General de Registro Civil, Identificación y Cedulación. Defunciones, enero-junio 2020. [Cited
 July 28, 2020]. https://www.registrocivil.gob.ec/cifras/

<sup>29</sup> Hanlon P, Chadwick F, Shah A et al. COVID-19 – Exploring the implications of long-term condition type and extent of multimorbidity on years of life fost: A modelling study. Welcome Open Res 2020; 5: 75. doi: 10.12688/wellcomeopenres.15849.1

<sup>30</sup> Riffe T & Acosta E. COVerAGE-DB: A database of COVID-19 cases and deaths by age. [Cited December 28, 2020]. https://osf.io/mpwjq/.

<sup>31</sup> O'Driscoll M, Ribeiro Dos Santos G, Wang L et al. Age-specific mortality and immunity patterns of SARS-CoV-2. *Nature* 2021; 590: 140-145. doi: 10.1038/s41586-020-2918-0

<sup>32</sup> Heuveline P & Tzen M. Beyond deaths per capita: Comparative CoViD-19 mortality indicators.

MedRxiv [preprint] 2021. doi: 10.1101/2020.04.29.20085506

<sup>33</sup> Aron J & Muellbauer J. A pandemic primer on excess mortality statistics and their comparability across

countries. Our world data. [Cited February 18, 2020]. https://ourworldindata.org/covid-excess-

mortality.

<sup>34</sup> Kontis V, Bennett JE, Rashid T et al. Magnitude, demographics and dynamics of the effect of the first wave of the COVID-19 pandemic on all-cause mortality in 21 industrialized countries," Nat. Med. 2020;
26: 1919-1928. doi: 10.1038/s41591-020-1112-0

<sup>35</sup> Rossen LM, Branum AM, Ahmed FB et al. Excess deaths associated with COVID-19, by age and race and ethnicity – United States, January 26 – October 3, 2020," Morb. Mortal. Wkly Rep. 2020; 69: 1522-1527. doi: 10.15585/mmwr.mm6942e2external icon.

<sup>36</sup> Dowd JB, Andriano L, Brazel DM, et al. Demographic science aids in understanding the spread and fatality rates of COVID-19. Proc. Natl Acad. Sci. 2020; 117: 9696-9698. doi: 10.1073/pnas.2004911117