# Bloody jello: Colloidal physics states acanthocyte sedimentation rate as a diagnostic biomarker for neuroacanthocytosis

Alexis Darras<sup>a,+,\*</sup>, Kevin Peikert<sup>b,c,+</sup>, Antonia Rabe<sup>a,d</sup>, François Yaya<sup>a,e</sup>, Greta Simionato<sup>a,f</sup>, Thomas John<sup>a</sup>, Anil Kumar Dasanna<sup>g</sup>, Semen Buvalyy<sup>g</sup>, Jürgen Geisel<sup>h</sup>, Andreas Hermann<sup>b,c,i,j</sup>, Dmitry A. Fedosov<sup>g</sup>, Adrian Danek<sup>k</sup>, Christian Wagner<sup>a,l</sup>, and Lars Kaestner<sup>a,d,\*</sup>

<sup>a</sup>Experimental Physics, Saarland University, 66123 Saarbruecken, Germany

<sup>b</sup>Translational Neurodegeneration Section "Albrecht-Kossel", Department of Neurology, University Medical Center Rostock, University of Rostock, Rostock, Germany

<sup>c</sup>Neurodegenerative Diseases, Department of Neurology, Technische Universität Dresden, Dresden, Germany <sup>d</sup>Theoretical Medicine and Biosciences, Saarland University, 66424 Homburg, Germany

<sup>e</sup>Laboratoire Interdisciplinaire de Physique, UMR 5588, 38402 Saint Martin d'Hères, France

<sup>f</sup>Institute for Clinical and Experimental Surgery, Saarland University, 66424 Homburg, Germany

<sup>g</sup>Institute of Biological Information Processing and Institute for Advanced Simulation, Forschungszentrum Jülich, 52425 Jülich, Germany

<sup>h</sup>Institute for Clinical and Experimental Surgery, Saarland University, 66424 Homburg, Germany <sup>i</sup>DZNE, German Center for Neurodegenerative Diseases, Research Site Rostock/Greifswald, Rostock, Germany <sup>j</sup>Center for Transdisciplinary Neurosciences Rostock (CTNR), University Medical Center Rostock, University of Rostock, Rostock, Germany

<sup>k</sup>Neurologische Klinik und Poliklinik, Ludwig-Maximilians-Universität, 81366 Munich, Germany <sup>1</sup>Physics and Materials Science Research Unit, University of Luxembourg, Luxembourg City, Luxembourg <sup>\*</sup>Corresponding authors. E-mail: alexis.darras@uni-saarland.de, lars\_kaestner@me.com

+AI.D. contributed equally to this work with K.P.

#### ABSTRACT

Chorea-acanthocytosis and McLeod syndrome are the core diseases among the group of rare neurodegenerative disorders called neuroacanthocytosis syndrome (NAS). NAS patients have irregularly spiky erythrocytes, so-called acanthocytes. Their detection is a crucial but error-prone parameter in the diagnosis of NAS, often leading to misdiagnosis. Based on the standard Westergren method, we show that the acanthocyte sedimentation rate (ASR) with a two-hour read-out is significantly prolonged in Chorea-acanthocytosis and McLeod syndrome without overlap compared to the erythrocyte sedimentation rate (ESR) of controls. Thus, the ASR/ESR is a clear, robust and easily obtained diagnostic marker. Mechanistically, through modern colloidal physics, we show that acanthocyte aggregation and plasma fibrinogen levels slow down the sedimentation. This study is also a hallmark of the physical view of the erythrocyte sedimentation by describing anticoagulated blood in stasis as a percolating gel, allowing the application of colloidal physics theory.

#### 1 Introduction

Neuroacanthocytosis syndromes (NASs) are a group of rare
 genetic neurodegenerative diseases, the two core disorders
 being chorea-acanthocytosis (ChAc, OMIM #200150) and
 McLeod syndrome (MLS, OMIM #300842)<sup>1-7</sup>. While both
 diseases are characterized by neurological symptoms such

7 as hyper- and hypokinesia, epileptic seizures and cognitive

<sup>8</sup> impairment, ChAc usually manifests much earlier in life (in

the twenties) than MLS (usually after 40 years of age). The
 mean life expectancy after diagnosis has been reported to

<sup>11</sup> be approximately 11 years for ChAc and 21 years for MLS, <sup>12</sup> leading to premature death in most cases<sup>8</sup>. The courses of

the diseases are heterogeneous, usually slowly progressive

<sup>14</sup> and often result in severe disability, while treatment options

remain purely symptomatic so far<sup>9</sup>. Diagnosis is confirmed 15 by genetic testing (VPS13A gene in ChAc, XK gene in MLS) 16 and chorein in Western blot (ChAc) or immunohematological 17 assessment  $(MLS)^5$ , although the initial detection often relies 18 on blood smears<sup>10</sup>. One of the common features of NAS is the 19 presence of acanthocytes among erythrocytes<sup>4–6</sup>. Acantho-20 cytes are deformed erythrocytes whose irregular membrane 21 presents disordered, asymmetric spikes. 22

For many of these rare diseases, diagnosis is challenging, and misdiagnosis is frequent, which additionally increases the patients' burden of disease. One of the reasons for delayed diagnosis is related to the difficult detection of acanthocytes in classical blood smears and routine laboratory testing: sensitive detection requires special isotonically diluted blood 28

<sup>29</sup> samples<sup>10</sup>, the number of acanthocytes varies among the pa-

30 tients, and their similarity to echinocytes often makes them

hard to identify<sup>6</sup>. Therefore, it would be desirable to have an

easier and more reliable initial diagnostic biomarker.

Here, we proposed the erythrocyte sedimentation rate (ESR)
as a convenient, inexpensive and objective parameter that
allows routine and systematic tests to be used to diagnose NAS.
At the same time, we investigated the physical cause of the
change in the acanthocyte sedimentation rate (ASR). In doing
so, we devised a novel physical view of the sedimentation
process based on colloidal physics that is in good agreement

<sup>40</sup> with the experimental data.

#### 41 2 Results

### 42 Significance of the ESR in Neuroacanthocytosis43 Syndrome Patients

Time-resolved measurements of the ESR following the stan-44 dard Westergren method were performed on 6 ChAc patients, 45 3 MLS patients and 8 healthy controls. The major hematologi-46 cal data as well as the neurological phenotypes of the individu-47 als are given in Table 1. An image of the sedimentation tubes, 48 the time curves of representative ASR/ESR measurements, 49 the statistical presentation of patient-based data at selected 50 time points and the overview of all measurements at selected 51 time points are summarized in Fig. 1A-D, respectively. 52 Although the number of patients was limited, at 30 min, the 53 difference in ESR was significantly lower for MLS patients 54 than for controls (p = 0.028). At 1 h, we found a significant 55 difference between both ChAc patients (p = 0.022) and MLS 56 patients (p = 0.017) and healthy controls. After 2 h, the p-57

value decreased to 0.006 for ChAc patients and to 0.005 for

<sup>59</sup> MLS patients relative to the healthy controls (Fig. 1C).

However, a clear systematic difference between the whole
range of measurements is only reached after 2 h. A sedimentation height of 10 mm can be regarded as a threshold (green
dashed line in Fig. 1D), i.e. only if the sedimentation phase
is less than 10 mm within 2h, ESR indicates ChAc and MLS.
These results are sex independent (cp. Supplemental Fig. 1A).

#### 66 Comparison of the Acanthocyte Percentage with 67 the ESR

To date the diagnosis of NAS has involved the detection of 68 acanthocytes. Consequently, here blood smears were prepared 69 based on the protocol from Storch et al.<sup>10</sup> and later stained 70 according to the Pappenheim method (see Supplemental Data). 71 An example is presented in Fig. 2A. The manual count of 72 cell shapes is prone to be subjective, biased and inaccurate 73 and therefore reproducible with only limited reliability, as 74 outlined in Supplemental Fig. 2. For a better assessment, 75 we fixed fresh cells in 0.1% glutaraldehyde and performed 76 confocal imaging of z-stacks to 3D render the erythrocytes. 77 See Supplemental Data for detailed protocol. Figure 2B shows 78 a number of representative acanthocytes in 3D, which were 79 used for further analysis. 80

We observed that the ESR tends to decrease when the frac-81 tion of acanthocytes increases. To relate the ESR to the num-82 ber of acanthocytes, the inverse of the ESR at 2 h is plotted 83 as a function of the number of acanthocytes (Fig. 2C). In a 84 more general approach, we plotted the inverse of the ESR 85 at 2 h against the number of all deformed cells (echinocytes, 86 acanthocytes and others; Fig. 2D). The analysis revealed a 87 significant correlation for both parameters ( $R^2$  coefficient of 88 0.61 and 0.59, respectively). Consistent with this finding, 89 the p-value decreased from 0.037 (acanthocytes) to 0.011 (all 90 deformed cells). 91

#### Explanations for Differences in the ESR

Because there was no significant difference in the ASR between ChAc and MLS patients (Fig. 1), we do not discriminate between ChAc and MLS patients for further statistical comparisons. Therefore, we subsume them into one NAS group.

The reasons for the differences in the ESR are varied and 98 can depend on the erythrocytes, the blood plasma or the 99 erythrocytes-plasma volume relation (i.e. the hematocrit or 100 erythrocyte volume fraction  $\phi$ ). The last reason can be ex-101 cluded since there was no significant difference in the hema-102 tocrit between the groups (cp. Table 1 and Supplemental 103 Fig. 1B). The primary physical parameters such as the den-104 sity of the individual erythrocytes, represented by the mean 105 cellular hemoglobin concentration (MCHC), as well as the 106 constitution of the plasma can also be part of the explanation. 107 However, the MCHC indicates a higher density of erythro-108 cytes in the NAS patients (Supplemental Fig. 3C), a feature 109 that would favor faster sedimentation and therefore rules it 110 out as an explanation. In contrast, the total plasma protein 111 analysis (Supplemental Fig. 4A), as a representative of the 112 plasma content, indicates a lower protein concentration and 113 therefore a first hint for a plasma contribution to the ESR 114 change. In the following sections, we present a more detailed 115 view of the influence of both blood components, erythrocytes 116 and plasma. 117

#### Role of Erythrocytes in Sedimentation

The sedimentation of erythrocytes in blood plasma has been 119 proposed, but was never established, to obey a so-called gel 120 sedimentation regime<sup>11,12</sup>. Gel sedimentation is efficiently 121 represented by several equations established for colloidal sys-122 tems<sup>13–16</sup>. In particular, gel sedimentation can exhibit a de-123 layed collapse<sup>17,18</sup>. Here, we apply the gel sedimentation 124 model for sedimenting erythrocytes. This means that erythro-125 cytes first experience only little sedimentation. Then, after 126 a characteristic time on the order of a few minutes, some 127 "cracks" or "channels" form within the aggregated structure 128 of the erythrocytes, as suggested by a previous study<sup>11</sup>. We 129 also observed this behavior in our samples (see Supplemen-130 tal Fig. 5). It is worthwhile to notice that, due to volume 131 conservation, the sedimentation of the erythocytes implies 132 an upward flow of the plasma. The channel structures then 133 enhance this upward flow of liquid. Here, we will apply for 134

92

93

94

95

96

97



**Figure 1.** Comparison of the ESR between neuroacanthocytosis patients and healthy controls. A: ESR measurement setup: standard Westergren tubes are filled with full blood and left to rest. The sedimentation height is measured over time. The picture was taken after 2 h. The first two tubes contain blood from an MLS patient (MLS-3), and the last two tubes contain blood from a healthy control donor. B: Representative time traces of the sedimentation height for the different conditions. C: Statistics on sedimentation height after typical times for healthy controls, ChAc patients and MLS patients. The bar height represents the average value, and the error bar depicts the range of individual values. Stars indicate the significance levels (Brown-Forsythe and Welch ANOVA test): n.s., not significant (p-value>0.05); \* p<0.05; and \*\* p<0.01. N refers to the number of patients. D: Span of all individual sedimentation height measurements (each blood sample was measured at least in duplicate, and the blood of some patients was measured at different time points). After 2 h, a  $\Delta h$  of 10 mm could be regarded as a threshold to differentiate the ChAc and MLS conditions. n refers to the number of individual measurements.



**Figure 2. Relation of acanthocyte counts to ESR.** A: A representative stained blood smear of a ChAc patient. B: Representative 3D-rendered acanthocytes from the freshly fixed blood of a ChAc patient. C: Plot of the number of acanthocytes vs. the inverse of the ESR at 2 h and the corresponding correlation analysis. D: Plot of the number of all deformed erythrocytes (echinocytes, acanthocytes and other unspecified cell shapes) vs. the inverse of the ESR at 2 h and the corresponding correlation analysis. D: Plot of the number of all deformed erythrocytes (echinocytes, acanthocytes and other unspecified cell shapes) vs. the inverse of the ESR at 2 h and the corresponding correlation analysis. Panels C and D contain measurements from 9 patients (cp. Table 1), where 1 patient was measured twice (independent blood sampling on different days). Since none of the controls presented with acanthocytes, they are summarized in one data point with their average ESR. The plotted line represents the linear regression. The analysis reveals a significant correlation for both parameters (panels C and D,  $R^2$  coefficient of 0.61 and 0.59, respectively).

and to sta			hematocrit	hemoglobin	<b>RBC number</b>	main clinical
subjects	sex	age range (y)	(%)	(g/dl)	$(10^{12}/l)$	characteristics
ChAc-1	male	50-55	43	15.1	4.67	Parkinsonim, dystonia, dysarthria,
						peripheral neuropathy,depression
ChAc-2	female	50-55	40	13.8	4.21	Epilepsy, parkinsonism dystonia,
						dysartifia, peripheral neuropatity,
						Enilance Darkinganiam dystania
ChAc-3	male	50-55	40	14.2	4.15	ducerthria ducerhagia parinharal
						nouropathy acquitive impeirment
						Drug registent enilency
ChAc-4	male	30-35	46	16.2	5.40	mild abores ties econitive
						mild chorea, tics, cognitive
						impairment, peripheral neuropatny,
						myopatny
ChAc-5	male	25-30	43	15.7	4.91	Drug resistant epilepsy,
						mild chorea, tics, cognitive
						impairment, irritability, anxiety,
						depression, psychosis
ChAc-6	male	40-45	41	15.2	4.96	Epilepsy, feeding dystonia,
						orofacial dyskinesia, chorea,
						peripheral neuropathy, myopathy,
						impulse control disorder
MLS-1	male	55-60	43	16.0	4.82	Myopathy
MLS-2	male	50-55	42	15.3	4.67	Epilepsy, peripheral neuropathy,
						myopathy
MLS-3	male	50-55	45	16.2	5.17	Neuropathy, myopathy
female controls		64+10	38 5+2 6	12.6+1.3	4 36+0 19	
(average, N=4)		04±10	50.5±2.0	12.0±1.5	4.50±0.17	
male controls		38+11	43.4+2.1	15.5+0.8	$5.17 \pm 0.1$	
(average, N=4)		J0±11		10.0 ± 0.0	0.17 ±0.1	

Table 1 : Overview of the patients and controls

the first time some quantitative scaling based on these observations. Indeed, the maximal sedimentation speed  $v_M$  following volume conservation and Darcy's law<sup>15</sup> can be approximated as:

$$v_M = \frac{k_0 \Delta \rho_g \phi_0}{\eta} \frac{2h_c}{\chi},\tag{1}$$

where  $k_0$  is the permeability of the aggregated erythrocyte 139 structure,  $\Delta \rho$  is the density difference between the erythro-140 cytes and the suspending liquid,  $\phi_0$  is the initial hematocrit, 141  $\eta$  is the liquid viscosity,  $h_c$  is the height of the channel, and 142  $\chi$  is the characteristic radius of the zone where this channel 143 influences the liquid flow<sup>15</sup>. Note that the permeability of 144 the aggregated erythrocyte structure  $k_0$  is proportional to the 145 characteristic area of the holes within the cross-section  $\langle H \rangle$  of 146 aggregated erythrocytes:  $k_0 \propto \langle H \rangle^{13}$ . Therefore, the maximal 147 sedimentation speed is also proportional to  $\langle H \rangle$ : 148

settle in a microscope chamber such that the final hematocrit 150 on the coverslip layer was close to the physiological value of 151 45% (Fig. 3A,B). This, when observing the chamber from 152 the bottom, allows a two-dimensional analysis of the hole 153 size *H* based on microscopy images as outlined in Fig. 3C-E. 154 The quantitative difference in average hole diameter (defined 155 as the square root of the average hole area) is exemplified 156 for one healthy control and one NAS patient by a probability 157 density distribution (Fig. 3E). Statistical analysis of the hole 158 diameters is given in Fig. 3F, revealing a larger hole size for 159 the blood of healthy donors. One can also note that the total 160 area covered by the erythrocytes is different between healthy 161 controls and NAS patients (e.g., in Fig. 3A-D, the total area 162 of the holes is 8.5  $10^4 \mu m^2$  under control conditions but 6.2 163  $10^4 \mu m^2$  for the patients). This is caused by the projection 164 area of the cells despite the absolute hematocrit being equal 165 between the groups. Under control conditions, almost all cells 166 form rouleaux and hence contribute to a side projection, while 167 acanthocytes and echinocytes have a larger projection area. 168

To estimate the hole size, we allowed the erythrocytes to

 $v_M \propto \langle H \rangle.$  (2)

To better understand the effect of acanthocyte shape on 169



## Figure 3. Characteristics of sedimenting erythrocyte aggregates in autologous

plasma. A, B: Sedimented layer of erythrocytes in plasma for a control subject and a patient, on the bottom of a microscopy chamber after 2 h. Pictures are taken from the bottom of the well. A: healthy control donor and B: patient (MLS-3). The volume fraction (hematocrit) in the microscope slide well (inner diameter 5 mm, height 1.5 mm) was initially 0.33% to reach a volume fraction close to 45% at the bottom of the well after sedimentation. C, D: Same pictures as previously, but with the holes detected in the percolating network of erythrocytes highlighted with random colors. The 50 µm scale is valid for panels A to D. E: Example of distributions of the square root of the hole areas, or hole diameter, shown as complementary cumulative density functions. The linear curve reflects an exponential distribution with the mean hole diameter as a single scale parameter. This scale parameter was estimated by the maximum likelihood method. F: Statistical comparison of the scale parameters for control and NAS erythrocytes in autologous plasma. The scale parameters are normally distributed among each population, and a t-test between the two populations indicates a significant difference (p = 0.005). G: Simulation snapshot of healthy conditions with 100% discoid erythrocytes shown in black color similar to experimental images. H: Simulated aggregate of a mixture of 80% normal erythrocytes (black area) and 20% acanthocytes (gray color) by area fraction. The scale bar is valid for panels G and H. I: Complementary cumulative density function of the square root of the hole areas from the simulations of healthy erythrocytes and erythrocyte-acanthocyte mixture exemplified in G and H.

the properties of sedimented aggregates (e.g., hole size), we 170 performed a number of two-dimensional simulations with dif-171 ferent compositions of healthy erythrocytes and acanthocytes; 172 see example snapshots in Fig. 3G,H (and see Supplemental 173 Data for simulation details). Figure 3I shows the correspond-174 ing cumulative density functions of hole diameter (defined 175 as the square root of the hole area), which compare favor-176 ably to the experimental data in Fig. 3E. The simulations 177 reveal that acanthocyte shape and rigidity effectively reduce 178 179 aggregation interactions, as they cannot maximize locally aggregated membrane areas between cells, which is achieved by 180 cell deformation in healthy erythrocytes. A lower effective 181 aggregation between cells leads to smaller hole sizes, which 182 we confirmed directly by reducing the aggregation strength in 183 the suspension of healthy cells. 184

To further assess the role of erythrocytes in the reduced sed-185 imentation rate and exclude the influence of individual plasma 186 properties, sedimentation experiments with washed erythro-187 cytes suspended in dextran (70 kDa) diluted in phosphate 188 buffered saline (PBS) were performed. A dextran concentra-189 tion of 55 mg/ml, which mimics erythrocyte aggregation in 190 blood samples of healthy controls, was empirically selected 191 (Fig. 4A,D). The aggregates formed in sedimented layers of 192 erythrocytes in dextran solution (Fig. 4A,B) indeed looked 193 very similar to the aggregates in autologous plasma (compare 194 with Fig. 3A,B). The hole size ratio  $H_{\text{control}}/H_{\text{NAS}}$  in the dex-195 tran solution was also very similar to that in plasma solution 196 (Fig. 4C). 197

For further comparison, the maximum sedimentation speed 198  $v_M$  was measured for each ESR test. Under the assumption 199 of a fixed hematocrit of 45%, only the parameters  $k_0$  and  $\Delta \rho$ , 200 related to the properties of erythrocytes, can imply a differ-201 ence in sedimentation speed. Standard blood parameters, i.e. 202 hemoglobin concentration (Table 1, Supplemental Fig. 3A) 203 and complete red blood cell count parameters, i.e. number 204 of erythrocytes (Table 1, Supplemental Fig. 3B) and mean 205 cell volume (MCV; Supplemental Fig. 3D) did not show any 206 significant difference that could decrease the ESR of NAS pa-207 tients through  $\Delta \rho$ . However, MCHC (Supplemental Fig. 3C) 208 was slightly increased in the NAS patients but, as previously 209 mentioned, would then increase the ESR and therefore cannot 210 explain the differences. Thus, different cases can be compared 211 by scaling the observed maximum sedimentation speed with 212 the hole size; see Eq. 2. If we denote patients with a subscript 213 NAS and the healthy controls with C, then we obtain 214

$$\frac{v_{\rm C}}{v_{\rm NAS}} = \frac{\langle H_{\rm C} \rangle}{\langle H_{\rm NAS} \rangle} \Rightarrow v_{\rm C} = \frac{\langle H_{\rm C} \rangle}{\langle H_{\rm NAS} \rangle} v_{\rm NAS}.$$
(3)

The  $v_M$  of control ( $v_C$ ) and NAS blood ( $v_{NAS}$ ) as well as the corresponding erythrocyte-dextran mixtures are shown in Fig. 4D (gray and orange columns, respectively). Additionally, the sedimentation speed was scaled according to Eq. 3. This scaling is plotted in Fig. 4D (green columns), along with the error bars corresponding to the standard deviation of the

scaled speed measurements. The scaling makes the velocities 221 comparable and indicates that the difference in the aggregate 222 geometry as characterized by hole area accounts for the ob-223 served difference in sedimentation velocity when erythrocytes 224 are suspended in a dextran-based medium. However, the scal-225 ing is not sufficient to explain the velocity ratio when the 226 sedimentation is measured in autologous plasma, as shown in 227 Fig. 1. This indicates that the aggregate geometry accounts 228 for some fraction of the ASR slowdown, while the remain-229 ing fraction must correspond to other contributions, such as 230 patient plasma composition. 231

#### Role of Plasma in the Sedimentation

As mentioned above, the total protein content likely con-233 tributes to the reduced ASR (Supplemental Fig. 4A). There-234 fore, we further analyzed plasma constituents, such as fibrino-235 gen (Fig. 5A), albumin (Supplemental Fig. 4B), C-reactive 236 protein (CRP, Supplemental Fig. 4C) and immunoglobulin G 237 (IgG, Supplemental Fig. 4D). Since the last three did not show 238 significant differences between NAS patients and healthy con-239 trols, we focused on fibrinogen, which was on average 23% 240 lower in the patients (p = 0.004). 241

To investigate the role of plasma in the increase in the 242 ESR, we performed experiments in which the erythrocytes and 243 plasma of patients were swapped with those of a healthy con-244 trol with a matching blood group. We observed that healthy 245 erythrocytes suspended in patient plasma sediment signifi-246 cantly faster than NAS erythrocytes in healthy plasma (which 247 confirms the importance of erythrocyte properties). However, 248 patient erythrocytes also sediment in healthy plasma signif-249 icantly faster than in patient plasma, which is a clear sign 250 that NAS plasma also slows sedimentation. Representative 251 time traces are shown in Supplemental Fig. 6. A statistical 252 presentation is given in Fig. 5B. 253

Previous reports have shown that an increase in fibrinogen 254 concentration is related to an increased sedimentation speed 255 and that fibringen has a larger impact on ESR than other pro-256 teins<sup>19,20</sup>. In particular, fibrinogen increases the interaction 257 energy between erythrocytes<sup>21</sup>. To test whether this differ-258 ence in interaction energy also occurs in NAS patients, we 259 measured the aggregation forces between erythrocytes. To 260 this end, we performed experiments using holographic optical 261 tweezers, as shown in Fig. 5C. As depicted in Fig. 5D, the ag-262 gregation forces of erythrocytes from acanthocytosis patients 263 in autologous plasma were significantly lower (p = 0.0005) 264 than those from healthy controls. 265

The details of the physical mechanisms leading to delayed 266 collapse due to the appearance of cracks in weakly aggregated 267 colloids are still under investigation<sup>11, 12, 16, 22–25</sup>. However, 268 recent studies highlighted that the aggregation energy can 269 modify the sedimentation speed in this regime, as well as the 270 time of onset<sup>22–25</sup>. Therefore, the difference in plasma fibrino-271 gen concentration is likely to explain the further difference 272 in sedimentation speed between NAS patients and healthy 273 controls. 274



**Figure 4.** Characteristics of sedimenting erythrocyte aggregates in various suspension media. A,B: Sedimented layer of erythrocytes in 55 mg/ml dextran from A: a healthy control donor and B: an NAS patient. The characteristic area of the holes in the network is visibly smaller in the network of NAS patient's erythrocytes. As the replacement medium, we used 70 kDa dextran diluted in PBS to a final mass concentration of 55 mg/ml. C: Measured ratio of mean hole size for the suspension media (autologous plasma and dextran diluted in PBS with a mass concentration of 55 mg/ml). The dextran concentration was empirically chosen in order to achieve a sedimentation speed as close as possible to the rate obtained in the plasma of the healthy samples. D: Characteristic (maximal) erythrocyte sedimentation velocities, with scaling based on aggregate geometries. The bars indicate the values of the characteristic speed of each patient's erythrocytes by the ratio of the characteristic hole areas, as justified by Eq. 3. While the scaled velocity in autologous plasma is significantly smaller than the characteristic speed of the control erythrocytes, the scaling lies within the same range when the autologous plasma is replaced by a dextran-based medium. n.s., not significant (p>0.05); \* p<0.05.



**Figure 5. Measurements related to the role of plasma in sedimentation.** A: Comparison of fibrinogen concentration. A significant difference consistent with the aggregation force difference is observed. B: Comparison of sedimentation velocities of control samples, NAS patient samples and samples with erythrocytes and plasma exchanged. C 1: Schematic of the forces acting on the erythrocytes trapped in optical tweezers. C 2: Microphotographs of the protocol for the measurement of the aggregating force. The purple circles show the location of external optical traps. From left to right: After being selected, erythorcytes were lifted 15  $\mu$ m from the microscope slide by 4 optical traps, one for each erythrocyte extremity. The optical traps have known trapping forces. The erythrocytes are then brought into contact. At equilibrium, the two inner traps are removed. The optical force holding the cells is stepwise decreased, and the overlap distance tends to increase in the same manner. Finally, spontaneous aggregation overcomes the optical forces, and the erythrocytes escape the trap. The trapping force at which the cells aggregate is considered to be the aggregation force. D: Comparison of aggregation force. A significant difference between patients and controls was observed. n.s., not significant (p>0.05); \*\* p<0.01; \*\*\* p<0.001; and \*\*\*\*

#### 275 **3 Discussion and Conclusions**

We herein not only present a decreased ESR in NAS but also 276 conceptualize the biophysical reasons for this observation 277 and highlight that measurement of the ESR needs to be im-278 plemented in routine clinical tests. Our measurements show 279 that a decreased ESR can be used as an efficient biomarker 280 to test for ChAc and MLS. For a sedimentation time of 2 h, 281 we could (with our test tubes) determine a clear diagnostic 282 threshold of 10 mm. In the context of a clinical workflow, 283 the ESR (manually or automatedly determined) is a robust 284 and cost-efficient parameter that could easily complement the 285 error-prone acanthocyte count. Mutations can still be con-286 firmed by molecular biology methods to differentiate among 287 NASs (cp. Supplemental Data). Since the ESR also depends 288 on the geometry of the tubes<sup>26,27</sup>, systems other than those 289 used here may need their own standardization. 290

Conventionally, ESR is used to detect general inflammation, 291 which is indicated by an increased ESR<sup>28,29</sup>; however, there 292 is no conflicting overlap because the change is in the opposite 293 direction in NAS. We would have liked to test whether the 294 ASR correlates to the severity of the disease, but there is no 295 validated index or score for NAS severity available. Similar 296 to the acanthocyte count, ASR cannot discriminate between 297 ChAc and MLS. Furthermore, we cannot make a statement 298 about other NASs (such as pantothenate kinase-associated 299 neurodegeneration (PKAN)), but we suppose a prolongation 300 of the ESR also occurs in blood samples of patients with other 301 diseases showing an increased number of acanthocytes. 302

The number of patients in our study may seem low (6 303 ChAc and 3 MLS patients), but given the ultrararity of these 304 diseases—the estimated prevalence is less than 1 to 5 per  $10^6$ 305 inhabitants each<sup>5</sup>—the number of participants can well be 306 regarded as acceptable. In particular, this holds true because 307 with such a low number of patients, we were able to obtain sig-308 nificant results. Recent studies have identified promising drug 309 targets, such as Lyn kinase, as a potential disease-modifying 310 311 therapy for ChAc<sup>5</sup>. Considering the rarity and high variability of the natural history of the disease, there is an urgent need 312 for a robust biomarker to achieve clinical trial readiness. The 313 ASR might represent an ideal biomarker candidate in this 314 context. 315

We have shown that both erythrocytes properties and blood 316 plasma composition of NAS patients influence the ASR. Both 317 components reduce the sedimentation speed, i.e. they are ad-318 ditive and provide a clear cut-off to differentiate from healthy 319 individuals. This is to some extent surprising since the ESR is 320 a measure that integrates numerous parameters, and its initial 321 intended use as a parameter for reporting inflammation has 322 lost importance to a certain degree within the last decades<sup>30</sup> 323 due to a lack of specificity. 324

Furthermore, we discussed physical mechanisms related to the erythrocyte and plasma properties that determine the ESR. Historical reports describe an altered aggregation behavior of acanthocytes<sup>31</sup> and the influence of erythrocyte shape on the ESR<sup>32</sup>. Here, we applied the principles of colloidal physics

to describe sedimenting erythrocytes in a quantitative man-330 ner. Thus, we clearly show that the geometric properties of 331 erythrocyte aggregates influence the porosity of sedimenting 332 aggregates, which has a direct impact on the sedimentation 333 velocity. Furthermore, a modeling approach provides evi-334 dence that the shapes of the erythrocytes (acanthocytes and 335 echinocytes) and their rigidity are the primary cause for the 336 differences in aggregate geometry. Regarding the plasma 337 composition, we highlighted a difference in fibrinogen con-338 centration, which modifies the interaction energy between 339 erythrocytes. While the detailed influence of this energy on 340 the sedimentation process is still under active discussion, it is 341 clearly a key feature in the sedimentation process. 342

Future investigations are required for the clinical explo-343 ration of the ASR in a broader context of acanthocyte-related 344 diseases and its longitudinal value in individual patients and a 345 deeper physical understanding of erythrocyte sedimentation. 346 The clinical application comprises a broader test for neurode-347 generative diseases as well as the verification of the ASR as 348 a biomarker for NAS treatments as proposed above. For the 349 mechanistic understanding of the process of erythrocyte sed-350 imentation based on colloidal physics, we have provided an 351 initial concept that also requires further research. 352

#### 4 Materials and Methods

#### Patients and Blood Sampling

Blood samples were collected from six ChAc and three MLS 355 patients. Patients were being treated at the neurologic de-356 partments of Technische Universität Dresden and the Univer-357 sity Hospital of the Ludwig-Maximilian-Universität Munich. 358 Blood sample collection was approved by the 'Ärztekammer 359 des Saarlandes', ethics votum 51/18, and performed after in-360 formed consent was obtained according to the declaration of 361 Helsinki. The diagnosis of ChAc and MLS was based on 362 the clinical phenotype and was confirmed by detection of the 363 VPS13A mutation and/or absence of the erythrocyte mem-364 brane chorein via Western blot (ChAc) or the detection of XK 365 mutations (MLS)<sup>33</sup>. 366

Blood was taken in the morning during routine patient visits 367 and immediately transported to Saarland University in Hom-368 burg and Saarbrücken for further analysis, which started 6 to 8 369 h after withdrawal. Because transportation can have a tremen-370 dous effect on blood parameters<sup>34</sup>, samples from relatives (if 371 available) or the investigators were taken as transportation 372 controls (in total, eight different individuals). This practice 373 excluded patients and controls matched in terms of sex, age or 374 weight. Furthermore, three patients came directly to Saarland 375 University, which allowed the investigation of fresh samples. 376 However, concerning the ASR, there was no significant ef-377 fect between fresh and transported samples, as outlined in 378 Supplemental Fig. 7. 379

#### Measurements of the ESR

ESR/ASR measurements were performed in standard Westergren tubes (Dispette original, REF GS1500) with 200 mm

380

353

height EDTA-blood. Color pictures of the tubes were auto-383 matically taken every minute for up to 50 h with a Canon EOS 384 500D camera to cover the whole range of sedimentation in 385 every case. For each tube, a custom written MATLAB algo-386 rithm extracted the position of the interface at the top of the 387 concentrated erythrocyte suspension with an accuracy of at 388 least 0.1 mm. Since the displacement of the interface between 389 two successive frames was close to the picture resolution, 390 instantaneous sedimentation velocities were extracted from 391 392 least squares fitting of the data points with cubic splines with 8 free interior knots, as allowed by the SLM (Shape Language 393 Modeling) package<sup>35</sup>. 394

<sup>395</sup> Further detailed measurement techniques are described in <sup>396</sup> the Supplemental Data.

#### 397 Data Sharing Statement

The datasets generated and/or analyzed in the current study are available from the corresponding author on reasonable request.

#### 401 Acknowledgements

This work was supported by the Deutsche Forschungsgemein-402 schaft (DFG) in the framework of the research unit FOR 2688 403 "Instabilities, Bifurcations and Migration in Pulsatile Flows", 404 by the European Union Horizon 2020 research and innovation 405 program under the Marie Skłodowska-Curie grant agreement 406 No 860436 – EVIDENCE and by the Saarland University 407 competitive funding (Forchungsausschuss). K.P. was sup-408 ported by the Else Kröner Clinician Scientist Program (TU 409 Dresden, Germany) and the Rostock Academy for Clinician 410 Scientists (RACS, University of Rostock, Germany), A.H. is 411 supported by the "Hermann und Lilly Schilling-Stiftung für 412 medizinische Forschung im Stifterverband". F.Y., T.J. and 413 C.W. acknowledge funding from French German University 414 (DFH / UFA). We are indebted to the Advocacy for Neuroa-415 canthocytosis Patients (www.naadvocacy.org) for financial 416 support for chorea Western blotting. We also gratefully ac-417 knowledge the computing time granted through JARA-HPC 418 on the supercomputer JURECA at Forschungszentrum Jülich. 419 We would like to thank Alexander Kihm and Dr. Hannes Glass 420 for help in obtaining blood samples, their handling and trans-421 portation. We are grateful to Prof. Joan-Lluis Vives-Corrons 422 and Jan de Zoeten for valuable discussion of the manuscript. 423

#### 424 Authors contribution

Ad. D. and K.P. suggested the joint study. K.P. and L.K. 425 defined the project. K.P., A.H. and Ad.D. performed the 426 diagnoses and identified the patients suitable for the study. 427 Al.D., T.J., C.W. and L.K. designed the physical measure-428 ments. Al.D., T.J. and L.K. performed the ESR measurements 429 430 and the microscope-based erythrocyte sedimenting experiments. Al.D. analyzed the sedimentation data and suggested 431 the physical mechanisms. A.R. and G.S. performed the cell 432

counts. F.Y. measured the aggregation forces. J.G. took responsibility for the measurement of the hematological parameters. D.A.F., A.K.D. and S.B. performed the numerical simulations. All authors interpreted the results and wrote or reviewed the manuscript.

#### **Competing Interests**

The authors declare no competing interest.

438

439

440

#### References

- Adam, M. P. *et al.* Mcleod neurocanthocytosis syndrom.
   In *GeneReviews*® (University of Washington, Seattle, 1993).
- 2. Danek, A. *et al.* Mcleod neuroacanthocytosis: genotype and phenotype. *Ann Neurol* 2001; 50:755–764. 445
- **3.** Baeza, A. V. *et al.* Chorea-acanthocytosis. In *GeneReviews*® (University of Washington, Seattle, 2019). 447
- Siegl, C. *et al.* Alterations of red cell membrane properties in nneuroacanthocytosis. *PLoS One* 2013; 8:e76715. 449
- Peikert, K., Danek, A. & Hermann, A. Current state of knowledge in chorea-acanthocytosis as core neuroacanthocytosis syndrome. *Eur J Med Genet*. 2018; 61:699– 705.
- Adjobo-Hermans, M. J., Cluitmans, J. C. & Bosman,
   G. J. Neuroacanthocytosis: observations, theories and perspectives on the origin and significance of acanthocytes. *Tremor Other Hyperkinet Mov* 2015; 5:328.
- Rubio, J. P. *et al.* Chorea-acanthocytosis: genetic linkage to chromosome 9q21. *Am J Hum Genet.* 1997; 61:899– 908.
- Walker, R. H., Miranda, M., Jung, H. H. & Danek, A. Life expectancy and mortality in chorea-acanthocytosis and mcleod syndrome. *Park. Relat Disord* 2019; 60:158–161.
- Walker, R. H. Management of neuroacanthocytosis syndromes. *Tremor Other Hyperkinet Mov* 2015; 5:346.
- Storch, A., Kornhass, M. & Schwarz, J. Testing for acanthocytosis. *J Neurol* 2005; 252:84–90.
- Pribush, A., Meyerstein, D. & Meyerstein, N. The mechanism of erythrocyte sedimentation. part 1: Channeling in sedimenting blood. *Colloids Surf B Biointerfaces* 2010; 470 75:214–223.
- Pribush, A., Meyerstein, D. & Meyerstein, N. The mechanism of erythrocyte sedimentation. part 2: The global collapse of settling erythrocyte network. *Colloids Surf B Biointerfaces* 2010; 75:224–229.
- Allain, C., Cloitre, M. & Wafra, M. Aggregation and sedimentation in colloidal suspensions. *Phys Rev Lett* 477 1995; 74:1478.
- Manley, S., Skotheim, J., Mahadevan, L. & Weitz, D. A. Gravitational collapse of colloidal gels. *Phys Rev Lett* 480 2005; 94:218302.

- <sup>482</sup> 15. Derec, C., Senis, D., Talini, L. & Allain, C. Rapid settling
  <sup>483</sup> of a colloidal gel. *Phys Rev E* 2003; 67:062401.
- 484
  16. Senis, D., Gorre-Talini, L. & Allain, C. Systematic study
  of the settling kinetics in an aggregating colloidal suspension. *Eur Phys J E Soft Matter* 2001; 4:59–68.
- **17.** Buscall, R., Choudhury, T. H., Faers, M. A., Goodwin,
  J. W., Luckham, P. A. & Partridge, S. J. Towards rationalising collapse times for the delayed sedimentation
  of weakly-aggregated colloidal gels. *Soft Matter* 2009;
  5:1345–1349.
- Lindström, S. B., Kodger, T. E., Sprakel, J. & Weitz, D. A.
  Structures, stresses, and fluctuations in the delayed failure of colloidal gels. *Soft Matter* 2012; 8:3657–3664.
- 495 **19.** Gray, S. J., Mitchell, E. B. & Dick, G. Effect of purified
   496 protein fractions on sedimentation rate of erythrocytes.
   497 *Proc. Soc. for Exp. Biol. Medicine* 1942; 51:403–404.
- 498 20. Flormann, D., Kuder, E., Lipp, P., Wagner, C. & Kaestner,
  499 L. Is there a role of c-reactive protein in red blood cell
  aggregation? *Int J Lab Hematol* 2015; 37:474–482.
- **21.** Brust, M. *et al.* The plasma protein fibrinogen stabilizes
   clusters of red blood cells in microcapillary flows. *Sci Rep* 2014; 4:1–6.
- Ghazali, M. E. B., Argo, Y., Kyotoh, H. & Adachi, Y.
   Effect of the concentration of NaCl and cylinder height
   on the sedimentation of flocculated suspension of Na montmorillonite in the semi-dilute regime. *Paddy Water Environ.* 2020; 18:309–316.
- 23. Padmanabhan, P. & Zia, R. Gravitational collapse of colloidal gels: non-equilibrium phase separation driven by osmotic pressure. *Soft Matter* 2018; 14:3265–3287.
- **24.** Razali, A., Fullerton, C. J., Turci, F., Hallett, J. E., Jack,
  R. L. & Royall, C. P. Effects of vertical confinement on
  gelation and sedimentation of colloids. *Soft Matter* 2017;
  13:3230–3239.
- **25.** Harich, R. *et al.* Gravitational collapse of depletioninduced colloidal gels. *Soft Matter* 2016; 12:4300–4308.
- Brigden, M. L. & Page, N. E. Three closed-tube methods
   for determining erythrocyte sedimentation rate. *Lab Med* 1993; 24:97–102.
- Patton, W., Meyer, P. & Stuart, J. Evaluation of sealed
   vacuum extraction method (seditainer) for measurement
   of erythrocyte sedimentation rate. *J Clin Pathol* 1989;
   42:313–317.
- Bedell, S. E. & Bush, B. T. Erythrocyte sedimentation
   rate. From folklore to facts. *Am J Med* 1985; 78:1001–
   1009.
- Kratz, A. *et al.* ICSH recommendations for modified and
   alternate methods measuring the erythrocyte sedimenta tion rate. *Int J Lab Hematol* 2017; 39:448–457.
- **30.** Brigden, M. L. Clinical utility of the erythrocyte sedimentation rate. *Am Fam Physician* 1999; 60:1443–1450.

- Salt, H. *et al.* On having no beta-lipoprotein. a syndrome comprising α-beta-lipoproteinaemia, acanthocytosis, and steatorrhoea. *Lancet* 1960; 2:325–329.
- Reinhart, W. H., Singh, A. & Straub, P. W. Red blood cell aggregation and sedimentation: the role of the cell shape. *Br J Haematol* 1989; 73:551–556.
- 33. Dobson-Stone, C. *et al.* Chorein detection for the diagnosis of chorea-acanthocytosis. *Ann Neurol* 2004; 56:299–302.
   540
- **34.** Makhro, A. *et al.* Red cell properties after different modes of blood transportation. *Front Physiol* 2016; 7:288. 543
- **35.** D'Errico, J.
   Slm-shape language modeling.
   544

   http://www.mathworks.com/matlabcentral/
   545

   fileexchange/24443-slm-shape-language-modeling:
   546

   Mathworks 2009;.
   547
- 36. Zhang, Z., Henry, E., Gompper, G. & Fedosov, D. A. Behavior of rigid and deformable particles in deterministic lateral displacement devices with different post shapes. *J Chem Phys* 2015; 143:243145.
- Cluitmans, J. C. *et al.* Abnormal red cell structure and function in neuroacanthocytosis. *PLoS One* 2015; 10:e0125580.
- 38. Steffen, P. *et al.* Stimulation of human red blood cells leads to Ca2+-mediated intercellular adhesion. *Cell calcium* 2011; 50:54–61.
- Svoboda, K. & Block, S. M. Biological applications of optical forces. Annu. Rev Biophys Biomol Struct 1994; 23:247–285.
- 40. Abay, A. *et al.* Glutaraldehyde–a subtle tool in the investigation of healthy and pathologic red blood cells. *Front Physiol* 2019; 10:514.

#### 564 Supplemental Data

#### 565 Blood Smear Preparation and Evaluation

Blood smears were prepared as described by Storch et al. 566  $2005^{10}$ . Whole EDTA-blood was suspended in a 1:1 ratio in 567 a 0.9% sodium chloride solution containing 5 IU/ml heparin 568 and incubated for 1 h at room temperature. Then, a drop of 569 this solution was smeared onto a glass coverslip by using a 570 second coverslip. Dried smears were stained according to 571 the Pappenheim method. First, they were placed in May-572 Grünwald solution for 3-4 min and then washed with distilled 573 water. This was followed by staining with Giemsa solution 574 for 20 min followed by another washing step with distilled 575 water. Images were visualized with an Olympus microscope 576 (BX60, Olympus, Japan) with a  $40 \times$  Plan Fluor Objective 577 and  $2 \times$  postmagnification and recorded with a CCD camera 578 (DP73, Olympus, Japan). At least 400 cells were counted by 579 visual inspection and classified into stomatocytes, discocytes, 580 echinocytes, acanthocytes and 'other' cell shapes. 58

#### Laboratory Parameters and Erythrocyte Preparation

When required, erythrocytes were washed by centrifugation 584 (3000 rcf for 7 min). The supernatant was then removed and 585 replaced by PBS before redispensing the erythrocytes. This 586 process was repeated three times. During the last iteration, 587 the PBS was replaced by an adequate preparation (e.g., dex-588 tran diluted in PBS). Complete erythrocyte count and plasma 589 parameters were measured by standard methods in the Clin-590 ical Chemistry Laboratory of Saarland University Hospital 591 (Homburg, Germany). 592

#### 593 Widefield Microscopy

Microscopic pictures were taken using an inverted microscope 594 in transmission mode with a bright field and a blue filter to 595 enhance erythrocyte contrast with the background. A  $20\times$ 596 objective was used to send pictures to a CCD camera. Samples 597 were made by filling 30 µl well slides from Ibidi (µ-Slide 18 598 Well, Catalog no 81826, inner diameter 5 mm, interior height 599 1.5 mm) with a hematocrit of 0.33%. The suspending liquid 600 was either plasma from a control subject or a patient or a 601 preparation of dextran in PBS. The volume fraction of 0.33 % 602 was determined empirically to reach the percolation threshold, 603 i.e., the smallest concentration that could produce percolating 604 aggregates of erythrocytes on the lower side of the well when 605 erythrocytes have sedimented. The final volume fraction at 606 the bottom of the well after sedimentation is close to 45 %. 607 Pictures of an area of  $610 \times 480 \ \mu\text{m}^2$  of the sedimented layer 608 were recorded and then analyzed through a custom written 609 MATLAB algorithm. 610

#### 611 **Two-dimensional Simulations**

Two-dimensional simulations were employed to study
changes in the hole areas of aggregated mixtures of healthy
erythrocytes and acanthocytes. Both healthy cells and acanthocytes were modeled as ring-like bead-spring chains, each

consisting of 50 vertices, with a discocyte and star-like (having 616 six rounded corners) shape, respectively. Even though the dis-617 cocyte and star shapes have different areas, they have the same 618 circumference, so that the aggregation interaction between 619 cells did not depend on cell type. The total potential energy of 620 the bead-spring chain model consists of three parts<sup>36</sup>. The first 621 part is elastic energy, represented by nonlinear springs that 622 connect neighboring vertices into a ring-like configuration. 623 The second part accounts for the cellular bending resistance 624 between two neighboring springs, with a bending modulus 625 set to  $50k_BT$ . Finally, the last contribution corresponds to an 626 area constraint, which forces the discocyte shape of healthy 627 erythrocytes with a fixed circumference. Note that acantho-628 cytes are modeled as rigid bodies in simulations, as they have 629 shown no significant deformation in previous experiments<sup>37</sup>. 630

The simulations were performed in a domain of size 631  $300 \times 300 \ \mu m^2$  with periodic boundary conditions in both 632 directions. Langevin dynamics with constant temperature 633 were employed for time integration of the simulated system. 634 Aggregation interactions between cells were modeled by the 635 Lennard-Jones potential,  $U(r) = 4\varepsilon((\sigma/r)^{12} - (\sigma/r)^6)$  for 636  $r < r_{\rm cut}$  with  $\sigma = 0.3 \ \mu m$  and  $r_{\rm cut} = 0.72 \ \mu m$ . The strength of 637 the potential,  $\varepsilon$ , was selected from the range of  $1.5 - 2.5 k_B T$ 638 for different simulations to reflect aggregation variability in 639 experiments. The total hematocrit  $\phi$  was set to 50%. Two 640 sets of simulations were performed, including one with only 641 healthy erythrocytes and one with a mixture of 80% healthy 642 cells and 20% acanthocytes (NAS case) by area fraction. Each 643 set consisted of 11 simulations with different parameter val-644 ues. Note that since the area of an acanthocyte is larger than 645 that of a healthy cell, the total number of cells in the NAS 646 case is smaller than that in the healthy case. 647

648

#### Measurement of Aggregation Forces

The aggregation forces between ervthrocytes were mea-649 sured through holographic optical tweezers as previously de-650 scribed<sup>38</sup>. This is a versatile technique for measuring forces 651 in the piconewton (pN) range and allows force measurement 652 at a single-cell level. A single infrared beam Nd:YAG laser 653 (1064 nm, 3 W, Ventus 1064, Laser Quantum, UK) is reflected 654 by a parallel aligned nematic liquid crystal spatial light mod-655 ulator (PAL-SLM, PPM X8267-15, Hamamatsu Photonics, 656 Japan) and focused with a large numerical aperture oil immer-657 sion objective ( $60 \times$ , Nikon, Japan) in an inverted microscope 658 (Nikon, TE 2000, Japan). The real-time manipulation of op-659 tical traps was made possible by imaging the sample with a 660 CMOS camera (ORCA Flash 4.0 V3, Hamamatsu, Japan) and 661 using a MATLAB routine to compute the phase. The optical 662 force holding cells can be tweaked by varying the initial laser 663 power. Measurements were carried out following the protocol 664 depicted in Fig. 5C. For each measurement, two erythrocytes 665 with a discocytic shape were manually selected. Each erythro-666 cyte was held by two optical traps placed at their extremity. 667 The lower erythrocyte was brought into contact with the up-668 per erythrocyte by changing the vertical position of the traps 669

holding the erythrocyte in steps of 1 µm/s. The overlapping 670 contact length between the erythrocytes was initially set to 671 4.5 µm for all the measurements. The aggregation force corre-672 sponds to the force that is insufficient to counterbalance the 673 spontaneous aggregation forces between the two erythrocytes. 674 The same protocol was repeated to measure at least 7 pairs 675 of cells in both the pathological and healthy samples. Optical 676 tweezers were calibrated, and forces were determined based 677 on Stokes' law as in<sup>39</sup>. 678

#### **Statistical Analysis**

Statistical analysis was performed in Prism8 (GraphPad Soft-680 ware, USA). All data sets were checked for normality of the 681 distribution by the Shapiro-Wilk test. The ESR for controls, 682 ChAc and MLS patients was evaluated by the Brown-Forsythe 683 and Welch ANOVA tests. Further significance between two 684 conditions was then performed by an unpaired t-test, except if 685 otherwise stated in the figure legend. Significance p-values are 686 abbreviated as n.s. (not significant) for p>0.05, \* for p<0.05, 687 \*\* for p<0.01, \*\*\* for p<0.001 and \*\*\*\* for p<0.0001. Cor-688 relation analysis was performed by simple linear regression, 689 and the p-value provides the slope difference from zero. 690

#### 691 Confocal Microscopy and 3D Rendering

Approximately 5  $\mu$ l of blood was placed into 1 ml of 0.1 % 692 glutaraldehyde (Sigma-Aldrich, USA) solution in PBS to fix 693 cells in the shape in which they occur in the circulation<sup>40</sup>. 694 Erythrocytes (5 µl in 1 ml PBS) were stained with 5 µl of 695 CellMaskTM Deep Red plasma membrane stain (0.5 mg/ml; 696 Thermo Fisher Scientific, USA) for 24 h at room tempera-697 ture. Then, the cells were washed 3 times by centrifugation 698 at 4000 rcf for 5 min (Eppendorf Micro Centrifuge 5415 C. 699 Brinkmann Instruments, USA) in 1 ml of PBS solution. Af-700 ter washing, the cells were resuspended in PBS and finally 701 placed on a glass slide for confocal microscopy. Each la-702 beled sample was placed between two glass slides for imaging 703 (VWR rectangular coverglass,  $24 \times 60 \text{ mm}^2$ ) by employing 704 a piezo stepper for a 20 µm z-range. Confocal image genera-705 tion was performed with a spinning disk-based confocal head 706 (CSU-W1, Yokogawa Electric Corporation, Japan). Image 707 sequences were acquired with a digital camera (Orca-Flash 708 4.0, Hamamatsu Photonics, Japan). A custom written MAT-709 LAB routine was used to crop single cells from each image 710 and perform 3D reconstruction to enable visualization of the 711 3D shape of the cell. Each single-cell 3D image contained 68 712 individual planes with an extent of 100 by 100 pixels and a 713 lateral (x/y) resolution of 0.11 µm/pixel. The piezo stepper 714 had a minimal step width of 0.3 µm, defining the z-resolution. 715 To compensate for the difference in resolution in the x/y and 716 z-directions, we modified the z-scale by means of linear in-717 terpolation. Thus, the obtained z-stack had dimensions of 718  $100 \times 100 \times 185$  voxels. The image stacks were then passed 719 to a custom written ImageJ script. By applying a fixed thresh-720 old for every image, the script binarized the confocal z-stack 721 to retrieve the cell membrane as an isosurface. 722



#### Supplemental Figure 1. Sex-specific comparisons. A:

Sex-specific comparison of the ESR after 2 h. In the control group, males and females are equally distributed, while in the patient group, the higher prevalence of the disease in males is reflected by the presence of only one female ChAc patient. We did not detect any significant sex-based difference between the controls and patients. B: Comparison of hematocrit. Since gender differences in the hematocrit are known, we compared this parameter separately for males and females between control and NAS blood samples. We did not find significant differences between healthy controls and NAS patients either in the gender-specific comparison as outlined here or when male and female subjects were pooled.



**Supplemental Figure 2. Analysis of blood smears.** The blood smear from patient ChAc-5 was analyzed by two skilled staff members familiar with erythrocyte shapes. They were blinded to the intended comparison when they performed the counts. Approximately 1000 erythrocytes were counted by each analyzer. While the number of discocytes was consistent, the number of echinocytes, acanthocytes and other erythrocyte shapes (including stomatocytes and other cell shapes of known or unknown categories) varied tremendously. These differences reflect the uncertainty of acanthocyte counting based on blood smears.



**Supplemental Figure 3. Comparison of red blood cells properties.** A: Comparison of the hemoglobin concentration. For the hemoglobin (Hb) concentration, there was no significant difference between healthy controls and NAS patients. B: Comparison of the number of erythrocytes. For the number of erythrocytes (RBCs) per volume, there was no significant difference between healthy controls and NAS patients. C: Comparison of the mean cellular hemoglobin concentration. The mean cellular hemoglobin concentration (MCHC) was significantly increased in NAS patients. This is surprising since there were no differences in hematocrit (Supplemental Figure 1B), hemoglobin (panel A), number of erythrocytes (panel B) or mean cellular volume (panel D). The difference cannot be explained by the unequal gender distribution in the two groups. D: Comparison of the mean erythrocyte volume. For the mean cellular volume (MCV) of erythrocytes, there was no significant difference between healthy controls and NAS patients.



**Supplemental Figure 4. Comparison of plasma parameters.** A: Comparison of total plasma protein. The total plasma protein was reduced in NAS patients (p = 0.023), providing an initial hint for the contribution of blood plasma to the reduced ESR. B: Comparison of albumin concentration. For the albumin concentration, there was no significant difference between healthy controls and NAS patients. C: Comparison of the C-reactive protein concentration. For the C-reactive protein (CRP) concentration, there was no significant difference between healthy controls and NAS patients. D: Comparison of the immunoglobulin G concentration. For the immunoglobulin G (IgG) concentration, there is no significant difference between healthy controls and NAS patients.



**Supplemental Figure 5. Channels that appear in sedimenting blood.** The cell glass is approximately 100 µm thick, which makes the cracks visible to the naked eye. The square panel is a magnified view of the squared area in the rectangular picture. These channels are characteristic of the transient gel sedimentation regime of colloidal suspensions.





**Supplemental Figure 6. Example of time traces obtained by switching control and patient plasma.** Erythrocytes from a control subject sediments slower in patient plasma. Erythrocytes from patient MLS-2 patient sediment faster in control plasma.

Supplemental Figure 7. Comparison of fresh and

**delayed observations.** A: Comparison of ESR time traces fresh blood vs. 6 h after withdrawal. Blood samples from one healthy control and patient ChAc-5 were measured immediately after blood withdrawal and 6 h later. The delayed curves are almost indistinguishable from the fresh measurements.B: Comparison of the ESR after 2 h - fresh measurements vs. approximately 6 h of transportation. A subpopulation of the patients traveled to our laboratory in Saarbrücken to allow blood draws for immediate measurement. For the other patients, blood was collected at their local hospitals. We detected no difference in the ESR between fresh and transported blood for either the healthy controls or the NAS patients.