

Social multipliers and the Covid-19 epidemic: Analysis through constrained maximum entropy modeling

Duncan K. Foley*

08/22/20

Leo Model Professor of Economics, New School for Social Research, 79 Fifth Avenue, New York, NY 10003 USA. email: foleyd@newschool.edu

Abstract

Social multipliers occur when individual actions influence other individual actions so as to lead to amplified aggregate effects. Epidemic infections offer a dramatic example of this phenomenon since individual actions such as social distancing and masking that have small effects on individuals' risk can have very large effects in reducing risk when they are widely adopted. This paper uses the info-metric method of constrained maximum entropy modeling to estimate the impact of social multiplier effects in the Covid-19 epidemic with a model that infers the length of infection, the rate of mortality, the base infection factor, and reductions in the infection factor due to changes in social behavior from data on daily infections and deaths. While patterns are not universal over the sample of country data, they strongly support the conclusion that changes in social behavior are the primary factor influencing the dynamics of epidemics.

1 Introduction

The phenomenon of social multipliers, social interactions in which the impact of individual behavioral decisions have amplified social effects due to their impact on other members of society, are a pervasive feature of economic and social life. Social multiplier effects dominate important social-economic phenomena such as the determination of aggregate demand and employment in capitalist economies, asset prices in speculative markets, financial and economic crises, revolutionary political movements, and traffic congestion. In viral epidemics the social multiplier effect appears because individuals' attempts to protect

*I'd like to thank Amos Golan, David A. Oliver, and Nicholas C. Foley for conversations concerning the logic of this model, methods, and important features of the disease.

themselves against infection through social isolation, reduced social contact, and masking have a multiplied impact on the social spread of infection by changing the probability of that an infectious individual will infect others. The case of Covid-19 is particularly instructive in this respect, because the mean length of the infectious period of the disease is relatively short, on the order of 5-10 days, but it is so infectious that its spread through aerosols and other mechanisms in public spaces can double the number of infected also in a matter of days. Under these circumstances even small changes in behavior will have greatly amplified effects on the spread an epidemic on a short time horizon through the social multiplier. Each individual can do very little to protect themselves from a widespread and rapidly growing epidemic, but the collective effect of changes in social behavior that reduce the probability of infections can rapidly bring the epidemic under control.

This paper presents a method for the empirical analysis of epidemic statistics consisting of daily reports of new infections and deaths, and applies the model to data on national Covid-19 epidemics from the European Center for Disease Prevention and Control database. The method is the constrained maximum entropy framework described by [1]. In the context of epidemic modeling, this framework provides a method for estimating the unobserved daily rate of infection together with estimates of average recovery and mortality rates from observed daily reports of new infections and deaths.

The results of this study confirm that the expected length of Covid-19 infectiousness is on the order of 5-10 days, and that in countries, regions, and cities where the Covid-19 epidemic has been brought under control the decisive factor has been changes in social behavior acting through the social multiplier to reduce the probability of one infected individual infecting another.

2 Epidemic modeling

2.1 SIR Model

On a given day of an epidemic $t = 1, \dots, T$, let $S_t \geq 0$ be the proportion of a population uninfected and therefore susceptible, $F_t \geq 0$ the proportion of the population infected and infectious, and $R_t = 1 - S_t - F_t$ the proportion of the population either recovered or died, and therefore unsusceptible. T is the number of days over which the model tracks the epidemic, starting from day 1.

These variables follow the deterministic laws:

$$S_{t+1} = (1 - r_t F_t) S_t, t = 1, \dots, T \quad (1)$$

$$F_{t+1} = (1 - (m + d) + r_t S_t) F_t, t = 1, \dots, T \quad (2)$$

Here r_t is the infection rate for an encounter between a susceptible and infected person given the pattern of social behavior at day t , and m and d are the recovery and death rates for infections, which apply uniformly to the current infected population. In this context m is the rate at which an infected individual

stops being infectious, not necessarily the rate at which infected individuals become symptom-free. This is a version of the well-known SIR dynamic model of epidemics [2].

This type of geometric recovery and death rate implies that the expected length of an infection is $\frac{1}{m+d}$.

This model assumes that recoveries and deaths occur uniformly among the stock of currently infected individuals, which is at best an approximation, since we know the likelihood of recovery or death in fact depend on the length of the infection. Given the limitations of the data we have to work with, this approximation seems unavoidable, and the model fits reported below indicate that it does not compromise the ability of the model to track epidemics.

2.1.1 Dynamic analysis

This stripped down SIR model is a two-dimensional discrete-time nonlinear dynamical system. The linear approximation to this dynamical system is:

$$\begin{pmatrix} S_{t+1} \\ F_{t+1} \end{pmatrix} = \begin{pmatrix} 1 - r_t F_t & -r_t S_t \\ r_t F_t & r_t S_t \end{pmatrix} \begin{pmatrix} S_t \\ F_t \end{pmatrix} \quad (3)$$

The eigenvalues of this system at $F_t = 0$ are inside the unit circle, and therefore stable, if $r_t S_t < m + d$ or $\frac{r_t S_t}{m+d} < 1$. When this condition is met, epidemics die out because the reproduction factor for infections is less than unity. The model parameters r_t are the daily rate of infections, and to translate them into the more frequently used infection factor it is necessary to multiply the daily infection rate by the expected length of an infection in days, $\frac{r_t S_t}{m+d}$. For much of the data analyzed the susceptible proportion of the population remains very close to 1.

2.2 Constrained maximum entropy modeling

Constrained maximum entropy (CME) modeling represents the theoretical principles defining the system that produced the data as constraints. These constraints restrict the set of joint probability distributions over the states of the system, but typically the constraints are not sufficient to determine the joint probability distribution. The resulting underdetermination is formally resolved by ranking the feasible joint probability distributions according to their Shannon information entropy. The feasible joint probability distribution with the highest informational entropy is the CME estimate of the system's unobserved states. Ranking probability distributions by their informational entropy ensures that all the information shaping the result enters through the explicit constraints.

In the case of epidemic modeling in this paper the explicit CME formalism is:

$$\max_{\{S_t, F_t, r_t, u_{tr}, v_{tr}\}, r_0, m, d \geq 0} \sum_{t=1}^T \left(\sum_{r=1}^M -(u_{tr} \log[u_{tr}] + v_{tr} \log[v_{tr}]) \right) \quad (4)$$

$$-\gamma \sum_{t=2}^{T-1} ((r_{t+1} - r_t) - (r_t - r_{t-1}))^2 - \frac{k}{2} \sum_{t=1}^T (r_0 - r_t)^2 \quad (5)$$

$$\text{subject to} \quad (6)$$

$$S_{t+1} = (1 - r_t F_t) S_t, t = 1, \dots, T \quad (7)$$

$$F_{t+1} = (1 - (m + d) + r_t S_t) F_t, t = 1, \dots, T \quad (8)$$

$$r_0 \geq r_t \geq 0, t = 1, \dots, T \quad (9)$$

$$\frac{x_t}{\text{Pop}} = r_t S_t F_t + \sum_{r=1}^M u_{tr} q_r, t = 1, \dots, T \quad (10)$$

$$\frac{y_t}{\text{Pop}} = d F_t + \sum_{r=1}^M v_{tr} s_r, t = 1, \dots, T \quad (11)$$

$$\sum_{r=1}^M u_{tr} = 1 \quad (12)$$

$$\sum_{r=1}^M v_{tr} = 1 \quad (13)$$

Here u_{tr}, v_{tr} are error frequency distributions over the M -dimensional supports q_r, s_r , the expectations of which constitute the errors in fitting the observed daily data on infections, x_t , and mortality, y_t . The supports are chosen symmetrically to include three (this factor appears in the documentation as the “supportwidth” parameter) standard deviations of the corresponding data series, so that the maximum entropy of the frequency distributions corresponds to zero errors, or, equivalently, errors require a reduction in the entropy of the frequency distributions. The maximization of entropy of the error frequency distributions implies minimization of the errors.

The objective function has two other terms.

The first penalizes the variation in r_t represented by the square of second differences. The weight on this factor in the simulations below is $\gamma = 1000$, which results in smoothing daily fluctuations in the estimated infection rates. Experimentation indicates that the qualitative results of the simulations are robust to variations in this parameter.

The second additional term penalizes the deviation of the daily infection rate, r_t below the base infection rate, r_0 . The idea is that changing social behavior to depress r_t below the base level r_0 requires social “force” similar to the physical force depressing a spring, and results in a potential energy $\frac{k}{2}(r_0 - r_t)^2$, which trades off against the entropy of the frequency distributions of error terms on the observed data. This off-setting force we call *resistance*. The reduction in the

daily infection rate, which turns out to be critical to the stabilization of most of the epidemics for which we have data in the source for this paper, is partly a spontaneous reaction of individuals to their recognition that an epidemic is spreading by reducing their exposure, and partly a social response mediated by legal guidelines and the development of social norms concerning masking, social distancing, and the like. Since these changes in behavior have substantial economic and other costs, it makes sense to include a parameter such as k to represent those costs. In the simulations reported below typically $k = \frac{1}{10}$, a value that is in a range that produces enough resistance to changing social behavior to require a reasonable level for the infection factor $\frac{r_0}{m+d}$.

The inclusion of this term is also necessary to allow the model to identify the base infection level, r_0 . The only evidence available on which to make an estimate of this base infection rate is the growth of new infections in the first few days of the epidemic, before changes in social behavior have begun to affect r_t . But the number of new infections is an inherently noisy variable, and estimates of r_0 through extrapolation of the number of new infections turns out to be highly unstable empirically. The inclusion of the resistance in the objective function has the effect of making the estimates of the daily infection rate in the days before the epidemic takes hold uniform over time, and of stabilizing these estimates.

The logic of the simulation method is as follows. We have data on observed new infections, x_t , and deaths, y_t , for T days, and want to infer the history of the epidemic, the vectors $\{S_t\}$ and $\{F_t\}$, the unobserved daily infection rates, $\{r_t\}$ and the parameters r_0, m, d , by finding the values of these parameters that maximize the sum of the entropy of the error distributions and the penalty functions subject to the constraints (4).

2.3 Example 1: United States

Figure

Figure 1 shows the results of the simulation for data from the United States. The model estimates the average duration of the disease as about 6 days, the proportion of cases resulting in death as 3%, the infection factor without changes in social behavior about 1.74, and the resulting herd immunity threshold as requiring 42% of the population to be immune to suppress the epidemic.

The plots visualize the modeling of the epidemic. The upper-left and upper-right plots show the actual daily observed new infections and deaths in blue with the model in orange and the resulting errors in green. The general fit, which has to accommodate both the infection and mortality data, is convincing. The lower-left plot shows the initial infection factor as a horizontal orange line, and the daily infection factors and rates (which are indistinguishable since the proportion of the population susceptible to infection remains very close to 1) in red and blue, with the infection risk to susceptibles plotted in green, scaled up by a factor of 1000 to show up on the plot. The lower-right plot summarizes the epidemic dynamics in the proportion infected in blue and the proportion removed through recovery or death in orange.



6




Figure 2: Simulation for the New York City. The report has the same format as Figure 1.

An important feature of this simulation is that it indicates that the epidemic in the U.S. was controlled, to the extent it was controlled, by changes in social behavior that reduced the infection factor, not by herd immunity exhausting the pool of individuals susceptible to the infection. The red and blue curves in the lower-left plot dramatically record the fall in the infection rate a few days after the infection began to take hold, as well as a rebound in the infection rate after the first peak of the epidemic.

2.4 Example 2: New York City

Data on infections and deaths from Covid-19 are also available for New York City, from <https://www1.nyc.gov/site/doh/covid/covid-19-data.page>, accessed August 23, 2020.

Figure 2 shows the results of the simulation for data from New York City. The model estimates the average duration of the disease as about 6 days, the pro-

Stat	Infection Weighted Median	50% Prob Interval
$\frac{1}{d+m}$	5.99	0.18 8.05
$\frac{d}{d+m}$	0.03	0.02 0.03
$\frac{r(0)}{d+m}$	1.74	1.02 1.89
$1 - \frac{d+m}{r(0)}$	0.42	0.02 0.47

{k → $\frac{1}{10}$, $\gamma \rightarrow 1000$, addedconstraints → {}, supportwidth → 3, M → 3}

Table 1: Infection-weighted median values of the epidemic parameters together with a 50% confidence interval for each, computed from the empirical distributions of the parameter estimates for 89 country epidemics reporting more than 10,000 total infections in the European Center for Disease Prevention and Control Covid-19 database for 31 December, 2019 to 25 August 2020.

portion of cases resulting in death as 10%, the infection factor without changes in social behavior about 1.95, and the resulting herd immunity threshold as requiring 49% of the population to be immune to suppress the epidemic.

An important feature of this simulation is that it indicates that the epidemic in the New York was controlled, to the extent it was controlled, by changes in social behavior that reduced the infection factor, not by herd immunity exhausting the pool of individuals susceptible to the infection. The red and blue curves in the lower-left plot dramatically record the fall in the infection rate a few days after the infection began to take hold, as well as a gradual rebound in the infection rate after the peak of the epidemic, though the risk of infection remains quite low due to the small number of infectious individuals.

3 Cross-country statistics

Data from the European Center for Disease Prevention and Control reports daily infection and mortality counts for 209 countries. The program reports errors or times out on some of these country data sets, but usable simulations result in countries. Some of these cases report small total numbers of infections. Summary statistics for the modeling of the 89 country epidemics reporting more than 10,000 infections are reported in Table 1.

The estimates, which depend on extrapolating early phases of epidemics, are predictably quite noisy, but the medians, 6 days for the expected infectious period, a 3% mortality rate among the infected, and a base infection factor 1.74, are within ranges frequently reported for these parameters. This method estimates a somewhat shorter, less infectious, but more lethal disease than some other estimates.

4 Country cases

Some national epidemics show basically the same pattern as the United States: an estimate of infection length between 4 and 12 days, an infection factor be-




Figure 3: China. The sharp spikes in the infection and mortality data represent outliers that are excluded from the model simulation constraints.

between 1.5 and 3, and a sharp drop in the daily infection factor presumably as a result of changes in social behavior such as isolation and masking motivated by a combination of individual protection and social norms backed up public policy. In many instances the model estimates that the infection factor rises again after a fall in infections, in some cases heralding a resurgence of the infection.

A selection of these cases appears here in Figures 3–8.

4.1 Anomalous cases

Some country epidemic data result in very short (less than one day) or very long (over fifty days) infection lengths. In some of these cases adding a constraint that the infection length lie between 2 and 30 days results in model estimates qualitatively more in line with the previous examples. Some of these cases are presented in Figures 9–12.

The data from India cannot be modeled successfully with the infection length constraint. The Indian case modeled without constraints is presented in Figure




Figure 4: Italy

5 Conclusion and discussion

The results of this modeling effort point to some conclusions both at the level of method and at the level of understanding the social aspects of an epidemic.

The model uses limited and noisy data, and reflects both the limitations of conclusions that can be drawn from the data and the potential of the info-metric constrained maximum entropy methods to extract whatever information a data set may contain. It is particularly striking that in many cases the constrained maximum entropy model allows us to draw fairly strong inferences about the behavior of unobserved, or not directly observed, variables like the daily infection factor without making strong assumptions such as linearity. There are undoubtedly numerous other ways to approach these same data sets with the CME methods, so it would be a mistake to argue that any particular CME model exhausts the information in the data, but given the constraints constituting any particular model the CME approach does promise to extract all the information available in that framework.

The epidemiological results of this exploration reflect limits not only of the quality and extent of the data but also of the specific constraints. In this case the greatest uncertainties arise because estimates of base infection factors and the length of the infectious period have to rest on extrapolation from limited observations of the initial stages of each epidemic and, as a result, are themselves fairly noisy. Nonetheless, the general thrust of the results seems to constitute




Figure 5: France




Figure 6: Spain




Figure 7: Germany




Figure 8: Russia




Figure 11: Sweden, modeled with infection length constraint.




Figure 12: United Kingdom, modeled with infection length constraint.




Figure 13: India. The estimated infection length and infection factor indicate insufficient variation in the data for the model to work.

a strong presumptive case that infections like Covid-19, which have a half-life of infectiousness measured in a few days and are infectious enough to create an epidemic, are extremely sensitive to changes in social behavior. The simulations strongly support the view that any measures that reduce infectiousness below the stability threshold can stabilize the epidemic in a relatively short time span due to the social multiplier effects outlined above. The other side of this coin, of course, is that relaxation of social distancing that allows the infection factor to rise above unity leaves the society vulnerable to a resurgence of the infection.

Covid-19 is a particularly nasty phenomenon precisely because it is very infective but it is quite lethal. Its infection rate is not high enough to establish herd immunity without tolerating loss of life on a scale unacceptable to many societies. Fortunately an understanding of the social multiplier dynamics governing the infection show that relatively low-cost measures to reduce the infection factor can be remarkably effective in suppressing the disease.

References

- [1] Amos Golan. *Foundations of Info-Metrics*. Modeling and Inference with Imperfect Information. Oxford University Press, 2017.
- [2] David Smith and Lang Moore. The sir model for spread of disease - the differential equation model. *Convergence*, 2004.