

Diabetic retinopathy environment-wide association study (EWAS) in NHANES 2005-8.

Kevin Blighe, Ph.D.¹, Sarega Gurudas, MSC.¹, Ying Lee, Ph.D.¹, Sobha Sivaprasad, FRCOphth.^{1,2,*}

¹University College London, Institute of Ophthalmology, London, EC1V 9EL, United Kingdom.

²National Institute for Health Research, Moorfields Biomedical Research Centre, London, United Kingdom.

* corresponding author:

Professor Sobha Sivaprasad

sobha.sivaprasad@nhs.net

+44 (0)20 7566 2039

16 SUMMARY

17 **Background:** Several circulating biomarkers are reported to be associated with diabetic
18 retinopathy (DR). However, their relative contributions to DR compared to known risk
19 factors, such as hyperglycemia, hypertension, and hyperlipidemia, remain unclear. In this
20 data driven study, we used novel models to evaluate the associations of over 400 laboratory
21 parameters with DR.

22 **Methods:** We performed an environment-wide association study (EWAS) of laboratory
23 parameters available in National Health and Nutrition Examination Survey (NHANES)
24 2007-8 in individuals with diabetes with DR as the outcome (test set). We employed
25 independent variable ('feature') selection approaches, including parallelized univariate
26 regression modeling, Principal Component Analysis (PCA), penalized regression, and
27 RandomForest™. These models were replicated in NHANES 2005-6 (replication set).

28 **Findings:** The test and replication set consisted of 1025 and 637 individuals with available
29 DR status and laboratory data respectively. Glycohemoglobin (HbA1c) was the strongest risk
30 factor for DR. Our PCA-based approach produced a model that incorporated 18 principal
31 components (PCs) that had AUC 0.796 (95% CI 0.761-0.832), while penalized regression
32 identified a 9-feature model with 78.51% accuracy and AUC 0.74 (95% CI 0.72-0.77).
33 RandomForest™ identified a 31-feature model with 78.4% accuracy and AUC 0.71 (95% CI
34 0.65-0.77). On grouping the selected variables in our RandomForest™, hyperglycemia alone
35 achieved AUC 0.72 (95% CI 0.68-0.76). The AUC increased to 0.84 (95% CI 0.78-0.9) when
36 the model also included hypertension, hypercholesterolemia, hematocrit, renal and liver
37 function tests.

38 **Interpretation:** All models showed that the contributions of established risk factors of DR
39 especially hyperglycemia outweigh other laboratory parameters available in NHANES.

40

41 RESEARCH IN CONTEXT

42 What is already known about this subject?

- 43 ▪ There are >500 publications that report associations of candidate circulating
- 44 biomarkers with diabetic retinopathy (DR).
- 45 ▪ Although hyperglycemia, hypertension, and hyperlipidemia are established risk
- 46 factors, they do not always explain the variance of this complication in people with
- 47 diabetes; DR also shares risk factors with other diabetes complications including
- 48 markers of renal and cardiovascular disease.
- 49 ▪ ‘Holistic’ studies that quantify risk across all of these parameters combined are
- 50 lacking.

51

52 What is the key question?

- 53 ▪ It is unclear whether risk models for DR may be improved by adding some of these
- 54 reported biomarkers - there is an unmet need to systematically evaluate as many
- 55 circulating biomarkers as possible to help rank their associations with DR.

56

57 What are the new findings?

- 58 ▪ We show that hyperglycemia is the strongest risk factor across all models.
- 59 ▪ We stratified the rest of the highest ranked parameters into groups related to diabetes
- 60 control, renal and liver function, and hematocrit changes.

61

62 How might this impact on clinical practice in the foreseeable future?

- 63 ▪ The importance of focusing on parameters beyond hyperglycemia control to reduce
- 64 risk of progression from diabetes to DR is emphasized.

65

66 INTRODUCTION

67 Diabetes represents the most common cause of microvascular changes in the retina. The
68 initial retinal lesions of diabetic retinopathy (DR) are microaneurysms but they can occur in
69 eyes with and without diabetes (1-3). With increasing duration of diabetes, other lesions
70 develop and co-exist in the retina such as retinal hemorrhages, exudates, intraretinal
71 microvascular abnormalities and neovascularization of the retina or optic disc. Based on the
72 presence of individual lesions or a constellation of them, DR severity level is graded from
73 mild, moderate and severe non-proliferative diabetic retinopathy (NPDR) to proliferative
74 diabetic retinopathy (PDR) (4, 5). Diabetic macular edema (DME) can occur in any stage of
75 DR (5). In population-based studies, approximately a third of people with diabetes have DR
76 (6, 7). The established systemic risk factors for DR are suboptimal control of hyperglycemia,
77 hypertension and hyperlipidemia (8, 9). Hypertension can also cause some of these retinal
78 lesions independent of diabetes (10).

79 There are several laboratory parameters that have been shown to be abnormal in people with
80 DR such as hyperuricemia (11), low vitamin D levels (12), low thyroxine levels (13), anemia
81 (14), oxidative stress and inflammatory markers (15). In addition, DR is also associated with
82 markers of diabetic kidney disease including microalbuminuria and serum creatinine (16, 17)
83 and cardiovascular disease markers such as raised C-reactive protein (CRP) (18). Most of
84 these associations and risks of DR are reported based on analysis of candidate laboratory-
85 based serum or urinary markers.

86 In addition to these risk factors, there are several other non-modifiable and modifiable
87 risk factors that have been attributed to the development and progression of DR. Some
88 of these include age of onset of diabetes, duration of diabetes, male sex, and ethnicity
89 (19-21).

90 There is an unmet need to rank these reported retinal, systemic and laboratory risk
91 factors to understand their relative contributions or associations with DR in people
92 with diabetes. The National Health and Nutrition Examination Survey (NHANES -
93 <https://wwwn.cdc.gov/Nchs/Nhanes/>) was initiated in the 1960s in order to examine the
94 health and nutritional status of US citizens and has been surveying the population up to
95 the present time. Since 1999, it has examined ~5000 citizens per year and includes
96 various topics, including cardiovascular disease, diabetes, environmental exposures,

97 eye diseases, hearing loss, infectious diseases, kidney disease, nutrition, etc. The data
98 also contains several laboratory markers including environmental toxins, allergens,
99 pollutants.

100 In this study, we used an environment wide association study (EWAS) methodology
101 (22-25) on NHANES 2007-8 to evaluate the rank order of systemic and laboratory
102 risks of DR among individuals identified as having diabetes to evaluate their relative
103 associations with DR. Our findings are then replicated in NHANES 2005/6. Our
104 objective is not to only use previously reported risk factors but also provide new
105 research avenues from this data driven agnostic modelling study.

106 **METHODS**

107 **Study data preparation**

108 We used National Health and Nutrition Examination Survey (NHANES) 2007-8 as our
109 primary cohort and 2005-6 as a replication cohort. Both datasets were prepared in the
110 same fashion, however, for ease of interpretation, the following methods describe
111 2007-8. Specifically, three main categories of data were used: examination data
112 (*Ophthalmology - Retinal Imaging data*; OPXRET_E), demographics data (DEMO_E),
113 and laboratory data (**Figure 1** footnote). The main outcome of interest in the
114 examination data was *4 levels retinopathy severity, worse eye* (OPDURL4) – this
115 variable was recoded as binary with levels: no retinopathy; retinopathy (including mild
116 NPR, moderate/severe NPR, and proliferative). All datasets were downloaded as SAS
117 XPORT (xpt) format and read into R (v4.0.2) via the *Hmisc* package.

118 Individuals with a missing value in the main outcome variable were removed before
119 aligning the examination, demographics, and laboratory data via each individual's
120 respondent sequence number (SEQN). This dataset was then further filtered for only
121 those individuals who had diabetes (**Figure 1**). Variables were removed from the data
122 that had 0 variance (i.e. constant values) (**Supplementary Table 1**). Prior to any
123 analysis, in addition, any variable that contained a single value occupying > 90% of
124 total values was removed, as were variables that had > 90% missingness. Further
125 specific filtering and encoding was then applied per dataset. [A] Examination data:
126 variables that were different encodings of the main outcome were removed; variables

that related to the status of the examination appointment were removed; OPDUHMA was removed, as it is a combination of 2 other variables that were retained (OPDUMA and OPDUHEM); variables related to glaucoma, for which there is already a single variable, were removed; variables related to the left or right eye where there was already a variable for ‘worse’ eye were removed; values encoded as missing were recoded as NA; and all other remaining variables were encoded as binary, with 0 representing the absence of the condition, and 1 representing any recorded presence (at any level) of the condition. [B] Demographic data: variables associated with interpreters and the language of the interview were removed; variables that were duplicates or different encoding of each were removed. [C] Laboratory data: categorical variables were removed and only continuous retained; duplicate variables related to the oral glucose tolerance test (OGTT_E) were removed; variables related to time since domestic activities (‘pump gas’, ‘shower’, etc.) were removed; variables that were duplicates or different encodings of each were removed; variables measured on the imperial system of weights and measures were removed if they had a corresponding variable in SI units. We focused only on continuous laboratory variables for the following reasons: 1, in NHANES, the majority of categorical variables are derived from the continuous variables; 2, our PCA-based approach can only work on continuous variables; 3, for RandomForest™, having continuous variables increases the number of splitting points in the data, and metrics of importance such as Gini are known to exhibit less bias on such data (26).

Diabetes status

To define the diabetes status for each individual, questionnaire data (DIQ_E) was used in addition to variables already included in the laboratory data. Diabetes status was then defined as an individual satisfying any of the following: Self-reported diabetes (DIQ010); on anti-diabetes drugs (DIQ070); taking insulin (DIQ050); fasting blood sugar (FBS) ≥ 6.1 (110mg/dl) (LBDGLUSI); random blood sugar (RBS) ≥ 11.1 (200mg/dl) (LBDSGLSI); oral glucose tolerance test (OGTT) ≥ 200 mg/dl (LBDGLTSI); Glycohemoglobin (HbA1c) $\geq 6.5\%$ (LBXGH).

Covariates

Age, ethnicity, and diabetes duration were used as covariates. Diabetes duration was calculated as age at screening minus the age at which the individual was first informed that he/she had diabetes.

Statistical analysis

Prior to statistical analysis, continuous laboratory variables were logged (\log_e) and then transformed into z-scores to ensure that these were on the same scale. In regression analysis, the complex sampling design of the NHANES dataset was accounted for through use of survey sampling weights via the *survey* package in R / CRAN. To do this, the following value-pairs were used with the *svydesign* function: (*id*, SDMVPSU; *strata*, SDMVSTRA; *weights*, WTMEC2YR; *nest*, TRUE).

Univariate analysis was performed on all candidate predictors using a survey-weighted compute-parallelized logistic regression model via the R / Bioconductor package *RegParallel*, adjusting for age, ethnicity, and duration of diabetes separately. The Benjamini-Hochberg (27) procedure was used to control the type I error false discovery rate (FDR). A customized Manhattan plot was generated using *ggplot2*, while pairwise scatter and correlation plots were generated via a customized pairs plot. Finally, a heatmap was generated via the R / Bioconductor package *ComplexHeatmap*.

As our study is also hypothesis-generating, multivariate approaches based on principal component analysis (PCA), penalized regression, and the RandomForest™ classification algorithm were additionally used. Variables were pre-filtered and prepared as per univariate testing. Principal component analysis was performed via the R / Bioconductor package *PCAtools*. After conducting PCA, each eigenvector was then independently regressed against retinopathy outcome via binary logistic regression and those that passed $p \leq 0.05$ were used to construct a multivariable model that was further tested in ROC analysis via the *pROC* package in R.

Separately, as model complexity and multi-collinearity can arise from a large number of predictors, elastic net regularization (penalized regression with L1 and L2 penalties of the Lasso and Ridge methods) was used to reduce the number of predictor variables using *glmnet* in R / CRAN. To fit the model, 100x cross-validation was used and alpha (α) set to 0.5. The final chosen variables were those whose coefficients were not

shrunk to zero – these were plot as violin plots with scatter overlays to show differences between non-DR and DR via *ggplot2*. To determine accuracy, model predictions were made on the data using the lambda (λ) one-standard-error rule using the *predict* function from the *stats* package in R.

Finally, the RandomForest™ (RF) model was fitted via the *randomForest* R / CRAN package. For this, the dataset was divided randomly into 50% training and 50% validation. Prior to model fitting, the initial model was tuned using functionality provided by the *caret* package in R / CRAN, as follows: 1), a 10x cross-validation control function was defined via *trainControl* function; 2) the best value for ‘mtry’, i.e., the ideal number of variables to randomly sample, was determined using the *train* function across a search / tuning grid ranging between 1-40 and with Kappa as the metric; and 3) using the selected value of ‘mtry’, the ideal number of trees, ‘ntrees’, was determined also via the *train* function with selection metric based on Kappa. After the initial model was fit, variables with mean decrease in accuracy $\leq 1\%$ were excluded and the model re-fit. This was then repeated in a recursive fashion until all variables with negative mean decrease accuracy were removed from the model.

Final risk models

Variables selected from RandomForest™ were grouped based on similarity of function or clinical use. Each group was then used to create independent univariate or multivariable binary logistic regression models with DR as the end-point. A single Wald test p-value was derived for each model using *wald.test* from the *aod* package. ROC analysis was performed using *pROC*. McFadden’s and Nagelkerke’s pseudo- R^2 were derived via the *pscl* and *rms* packages, respectively.

Role of the funding source

The funders had no role in study design, data collection, data analysis, data interpretation, writing, editing the report, or the decision to submit for publication.

RESULTS

Study cohort

In NHANES 2007-8, retinal imaging data is available for 3863 individuals, demographics data is available for 10149 individuals, and laboratory data is available for between 394 and

9307 individuals, depending on the individual laboratory dataset in NHANES (see **Figure 1** footnote). After aligning all data and filtering for those who had diabetes by our classification, 1025 individuals remained in our dataset. The selection process is illustrated in **Figure 1**, while **Table 1** provides an overview of the demographics of these individuals.

For our replication cohort, NHANES 2005-6, we prepared laboratory data following the same filter criteria as NHANES 2007-8 and produced a final dataset of 2459 individuals, among which 637 (with retinopathy, 176; no retinopathy, 461) had diabetes.

Retinal lesions of diabetic retinopathy

To help validate our methodology and cohort selection, we aimed to determine retinal lesions that define DR. To this end, we identified 9 retinal lesions in NHANES 2007-8 that were statistically significantly associated with DR and survived to p-value adjustment for false discovery (**Table 2**). The top lesions were retinal microaneurysms ($p \leq 0.0001$), followed by retinal hard exudates (typically due to lipoprotein deposition in the retina and may be associated with macular edema) ($p \leq 0.0001$). Other key lesions at $p \leq 0.0001$ were retinal soft exudate (now termed cotton wool spots), retinal blot hemorrhages, intraretinal microvascular abnormalities (IRMA), and macular edema. In NHANES, retinal microaneurysms and retinal blot hemorrhages are encoded to be mutually exclusive, i.e., an individual is recorded as having retinal microaneurysms only when not accompanied with retinal blot hemorrhages, and *vice-versa* (**Table 2**).

Univariate logistic regression analysis

In total, 6 variables reached statistical significance in the unadjusted univariate analysis, 11 after adjustment for age, 2 after adjustment for ethnicity, and 7 for diabetes duration (**Supplementary Figure 1; Table 3**). Glycohemoglobin (HbA1c) was the only variable that was statistically significant in both the unadjusted and adjusted analyses. Other risk variables of note that reached statistical significance in the unadjusted analysis included serum glucose (mmol/L) (i.e., RBS), osmolality (mmol/Kg), urinary albumin (mg/L), and fasting glucose (mmol/L) (i.e., FBS). The only protective variable, i.e., negatively associated, was hemoglobin (g/dL). These variables indicate suboptimal diabetes control, abnormal kidney function and presence of anemia as risk factors for DR. There was evidence of co-correlation among these statistically significant variables from the unadjusted analysis (**Supplementary Figure 2**).

Interestingly, after adjustment for diabetes duration, the following variables reached statistical significance: HbA1c (%), osmolality (mmol/Kg), urinary iodine ($\mu\text{g/L}$), urinary cobalt ($\mu\text{g/L}$), urinary triclosan (ng/mL), urinary creatinine ($\mu\text{mol/L}$), and urinary barium ($\mu\text{g/L}$).

Principal component analysis

Unsupervised PCA using all laboratory variables revealed that 59 PCs could account for 80% or more variation in the dataset. Eighteen PCs were statistically significantly associated with DR at $p \leq 0.05$ via independent binomial regression models testing each PC (**Supplementary Table 2**). The top variables responsible for variation along these PCs included measures of blood glucose (HbA1c, random blood glucose and fasting blood glucose), kidney function markers (urinary albumin, blood urea nitrogen [BUN]), hematological markers (hematocrit, hemoglobin, red blood cell distribution width), inflammatory markers (CRP), white blood cell count, urinary nitrates, segmented neutrophil count), and toxic elements (urinary beryllium and cotinine) among others – these PCs were also statistically significantly correlated to microaneurysms, the previously-identified top retinal lesion, and the covariates used during univariate testing (**Supplementary Figure 3**). Through ROC analysis, these 18 PCs achieved AUC 0.796 (95% CI: 0.761-0.832).

Penalized regression model

We fitted an unbiased elastic-net penalized regression model to the laboratory variables and cross-validated it 100x. The model selected 9 variables whose coefficients were not shrunk to zero: urinary albumin, BUN, urinary cobalt, CRP, HbA1c, blood osmolality, serum potassium, systolic blood pressure, and urinary nitrate (**Figure 2**). Of note, these measurements mainly represent diabetes and blood pressure control and kidney function. This model had an accuracy of 78.51% and AUC 0.74 (95% CI: 0.72-0.77) when predicted on the same dataset on which the model was produced.

RandomForest™ classification model

From our RandomForest™ model, HbA1c was the single best predictor of DR (mean decrease accuracy, 31.94%; Gini, 21.75) (**Table 4**). However, other notable variables of appreciable accuracy were markers of diabetes control (FBS, RBS) inflammation (CRP), kidney function (potassium, BUN, creatinine and urinary albumin), hematological markers

(hematocrit), systolic blood pressure, among others. The overall accuracy of the model on the validation cohort was 78.4% and AUC 0.71 (95% CI: 0.65-0.77).

Replication cohort

In the NHANES 2005-6 replication cohort, we performed penalized regression and RandomForest™ in the same way as per the 2007-8 cohort. Our penalized regression model identified urinary albumin (mg/L), cockroach IgE antibody (kU/L), HbA1c (%), hemoglobin (g/dL), and urinary nitrate (ng/mL), with a model accuracy of 73.16% and AUC 0.76 (95% CI: 0.73-0.78). RandomForest™ identified HbA1c (%) as the variable contributing most to accuracy (mean decrease 16.96%), with many other variables contributing appreciable accuracy to the overall model (**Supplementary Table 3**) - the overall model accuracy was 72.98% and AUC 0.68 (95% CI: 0.61-0.75).

Final clinical risk models

The 31 features identified by RandomForest™ (**Table 4**) were grouped into different categories of blood tests according to diabetes status, hematocrit values, blood pressure (BP), immune markers, renal function, sterols, toxins and metals, and liver function. When modeled against DR outcome, each group varied in performance; diabetes tests alone achieved AUC 0.72 (95% CI: 0.68-0.76). A final clinical risk model comprising diabetes tests, BP, renal and liver function tests, hematocrit values, circulating sterols and immune markers achieved AUC 0.84 (95% CI: 0.78-0.9) ($p=0.00013$) (Nagelkerke R^2 0.36) (**Table 5; Figure 3**).

DISCUSSION

This EWAS of NHANES 2007-8 data with DR outcomes in individuals with diabetes included an unbiased feature selection approach based on a rudimentary univariate regression enabled for compute parallelization, PCA, penalized regression, and RandomForest™ of a large number of laboratory parameters. In contrast, epidemiological studies are typically conducted based on pre-conceived hypotheses and involve a single or just a few variables. These methods can be scaled to datasets of any size and therefore provide ways of working with large clinical and epidemiological datasets for the purpose of searching for novel hypotheses that could then lead to further focused investigations.

In our rudimentary approach, which is ultimately running many univariate models in a parallelized fashion, HbA1c was the only variable to reach statistical significance after adjusted for age, ethnicity, and diabetes duration. The relationship between HbA1c and DR has been explored extensively and was selected as the strongest risk factor in every approach we undertook, with a mean decrease accuracy of 31.94% via RandomForest™.

Our penalized regression and RandomForest™ algorithms also identified an association between elevated systolic blood pressure—but not diastolic—and DR (mean decrease accuracy, 1.9%), again confirming literature (10, 28-30). Further risk variables identified by both penalized regression and RandomForest™ were renal function tests including BUN, urinary albumin, potassium, osmolality, and urinary nitrate. These confirm the strong association of DR with markers of impaired kidney function. Other known risk factors that contributed higher up in the ranking order include hematocrit (%) and cholesterol. Although HbA1c has the strongest association with DR, our study highlight how the addition of other clinical parameters, e.g., from renal and liver function and hematocrit can increase the sensitivity and specificity of predicting DR outcome, with our final clinical risk model achieving AUC 0.83 (95% CI: 0.77-0.89) ($p=0.00012$) (Nagelkerke R^2 0.33), higher than any traditional diabetes control parameter in isolation or in combination.

The EWAS methodology and our RandomForest™ approach of non-targeted recursive feature also indicates a small contribution from toxins and metals, including 3-hydroxyphenanthrene, 9-hydroxyfluorene, phthalates, blood o-Xylene, and blood nitromethane. Therefore, retina may be a target tissue for environmental contamination. Some of the associations provide directions to future mechanistic research in DR. For example, we found that retinal microaneurysms (FDR-adjusted $p \leq 0.0001$), the most statistically significant retinal lesions in individuals with DR, is already correlated with some of the variables such as HbA1c, CRP, BUN, beryllium, and hematocrit, suggesting early effects. In contrast, increased urinary cobalt, triclosan and barium became significant only when adjusted for duration of diabetes. Most of these parameters are also linked to risk of allergies and lung disease, an association that has not been previously explored systematically.

As this is a cross-sectional study, a cause-effect relation cannot be established. Moreover, we are unable to rule out any confounding effects of any unmeasured factors. On the other hand, the main strength of the study is the use of the well characterised NHANES cohort in whom standardised protocols were used to measure laboratory parameters. We are not aware of any

other association studies in DR where over 400 laboratory parameters were analysed simultaneously to develop multiple models. As the top variables of all four data driven agnostic models were similar, we also believe our findings are generalisable.

CONCLUSION

We confirm that DR is a complex disease and that the already established risk factors contribute significantly to the risk models of DR, with HbA1c being the strongest risk factor. Although our model provides an accuracy of approximately 80%, it also provides mechanistic insights into future research on DR including interrogating the interaction of low-ranking risk factors with more established factors in the models and highlights need to explore epigenetic screens to gauge better how risk factors influence gene expression. Most importantly, the study reinforces the need to control known risk factors of DR, especially hyperglycemia.

ACKNOWLEDGEMENTS

We thank the many thousands of NHANES study participants who, over the course of decades, have provided valuable information for epidemiological studies.

FUNDING

This work was funded by Global Challenges Research Fund and UK Research and Innovation through the Medical Research Council grant number MR/P027881/1. The research was supported] by the National Institute for Health Research (NIHR) Biomedical Research Centre based at Moorfields Eye Hospital NHS Foundation Trust and UCL Institute of Ophthalmology. The views expressed are those of the author(s) and not necessarily those of the NHS, the NIHR or the Department of Health.

AUTHOR CONTRIBUTIONS

KB, SG and SS conceived and designed the study. KB, SG and YL analysed the data. KB is the study data guarantor. All authors interpreted the results and reviewed the manuscript. KB had full access to all of the data and takes responsibility for the integrity of the data and the accuracy of data analysis. All authors read and approved the final manuscript.

TABLES

Table 1. Demographic overview of study cohort.

Characteristics mean (\pm SD) n (%)		Diabetes with no diabetic retinopathy (n=787)	Diabetic retinopathy (n=238)	p-value	β -coefficient	OR (95% CI)
Age	-	62.39 (\pm 11.02)	63.53 (\pm 10.54)	0.16	0.01	1.00 (1.00-1.02)
Sex	Male	420 (53.37)	126 (52.94)	-	-	-
	Female	367 (46.63)	112 (47.06)	0.91	0.02	1.02 (0.76-1.36)
Ethnicity	Non-hispanic white	369 (46.88)	91 (38.23)	-	-	-
	Non-hispanic black	166 (21.09)	74 (31.09)	0.0012	0.59	1.81 (1.27-2.58)
	Mexican-American	138 (17.54)	41 (17.23)	0.38	0.19	1.21 (0.79-1.83)
	Other Hispanic	89 (11.31)	28 (11.77)	0.32	0.24	1.28 (0.79-2.07)
	Other race - including multi-racial	25 (3.18)	4 (1.68)	0.43	-0.43	0.65 (0.22-1.91)
Education	Less than 9 th grade	148 (18.8)	54 (22.69)	-	-	-
	9-11 th grade [±]	148 (18.8)	51 (21.43)	0.8	-0.06	0.94 (0.61-1.47)
	High school graduate / GED or equivalent	199 (25.29)	56 (23.53)	0.24	-0.26	0.77 (0.5-1.19)
	Some college or AA degree	172 (21.86)	55 (23.11)	0.55	-0.13	0.88 (0.57-1.35)
	College graduate or above	120 (15.25)	22 (9.24)	0.014	-0.69	0.5 (0.29-0.87)
Marital status	Married	464 (58.96)	133 (55.88)	-	-	-
	Widowed	119 (15.12)	36 (15.13)	0.8	0.05	1.06 (0.69-1.61)
	Divorced	90 (11.44)	37 (15.55)	0.099	0.36	1.43 (0.94-2.2)
	Separated	32 (4.07)	7 (2.94)	0.53	-0.27	0.76 (0.33-1.77)
	Never married	53 (6.73)	18 (7.56)	0.56	0.17	1.19 (0.67-2.09)
	Living with partner	29 (3.68)	7 (2.94)	0.69	-0.17	0.84 (0.36-1.97)
Family income	\$0-\$4999	11 (1.4)	3 (1.26)	-	-	-
	\$5000-\$9999	47 (5.97)	12 (5.04)	0.93	-0.07	0.94 (0.23-3.89)
	\$10000-\$14999	82 (10.42)	22 (9.25)	0.98	-0.02	0.98 (0.25-3.84)
	\$15000-\$19999	68 (8.64)	22 (9.25)	0.81	0.17	1.19 (0.30-4.64)
	\$20000-\$24999	69 (8.77)	25 (10.5)	0.68	0.28	1.33 (0.34-5.16)
	\$25000-\$34999	103 (13.09)	42 (17.65)	0.55	0.40	1.50 (0.40-5.63)
	\$35000-\$44999	68 (8.64)	23 (9.66)	0.76	0.22	1.24 (0.32-4.84)
	\$45000-\$54999	55 (6.99)	12 (5.04)	0.76	-0.22	0.80 (0.19-3.31)
	\$55000-\$64999	41 (5.21)	15 (6.3)	0.68	0.29	1.34 (0.33-5.48)
	\$65000-\$74999	36 (4.57)	6 (2.52)	0.53	-0.49	0.61 (0.13-2.86)
	\$75000-\$99999	44 (5.59)	12 (5.04)	1.00	0.00	1.00 (0.24-4.17)
	\geq \$100000	88 (11.18)	17 (7.14)	0.62	-0.35	0.71 (0.18-2.81)
	Over \$20000	31 (3.94)	9 (3.78)	0.93	0.06	1.06 (0.24-4.66)
	Under \$20000	17 (2.16)	2 (0.84)	0.4	-0.84	0.43 (0.06-3.01)
	Missing	27 (3.43)	16 (6.73)	-	-	-
Diabetes duration	-	9.05 (\pm 11.05)	16.33 (\pm 12.57)	\leq 0.0001	0.05	1.05 (1.04-1.07)

Notes: This table only relates to those individuals who have been determined as having diabetes by our selection criteria: 1, Self-reported diabetes (DIQ010); 2, On anti-diabetes drugs (DIQ070); 3, Taking insulin (DIQ050); 4, Fasting blood sugar (FBS) \geq 6.1 (110mg/dl) (LBDGLUSI); 5, Random blood sugar (RBS) \geq 11.1 (200mg/dl)

(LBDSGLSI); 6, Oral glucose tolerance test (OGTT) $\geq 200\text{mg/dl}$ (LBDGLTSI); 7, Glycohemoglobin (HbA1c) $\geq 6.5\%$ (LBXGH). Ethnicity, education, and diabetes duration contain at least one term that is statistically significant. \pm includes 12th grade with no diploma

Table 2. Retinal lesions that constitute diabetic retinopathy outcome.

Retinal co-morbidity	n (%)	β -coefficient	OR (95% CI)	p-value	FDR-adjusted p-value
Retinal microaneurysms only, worse eye	129 (12.59)	5.67	288.87 (127.66-653.68)	≤ 0.0001	≤ 0.0001
Retinal hard exudate, worse eye	86 (8.39)	3.72	41.34 (20.31-84.17)	≤ 0.0001	≤ 0.0001
Retinal blot hemorrhages, worse eye	47 (4.59)	3.36	28.71 (14.11-58.42)	≤ 0.0001	≤ 0.0001
Retinal soft exudate, worse eye	76 (7.41)	4.2	66.49 (26.44-167.21)	≤ 0.0001	≤ 0.0001
IRMA, worse eye	62 (6.05)	2.85	17.36 (9.06-33.26)	≤ 0.0001	≤ 0.0001
Macular edema, worse eye	51 (4.98)	3.88	48.24 (17.18-135.44)	≤ 0.0001	≤ 0.0001
Retinal fibrous proliferation, worse eye	19 (1.85)	3.41	30.19 (6.92-131.67)	≤ 0.0001	≤ 0.0001
Macular edema in center, worse eye	26 (2.54)	4.53	92.33 (12.45-684.9)	≤ 0.0001	≤ 0.0001
Retinal new vessels elsewhere, worse eye	15 (1.46)	3.12	22.68 (5.08-101.23)	≤ 0.0001	0.0003

Notes: Lesions are taken from the NHANES *ophthalmology - retinal imaging* (OPXRET_E) dataset. Only lesions with FDR-adjusted $p \leq 0.05$ are listed. Soft exudate is now termed cotton wool spots.

Table 3. Laboratory variables associated with retinopathy in individuals with diabetes.

Description	Unadjusted / Non-covariate adjusted				Age-adjusted				Ethnicity-adjusted				Diabetes duration-adjusted			
	β -coefficient	OR (95% CI)	p-value	FDR-adjusted p-value	β -coefficient	OR (95% CI)	p-value	FDR-adjusted p-value	β -coefficient	OR (95% CI)	p-value	FDR-adjusted p-value	β -coefficient	OR (95% CI)	p-value	FDR-adjusted p-value
Glycohemoglobin (%)	0.82	2.27 (1.84-2.8)	0.000	0.0003	0.85	2.34 (1.87-2.92)	0.000	0.0011	0.83	2.28 (1.82-2.87)	0.000	0.0036	0.72	2.05 (1.55-2.73)	0.002	0.0341
Glucose, serum (mmol/L)	0.54	1.72 (1.42-2.08)	0.001	0.0047	0.57	1.77 (1.45-2.15)	0.001	0.0115	0.54	1.71 (1.41-2.09)	0.002	0.0334	0.36	1.43 (1.1-1.86)	0.0180	0.0520
Osmolality (mmol/Kg)	0.49	1.63 (1.34-1.99)	0.002	0.0143	0.45	1.57 (1.3-1.9)	0.004	0.0127	0.49	1.63 (1.34-1.99)	0.005	0.0707	0.39	1.48 (1.12-1.94)	0.0140	0.0415
Albumin, urine (mg/L)	0.45	1.57 (1.28-1.93)	0.006	0.0288	0.43	1.53 (1.24-1.88)	0.012	0.0127	0.42	1.53 (1.25-1.87)	0.017	0.1418	0.25	1.28 (0.98-1.68)	0.0905	0.1617
Hemoglobin (g/dL)	-0.33	0.72 (0.61-0.85)	0.013	0.0387	-0.30	0.74 (0.63-0.88)	0.042	0.0234	-0.29	0.75 (0.62-0.9)	0.094	0.2638	0.00	1 (0.73-1.37)	0.9784	0.9883
Fasting Glucose (mmol/L)	0.49	1.63 (1.28-2.07)	0.011	0.0387	0.51	1.66 (1.3-2.13)	0.013	0.0127	0.46	1.59 (1.25-2.02)	0.031	0.2290	0.22	1.24 (0.93-1.66)	0.1687	0.2731
4-(methylnitrosamino)-1-(3-pyridyl)-1-butanol (NNAL), urine (ng/mL)	-0.26	0.77 (0.67-0.88)	0.021	0.0539	-0.20	0.82 (0.71-0.94)	0.133	0.0552	-0.28	0.75 (0.65-0.87)	0.033	0.2290	-0.25	0.78 (0.52-1.19)	0.2655	0.4061
Iodine, urine (ug/L)	-0.17	0.84 (0.76-0.93)	0.039	0.0873	-0.21	0.81 (0.73-0.9)	0.022	0.0156	-0.15	0.86 (0.77-0.96)	0.181	0.2638	-0.29	0.75 (0.63-0.89)	0.054	0.0341
Cobalt, urine (ug/L)	-0.51	0.6 (0.44-0.82)	0.052	0.1033	-0.50	0.6 (0.45-0.82)	0.058	0.0300	-0.51	0.6 (0.44-0.81)	0.073	0.2638	-0.52	0.59 (0.45-0.78)	0.021	0.0341
Hematocrit (%)	-0.31	0.73 (0.6-0.89)	0.061	0.1089	-0.28	0.76 (0.62-0.92)	0.148	0.0595	-0.28	0.75 (0.62-0.92)	0.202	0.2638	-0.03	0.97 (0.7-1.35)	0.8787	0.9329
Blood urea nitrogen (mmol/L)	0.33	1.4 (1.13-1.73)	0.075	0.1225	0.27	1.31 (1.01-1.71)	0.0645	0.1584	0.35	1.43 (1.16-1.75)	0.066	0.2638	0.15	1.17 (0.87-1.57)	0.3233	0.4793
Albumin (g/L)	-0.22	0.8 (0.69-0.93)	0.116	0.1736	-0.21	0.81 (0.69-0.94)	0.183	0.0687	-0.19	0.83 (0.7-0.99)	0.575	0.3225	-0.09	0.92 (0.74-1.14)	0.4412	0.5899
Urinary Triclosan (ng/mL)	-0.42	0.65 (0.49-0.88)	0.126	0.1743	-0.40	0.67 (0.5-0.89)	0.165	0.0639	-0.42	0.66 (0.49-0.89)	0.195	0.2638	-0.60	0.55 (0.36-0.83)	0.0135	0.0412
Mean cell hemoglobin (pg)	-0.24	0.79 (0.66-	0.159	0.2046	-0.27	0.77 (0.65-	0.076	0.0368	-0.17	0.84 (0.7-	0.1069	0.4855	-0.01	0.99 (0.76-	0.9174	0.9489

		0.93)				0.91)				1.02)				1.28)		
Lead, urine (µg/L)	-0.40	0.67 (0.49- 0.91)	0.0 222	0.2668	-0.40	0.67 (0.49- 0.91)	0.0 230	0.0831	-0.41	0.66 (0.49- 0.9)	0.0 228	0.2638	-0.43	0.65 (0.43- 0.99)	0.0 658	0.1312
Creatinine, urine (µmol/L)	-0.22	0.8 (0.67- 0.97)	0.0 348	0.2723	-0.19	0.83 (0.68-1)	0.0 710	0.1664	-0.27	0.77 (0.64- 0.92)	0.0 156	0.2638	-0.28	0.76 (0.64- 0.9)	0.0 070	0.0341
Alanine aminotransferase (ALT) (U/L)	-0.25	0.78 (0.62- 0.97)	0.0 380	0.2723	-0.20	0.82 (0.66- 1.02)	0.0 965	0.2003	-0.22	0.81 (0.64- 1.01)	0.0 869	0.4259	-0.09	0.92 (0.68- 1.23)	0.5 656	0.6993
Creatinine (µmol/L)	0.24	1.27 (1.04- 1.54)	0.0 335	0.2723	0.18	1.19 (0.96- 1.48)	0.1 291	0.2529	0.20	1.23 (0.99- 1.53)	0.0 933	0.4362	0.13	1.14 (0.89- 1.45)	0.3 111	0.4699
Red blood cell count (million cells/µL)	-0.19	0.82 (0.7- 0.97)	0.0 372	0.2723	-0.13	0.88 (0.74- 1.04)	0.1 439	0.2740	-0.19	0.83 (0.7- 0.98)	0.0 540	0.3172	0.01	1.01 (0.74- 1.37)	0.9 623	0.9769
Mean cell volume (fL)	-0.21	0.81 (0.68- 0.98)	0.0 463	0.2723	-0.25	0.78 (0.65- 0.93)	0.0 159	0.0623	-0.15	0.86 (0.71- 1.06)	0.1 865	0.6799	-0.05	0.95 (0.72- 1.26)	0.7 361	0.8368
Platelet count (1000 cells/µL)	-0.21	0.81 (0.68- 0.98)	0.0 435	0.2723	-0.18	0.83 (0.68- 1.02)	0.1 005	0.2049	-0.23	0.79 (0.67- 0.94)	0.0 216	0.2638	-0.29	0.75 (0.57- 0.99)	0.0 616	0.1250
Mean platelet volume (fL)	0.24	1.27 (1.04- 1.55)	0.0 352	0.2723	0.26	1.3 (1.06- 1.59)	0.0 243	0.0835	0.22	1.25 (1.02- 1.53)	0.0 564	0.3225	0.09	1.09 (0.83- 1.43)	0.5 336	0.6642
Cotinine (ng/mL)	-0.15	0.86 (0.76- 0.99)	0.0 451	0.2723	-0.09	0.91 (0.79- 1.04)	0.1 994	0.3443	-0.18	0.84 (0.74- 0.95)	0.0 221	0.2638	-0.06	0.94 (0.66- 1.34)	0.7 352	0.8368
Insulin (pmol/L)	-0.32	0.73 (0.55- 0.97)	0.0 469	0.2723	-0.30	0.74 (0.55-1)	0.0 699	0.1660	-0.29	0.75 (0.56-1)	0.0 734	0.3816	-0.22	0.8 (0.51- 1.26)	0.3 520	0.5062
Blood cadmium (nmol/L)	-0.23	0.8 (0.65- 0.97)	0.0 429	0.2723	-0.25	0.78 (0.64- 0.95)	0.0 273	0.0866	-0.23	0.79 (0.64- 0.98)	0.0 589	0.3225	-0.28	0.76 (0.5- 1.16)	0.2 224	0.3528
Urinary perchlorate (ng/mL)	-0.25	0.78 (0.63- 0.95)	0.0 294	0.2723	-0.24	0.78 (0.63- 0.98)	0.0 495	0.1267	-0.24	0.78 (0.64- 0.96)	0.0 399	0.2668	-0.31	0.73 (0.53- 1.01)	0.0 818	0.1525
Urinary nitrate (ng/mL)	-0.28	0.75 (0.6- 0.94)	0.0 264	0.2723	-0.23	0.79 (0.63-1)	0.0 675	0.1632	-0.27	0.76 (0.6- 0.96)	0.0 422	0.2688	-0.34	0.71 (0.55- 0.91)	0.0 186	0.0528
Cesium, urine (µg/L)	-0.33	0.72 (0.54- 0.95)	0.0 339	0.2723	-0.32	0.72 (0.55- 0.96)	0.0 390	0.1059	-0.33	0.72 (0.54- 0.95)	0.0 389	0.2665	-0.34	0.71 (0.47- 1.06)	0.1 161	0.1950
Thallium, urine (µg/L)	-0.40	0.67 (0.48- 0.95)	0.0 417	0.2723	-0.39	0.68 (0.48- 0.96)	0.0 488	0.1266	-0.42	0.66 (0.47- 0.92)	0.0 307	0.2638	-0.48	0.62 (0.4- 0.94)	0.0 414	0.0923
25OHD2+25OHD3 (nmol/L)	-0.24	0.79	0.0	0.2723	-0.25	0.78	0.0	0.0831	-0.18	0.83	0.1	0.5685	-0.20	0.82	0.2	0.3401

		(0.65-0.95)	279			(0.65-0.94)	230			(0.66-1.04)	390			(0.61-1.1)	129	
Blood Toluene (ng/mL)	-0.21	0.81 (0.68-0.96)	0.0 304	0.2723	-0.20	0.82 (0.69-0.97)	0.0 366	0.1029	-0.22	0.8 (0.65-0.98)	0.0 578	0.3225	-0.30	0.74 (0.45-1.21)	0.2 536	0.3896
C-reactive protein (mg/dL)	-0.19	0.82 (0.69-0.99)	0.0 551	0.3102	-0.18	0.83 (0.69-1.01)	0.0 827	0.1820	-0.25	0.78 (0.66-0.92)	0.0 153	0.2638	-0.27	0.77 (0.6-0.98)	0.0 489	0.1063
Barium, urine (µg/L)	-0.39	0.68 (0.47-0.99)	0.0 600	0.3114	-0.38	0.69 (0.48-0.99)	0.0 607	0.1525	-0.38	0.68 (0.47-1)	0.0 756	0.3885	-0.44	0.64 (0.47-0.87)	0.0 140	0.0415
Urinary 4-tert-octylphenol (ng/mL)	-0.34	0.71 (0.45-1.12)	0.1 607	0.5259	-0.33	0.72 (0.46-1.12)	0.1 682	0.3007	-0.42	0.66 (0.43-1.01)	0.0 846	0.4203	-0.82	0.44 (0.23-0.87)	0.0 331	0.0794
Dimethyldithiophosphate (µg/L)	-0.31	0.74 (0.51-1.06)	0.1 173	0.4800	-0.35	0.71 (0.49-1.01)	0.0 783	0.1782	-0.21	0.81 (0.59-1.12)	0.2 319	0.7382	-0.67	0.51 (0.29-0.91)	0.0 388	0.0887

Variables were first tested in an unadjusted / non-covariate adjusted analysis, and then again adjusting for age, ethnicity, and diabetes duration. To provide a broad overview, any variable passing nominal (i.e. prior to FDR-correction) $p \leq 0.05$ from either the non-covariate adjusted or any of the covariate-adjusted analyses are listed.

Table 4. RandomForest™-selected variables (features).

Marker	Group	Mean decrease accuracy	Mean decrease Gini
Glycohemoglobin (%)	Diabetes status	31.93719324	21.75225602
C-reactive protein (mg/dL)	Immune markers	11.51047187	10.1975135
Potassium (mmol/L)	Renal function	11.44142126	8.198116194
Albumin, urine (mg/L)	Renal function	8.220187056	10.21346621
Monocyte number (1000 cells/uL)	Immune markers	7.663589246	3.957711466
Osmolality (mmol/Kg)	Renal function	7.510989556	5.33445026
White blood cell count (1000 cells/uL)	Immune markers	7.440157644	4.405168072
Blood urea nitrogen (mmol/L)	Renal function	7.224789174	5.073947519
Segmented neutrophils num (1000 cell/uL)	Immune markers	7.020397225	4.067402271
Fasting Glucose (mmol/L)	Diabetes status	6.563694988	2.826827797
Red cell distribution width (%)	Hematocrit	6.138538515	4.947185792
Urinary nitrate (ng/mL)	Renal function	5.899174386	5.844258103
Glucose, serum (mmol/L)	Diabetes status	5.85339684	4.77357191
2-hydroxyphenanthrene (ng/L)	Toxins / Metals	4.028082549	2.307374348
MCHC (g/dL)	Hematocrit	3.936015913	3.896804592
Creatinine (μmol/L)	Renal function	3.530916132	3.747500758
Mono-2-ethyl-5-carboxypentyl phthalate	Toxins / Metals	2.996581682	2.212151134
Blood Nitromethane (pg/mL)	Toxins / Metals	2.936160917	3.31634662
Phosphorus (mmol/L)	Toxins / Metals	2.819084205	3.768671211
Total Cholesterol (mmol/L)	Sterols	2.448578413	3.833301666
Enterodiol (ng/mL)	Sterols	2.401651721	2.722762228
Hematocrit (%)	Hematocrit	2.364741874	4.841281382
Mono-n-octyl phthalate (ng/mL)	Toxins / Metals	2.215508752	0.231647322
Mean cell hemoglobin (pg)	Hematocrit	2.183482824	4.292282119
Gamma glutamyl transferase (U/L)	Liver Function	1.989695745	4.015070221
Systolic blood pressure	Blood pressure	1.892815461	8.85430752
Blood o-Xylene (ng/mL)	Toxins / Metals	1.670964308	3.708248225
Lactate dehydrogenase LDH (U/L)	Liver Function	1.593869272	4.231659273
9-hydroxyfluorene (ng/L)	Toxins / Metals	1.430814333	2.206778568
Cholesterol (mmol/L)	Sterols	1.201366974	4.025207689
3-hydroxyphenanthrene (ng/L)	Toxins / Metals	1.00569817	1.63630888

Notes: The model was initially trained on all laboratory variables in an unsupervised fashion, with Kappa-based model tuning to select the optimum values for ‘mtry’ (the ideal number of variables to randomly sample) and ‘ntrees’ (the ideal number of trees). Only variables contributing >1% mean decrease in accuracy from the initial model were retained, followed by recursive steps to remove low-informative variables. Variables are manually assigned to groups based on similar organ function or other characteristic. The Gini importance measure relates to the ‘splitting’ criterion that is employed in classification trees, and it is known to be less biased for continuous variables (26), which naturally have more splitting points compared to categorical variables. ‘Group’ is manually curated.

Table 5. Final clinical risk models.

Model	Wald test p-value	McFadden R ²	Nagelkerke R ²	AUC (95% CI)
Diabetes Status	≤ 0.0001	0.102	0.142	0.72 (0.677-0.763)
Hematocrit	0.0023	0.007	0.009	0.57 (0.539-0.601)
Blood Pressure (BP)	≤ 0.0001	0.017	0.024	0.586 (0.554-0.618)
Immune Markers	0.42	0.002	0.002	0.527 (0.495-0.559)
Renal function tests (renal)	≤ 0.0001	0.039	0.054	0.636 (0.604-0.669)
Sterols (include cholesterol)	0.055	0.012	0.017	0.582 (0.522-0.642)

Toxins / Metals	0.94	0.029	0.041	0.589 (0.478-0.7)
Liver function tests	0.07	0.002	0.003	0.525 (0.494-0.557)
Diabetes control + BP + Renal function	≤ 0.0001	0.13	0.18	0.73 (0.686-0.774)
Diabetes control + BP + renal function + Hematocrit	≤ 0.0001	0.135	0.184	0.737 (0.694-0.78)
Diabetes control + BP + renal function + Hematocrit + Sterols + Liver function	≤ 0.0001	0.238	0.315	0.823 (0.765-0.881)
All groups \pm	0.00013	0.272	0.355	0.84 (0.783-0.897)

Features from RandomForest™ were grouped logically based on similar function or clinical use (**Table 4**) and then tested independently in a univariate or multivariate regression model against DR outcome.

\pm The only toxins / metal included was Phosphorus (mmol/L) – others filtered out due to high missingness (>50%), resulting in difficulty fitting model.

FIGURE LEGENDS

Figure 1. Cohort selection process for NHANES 2007-8.

Figure 2.

Penalized regression-selected variables.

Variables were selected from a 100x cross-validated model with $\alpha=0.5$. Final variable selection was based on coefficients not shrunk to 0. Model accuracy was determined to be 78.4% accuracy and AUC 0.71 (95% CI: 0.65-0.77).

Figure 3. Final clinical risk models.


Features from RandomForest™ were grouped logically based on similar function or clinical use (**Table 4**) and then tested independently in a univariate or multivariate regression model against DR outcome. A final risk model including markers of hypertension, hypercholesterolemia, renal and liver function tests, and hematocrit achieved AUC 0.84 (0.78-0.9).

REFERENCES

1. Feman SS. The natural history of the first clinically visible features of diabetic retinopathy. *Trans Am Ophthalmol Soc.* 1994;92:745-73.
2. Chao JR, Lai M-Y, Azen SP, Klein R, Varma R, Group tLALES. Retinopathy in Persons without Diabetes: The Los Angeles Latino Eye Study. *Investigative Ophthalmology & Visual Science.* 2007;48(9):4019-25.
3. Venkatramani J, Mitchell P. Ocular and systemic causes of retinopathy in patients without diabetes mellitus. *BMJ (Clinical research ed).* 2004;328(7440):625-9.
4. Singh R, Ramasamy K, Abraham C, Gupta V, Gupta A. Diabetic retinopathy: an update. *Indian J Ophthalmol.* 2008;56(3):178-88.
5. Wang W, Lo ACY. Diabetic Retinopathy: Pathophysiology and Treatments. *Int J Mol Sci.* 2018;19(6):1816.

6. Yau JW, Rogers SL, Kawasaki R, Lamoureux EL, Kowalski JW, Bek T, et al. Global prevalence and major risk factors of diabetic retinopathy. *Diabetes Care*. 2012;35(3):556-64.
7. Heintz E, Wiréhn AB, Peebo BB, Rosenqvist U, Levin LA. Prevalence and healthcare costs of diabetic retinopathy: a population-based register study in Sweden. *Diabetologia*. 2010;53(10):2147-54.
8. Group UPDSU. Intensive blood-glucose control with sulphonylureas or insulin compared with conventional treatment and risk of complications in patients with type 2 diabetes (UKPDS 33). UK Prospective Diabetes Study (UKPDS) Group. *Lancet*. 1998;352(9131):837-53.
9. Chatziralli IP. The Role of Dyslipidemia Control in the Progression of Diabetic Retinopathy in Patients with Type 2 Diabetes Mellitus. *Diabetes Ther*. 2017;8(2):209-12.
10. Liu L, Quang ND, Banu R, Kumar H, Tham Y-C, Cheng C-Y, et al. Hypertension, blood pressure control and diabetic retinopathy in a large population-based study. *PLoS One*. 2020;15(3):e0229665-e.
11. Chen X, Meng Y, Li J, She H, Zhao L, Zhang J, et al. Serum uric acid concentration is associated with hypertensive retinopathy in hypertensive chinese adults. *BMC Ophthalmol*. 2017;17(1):83-.
12. Luo B-A, Gao F, Qin L-L. The Association between Vitamin D Deficiency and Diabetic Retinopathy in Type 2 Diabetes: A Meta-Analysis of Observational Studies. *Nutrients*. 2017;9(3):307.
13. Kong X, Wang J, Gao G, Tan M, Ding B, Li H, et al. Association between Free Thyroxine Levels and Diabetic Retinopathy in Euthyroid Patients with Type 2 Diabetes Mellitus. *Endocr Res*. 2020;45(2):111-8.
14. Merin S, Freund M. Retinopathy in severe anemia. *Am J Ophthalmol*. 1968;66(6):1102-6.
15. Khan AA, Rahmani AH, Aldebasi YH. Diabetic Retinopathy: Recent Updates on Different Biomarkers and Some Therapeutic Agents. *Curr Diabetes Rev*. 2018;14(6):523-33.
16. Moriya T, Tanaka S, Kawasaki R, Ohashi Y, Akanuma Y, Yamada N, et al. Diabetic retinopathy and microalbuminuria can predict macroalbuminuria and renal function decline in Japanese type 2 diabetic patients: Japan Diabetes Complications Study. *Diabetes Care*. 2013;36(9):2803-9.
17. Chen YH, Chen HS, Tarng DC. More impact of microalbuminuria on retinopathy than moderately reduced GFR among type 2 diabetic patients. *Diabetes Care*. 2012;35(4):803-8.
18. Lim LS, Tai ES, Mitchell P, Wang JJ, Tay WT, Lamoureux E, et al. C-reactive Protein, Body Mass Index, and Diabetic Retinopathy. *Investigative Ophthalmology & Visual Science*. 2010;51(9):4458-63.
19. Raymond NT, Varadhan L, Reynold DR, Bush K, Sankaranarayanan S, Bellary S, et al. Higher prevalence of retinopathy in diabetic patients of South Asian ethnicity compared with white Europeans in the community: a cross-sectional study. *Diabetes care*. 2009;32(3):410-5.
20. Spanakis EK, Golden SH. Race/ethnic difference in diabetes and diabetic complications. *Current diabetes reports*. 2013;13(6):814-23.
21. Wong TY, Klein R, Islam FMA, Cotch MF, Folsom AR, Klein BEK, et al. Diabetic retinopathy in a multi-ethnic cohort in the United States. *American journal of ophthalmology*. 2006;141(3):446-55.

22. McGinnis DP, Brownstein JS, Patel CJ. Environment-Wide Association Study of Blood Pressure in the National Health and Nutrition Examination Survey (1999–2012). *Scientific Reports*. 2016;6(1):30373.
23. Zhuang X, Guo Y, Ni A, Yang D, Liao L, Zhang S, et al. Toward a panoramic perspective of the association between environmental factors and cardiovascular disease: An environment-wide association study from National Health and Nutrition Examination Survey 1999-2014. *Environ Int*. 2018;118:146-53.
24. Hall MA, Dudek SM, Goodloe R, Crawford DC, Pendergrass SA, Peissig P, et al. Environment-wide association study (EWAS) for type 2 diabetes in the Marshfield Personalized Medicine Research Project Biobank. *Pac Symp Biocomput*. 2014:200-11.
25. Patel CJ, Rehkopf DH, Leppert JT, Bortz WM, Cullen MR, Chertow GM, et al. Systematic evaluation of environmental and behavioural factors associated with all-cause mortality in the United States national health and nutrition examination survey. *Int J Epidemiol*. 2013;42(6):1795-810.
26. Nembrini S, König IR, Wright MN. The revival of the Gini importance? *Bioinformatics*. 2018;34(21):3711-8.
27. Benjamini Y, Hochberg Y. Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing. *Journal of the Royal Statistical Society Series B (Methodological)*. 1995;57(1):289-300.
28. Stratton IM, Kohner EM, Aldington SJ, Turner RC, Holman RR, Manley SE, et al. UKPDS 50: risk factors for incidence and progression of retinopathy in Type II diabetes over 6 years from diagnosis. *Diabetologia*. 2001;44(2):156-63.
29. Klein R, Klein BEK. Blood pressure control and diabetic retinopathy. *Br J Ophthalmol*. 2002;86(4):365-7.
30. Zheng Y, Lamoureux EL, Lavanya R, Wu R, Ikram MK, Wang JJ, et al. Prevalence and risk factors of diabetic retinopathy in migrant Indians in an urbanized society in Asia: the Singapore Indian eye study. *Ophthalmology*. 2012;119(10):2119-24.



Footnotes:

¹ Laboratory datasets (2007-8): ALB_CR_E; APOB_E; BIOPRO_E; BPX_E; CARB_E; CBC_E; COTNAL_E; CRP_E; DEET_E; ENX_E; EPH_E; FASTQX_E; FERTIN_E; FOLATE_E; FOLFMS_E; GHB_E; GLU_E; HDL_E; HEPA_E; HEPBD_E; HEPB_S_E; HEPC_E; HIV_E; HPVSE_E; HPVSE_E; HSV_E; IHG_E; OGTT_E; OPD_E; PAH_E; PBCD_E; PERNT_E; PFC_E; PHTHE_E; PHYTO_E; POOLTF_E; PP_E; PSA_E; SSHCV_E; SSUSG_E; TCHOL_E; TFR_E; THYROD_E; TRIGLY_E; UAM_E; UAS_E; UCPREG_E; UHG_E; UHM_E; UIO_E; UPHOPM_E; UPP_E; VID_E; VIT_B6_E; VOC_E; VOCMWB_E; VOCWB_E.

Model accuracy: 78.51%

OPDURL4  0  1

