1	Socio-educational Impact and Psychological Distress of Medical Students amid the
2	COVID-19 Pandemic: A Japanese Cross-Sectional Survey
3	
4	Yoshito, Nishimura ^{1*} , Kanako, Ochi ^{1,2} , Kazuki, Tokumasu ¹ , Mikako, Obika ¹ , Hideharu,
5	Hagiya ¹ , Hitomi, Kataoka ^{1,3} , Fumio, Otsuka ¹
6	
7	¹ Department of General Medicine, Okayama University Graduate School of Medicine, Dentistry
8	and Pharmaceutical Sciences, Okayama, 7008558, Japan
9	² Center for Developing Medical and Health Care Education, Okayama University Medical
10	School, Okayama, 7008558, Japan
11	³ Center for Diversity and Inclusion, Okayama University Hospital, Okayama, 7008558, Japan
12	
13	*Corresponding author
14	Yoshito, NISHIMURA MD, PhD, MPH
15	Department of General Medicine, Okayama University Graduate School of Medicine, Dentistry
16	and Pharmaceutical Sciences, 2-5-1 Shikata-cho, Kita-ku, Okayama, Japan.
17	Tel.: +81-86-235-7342; Fax: +81-86-235-7345
18	E-mail: nishimura-yoshito@okayama-u.ac.jp

19 Abstract

20	The COVID-19 pandemic has negatively affected medical education. However, few data are
21	available about medical students' distress during the pandemic. This study aimed to provide
22	details on how medical students had been affected by the pandemic. In this cross-sectional study,
23	717 medical students participated in the web-based survey. The questions included how their
24	mental status had changed before and after the Japanese nationwide state of emergency (SOE).
25	65.9% (473/717) participated in the study. 29.8% (141/473) reported concerns about the shift
26	toward online education, mostly because they thought online education could have been
27	ineffective compared with in-person learning. Participants' subjective mental health status
28	significantly worsened after the SOE was lifted (p <.001). Those who had concerns about a shift
29	toward online education had higher odds of having generalized anxiety and being depressed (OR
30	1.97, 95% CI 1.19 – 3.28), as did those who requested food aid and mental health care resources
31	(OR 1.99, 95% CI 1.16 – 3.44; OR 3.56, 95% CI 2.07 – 6.15, respectively). Given our findings,
32	the sudden shift to online education might have overwhelmed medical students. Thus, we
33	recommend educators to inform learners that online learning is non-inferior to in-person learning,
34	which could attenuate potential depression and anxiety.

35



37 Introduction

38 The coronavirus disease 2019 (COVID-19) global pandemic has drastically changed our lives, 39 with more than eight million cases and 400,000 deaths reported globally as of June 18, 2020, 40 according to the World Health Organization statistics(1). In Japan, 17,668 cases have been 41 confirmed as of June 18, 2020, with an explosion in the number of cases in early April(2). While 42 Japan seems to be bringing the outbreak under control through active cluster tracing, restriction 43 of mass gatherings, and advocating universal masking and hand hygiene, there were no simple 44 solutions to this issue(3). After the surge in COVID-19 cases, on April 16, 2020, the Government 45 of Japan (GoJ) declared a state of emergency (SOE) in all 47 prefectures, which lasted until May 46 25, 2020(4). In Okayama, a prefecture in the western part of mainland Japan(5) with 47 approximately 1.9 million people (ranked 20/47 among the 47 prefectures), only 25 confirmed 48 cases had been reported by the end of May 2020. Behind the scenes of successful COVID-19 49 mitigation, however, those in education-related jobs and medical students struggled with the 50 rapid change in the socio-educational system.

51

Although the SOE was lifted on May 14 in Okayama based on the low incidence of COVID-19,
even before the SOE, from March 25, Okayama University School of Medicine (OUSM), one of
the largest national universities in Japan, requested its students to stay home to prevent the

55	possible community spread of COVID-19 within the medical school and hospital. Until the
56	stay-home request was lifted on May 22, medical students were required to cope with this sudden
57	change in their lifestyle, and shifted to online education amid the fear and considerable
58	uncertainty surrounding COVID-19. In the early 2000s, the severe acute respiratory syndrome
59	coronavirus (SARS-CoV) outbreak had a devastating impact on academic education, including
60	sudden curriculum changes and rapid integration of information technology(6-8). Similarly, the
61	current COVID-19 pandemic has provoked significant turmoil in society. In particular, mental
62	health problems due to the pandemic have drawn attention worldwide as studies have suggested
63	the need for mental health care interventions during the outbreak(9-14). Medical students, who
64	essentially need clinical exposure, may have been impacted even further by the pandemic.
65	

In Japan, medical schools have different admission practices and curricula than in the United States. Students typically enter medical school immediately after high school graduation, often at 18 years old, and they go through six years of medical education before graduation. Despite the differences in education system, the fundamental philosophy of academic medicine is the same; to provide quality educational experience. To date, few published studies have investigated the status of medical students' living environment, socio-educational, and mental health status during the COVID-19 pandemic. To address this, we conducted a total population survey of OUSM

73 medical students to comprehensively clarify what they require and how they have been mentally

74 affected by COVID-19.

75

- 76 Methods
- 77 Study Design, setting, and participants

We performed a cross-sectional study that employed an anonymous, self-administered voluntary web-based survey. Participants comprised medical students in all years of study at OUSM. We used purposive sampling to conduct a total population sample for the 717 medical students who belonged to the OUSM as of April 1, 2020 (the first day of the academic year in Japan). The participants' consent was implied by their completion of the survey.

83

The survey was developed through consultation with a medical education expert panel at OUSM and piloting, and administered with Qualtrics (Qualtrics International Inc., Provo, Utah), a web-based survey platform. We provided survey instructions and instruments in Japanese. We distributed survey links to the students using OUSM official mailing lists. All participants were invited to complete the survey within one week (June 8–14, 2020, in Japan Standard Time). No financial incentives were provided for their participation in the survey. In the survey, we included entries on demographics (age, gender, education before entering medical school, employment

91	status on date of response, changes in employment status due to the COVID-19 pandemic,
92	marital status, living environment, household size, and comorbidities) as well as COVID-19
93	related items (e.g., "Chance of contracting COVID-19 during the current pandemic"),
94	self-learning associated activities (e.g., Average time of self-learning/day), validated depression
95	and anxiety scale instruments, and financial situations. To protect participants' anonymity as
96	much as possible, respondents were not prompted to enter their year of study.
97	
98	Measurements
99	COVID-19-related questions
100	To evaluate the extent of participants' concern and preparedness for COVID-19, they were
101	asked: "What do you think the chances are that you will contract COVID-19 during the current
102	pandemic?"; "What is your degree of concern about the health status of your family?"; "Do you
103	think you have enough information about the symptoms of COVID-19?"; "Do you think you
104	have enough information about prevention and treatment of COVID-19?"; and "Do you feel
105	worried about COVID-19?" To further assess students' concerns, the participants were prompted
106	to answer "I am concerned because my future career formation may be negatively affected due to
107	the COVID-19 pandemic"; "I am concerned because the COVID-19 pandemic may attenuate our
108	relationship to teachers"; "I am concerned because of the disruption to ongoing research or

109	extracurricular activities"; and "I am concerned about the shift toward online education." All
110	questions were evaluated on a 5-point Likert scale except for the item on the health status of the
111	family (3-point Likert scale). Those who answered "Very concerned" or "Concerned" in the
112	entry on concern about the shift toward online education were prompted to provide reasons. To
113	describe participants' needs, they were asked to note the types of support they wish to receive
114	from the university if there is a resurgence of COVID-19. These responses were mandatory.
115	
116	Self-learning and related activity
117	For subjective mental health status and the average time per day that participants stayed at home,
118	read books, played video games, and learned by themselves, respondents were prompted to
119	answer "How many hours a day did you / stay at home / read books / play video games /
120	self-learn?" before the SOE order (April 16, 2020) or during the last two weeks (with the base
121	date of when participants completed the survey).
122	
123	Depression and anxiety disorders
124	We assessed the presence of depression using the 9-item Patient Health Questionnaire (PHQ-9),
125	a common screening tool for mood disorders. We used the validated Japanese translation of the
126	scale(15). The total scores of PHQ-9 range from 0 to 27 and we defined scores of 10 or more as

127	having "depression." We screened for anxiety disorders using the Japanese version of the 7-item
128	Generalized Anxiety Disorder Scale (GAD-7), which was validated in 2010(16). The score
129	ranges from 0 to 21, and we defined scores of 10 or more as having "anxiety." Both instruments
130	ask respondents about their mental health status during the last two weeks.

131

132 **Financial situation**

In Japan, university students typically live on a monthly allowance from parents. According to the latest statistics by the National Federation of University Co-operative Associations, students receive 72,810 JPY (approximately 680 USD) a month on average(17). Participants were prompted to give their monthly allowance from the following options: "None," "0," "< 30,000 JPY (280 USD)," "30,000 – 49,999 JPY (280 – 467 USD)," "50,000 – 69,999 JPY (467 – 654 USD)," "70,000 – 99,999 JPY (654 – 935 USD)," and " \geq 100,000 JPY (\geq 935 USD)." Respondents were also asked to answer if they were on scholarship or student loan.

141 Statistical analysis

We analyzed the data using JMP version 13.1.0 (SAS Institute Inc., Cary, North Carolina). We used the Wilcoxon signed-rank test to examine differences in the time participants spent on self-learning related activities based on non-normal distribution. For associations between

145 categorical variables containing small sample sizes, we employed Fisher's exact	et test. To examine
---	---------------------

- 146 the predictive factors of categorical dependent variables, we used univariate logistic regression
- 147 analyses. The threshold for significance was defined as p < 0.05.

148

149 **Ethical approval**

150 This study protocol was approved by the institutional review board of Okayama University

151 Hospital (reference number 2006-029, approved on June 5, 2020).

152

153 Data Availability Statement

154 The datasets generated and analysed during the current study are available from the 155 corresponding author on reasonable request.

156

157 **Results**

Response rates to the survey were 473 out of 717 (65.9%) OUSM students in all six years combined. Participants' demographic characteristics are summarized in **Table 1**. Of note, 250 (52.9%) respondents reported that they were engaged in part-time work, while 44 (9.3%) reported having resigned or been fired due to the COVID-19 pandemic. Eight (1.7%) and 6 (1.3%) noted that they had a past medical history of anxiety disorders and depression,

163 respectively.

164

165 **COVID-19 related survey items**

166	Table 2 and Supplementary Figure 1 show respondents' answers to the COVID-19 related
167	survey items. Eighty-one (17.2%) responded that they were either "likely" or "very likely" to
168	contract COVID-19 during the ongoing pandemic; 275 (58.1%) agreed or strongly agreed to a
169	prompt "I feel worried about COVID-19." Regarding the breakdown of students' concerns about
170	COVID-19, 182 (38.5%), 121 (25.6%), and 235 (49.7%) acknowledged that they were concerned
171	about the negative impacts of COVID-19 on their future career formation, relationship with
172	teachers, and ongoing research or extracurricular activities, respectively, while 141 (29.8%) also
173	reported concern about a shift toward online education. The reasons for concerns included
174	ineffective educational effect compared with on-site education ($n = 92, 65.2\%$), possible
175	resurgence of the COVID-19 outbreak leading to a sudden change in the curriculum ($n = 74$,
176	52.5%), and decreased clinical exposure ($n = 92, 65.2\%$).

177

178 Of the participants, 298 (63.0%) answered that they would request financial aid if a stay-home 179 order recurred due to a COVID-19 resurgence, followed by request for food aid (n = 100, 21.1%), 180 technical support for online education (n = 100, 21.1%), and mental health care resources (n = 85,

181 18.0%).

182

183 Change in self-learning related time before and after the SOE

- 184 We compared the change in average time respondents spent at home, reading books, playing
- 185 video games, and self-learning per day before the SOE and within two weeks prior to the survey
- 186 completion (after the SOE). As shown in **Table 3** and **Supplementary Figure 2**, the participants
- 187 spent significantly longer on all the activities mentioned above after the SOE than before the
- 188 SOE (p < .001). There were also significant differences in their subjective mental health status
- 189 before and after the SOE based on Wilcoxon signed-rank test (p < .001).
- 190

191 **Regression analyses of factors associated with depression and anxiety**

Of the participants, 75 (15.8%) had PHQ-9 scores of 10 or more, and 34 (7.2%) had GAD-7 scores of 10 or more. **Table 4** presents all the results of univariate regression analyses. The odds of being depressed were significantly higher in those who had concerns about a shift toward online education (OR 1.97, 95% CI 1.19 – 3.28), and in those who requested food aid (OR 1.99, 95% CI 1.16 – 3.44) and mental health care resources (OR 3.56, 95% CI 2.07 – 6.15) from the university in the event of resurgence of COVID. Regarding generalized anxiety, the odds were higher in respondents who requested food aid (OR 2.50, 95% CI 1.21 – 5.20) and mental health

199 care resources (OR 3.16, 95% CI 1.51 – 6.59).

200

201 **Prevalence of financial hardship**

Table 5 and Supplementary Figure 3 show the self-reported amounts of monthly allowance from parents and situations regarding financial aid. A total of 332 (70.2%) respondents had a monthly allowance that was lower than the national average (72,810 JPY, approximately 680 USD)(17), while 141 (27.7%) received some sort of financial aid for their living and education.

206

207 **Discussion**

To the best of our knowledge, this study is the first survey of Japanese medical students 208 209 regarding their life circumstances and challenges due to COVID-19. Although Japan has had 210 fewer COVID-19 cases than the US, and Okayama has had comparatively fewer cases than other 211 Japanese prefectures, the study results underscore that the pandemic caused a profound negative 212 socio-educational impact on our society and medical students. We found that approximately 10% 213 of the students on part-time jobs had been fired due to the pandemic. Furthermore, a considerable 214 number of medical students had concerns around their basic life security, demanding support 215 from the university in the form of financial aid, food aid, technical support, and mental health care resources due to the SOE. Regarding students' subjective psychological distress, those who 216

expressed concerns about the rapid shift toward online education and fear around basic lifesecurity were more likely to be depressed and anxious after the SOE.

219

220	Concerns around future career disruption, attenuated relationship to medical teachers, and
221	disruption of ongoing extracurricular activities were prevalent among the participants, which
222	underlines the considerable uncertainty amid the COVID-19 pandemic (Table 2). In particular,
223	63% of respondents reported the need for financial aid in the event of a second wave of the
224	pandemic. These data correspond to the fact that more than 70% of participants received no more
225	than the national average monthly allowance from their parents (Table 5). Contrary to the general
226	notion in Japan that "medical students are financially well-off," many may have experienced
227	financial hardship. Recently, the theory of willpower has gathered public attention. Financial
228	stability is considered a prerequisite of willpower, which aids appropriate decision-making(18).
229	Combined with previous research findings that financial instability could lead to worse mental
230	health outcome(19, 20), educators are expected to develop a strategy to offer financial support to
231	those in need.

232

233 Regarding lifestyle changes and psychological distress, after the SOE was lifted, the participants

234 experienced significantly worse mental health status, and spent significantly longer at home,

235	reading books, playing video games, and learning by themselves (Table 3) than before the
236	announcement of SOE. Our results are consistent with prior longitudinal and cross-sectional
237	studies that the COVID-19 pandemic led to a sedentary life and psychological distress(9, 12, 13,
238	21). As suggested by a previous study, the increase in gaming could be a coping behavior against
239	psychological stress(22).

240

241 As for regression analyses of factors associated with depression and anxiety, in our study 242 population, those who requested food aid and mental health care resources from the university 243 upon the future resurgence of the COVID-19 outbreak had significantly higher odds of depression and anxiety. Furthermore, concerns around the shift toward online education were 244 245 identified as a factor associated with depression (Table 4). Surprisingly, 65.2% of those who had 246 concerns about the shift toward online education thought online education was less effective than 247 in-person education. While previous studies have reported the utility and noninferiority of online 248 learning compared to offline(23, 24), in-person learning, the results of the current study revealed 249 a potential gap in perception regarding the effectiveness of online education between medical 250 students and educators. Educators should not assume that students know the potential benefits of 251 online learning, and it is essential to inform learners that online learning is non-inferior to 252 in-person learning, which could attenuate potential depression and anxiety. While in-person communication has become difficult nowadays due to the fear for COVID-19, medical schools
may need to reach out to students to find out those suffering from underlying life insecurity and
to provide multilateral support, including early mental health care interventions.

256

257 Several limitations to this study should be noted. First, due to the single-center cross-sectional 258 survey design, we may not conclude causal relationships. Second, the cross-sectional research 259 contains a limitation in terms of addressing changes over time. While we illustrated the changes 260 in students' average time spent on self-learning related activities before and after the SOE in 261 Table 3, a longitudinal study design would be more appropriate to examine the differences in the 262 study cohort. Third, we asked participants to provide their mental health status and time spent on 263 self-learning related activities before the SOE (approximately six weeks before the survey 264 implementation), both of which are subject to recall bias. Also, PHQ-9 and GAD-7 scores were 265 obtained only after the SOE. Thus, it is possible that they might have not changed during the period. Lastly, due to the nature of the survey topic, those who were interested in the public 266 health emergency or mental health may have been more likely to respond, which would lead to 267 268 self-selection bias. Despite these limitations, a total population sampling strategy coupled with higher-than-usual response rates(25) contributed to high internal validity. 269

270

271 In conclusion, through the study, we have provided graphical data and evidence regarding the 272 socio-educational impacts of the COVID-19 pandemic on medical students. In circumstances of 273 considerable uncertainty, both educators and medical students need to be flexible, patient, and 274 resilient. The uncertainty and drastic change triggered substantial psychological distress in 275 students, which was greater than we assumed. As discussed, Japan has had a relatively small 276 number of COVID-19 cases compared to the US and other European countries. Although 277 Okayama prefecture survived the pandemic with fewer confirmed COVID-19 cases than other 278 prefectures, medical students experienced significant impacts due to the public health emergency. 279 The educational institution should recognize the prevalence of basic needs insecurity, such as 280 financial difficulties and a shortage of staples, including food, etc. As medical educators, we 281 need to be accountable for the advantages of online education in the field of medicine to alleviate 282 students' psychological distress, in addition to providing multilateral support to those in need, 283 including early mental health care interventions. While we targeted medical students in a single Japanese national university, the survey results warrant further research and analysis if the 284 distress is amplified by existing anxiety, depression, burnout, etc., or is a wholly 285 286 COVID-19-related phenomenon. We call for increased research in populations with more 287 COVID-19 cases than Japan, to figure out the challenges medical students have been facing amid 288 the pandemic in different cultures and backgrounds.

289

290 **References**

- 291 1. World Health Organization. Coronavirus disease (COVID-19) Situation Report 150
- 292 2020 [Available from:
- 293 https://www.who.int/docs/default-source/coronaviruse/situation-reports/20200618-covid-19-sitre
- 294 <u>p-150.pdf?sfvrsn=aa9fe9cf_2</u>.
- 295 2. Ministry of Health LaWJ. About Coronavirus Disease 2019 (COVID-19) 2020
- 296 [Available from: <u>https://www.mhlw.go.jp/stf/seisakunitsuite/bunya/newpage_00032.html</u>.
- 297 3. Looi MK. Covid-19: Japan ends state of emergency but warns of "new normal". BMJ.
- 298 2020;369:m2100.
- 299 4. Government of Japan. Basic Policies for Novel Coronavirus Disease Control by the
- 300 Government of Japan (Summary) 2020 [Available from: 301 https://www.mhlw.go.jp/content/10900000/000634753.pdf.
- 302 5. Statistics Bureau of Japan. Current Population Estimates as of October 1, 2019 2019
- 303 [Available from: https://www.stat.go.jp/english/data/jinsui/2019np/index.html.
- 304 6. Clark J. Fear of SARS thwarts medical education in Toronto. BMJ.
 305 2003;326(7393):784.
- 306 7. Patil NG, Chan Y, Yan H. SARS and its effect on medical education in Hong Kong. Med

307 Educ. 2003;37(12):1127-8.

308	8.	Sherbino J, Atzema C. "SARS-Ed": severe acute respiratory syndrome and the impact
309	on medic	cal education. Ann Emerg Med. 2004;44(3):229-31.
310	9.	Huckins JF, DaSilva AW, Wang W, Hedlund E, Rogers C, Nepal SK, et al. Mental
311	Health a	and Behavior During the Early Phases of the COVID-19 Pandemic: A Longitudinal
312	Mobile S	Smartphone and Ecological Momentary Assessment Study in College Students. J Med
313	Internet	Res. 2020.
314	10.	Liu S, Yang L, Zhang C, Xiang YT, Liu Z, Hu S, et al. Online mental health services in
315	China du	uring the COVID-19 outbreak. Lancet Psychiatry. 2020;7(4):e17-e8.
316	11.	Shigemura J, Kurosawa M. Mental health impact of the COVID-19 pandemic in Japan.
317	Psychol	Trauma. 2020.
318	12.	Talevi D, Socci V, Carai M, Carnaghi G, Faleri S, Trebbi E, et al. Mental health
319	outcome	s of the CoViD-19 pandemic. Riv Psichiatr. 2020;55(3):137-44.
320	13.	Wang C, Pan R, Wan X, Tan Y, Xu L, Ho CS, et al. Immediate Psychological Responses
321	and Asso	ociated Factors during the Initial Stage of the 2019 Coronavirus Disease (COVID-19)
322	Epidemi	c among the General Population in China. Int J Environ Res Public Health. 2020;17(5).
323	14.	Zhang Y, Zhang H, Ma X, Di Q. Mental Health Problems during the COVID-19
324	Pandemi	cs and the Mitigation Effects of Exercise: A Longitudinal Study of College Students in

325 China. Int J Environ Res Public Health. 2020;17(10).

- Muramatsu K, Miyaoka H, Kamijima K, Muramatsu Y, Tanaka Y, Hosaka M, et al. 326 15. Performance of the Japanese version of the Patient Health Questionnaire-9 (J-PHQ-9) for 327 328 depression in primary care. Gen Hosp Psychiatry. 2018;52:64-9. 329 16. Muramatsu K, Miyaoka H, Ueshima K, Muramatsu Y, Fuse K, Yoshimine H, et al. 330 Validation and utility of a Japanese version of the GAD-7. Jpn J Psychosom Med. 331 2010;50(6):592. 332 17. National Federation of University Co-operative Associations. Summary of Campus Life 333 Data 2019 [Article] in Japanese] 2019 [Available from: 334 https://www.univcoop.or.jp/press/life/report.html. 335 18. Baumeister RF, Tice DM, Vohs KD. The Strength Model of Self-Regulation: Conclusions From the Second Decade of Willpower Research. Perspect Psychol Sci. 336 337 2018;13(2):141-5. 338 Richardson T, Elliott P, Roberts R, Jansen M. A Longitudinal Study of Financial 19. Difficulties and Mental Health in a National Sample of British Undergraduate Students. 339 340 Community Ment Health J. 2017;53(3):344-52.
- 341 20. McCloud T, Bann D. Financial stress and mental health among higher education
 342 students in the UK up to 2018: rapid review of evidence. J Epidemiol Community Health.

343 2019;73(10):977-84.

344	21. Brooks SK, Webster RK, Smith LE, Woodland L, Wessely S, Greenberg N, et al. The
345	psychological impact of quarantine and how to reduce it: rapid review of the evidence. Lancet.
346	2020;395(10227):912-20.
347	22. Balhara YPS, Kattula D, Singh S, Chukkali S, Bhargava R. Impact of lockdown
348	following COVID-19 on the gaming behavior of college students. Indian J Public Health.
349	2020;64(Supplement):S172-S6.
350	23. Pei L, Wu H. Does online learning work better than offline learning in undergraduate
351	medical education? A systematic review and meta-analysis. Med Educ Online.
352	2019;24(1):1666538.
353	24. Seymour-Walsh AE, Weber A, Bell A. Practical approaches to pedagogically rich online
354	tutorials in health professions education. Rural Remote Health. 2020;20(2):6045.
355	25. Cunningham CT, Quan H, Hemmelgarn B, Noseworthy T, Beck CA, Dixon E, et al.
356	Exploring physician specialist response rates to web-based surveys. BMC Med Res Methodol.
357	2015;15:32.
358	
359	

360

20

361	
362	
363	
364	
365	
366	Acknowledgments
367	We would like to thank the Academic Affairs Division of Okayama University Medical School
368	for its cooperation in conducting the survey.
369	
370	Author contributions
371	YN wrote the manuscript, designed the study, and analysed the data. KO, KT, MO, HH, and HK
372	analyzed the data and revised the manuscript. FO designed and supervised the research.
373	
374	Conflicts of interest
375	The authors declare no conflicts of interest in association with the present study.
376	
377	Funding
378	None. 21

379	
380	
381	
382	
383	
384	Figure Legend
385	Supplementary Figure 1
386	(A) The breakdown of students' concerns about COVID-19. 182 (38.5%), 121 (25.6%), and 235
387	(49.7%) acknowledged that they were concerned about the negative impacts of COVID-19 on
388	their future career formation, relationship with teachers, and ongoing research or extracurricular
389	activities, respectively, while 141 (29.8%) also reported concern about a shift toward online
390	education. (B) Of the participants, 298 (63.0%) answered that they would request financial aid if
391	a stay-home order recurred due to a COVID-19 resurgence, followed by request for food aid (n =
392	100, 21.1%), technical support for online education ($n = 100, 21.1\%$), and mental health care
393	resources $(n = 85, 18.0\%)$.
394	

395 Supplementary Figure 2

396 The bar charts show the change in average time respondents spent at home, reading books,

- 397 playing video games, and self-learning per day before the SOE and within two weeks prior to the
- 398 survey completion (after the SOE). Error bars show the 95% confidence intervals.

399

- 400 Supplementary Figure 3
- 401 The self-reported amounts of monthly allowance from parents and situations regarding financial
- 402 aid are shown.

Table 1. Demographic Characteristics of the Study Participants Comprising 473

Characteristic	Value	SD	95% CI
Age in years ^a			
Mean	22.0	3.3	21.7-22.3
Gender, no. (%)			
Female	161 (34.0)		
Male	311 (65.8)		
Non-confirming	1 (0.2)		
Education before medical school, no. (%)			
High school	434 (91.8)		
Career college	1 (0.2)		
4-year university	31 (6.6)		
Master's degree	7 (1.5)		
Employment status, no. (%)			
Part-time job	250 (52.9)		
Schoolwork only: no extra work	221 (46.7)		
Self-owned business	2 (0.4)		

Japanese Medical Students

Resigned or fired due to the pandemic	44 (9.3)		
Marital status, no. (%)			
Single	467 (98.7)		
Married	5 (1.1)		
Missing	1 (0.2)		
Living environment (%)			
Living alone	308 (65.1)		
Living with family	127 (26.9)		
Living with a partner	20 (4.2)		
Living alone: family or relatives nearby	18 (3.8)		
Household size			
Mean	1.7	1.3	1.6-1.8
Comorbidity, no. (%)			
None	401 (84.8)		
Asthma	37 (7.8)		
Anxiety disorder	8 (1.7)		
Depression	6 (1.3)		
Hypertension	5 (1.1)		

Total number of participants	473	
Other	24 (5.1)	
Dyslipidemia	2 (0.4)	
Heart diseases	3 (0.6)	

Abbreviations: CI, confidence interval; SD, standard deviation.

^a Of the 473 respondents, 2 did not specify their age.

Table 2. Results of the COVID-19-related Survey Items

Survey Entry	No. (%)
"Chance of contracting COVID-19 during the current pandemic"	
Very likely	15 (3.2)
Likely	66 (14.0)
Neutral	238 (50.3)
Unlikely	121 (25.6)
Very unlikely	33 (6.9)
"Degree of concern about the health status of your family"	
Very concerned	198 (41.9)
Somewhat concerned	237 (50.1)
Not concerned	38 (8.0)
"I have enough knowledge about symptoms of COVID-19"	
Strongly agree	8 (1.7)
Agree	158 (33.4)
Neutral	169 (35.7)
Disagree	116 (24.5)
Strongly disagree	22 (4.7)

"I have enough knowledge about the treatment of COVID-19"

Strongly agree	6 (1.3)
Agree	65 (13.7)
Neutral	138 (29.2)
Disagree	196 (41.4)
Strongly disagree	68 (14.4)

"I feel worried about COVID-19"

Strongly agree	46 (9.7)
Agree	229 (48.4)
Neutral	106 (22.4)
Disagree	77 (16.3)

15 (3.2)

Students' concern	Future Career	Attenuated	Disruption of	Shift towards
	Formation due	Relationship to	Ongoing	Online
	to COVID-19	Teachers	Research or	Education
			Extracurricular	
			Activities	

	No. (%)	No. (%)	No. (%)	No. (%)
Very concerned	46 (9.7)	20 (4.2)	66 (14.0)	18 (3.8)
Concerned	136 (28.8)	101 (21.4)	169 (35.7)	123 (26.0)
Neutral	130 (27.5)	136 (28.7)	91 (19.2)	115 (24.3)
Not concerned	85 (18.0)	129 (27.3)	67 (14.2)	121 (25.6)
Not at all concerned	76 (16.0)	87 (18.4)	80 (16.9)	96 (20.3)

"The Reasons for Concern about Shift toward Online Education" ^a

Online education may not be as effective as on-site education	92 (65.2)
Fear of possible resurgence of COVID-19 outbreak, which would cause a	74 (52.5)
sudden change in the curriculum	
Less clinical exposure	92 (65.2)
Other	16 (11.3)

"What supports do you want from the university if a stay-home order takes

place again due to the resurgence of COVID-19?"^b

Financial aid (e.g., tuition waiver, no-interest student loan)	298 (63.0)
Food aid	100 (21.1)
Resources for mental health care	85 (18.0)
Technical support (e.g., free rental of PC, pocket WiFi)	100 (21.1)

Basic livelihood support (e.g., resources for health maintenance	57 (12.1)
or preventing DV)	
Parenting support	6 (1.3)
Other	55 (11.6)

Abbreviations: COVID-19, coronavirus disease 2019; PC, personal computer.

^a The survey entry was explicitly answered by participants who answered "Very concerned" or "Concerned" in the previous entry ("Concerned about Shift toward Online Education," n = 141). Multiple answers permitted.

^b Multiple answers permitted.

Table 3. Comparison of Age-weighted Self-learning and Related Activity Time of

	Befor	e SOE	Last 2	Weeks ^a	
	Time (h)	95% CI	Time (h)	95% CI	
Measure					P value ^b
Average Time at Home / Day	16.1	15.7-16.6	18.8	18.5-19.2	< .001
Average Time Reading Books / Day	0.70	0.58-0.82	1.1	0.96-1.2	< .001
Average Time Playing Video Games / Day	1.5	1.3-1.7	2.1	1.9-2.3	< .001
Average Time Self-learning / Day	2.6	2.4-2.8	4.0	3.7-4.2	< .001
	Befor	e SOE	Last 2	Weeks	
Measure	No.	(%)	No.	(%)	
Subjective Mental Health Status ^c					
Very good	85 (18.0)	39 ((8.2)	
Good	171	(36.1)	58 (12.2)	
Fair	193	(40.8)	286	(60.5)	
Bad	18	(3.8)	76 (16.1)	
Very bad	6 (1.3)	14 ((3.0)	

Medical Students Before and After the National State of Emergency in Japan

Abbreviations: CI, confidence interval; h, hour; SOE, state of emergency.

^a The base date of the "last 2 weeks" is the day when the participants answered the

survey. The survey period was from June 8 to 14, 2020.

^b *P* value is calculated with Wilcoxon signed-rank test.

^c P < .001 with Wilcoxon signed-rank test. There is a significant difference in the subjective mental health status of participants before the Japanese national state of emergency and the last two weeks prior to the date of answering the survey.

Table 4. Results of Univariate Regression Analyses of Factors Associated with

Depression and Generalized Anxiety in Japanese Medical Students

Dependent Variable	Independent Variable	OR (95% CI) ^a	P value
Depression $(n = 75)$	Female (vs. Male)	1.08 (0.58 – 2.01)	.81
(PHQ-9 score ≥ 10)	Age	1.06 (0.96 – 1.16)	.45
	Concerns around Future Career Formation ^b	1.40 (0.85 – 2.30)	.18
	Concerns around Attenuated Relationship to	1.46 (0.85 – 2.50)	.17
	Teachers ^b		
	Concerns around Disruption of Ongoing Research	0.81 (0.50 – 1.33)	.41
	or Extracurricular Activities ^b		
	Concerns around Shift toward Online Education ^b	1.97 (1.19 – 3.28)	.009
	Request for Financial Aid	0.92 (0.55 – 1.53)	.74
	Request for Food Aid	1.99 (1.16 – 3.44)	.013
	Request for Mental Health Care Resources	3.56 (2.07 – 6.15)	<.0001
Generalized Anxiety (n	Female (vs. Male)	0.81 (0.37 – 1.74)	.58
= 34)	Age	0.96 (0.85 - 1.09)	.57
(GAD-7 score ≥ 10)	Concerns around Future Career Formation ^b	0.75 (0.36 – 1.58)	.45
	Concerns around Attenuated Relationship to	1.43 (0.68 – 3.03)	.35
	Teachers ^b		

Concerns around Disruption of Ongoing Research	0.53 (0.26 - 1.09)	.09
or Extracurricular Activities ^b		
Concerns around Shift toward Online Education ^b	1.50 (0.73 – 3.10)	.27
Request for Financial Aid	1.25 (0.59 – 2.62)	.56
Request for Food Aid	2.50 (1.21 – 5.20)	.014
Request for Mental Health Care Resources	3.16 (1.51 - 6.59)	.002

Abbreviations: CI, confidence interval; GAD-7, Generalized Anxiety Disorder-7; OR,

odds ratio; PHQ-9, Patient Health Questionnaire-9.

^a Univariate regression analysis was performed for each dependent variable.

^b Participants who answered, "Very concerned" or "Concerned" in the respective survey entries listed in Table 2 were considered the "Positive concerns group." On the other hand, those who answered "Neutral," "Not concerned," or "Not at all concerned" were considered the "No concerns group" in this analysis.

Table 5. Description of	f the Financial Situation	of Japanese Medical Students
-------------------------	---------------------------	------------------------------

Characteristic	No. (%)
Monthly allowance from parents (JPY) ^a	
No allowance	127 (26.9)

< 30,000 JPY	43 (9.1)
30,000 – 49,999 JPY	81 (17.1)
50,000 – 69,999 JPY	81 (17.1)
70,000 – 99,999 JPY	78 (16.5)
≥ 100,000 JPY	63 (13.3)
Financial Aid	
None	342 (72.3)
Student loan only	74 (15.6)
Scholarship only	44 (9.3)
Both student loan and scholarship	13 (2.8)

Abbreviations: JPY, Japanese Yen.

^a According to the Bank of Japan, the exchange rates of 1 US dollar to JPY was

107.39-41 on the morning of June 17, 2020.