1	Active transport of brilliant blue FCF across the Drosophila
2	midgut and Malpighian tubule epithelia
3	

5	
6	Dawson B.H. Livingston ¹ , Hirva Patel ¹ , Andrew Donini ² , and Heath A. MacMillan ¹
7	
8	¹ Department of Biology, Carleton University, Ottawa, Canada, K1S 5B6
9	² Department of Biology, York University, Toronto, Canada, M3J 1P3
10	
11	
12	
13	
14	
15	
16	
17	
18	Keywords: Organic anion; paracellular barriers; barrier dysfunction; renal function; cold
19	tolerance; FD&C blue dye no. 1
20	
21	
22	
23	
24	
25	Commence Statements D is 11 and 11
20	Summary Statement: Drosophila are able to actively secrete Brilliant blue FCF, a commonly
27	used marker of barrier dysfunction
28	

29 Abstract:

30 Under conditions of stress, many animals suffer from epithelial barrier disruption that can cause 31 molecules to leak down their concentration gradients, potentially causing a loss of organismal 32 homeostasis, further injury or death. Drosophila is a common insect model, used to study barrier 33 disruption related to aging, traumatic injury, or environmental stress. Net leak of a non-toxic dye 34 (Brilliant blue FCF) from the gut lumen to the hemolymph is often used to identify barrier failure 35 under these conditions, but *Drosophila* are capable of actively transporting structurally-similar 36 compounds. Here, we examined whether cold stress (like other stresses) causes Brilliant blue 37 FCF (BB-FCF) to appear in the hemolymph of flies fed the dye, and if so whether *Drosophila* 38 are capable of clearing this dye from their body following chilling. Using *in situ* midgut leak and 39 transport assays as well as Ramsay assays of Malpighian tubule transport, we tested whether 40 these ionoregulatory epithelia can actively transport BB-FCF. In doing so, we found that the 41 Drosophila midgut and Malpighian tubules can mobilize BB-FCF via an active transcellular 42 pathway, suggesting that elevated concentrations of the dye in the hemolymph may occur from 43 increased paracellular permeability, reduced transcellular clearance, or both.

44 Introduction

Epithelial barriers are an important constituent of all metazoan life on earth and are critical in the
maintenance of homeostasis. In the gut, epithelial barriers play a major role in regulating rates of
nutrient absorption; ion and water secretion and absorption; the clearance of wastes (Salvo
Romero et al., 2015); and protect organisms from potential threats such as infection and
dehydration (Presland and Jurevic, 2002).

50

51 Drosophila are a common model of digestive and renal physiology (Dow & Romero, 2010; 52 Buchon et al., 2013; Miguel-Aliaga et al., 2018). The gut of *Drosophila* is divided into three 53 distinct regions; the foregut, midgut and hindgut. The foregut primarily functions in food storage 54 and regulates the passage of food into the midgut. The insect midgut is largely analogous to the 55 small intestine in humans, and is the primary site of food digestion, and nutrient absorption 56 (Buchon et al., 2013). From the midgut, digested material enters the hindgut, which functions in 57 water resorption prior to defecation. The Malpighian tubules are blind-ended tubes that branch 58 from the midgut-hindgut junction, and are a single cell layer thick (Demerec, 1950) The Malpighian tubules are responsible for the production of primary urine containing water, ions, 59 60 toxins, and waste products (Dow, 2009; Buchon et al., 2013). Drosophila have four Malpighian 61 tubules; two anterior and two posterior, which all function in ion and water transport (Denholm 62 and Skaer, 2009; Chintapalli et al., 2012).

63

64 In addition to its critical role in facilitating our understanding of gut and renal physiology,

65 *Drosophila* are also used to probe the roles of epithelial barriers in aging and disease (Dow et al.,

66 1994; Rera et al., 2012; Rodan et al., 2012; He & Jasper, 2014; Clark et al., 2015; Dambroise et

67 al., 2016). Advanced age, infections and thermal stress, for example, are all known to cause a 68 loss or decrease in epithelial barrier function (Jasper, 2015; MacMillan et al., 2017), and this 69 barrier dysfunction is thought to contribute to ageing phenotypes and/or stress-induced injury or 70 death (Katzenberger et al., 2015; Cerqueira César Machado and Pinheiro da Silva, 2016). In most 71 of these studies, barrier dysfunction in fruit flies is identified and quantified using the "Smurf 72 assay" (Rera et al., 2011, Martins et al., 2018). In this assay, flies are fed a food containing a 73 high concentration of a non-toxic organic dye called Brilliant blue FCF (BB-FCF). Flies that are 74 healthy appear to retain the dye in their gut lumen, but those that have some form of intestinal 75 epithelial dysfunction turn blue (Rera et al., 2011). It is presumed that the presence of these 76 Smurf phenotypes is a direct result of BB-FCF leaking down its concentration gradient (i.e. 77 passively diffusing) from the gut lumen into the hemolymph because of barrier disruption, which 78 will eventually make the fly appear blue (Rera et al., 2011).

79

80 When exposed to cold temperatures, D. melanogaster become less capable at regulating 81 hemolymph osmolarity and ion balance, and as a result, hyperkalemia causes cell depolarization 82 and cell death (MacMillan et al., 2015a). This inability to maintain ion and water balance in the 83 cold has been attributed, in part, to a susceptibility of the renal system to chilling (Overgaard and 84 MacMillan, 2017). Specifically, low temperatures suppress ionomotive enzyme activity in the Malpighian tubules and gut epithelia, such that a net leak of Na⁺ and water from the hemolymph 85 to the gut causes a toxic increase in concentration of K⁺ in the hemolymph (Overgaard and 86 87 MacMillan, 2017), This cascade of issues can be largely prevented in *Drosophila* given a period 88 of cold acclimation (MacMillan et al., 2015b; Yerushalmi et al., 2018). When ion balance is 89 disturbed, recovery of K^+ balance (and thus cell membrane potential) is thought to be particularly

90	important to the recovery of neuromuscular function following rewarming (MacMillan et al.,
91	2012; Andersen and Overgaard, 2019). The ability of insects to recover can be quantified by
92	measuring the time required for an insect to stand following chilling (termed chill coma recovery
93	time; CCRT) (Macdonald et al., 2004; Findsen et al., 2014). Like other stressful conditions, cold
94	exposures also cause gut barrier dysfunction (MacMillan et al., 2017), but this cold-induced
95	barrier failure was described using another common paracellular probe (dextran) and has
96	previously not been investigated using BB-FCF.
97	
98	Some solutes are able to pass through the epithelial cells of an insect gut, while others can't
99	(Huang et al., 2015). Large polar organic molecules (like BB-FCF or dextran) cannot easily pass
100	through cell membranes, but there are two primary routes by which they could potentially cross

101 epithelial barriers; transcellular and paracellular transport. Transcellular transport occurs across

102 cell membranes via active (i.e. energy-consuming) or passive routes (O'Donnell & Maddrell,

103 1983). Many complex organic compounds have been shown to be actively (e.g. para-

aminohippuric acid, tetraethylammonium) (Linton & O'Donnell, 2000; Rheault & O'Donnell,

105 2004), or passively (e.g. urea, sucrose) (Maddrell & Gardiner, 1974) transported across

106 Drosophila epithelia. Many inorganic ions and compounds are actively transported at high rates

107 by the gut and renal epithelia of insects (e.g. phosphate, Mg^{2+}) (Maddrell & O'Donnell, 1992).

108 Passive diffusion of a molecule down its concentration gradient can occur in the transcellular

109 manner (e.g. through suitable channels) provided the molecule has a low molecular weight and is

soluble in the hemolymph. By contrast, primary active transport occurs transcellularly through

111 dedicated transporters and secondary active transport can mobilize molecules up or down a

112 concentration gradient by coupling concentration gradients through exchangers (Maddrell &

113 O'Donnell, 1992; Rheault & O'Donnell, 2004).

114Paracellular transport is defined as the transport of molecules/solutes across the epithelial cell

115 layer, specifically between the cells via the intercellular spaces (O'Donnell & Maddrell, 1983).

116 In insects, paracellular permeability is regulated by the presence of septate junctions (SJ), protein

117 complexes analogous to vertebrate tight junctions that play an important role in selectivity of

passage (O'Donnell & Maddrell, 1983; Jonusaite et al., 2017). SJs act as permeability barriers to

the free diffusion of molecules, and thereby control the rate at which molecules can leak down

their concentration gradient from one side of an epithelium to the other (Banerjee et al., 2006;

121 Ganot et al., 2014; Jonusaite et al., 2016). There are two types of these junctions in insects,

122 pleated and smooth SJs; while pleated SJs are typically found in the salivary glands, epidermis

and the photoreceptors of the flies, smooth SJs are primarily found in endodermal tissues like the

midgut and Malpighian tubules (Hall and Ward, 2016; Jonusaite et al., 2017).

125

126 The mode of transport of BB-FCF through the Drosophila gut epithelia is unknown, and despite 127 its use in understanding epithelial health and disease, whether or not flies can actively transport 128 this dye remains untested. Importantly, other organic compounds can be actively transported by 129 the Drosophila midgut and Malpighian tubules, including TEA, Texas red and methotrexate 130 (Rheault & O'Donnell, 2004; O'Donnell & Leader, 2006; O'Donnell, 2009). Texas red (625.12) g mol⁻¹), for example, is of a similar size to brilliant blue (792.85 g mol⁻¹), and is transported 131 132 across the Malpighian tubule epithelia via the action of a sodium-independent active transport 133 mechanism (Leader and O'Donnell, 2005). This active transport has previously been confirmed

by experimentally saturating the putative transporter with high concentrations of Texas red(Leader and O'Donnell, 2005).

136

137 In the present study, we hypothesized that 1) Like other stressors, cold stress can cause the Smurf 138 phenotype in *Drosophila*, but that this outcome could be due to either barrier failure or a reduced capacity for active transport in the cold because 2) the gut and Malpighian tubule epithelia of 139 140 Drosophila are innately capable of actively secreting BB-FCF. We used dye leak and clearance 141 assays using whole animals and isolated organs to test whether and how flies transport the dye 142 and found that cold stress does indeed cause the Smurf phenotype in *Drosophila* and that Smurf 143 flies are somewhat more likely to die from cold stress. We argue that Smurf assays, however, 144 cannot distinguish between a decrease of dye transport rates and a failure to maintain epithelial barriers under conditions of stress, as the midgut and tubules are continuously clearing the dye 145 146 from the hemolymph via active transcellular transport.

147

148 Materials and Methods

149 Animal Husbandry

150 The line of *Drosophila melanogaster* used in this study were derived from isofemale lines

151 established from collections from London, Ontario and Niagara on the Lake, Ontario in 2007

152 (Marshall and Sinclair, 2009). D. melanogaster were reared in 200 mL bottles that contained ~50

153 mL of Bloomington *Drosophila* medium (a cornmeal, corn syrup, agar and yeast-based diet).

154 FD&C Blue dye #1 (brilliant blue FCF, BB-FCF) was a generous gift from Calico Food

155 Ingredients Ltd. (Kingston, ON). The flies used in our cold tolerance assays were kept at 25°C

156 with a 14 hr: 10 hr light:dark cycle in a MIR-154-PA incubator (Panasonic Corporation,

157	Kadoma, Japan). Adult flies were transferred to a fresh bottle with food to lay eggs for two hours
158	to ensure a consistent number (~200) of progeny per bottle. Following adult emergence
159	(approximately ten days after the eggs were laid), the flies were collected in 40 mL vials
160	containing 7 mL of food medium. The flies were sexed on a CO ₂ pad three days post-adult-
161	emergence (day 13 of their life cycle), this was done in less than 5 min to reduce the time flies
162	are exposed to the CO ₂ . All of the experiments were completed on adult females, seven days
163	after final ecdysis.
164	
165	Survival Assay
166	A total of 250 flies were used to quantify survival rates following cold stress. Dyed medium was
167	prepared using the standard rearing medium, but with BB-FCF added at a concentration of 2.5%
168	(w/v) (Rera et al., 2012). Flies were maintained on dyed medium for nine hours and then
169	transferred into empty 40 mL vials. Control flies (n=50; ~10 per vial) were observed at this point
170	for the Smurf phenotype. A fly was counted as Smurf when dye coloration was observed outside
171	of the digestive tract (observed using a microscope, without dissection), while they lay in ventral
172	position. After observation the control flies were transferred to new vials containing regular
173	Bloomington food for 24 h before survival was assessed. The rest of the vials were put in a
174	Styrofoam container with a mixture of water and ice (0°C). After 6, 12, 18 and 24h at 0°C,
175	subsamples of 50 flies were removed from the cold stress. After observation for the Smurf/non-
176	Smurf phenotypes, the flies were transferred to new vials containing standard Bloomington food
177	while separating the Smurf flies from the Non-Smurf flies into groups of ~10 flies/vial. The vials
178	with the recovering flies were left on their side for 24h before survival was assessed. Flies then

able to stand and walk were counted as alive, and flies not moving or moving but unable to standand walk counted as dead.

181

182 *Chill coma recovery time (CCRT)*

183 Flies were maintained on dyed medium (prepared as described above) for 9 h before individual 184 flies were transferred into empty 4 mL glass screw-top vials. The vials were put into a plastic bag 185 (10 vials/bag) and submerged in an ice-water mixture (0°C). After 12 h, all of the vials were 186 removed from the ice-water bath. Each fly was immediately assessed to determine whether or 187 not they had a Smurf phenotype. All of the vials were then arranged on a sheet of paper on a 188 laboratory bench at room temperature (23°C). All of the flies were observed for 1 h to record 189 CCRT. The CCRT was recorded as the time it took for the fly to stand up on all six legs, and the 190 flies that did not stand within the hour were considered to have suffered chilling injury.

191

192 *Cold acclimation:*

193 Thermal acclimation over a period of days can strongly influence cold tolerance in D. 194 melanogaster, and cold acclimated flies better maintain barrier function in the cold (MacMillan et al., 2017). To test whether cold acclimation prevented flies from having a Smurf phenotype, 195 196 we acclimated flies to one of two temperatures and examined their ability to retain the dye in 197 their gut. Flies (reared as described above) were collected on the day of adult emergence and 198 sorted by sex under short CO₂ anaesthesia (< 5 minutes) before being transferred to either 10°C 199 or 25°C. At six days of age (post adult ecdysis), warm- and cold-acclimated flies were 200 transferred to fresh vials of their standard medium containing 2.5% (w/v) of BB-FCF. The flies 201 were separated (without anaesthesia) into groups of 20-26 flies per vial (four vials for each

202 acclimation group) and the vials were placed in an ice-water mixture (at 0°C) for 16 h. After 203 removal, and while the flies were still in chill coma, they were placed under the view of a 204 dissecting microscope and the number of Smurf flies and non-Smurf flies were counted. 205 Dye clearance assay: 206 Dyed food medium was prepared as described above, and flies were maintained on the medium 207 for 9 h after which they were then transferred to empty 40 mL vials. Control flies were observed 208 at this point and scored for the Smurf phenotype. After observation, these control flies were 209 transferred to 40 mL vials with 7 mL of regular Bloomington media for 12 hours before the 210 ability of the flies to clear the dye from the gut could be seen. The remainder of the flies in the 211 empty vials were placed in an ice bath at 0°C for 12 h. After examination of the flies for Smurf 212 phenotype (observed using a microscope without dissection), the Smurfs were then placed into 213 40 mL vials with 7 mL of standard Bloomington media for 30 h, after which they were re-scored 214 for the Smurf phenotype to assess their ability to clear the dye from the hemolymph. Photos of 215 representative Smurf flies were taken directly following the cold stress using an Axiocam 105 216 color microscope camera (Zeiss, Oberkochen, Germany). Photos of the same flies were then 217 taken for direct phenotypic comparison.

218

219 Midgut dye leak and transport assay

To quantify the effects of cold exposure on the leak of BB-FCF across the *Drosophila* midgut, and whether the midgut is capable of transporting this dye, an *in situ* barrier function/transport assay was developed. Flies that had been fed on diet containing BB-FCF (as above) were dissected under paraffin oil such that the gut was surrounded by the oil but remained attached to the anterior (head and thorax) and posterior (abdomen) ends of the animal (both of which were

225	held in place using minute dissecting pins). A single 2 µL droplet of <i>Drosophila</i> saline (117.5
226	mM NaCl; 20 mM KCl; 2 mM CaCl ₂ ; 8.5 mM MgCl ₂ ; 10.2 mM NaHCO ₃ ; 4.3 mM NaH ₂ PO ₄ ; 15
227	mM MOPS; 20 mM glucose; 10 mM glutamine; pH 7.0) was placed on the posterior midgut. The
228	preparation was then either left at 23°C, or was transferred to 0°C by placing it on the surface of
229	an ice water mixture within an insulated container. After 3h, a 1 μ L sample of the droplet on the
230	gut was taken and mixed with 100 μ L of milliQ water in a 96-well microplate. Absorbance was
231	read at 630 nm and concentrations of BB-FCF that had leaked into the droplet from the gut were
232	quantified by reference to an appropriately ranged BB-FCF standard curve. These values were
233	then expressed relative to the length of the midgut inside the droplet in each preparation. To
234	determine whether the Drosophila midgut could actively transport BB-FCF up a concentration
235	gradient, we conducted the same assay at 23°C, but with a 2 μ L droplet of saline containing a
236	known concentration of the dye (75 μ M), well below that of the diet.

237

238 Ramsay Assays

239 To test whether the Malpighian tubules can transport BB-FCF, and whether this transport is 240 likely to be carrier-mediated, we used a modified Ramsay assay procedure, similar to that 241 described by Rheault and O'Donnell (Rheault and O'Donnell, 2004). The flies used for the 242 Ramsay assays were reared under the same conditions with the exception that the light:dark 243 cycle was 12 hr:12 hr (simply because of available rearing space). Individual flies were briefly 244 aspirated into ethanol and the exoskeleton was carefully dissected away to reveal the gut and 245 Malpighian tubules. The Malpighian tubules were excised by cutting at the base of the ureter 246 using a 3G needle. Using a thin pulled glass rod, the pair of tubules was transferred to a saline 247 droplet in a dish containing heavy paraffin oil. The control droplets submerged in the oil were 10

248	μ L of 1:1 mixture of <i>Drosophila</i> saline (described above) and Schneider's Insect Medium
249	(Sigma-Aldrich S0146, St. Louis, USA). The experimental droplets contained the same 1:1
250	mixture mentioned above, but had BB-FCF added to obtain a range of dye concentrations
251	between 0.0126 μM and 25.2 μM (n=5-15 tubules per concentration). One tubule was wrapped
252	around a pin outside of the droplet, such that the ureter was half-way between the droplet and the
253	pin, and the other tubule remained in the droplet. The droplet that formed at the ureter was
254	removed after 15 min, and droplet volumes were approximated using droplet diameter, which
255	was measured using an ocular micrometer (assuming a sphere when the droplet was suspended in
256	the oil). The droplets were then collected using a 2 μ L micropipette and transferred to an empty
257	microcentrifuge tube. A 3 μ L volume of <i>Drosophila</i> saline was added to each sample in order to
258	separate the aqueous fluid from any oil collected along with the droplet. After brief
259	centrifugation, a 2 μ L aliquot of aqueous layer was transferred to a Take3 microwell plate
260	(BioTek, Winooski, USA) and the absorbance was read using the Biotek Cytation 5 Imaging
261	Reader at 630 nm. The absorbance reading was then compared to a standard curve to determine
262	the concentration of BB-FCF, which was then used to calculate the original concentration in the
263	primary urine.

264

265 Data Analysis

All data analysis was completed using R (version 3.5.2). Linear regression was used to assess
whether the tendency for flies to adopt a Smurf phenotype increased with cold exposure
duration, and survival following cold stress was analyzed using a generalized linear model with a
binomial error distribution (with phenotype and time included as factors). A Welch's two sample
t-test were used to determine whether there was a difference in CCRT between Smurfs and non-

271	Smurfs, and whether cold acclimation altered the tendency of flies to become Smurfs. The effect
272	of temperature on the leak of BB-FCF from the midgut was examined using a one-way ANOVA
273	To analyze our Ramsay assay data, we fit a Michaelis-Menton type model to the relationship
274	between the concentration of BB-FCF bathing the Malpighian tubules and the rate of dye flux
275	(nmol min ⁻¹) through the tubules using the nls() function. This model was used to determine the
276	maximal rate of dye clearance (J_{max}) and the saline concentration at which dye transport rates
277	were $\frac{1}{2}$ the maximal rate (K _t).
278	

279 Results:

280 Low temperature survival and chill coma recovery

Increased cold exposure times increased the likelihood of a fly becoming a Smurf, and decreased the probability of survival following chilling. We found a significant linear relationship between the duration of exposure to 0°C and the proportion of the flies that turned blue (i.e. became Smurfs; P = 0.049; Fig. 1A). Longer exposures to 0°C also resulted in lower rates of survival $P_{\text{time}} < 0.001$; Fig. 1B), and flies scored as Smurfs were slightly, but significantly, more likely to die from a cold stress ($P_{\text{Smurf}} = 0.024$; Fig. 1B).

287

The chill coma recovery time (CCRT) of both Smurf and non-Smurfs was scored following 12 h at 0°C. Although the tendency to turn blue was related to low temperature survival, it was not significantly related to CCRT. The Smurf flies had an average recovery time of 44.2 ± 10.9 minutes, and the non-Smurf flies had an average recovery time of 42.6 ± 8.07 minutes (t₂₃ = 0.530, *P* = 0.6010; Fig. 1C).

293 *Effects of acclimation on barrier disruption*

294 To examine whether improvements in cold tolerance associated with cold acclimation may be 295 related to the ability of the flies to avoid barrier disruption in the cold, we quantified the number 296 of warm- and cold-acclimated flies with a Smurf phenotype following cold stress. Both warm-297 and cold- acclimated flies fed Bloomington food containing BB-FCF retained the dye in the gut 298 lumen prior to any cold stress, but exposure to 0°C for 16h caused flies to lose barrier function 299 and turn blue (Fig. 1C-E). The tendency to turn blue during cold stress, however, was 300 significantly associated with the acclimation history of the flies ($t_4 = 39.2$, P < 0.001); the cold 301 exposure caused $\sim 90\%$ of warm-acclimated flies to turn blue, only $\sim 10\%$ of cold-acclimated flies 302 turned blue after the cold stress (Fig. 1E). 303

304 *Smurf recovery assay*

305 The Smurf recovery assay was used to test whether Drosophila with the Smurf phenotype could 306 recover, and clear the BB-FCF from their hemolymph. If possible, we expected such recovery to 307 change flies from a cold-induced Smurf to a non-Smurf phenotype within 30 h at benign 308 temperatures. None of the control flies (i.e. those not given any cold exposure) became Smurf 309 flies (n=25), and consequently, all of these control flies (n=25) were still non-Smurfs after the 310 12-hr recovery from the blue food, with 7 flies showing some residual blue dye inside the 311 rectum. Of the total Smurf flies (n = 40), only 16 survived the cold exposure, however, of the 16 312 Smurf flies that survived, 14 resulted in non-Smurf phenotypes following the 30 h recovery 313 period (Fig. 2). The remaining flies (n=2) had some residual blue within their hemolymph and 314 rectum.

316 *Midgut dye leak and transport assay*

317 To test whether the midgut leaks BB-FCF in situ, we measured the appearance of the dye in a 318 droplet surrounding the posterior midgut of flies. Notably, midguts of flies fed on the dye had measurable leak $(3.32 \pm 0.49 \text{ pmol mm}^{-1} \text{ h}^{-1}$ in the mucosal to serosal direction) even at room 319 temperature (Fig. 3). Midguts also leaked the dye significantly faster at 0°C than at 23°C ($F_{14} =$ 320 321 7.7, P = 0.015). To test whether control (room temperature) midguts could also transport the dye 322 up a concentration gradient (despite concurrent leak), we added a known concentration of the 323 dye to the bathing saline, and quantified disappearance of the dye from the saline. At 23°C, 324 midguts had a net transport rate of the dye (in the serosal to mucosal direction) of 6.10 ± 0.66 $pmol mm^{-1} h^{-1}$ (Fig. 3). 325 326 327 Ramsay Assays 328 The Malpighian tubules of *D. melanogaster* were used to determine whether or not the tubules 329 could actively transport BB-FCF. The Michaelis-Menten model closely fit the data and estimated a J_{max} value (highest possible rate of flux) of 232 fmol min⁻¹, and a K_t value (point where rate of 330 flux reached half of J_{max}) of 760 µmol L⁻¹ (Fig. 4). These results are also demonstrated 331 332 qualitatively with example images of the Ramsay assays at each initial saline concentration in the

supplementary material (Fig. S1).

334 Discussion

335 Here, we demonstrate for the first time that cold stress (like advanced age or traumatic injury) 336 causes the Smurf phenotype in *Drosophila*, and that the tendency to turn blue is related to cold 337 tolerance. We also show, however, that *Drosophila* appears to be able to transport BB-FCF, 338 which complicates its use as a marker of barrier disruption. 339 340 Like other stresses, cold exposure caused flies to adopt a Smurf phenotype (turn blue), and 341 longer cold stresses increased the proportion of Smurf flies (Fig. 1A). We noted an increase in 342 abundance of Smurf phenotypes as the duration of cold exposure increased and as survival rates 343 declined, but there was also a drastic decrease in survival as well (Fig. 1B). With these findings 344 alone, one could presume that barrier failure was being well characterized by this blue dye, 345 especially considering that barrier dysfunction occurs in the cold when measured using dextran 346 (MacMillan et al., 2017), and that other large polar molecules like glucose can leak into the 347 hemolymph under conditions of stress (Katzenberger et al., 2015). 348 349 Insects, including Drosophila have an acute ability to recover from cold exposure (Hoback and 350 Stanley, 2001). Upon chilling, *Drosophila* will enter a comatose state (chill coma), wherein their

neural and muscular systems are silenced (Rodgers et al., 2010). After they return to a normal
homeostatic temperature, the flies can revert from the comatose state back to a functional state
(David et al., 1998). The time required to return from chill coma is a common measure of cold
tolerance, and is related to the degree to which ion and water balance have been disturbed, both
in the nervous system and the hemolymph (Andersen and Overgaard, 2019). We performed a
CCRT experiment using the Smurf flies and the non-Smurf flies, and found no difference in

357 CCRT between the two phenotypic groups (Fig. 1C). This suggests that the severity of ion 358 balance disruption is not related to the tendency of BB-FCF to leak across the epithelial barrier. 359 Therefore, the cold stress appeared to make the epithelial barriers leaky, and this leak was 360 associated with low temperature, death and injury, but not with chill coma recovery. 361 362 We demonstrated that under normal conditions, there was no evidence of BB-FCF leaking/being 363 transported into the hemolymph, however after a stress is applied, there is strong evidence of net 364 leak of BB-FCF. Thermal acclimation can drastically alter cold tolerance phenotypes in adult 365 Drosophila (Ransberry et al., 2011; Colinet and Hoffmann, 2012). To examine whether or not 366 the tendency for flies to turn blue was related to thermal plasticity, we then investigated whether 367 or not acclimation would similarly impact the tendency of flies to turn blue during a cold stress. 368 Using this approach, we found a striking reduction in the tendency for cold-acclimated flies to 369 accumulate the blue dye in their hemolymph, relative to warm-acclimated flies (Fig. 1D-E). We 370 suggest two possible explanations for a reduction in Smurf phenotypes given a cold acclimation; 371 1) cold acclimation prevents epithelial barrier failure (MacMillan et al., 2017), or 2) cold 372 acclimation improves the ability of transporters to clear the dye from the hemolymph in the cold. 373 Given these two possibilities we opted to investigate whether or not flies were capable of 374 clearing BB-FCF from their hemolymph.

375

Our Smurf recovery assay demonstrated the ability of the flies to revert their phenotype from
Smurf to non-Smurf, which suggests that the flies can either transport and excrete BB-FCF out of
their hemolymph, or they are able to effectively metabolize it. In order to clarify whether flies
are capable of transporting BB-FCF, we tested for the capacity for transport of this dye through

380 the midgut and/or the Malpighian tubule epithelia using *in situ* assays. We took this approach 381 because BB-FCF dye is an organic anion which is similar in structure to other sulphonates (e.g. 382 amaranth, indigo carmine) that are already known to be transported by the Malpighian tubules 383 and midgut of *D. melanogaster* at relatively high rates (Linton & O'Donnell, 2000; O'Donnell et 384 al., 2003). We found that the Drosophila midgut is capable of effectively transporting this dye 385 against a concentration gradient (Fig. 3). This is similar to other large organics, which have also 386 been found to be transported across the midgut epithelium (Rheault & O'Donnell, 2004, Bijelic 387 et al., 2005; Chahine & O'Donnell, 2009). The Malpighian tubules function in ion and water 388 regulation but are also critical to the clearance of wastes and toxic dietary compounds from the 389 hemocoel (Dow, 2009; Denholm, 2013). We investigated the ability of the Malpighian tubules to 390 transport BB-FCF via modified Ramsay assays (Rheault and O'Donnell, 2004). The rate of flux 391 of BB-FCF into the Malpighian tubules was seen to increase along with the bathing 392 concentration at low concentrations (Fig. 4), but plateaued at high concentrations. A Michaelis-393 Menten model was fit to the relationship between the concentration of the dye in the bathing 394 saline and the rate of dye flux into the primary urine (Fig. 4). We found that the J_{max} of BB-FCF is approximately 232 fmol min⁻¹, which is comparable to other organic compounds such as 395 methotrexate and Texas red which have J_{max} values of 118.4 and 1028 fmol min⁻¹ respectively 396 397 (Leader and O'Donnell, 2005; Chahine and O'Donnell, 2009). Tetraethylammonium (TEA) is 398 similarly sized but a cation (and is thus mobilized by different transporters) but has a much higher J_{max} of 1.52 pmol min⁻¹ tubule⁻¹ (Rheault and O'Donnell, 2004). The K_t of BB-FCF (760 399 400 μ M) was higher than those of methotrexate and TEA in the Malpighian tubules of 401 D. melanogaster, as the K_t values for these two molecules were 12.2 μ M and 220 μ M 402 respectively (Rheault and O'Donnell, 2004; Chahine and O'Donnell, 2009). Overall, our Ramsay 403 assay results indicate that transport of BB-FCF is saturable, like several other similarly-sized404 organic compounds.

405

406	There are two known types of organic anion transporters in the Drosophila Malpighian tubules;
407	MRP1 and MRP2 (Multidrug resistant proteins) that transport either type I or type II organic
408	anions (OAs) (Leader & O'Donnell, 2005; Chahine et al., 2012). Type I OAs are generally
409	smaller molecules and are transported by Na ⁺ -dependent processes (Bresler et al., 1990; Linton
410	& O'Donnell, 2000). Type II OAs are typically larger molecules and are transported Na^+ -
411	independently (Leader and O'Donnell, 2005; Chahine and O'Donnell, 2009). Texas red, which is
412	an organic anion of a similar size to BB-FCF, is transported by MRP2 (Leader and O'Donnell,
413	2005). Based on similarity to Texas red, we suggest that BB-FCF is actively transported in a
414	Na ⁺ -independent manner by MRP2. In support of this hypothesis, BB-FCF has recently been
415	shown to inhibit MRP2 transport in human intestinal membrane vesicles in substrate competition
416	experiments, and MRP2 is capable of transporting BB-FCF into vesicles (Sjöstedt et al., 2017).
417	Future studies could verify whether MRP2 is responsible for the transport of BB-FCF into the
418	Malpighian tubules, such as transgenic knockdown of MRP2 or substrate competition
419	experiments.

420

421 A good marker of paracellular permeability requires a few important qualities; it should 1) have 422 a distinct colorimetric or radioactive attribute (allowing the marker to be quantified), 2) be non-423 lethal to the organism, and 3) not be able to be transported by the tissues and organs of the 424 organism under study. We argue that brilliant blue FCF, while convenient in studies of barrier 425 disruption for the first two requisite attributes, violates the third requirement. In most of the 426 present studies using BB-FCF to characterize barrier dysfunction, the non-Smurfs are presumed 427 to maintain all of the brilliant blue dye in the gut unless gut failure occurs, but we show here that 428 this may not be the case. Based on our findings, the midgut appears instead to leak the dye out of 429 the gut into the hemolymph, and the gut and Malpighian tubules secrete the dye back into the gut 430 in a constant cycle. Thus, even if a fly fed the dye is not a Smurf, there is still some brilliant blue 431 in the hemolymph at all times. The dye is simply being actively transported back into the gut at 432 high enough rates so as to keep the concentration of BB-FCF in the hemolymph low enough to 433 evade visual detection (i.e. all the flies are Smurfs, but some are more Smurfy than others). 434 Based on this model, any break in the cycle (e.g. via an inhibition of transport, paracellular barrier failure, or physical damage to the gut) could induce the Smurf phenotype, and 435 436 observation of the organismal phenotype is not indicative of any one form of failure. In the case 437 of cold exposure, low temperatures supress active transport rates by a factor of 2-3 for every 10°C drop in temperature (the Q₁₀ effect), and cold causes paracellular barrier disruption 438 439 (MacMillan et al., 2017), so at least two factors likely contribute to the cold-induced Smurf 440 phenotype described here.

441

We examined the practical use of brilliant blue dye (BB-FCF), and whether or not *Drosophila* could transport it via their midgut and/or Malpighian tubules. We concluded that both the midgut and Malpighian tubules have the ability to transport the dye. These results together suggest that BB-FCF is not an effective tool for measuring paracellular barrier dysfunction, as *Drosophila* can and do effectively transport BB-FCF across their intestinal epithelial barriers, possibly via MRP2. Importantly, these findings should not be interpreted as undermining the importance of the Smurf assay or what has been learned from it. Instead they should be seen as allowing for

449	new hypotheses on the precise mechanisms that cause a fly to turn blue during stress or at an
450	advanced age. In addition, we argue that the insect community should actively test other methods
451	of assaying paracellular barrier disruption, and be careful to avoid the assumption that markers
452	used in vertebrates (e.g. dextran, inulin, or polyethylene glycol) are inherently good markers for
453	insects.
454	
455	Data Accessibility:
456	All data is provided as a supplementary file for review and the same file will be uploaded to a
457	data repository (e.g. Dryad) should the manuscript be accepted for publication.
458	
459	Author Contributions:
460	All authors contributed to the conception and design of the study, D.L. H.P., and H.M. conducted
461	the experiments. D.L. and H.M. analyzed the data, D.L. drafted the manuscript, and all authors
462	edited the manuscript.
463	
464	Funding:
465	This research was supported by Natural Sciences and Engineering Research Council of Canada
466	Discovery Grants to both H.M. (grant RGPIN-2018-05322) and A.D. (grant RGPIN-2018-
467	05841).

469 **References:**

- Andersen, M.K., Overgaard, J., 2019. The central nervous system and muscular system play
 different roles for chill coma onset and recovery in insects. Comp. Biochem. Physiol. Part A
 Mol. Integr. Physiol. 233, 10–16. https://doi.org/10.1016/j.cbpa.2019.03.015
- Banerjee, S., Sousa, A.D., Bhat, M.A., 2006. Organization and function of septate junctions. Cell
 Biochem. Biophys. 46, 65–77. https://doi.org/10.1385/CBB:46:1:65
- Bijelic, G., Kim, N.R., O'Donnell, M.J., 2005. Effects of dietary or injected organic cations on
 larval *Drosophila melanogaster*: mortality and elimination of tetraethylammonium from the
 hemolymph. Arch. Insect Biochem. Physiol. Publ. Collab. with Entomol. Soc. Am. 60, 93–
 103.
- Bresler, V.M., Belyaeva, E.A., Mozhayeva, M.G., 1990. A comparative study on the system of
 active transport of organic acids in Malpighian tubules of insects. J. Insect Physiol. 36, 259–
 270. https://doi.org/10.1016/0022-1910(90)90110-2
- Buchon, N., Broderick, N.A., Lemaitre, B., 2013. Gut homeostasis in a microbial world: insights
 from *Drosophila melanogaster*. Nat. Rev. Microbiol. 11, 615.
- 484 Cerqueira César Machado, M., Pinheiro da Silva, F., 2016. Intestinal barrier dysfunction in
 485 human pathology and aging. Curr. Pharm. Des. 22, 4645–4650.
- Chahine, S., Campos, A., O'Donnell, M.J., 2012. Genetic knockdown of a single organic anion
 transporter alters the expression of functionally related genes in Malpighian tubules of *Drosophila melanogaster*. J. Exp. Biol. 215, 2601–2610.
- Chahine, S., O'Donnell, M.J., 2009. Physiological and molecular characterization of
 methotrexate transport by Malpighian tubules of adult *Drosophila melanogaster*. J. Insect
 Physiol. 55, 927–935. https://doi.org/10.1016/j.jinsphys.2009.06.005
- Chintapalli, V.R., Terhzaz, S., Wang, J., Al Bratty, M., Watson, D.G., Herzyk, P., Davies, S.A.,
 Dow, J.A.T., 2012. Functional correlates of positional and gender-specific renal asymmetry
 in *Drosophila*. PLoS One 7, e32577–e32577. https://doi.org/10.1371/journal.pone.0032577
- Clark, R.I., Salazar, A., Yamada, R., Fitz-Gibbon, S., Morselli, M., Alcaraz, J., Rana, A., Rera,
 M., Pellegrini, M., William, W.J., 2015. Distinct shifts in microbiota composition during *Drosophila* aging impair intestinal function and drive mortality. Cell Rep. 12, 1656–1667.
- 498 Colinet, H., Hoffmann, A.A., 2012. Comparing phenotypic effects and molecular correlates of
- developmental, gradual and rapid cold acclimation responses in *Drosophila melanogaster*.
 Funct. Ecol. 26, 84–93.
- Dambroise, E., Monnier, L., Ruisheng, L., Aguilaniu, H., Joly, J.-S., Tricoire, H., Rera, M.,
 2016. Two phases of aging separated by the Smurf transition as a public path to death. Sci.
 Rep. 6, 23523.
- David, R.J., Gibert, P., Pla, E., Petavy, G., Karan, D., Moreteau, B., 1998. Cold stress tolerance
 in *Drosophila*: analysis of chill coma recovery in *D. melanogaster*. J. Therm. Biol. 23, 291–
 299.
- Demerec, M., 1950. The Biology of *Drosophila*. Ann. Entomol. Soc. Am. 43, 522.
 https://doi.org/10.1093/aesa/43.4.522
- Denholm, B., 2013. Shaping up for action: the path to physiological maturation in the renal
 tubules of *Drosophila*. Organogenesis 9, 40–54. https://doi.org/10.4161/org.24107
- 511 Denholm, B., Skaer, H., 2009. Bringing together components of the fly renal system. Curr. Opin.
 512 Genet. Dev. 19, 526–532. https://doi.org/10.1016/j.gde.2009.08.006
- 513 Dow, J.A., Maddrell, S.H., Görtz, A., Skaer, N.J., Brogan, S., Kaiser, K., 1994. The Malpighian

- tubules of *Drosophila melanogaster*: a novel phenotype for studies of fluid secretion and its
 control. J. Exp. Biol. 197, 421–428.
- 516 Dow, J.A.T., 2009. Insights into the Malpighian tubule from functional genomics. J. Exp. Biol.
 517 212, 435 LP 445. https://doi.org/10.1242/jeb.024224
- 518 Dow, J.A.T., Romero, M.F., 2010. *Drosophila* provides rapid modeling of renal development,
 519 function, and disease. Am. J. Physiol. Physiol. 299, F1237–F1244.
- 520 https://doi.org/10.1152/ajprenal.00521.2010
- Findsen, A., Pedersen, T.H., Petersen, A.G., Nielsen, O.B., Overgaard, J., 2014. Why do insects
 enter and recover from chill coma? Low temperature and high extracellular potassium
 compromise muscle function in *Locusta migratoria*. J. Exp. Biol. 217, 1297 LP 1306.
 https://doi.org/10.1242/jeb.098442
- Ganot, P., Zoccola, D., Tambutté, E., Voolstra, C.R., Aranda, M., Allemand, D., Tambutté, S.,
 2014. Structural Molecular Components of Septate Junctions in Cnidarians Point to the
 Origin of Epithelial Junctions in Eukaryotes. Mol. Biol. Evol. 32, 44–62.
- 528 https://doi.org/10.1093/molbev/msu265
- Hall, S., Ward, R.E., 2016. Septate junction proteins play essential roles in morphogenesis
 throughout embryonic development in *Drosophila*. G3 Genes, Genomes, Genet. 6, 2375–
 2384.
- He, Y., Jasper, H., 2014. Studying aging in *Drosophila*. Methods 68, 129–133.
- Hoback, W.W., Stanley, D.W., 2001. Insects in hypoxia. J. Insect Physiol. 47, 533–542.
- Huang, J.-H., Jing, X., Douglas, A.E., 2015. The multi-tasking gut epithelium of insects. Insect
 Biochem. Mol. Biol. 67, 15–20. https://doi.org/10.1016/j.ibmb.2015.05.004
- Jasper, H., 2015. Exploring the physiology and pathology of aging in the intestine of *Drosophila melanogaster*. Invertebr. Reprod. Dev. 59, 51–58.
- Jonusaite, S., Donini, A., Kelly, S.P., 2016. Occluding junctions of invertebrate epithelia. J.
 Comp. Physiol. B 186, 17–43.
- Jonusaite, S., Kelly, S.P., Donini, A., 2017. Identification of the septate junction protein
 gliotactin in the mosquito *Aedes aegypti*: evidence for a role in increased paracellular
 permeability in larvae. J. Exp. Biol. 220, 2354 LP 2363.
- 543 https://doi.org/10.1242/jeb.156125
- Katzenberger, R.J., Chtarbanova, S., Rimkus, S.A., Fischer, J.A., Kaur, G., Seppala, J.M.,
 Swanson, L.C., Zajac, J.E., Ganetzky, B., Wassarman, D.A., 2015. Death following
 traumatic brain injury in *Drosophila* is associated with intestinal barrier dysfunction. Elife
 4, e04790.
- Leader, J.P., O'Donnell, M.J., 2005. Transepithelial transport of fluorescent p-glycoprotein and
 MRP2 substrates by insect Malpighian tubules: confocal microscopic analysis of secreted
 fluid droplets. J. Exp. Biol. 208, 4363–4376.
- Linton, S., O'Donnell, M.J., 2000. Novel aspects of the transport of organic anions by the malpighian tubules of *Drosophila melanogaster*. J. Exp. Biol. 203, 3575–3584.
- Macdonald, S.S., Rako, L., Batterham, P., Hoffmann, A.A., 2004. Dissecting chill coma
 recovery as a measure of cold resistance: evidence for a biphasic response in *Drosophila melanogaster*. J. Insect Physiol. 50, 695–700.
- 556 https://doi.org/https://doi.org/10.1016/j.jinsphys.2004.05.004
- 557 MacMillan, H.A., Baatrup, E., Overgaard, J., 2015a. Concurrent effects of cold and
- 558 hyperkalaemia cause insect chilling injury. Proc. R. Soc. B Biol. Sci. 282, 20151483.
- 559 https://doi.org/10.1098/rspb.2015.1483

- MacMillan, H.A., Andersen, J.L., Loeschcke, V., Overgaard, J., 2015b. Sodium distribution
 predicts the chill tolerance of *Drosophila melanogaster* raised in different thermal
- 562 conditions. Am. J. Physiol. Regul. Integr. Comp. Physiol. 308, 823–831.
- MacMillan, H.A., Williams, C.M., Staples, J.F., Sinclair, B.J., 2012. Reestablishment of ion
 homeostasis during chill-coma recovery in the cricket *Gryllus pennsylvanicus*. Proc. Natl.
 Acad. Sci. U. S. A. 109, 20750–20755. https://doi.org/10.1073/pnas.1212788109
- 566 MacMillan, H.A., Yerushalmi, G.Y., Jonusaite, S., Kelly, S.P., Donini, A., 2017. Thermal
- acclimation mitigates cold-induced paracellular leak from the *Drosophila* gut. Sci. Rep. 7,
 8807. https://doi.org/10.1038/s41598-017-08926-7
- Maddrell, S.H., O'Donnell, M.J., 1992. Insect Malpighian tubules: V-ATPase action in ion and
 fluid transport. J. Exp. Biol. 172, 417–429.
- Maddrell, S.H., Gardiner, B.O.C., 1974. The passive permeability of insect Malpighian tubules
 to organic solutes. J. Exp. Biol. 60, 641–652.
- Martins, R., McCracken, A., Simons, M., Henriques, C., Rera, M., 2018. How to catch a Smurf?
 Ageing and beyond...*in vivo* assessment of intestinal permeability in multiple model
 organisms. Bio-Protocol 8. doi:10.21769/BioProtoc.2722
- Miguel-Aliaga, I., Jasper, H., Lemaitre, B., 2018. Anatomy and physiology of the digestive tract
 of *Drosophila melanogaster*. Genetics 210, 357–396.
- 578 https://doi.org/10.1534/genetics.118.300224
- 579 O'Donnell, M.J., 2009. Too much of a good thing: how insects cope with excess ions or toxins in
 580 the diet. J. Exp. Biol. 212, 363 LP 372. https://doi.org/10.1242/jeb.023739
- O'Donnell, M.J., Ianowski, J.P., Linton, S.M., Rheault, M.R., 2003. Inorganic and organic anion
 transport by insect renal epithelia. Biochim. Biophys. Acta Biomembranes 1618, 194–206.
- 583 O'Donnell, M.J., Leader, J.P., 2006. Changes in fluid secretion rate alter net transpithelial
 584 transport of MRP2 and P□glycoprotein substrates in Malpighian tubules of *Drosophila*585 *melanogaster*. Arch. Insect Biochem. Physiol. Publ. Collab. with Entomol. Soc. Am. 63,
 586 123–134.
- O'Donnell, M.J., Maddrell, S.H., 1983. Paracellular and transcellular routes for water and solute
 movements across insect epithelia. J. Exp. Biol. 106, 231-253.
- 589 Overgaard, J., MacMillan, H.A., 2017. The integrative physiology of insect chill tolerance.
 590 Annu. Rev. Physiol. 79, 187–208.
- 591 Presland, R.B., Jurevic, R.J., 2002. Making sense of the epithelial barrier: what molecular
 592 biology and genetics tell us about the functions of oral mucosal and epidermal tissues. J.
 593 Dent. Educ. 66, 564–574.
- Ransberry, V.E., MacMillan, H.A., Sinclair, B.J., 2011. The relationship between chill-coma
 onset and recovery at the extremes of the thermal window of *Drosophila melanogaster*.
 Physiol. Biochem. Zool. 84, 553–559.
- Rera, M., Bahadorani, S., Cho, J., Koehler, C.L., Ulgherait, M., Hur, J.H., Ansari, W.S., Lo Jr,
 T., Jones, D.L., Walker, D.W., 2011. Modulation of longevity and tissue homeostasis by the *Drosophila* PGC-1 homolog. Cell Metab. 14, 623–634.
- Rera, M., Clark, R.I., Walker, D.W., 2012. Intestinal barrier dysfunction links metabolic and
 inflammatory markers of aging to death in *Drosophila*. Proc. Natl. Acad. Sci. U. S. A. 109,
 21528–21533. https://doi.org/10.1073/pnas.1215849110
- Rheault, M.R., O'Donnell, M.J., 2004. Organic cation transport by Malpighian tubules of
 Drosophila melanogaster: application of two novel electrophysiological methods. J. Exp.
- 605 Biol. 207, 2173–2184.

- Rodan, A.R., Baum, M., Huang, C.-L., 2012. The *Drosophila* NKCC Ncc69 is required for
 normal renal tubule function. Am. J. Physiol. Physiol. 303, C883–C894.
 https://doi.org/10.1152/ajpcell.00201.2012
- Rodgers, C.I., Armstrong, G.A.B., Robertson, R.M., 2010. Coma in response to environmental
 stress in the locust: a model for cortical spreading depression. J. Insect Physiol. 56, 980–
 990.
- Salvo Romero, E., Alonso Cotoner, C., Pardo Camacho, C., Casado Bedmar, M., Vicario, M.,
 2015. The intestinal barrier function and its involvement in digestive disease. Rev Esp
 Enferm Dig 107, 686–696.
- Sjöstedt, N., Deng, F., Rauvala, O., Tepponen, T., Kidron, H., 2017. Interaction of food additives
 with intestinal efflux transporters. Mol. Pharm. 14, 3824–3833.
- 617 https://doi.org/10.1021/acs.molpharmaceut.7b00563
- Yerushalmi, G.Y., Misyura, L., MacMillan, H.A., Donini, A., 2018. Functional plasticity of the
 gut and the Malpighian tubules underlies cold acclimation and mitigates cold-induced
- 620 hyperkalemia in *Drosophila melanogaster*; J. Exp. Biol. 221, jeb174904.
- 621 https://doi.org/10.1242/jeb.174904



Figure 1. Cold stress causes the Smurf phenotype. A) Proportion of flies with a Smurf

- phenotype following exposure to 0°C for up to 24 h (n=49-50 flies per time period). B) Rates of
 survival of cold exposed flies that did, and did not adopt the Smurf phenotype during cold
- exposure. The time to cause 50% mortality at 0°C (Lt₅₀) was slightly lower in Smurfs (13.8 \pm
- 630 1.53 h) than in non-Smurfs (17.8 \pm 1.18h). C) Chill coma recovery times (CCRT) of Smurf and
- 631 Non-Smurf flies following 12 h at 0° C (n=56 flies in total; 17 Smurf, 39 non-Smurf). Open
- 632 circles represent CCRT of individual flies and closed circles represent the mean \pm sem. Recovery
- from chill coma was not related to the Smurf phenotype. D) Representative cold- $(10^{\circ}C)$ and
- 634 warm-acclimated (25°C) flies following 16 h at 0°C. E) Proportions of cold- and warm-
- acclimated flies with a Smurf phenotype following 16 h at 0°C. Cold acclimation strongly
- reduces the probability of flies exhibiting a Smurf phenotype. Open circles represent proportions
- 637 from n=4 replicates per acclimation temperature, each with 20-26 flies. Closed circles represent
- 638 the mean \pm sem. Error bars that are not visible are obscured by the symbols.



- 640 Figure 2. *D. melanogaster* can clear brilliant blue FCF from their hemolymph, reversing
- 641 **the Smurf phenotype.** A) Smurf fly directly after removal from the cold stress. B) Smurf fly
- shown after recovery. Flies in 40 mL vials were submerged in an ice bath at 0°C for 12 hours to
- allow for the onset of the Smurf phenotype. The flies were then removed from the ice bath, and
- after inspection for the Smurf phenotype, they were put into an incubator at 25°C for 30 hours to
- allow them to clear the dye from the hemolymph.
- 646



648 649

Figure 3. The midgut of *D. melanogaster* can transport brilliant blue FCF up a steep

651 concentration gradient and cold causes net leak of the dye down its concentration gradient.
 652 Midguts dissected from adult female flies that had fed on 2.5% w/v brilliant blue FCF were

653 submerged under oil and a droplet of saline containing no dye was placed on the posterior

midgut and left for 3 h at either 23°C or 0°C. Net leak rates of the dye (solid circles; mean \pm

sem) were calculated from the appearance of the dye in the saline relative to the length of midgut

tissue suspended in the droplet. These values are compared to the rate of net dye transport at

657 23°C, calculated from the rate of disappearance of the dye from a droplet containing a known

658 concentration of brilliant blue FCF (75 μ M; white dashed line and grey area; mean \pm sem).



659

660 Figure 4. Rates of brilliant blue FCF transport through the Malpighian tubules of adult

661 **female** *Drosophila melanogaster*. Modified Ramsay assays were used to characterize rates of

transport of BB-FCF within 15 min of exposure to the dye. The black line denotes the fit of a Michaelis-Menten model. Solid symbols represent the mean \pm sem and open circles represent

663 Michaelis-Menten model. Solid symbols represent the mean \pm sem and open circles rep 664 values from individual tubules. Insert shows the ratio between the secreted BB-FCF

665 concentration (S) and the initial bathing solution of the BB-FCF (M).



666

Figure S1. The *Drosophila* **Malpighian tubules can transport brilliant blue FCF.** Modified Ramsay assays were performed to examine the effects of increasing BB-FCF concentrations in the bathing saline on the concentration of the resultant primary urine. A = Control (0 μ M), B = 0.0126 μ M, C = 0.126 μ M, D = 1.26 μ M, E = 3.15 μ M, and F = 12.6 μ M.