

Changes in eating habits and lifestyles in Peruvian population during social isolation by the COVID-19 pandemic

Salomón Huancahuire-Vega ^{1,*}, Edda E. Newball-Noriega ¹, Ricardo Rojas-Humpire ¹, Jacksaint Saintila ², Mery Rodriguez Vásquez ², Percy. G. Ruiz-Mamani ³, Wilter C. Morales-García ⁴, Michael White ⁵

¹ Departamento de Ciencias Básicas, Escuela de Medicina Humana, Facultad de Ciencias de la Salud, Universidad Peruana Unión - Lima, Perú;

² Departamento de Nutrición, Escuela de Nutrición Humana, Facultad de Ciencias de la Salud, Universidad Peruana Unión – Lima, Perú;

³ Universidad Privada San Juan Bautista, Lima, Perú;

⁴ Escuela de Posgrado, Universidad Peruana Unión – Lima, Perú;

⁵ Dirección General de Investigación, Universidad Peruana Unión – Lima, Perú

* Correspondence: Dr. Salomón Huancahuire-Vega

salomonhuancahuire@upeu.edu.pe;

Carretera central Km 19.5 Ñaña Chosica Lima 15 – Peru

Tel.: +51-997-574-011

Competing interests: The authors have no competing interests to declare.

Author contribution: S.H-V., E.E.N-N., and R.R-H., have contributed to conceptualization, formal analysis and writing the manuscript. J.S., M.R.V., and P.G.R-M. helped in data curation and writing—original draft preparation. W.C.M-G. and M.W. contributed to the investigation and manuscript writing. All authors have read and agreed to the published version of the manuscript.

Article type: Original Research

Keywords: COVID-19, lifestyle, feeding behavior, Peru, social isolation.

Running head: Eating habits and lifestyles in Peruvian population due to COVID-19

Abstract:

Background: The COVID-19 pandemic caused that some governments have implemented house confinement measures with probable consequences on lifestyle, particularly affecting eating habits, physical activity, sleep quality, and mental health.

Objectives: The aim of this study was to assess the frequency of lifestyles, physical activity and sleep characteristics, as well as changes in eating habits in the Peruvian population during COVID-19 pandemic.

Methods: A Cross-sectional descriptive study was performed. We analyzed adults from Peru between July to August 2020 based on an online self-administered questionnaire divided into sociodemographic, anthropometrics and COVID-19 diagnostic reported, lifestyle habits and frequency of consumption of foods.

Findings: During confinement by COVID-19, 1176 participants were studied, 39% were student, 37.5% were workers and 46% were assert not to work. The population asserted gain weight (1 to 3 Kg) and 35.7% were overweight. The lifestyles habits showed that 54.8% affirmed to doing physical activity and a large proportion (37.2%) asserted sleep less. The Peruvian population presented a main feeding patter of breakfast (95.7%), lunch (97.5%), dinner (89.1%) and brunch (44.9%). Likewise, feeding habits before and during COVID-19 pandemic showed that vegetables (OR:1.56, CI95% 1.21 - 200), fruit (OR: 1.42, CI95% 1.10 – 1.81), legumes (OR:1.67, CI95% 1.23 – 2.28) and eggs (OR: 2.00, CI95% 1.52 – 2.65) presented significantly consumption increase during social isolation, while bakery products (OR: 0.74, CI95% 0.56 – 0.97), meat, snack, refreshment and fast-food decrease consumption. Other food no significant differences were presented.

Conclusion: This study in a Peruvian population showed an important frequency of overweight and sleep disorders. There was a slight increase in physical activity despite the social isolation measures and an increase in health eating habits, nevertheless a majority reported gaining weight.

Keywords: COVID-19, lifestyle, feeding behavior, Peru, social isolation.

1. Introduction

An announcement made by the WHO on January 30, 2020 declared COVID-19 a global health emergency due to the exponential growth of cases in China and other countries of the world.¹ As of October 18, 2020, according to the Coronavirus Resource Center at Johns Hopkins University, more than 39 million cases have been confirmed, while the number of deaths has risen to more than 1 million.² Given the circumstances measures needed to control the spread of the disease, government authorities were forced to take decisive actions such as social distancing and mandatory isolation in homes, which have led to restrictions on various daily activities, with probable consequences on lifestyle, particularly affecting eating habits, physical activity, sleep quality, and mental health.³

In Peru, the first case of COVID-19 infection was reported by the government on March 6, 2020. Since then, the government has implemented a series of preventive measures in order to limit and contain the spread of the disease. On March 16 the government, through Supreme Decree No. 008-2020-SA, declared a nationwide state of health emergency for a period of ninety (90) calendar days, dictating prevention measures and mandatory social isolation (national quarantine) for the control of COVID-19.⁴ As of June 26, the government announced an extension that modified the

restriction to a targeted social isolation, but the state of health emergency and social distancing measures were extended until December 31.⁵

The period of social isolation due to COVID-19 has caused a series of changes in people's daily routine and lifestyle.⁶ Available evidence has shown that during the pandemic, physical activity decreased in the population, while the amount of screen time increased significantly.⁶ In fact, prior to the pandemic, these factors, along with obesity, were described as a public health problem in the Peruvian population, and COVID-19 may make this situation even worse.⁷ In addition, studies have shown the negative impact of sedentary behaviors on physical and mental health.⁸ Likewise, the fear caused by COVID-19 has an indirect effect, causing negative emotions by affecting the quality of sleep, contributing to an increase in obesity.⁹

Additionally, evidence has shown that the risks of choosing inappropriate eating behaviors can increase as a psychological and emotional response to COVID-19.¹⁰ In fact, it has already been shown that negative emotional experiences can lead to overeating.¹¹ Eating habits characterized by a high consumption of ultra-processed, high-calorie-density foods and low intake of fiber, antioxidants and other bioactive elements can affect the immune system, causing chronic activation of the innate system and inhibition of the adaptive immune system response by increasing oxidative stress, thus delaying defensive intervention against pathogens.¹²

Previous studies have demonstrated the beneficial effects of physical activity and sleep quality on mental health.¹³ In addition, adequate physical activity can improve the quality of sleep, reducing the risk of morbidity from chronic non-communicable diseases.¹⁴ Furthermore, appropriate eating habits can be a protective factor for physical and mental health, disease prevention and strengthening of the immune system.¹⁵ Indeed, improved lifestyle, as a possible strategy that includes regular

physical activity, less screen time, adequate rest and proper dietary habits through a diet based on consumption of minimally processed plant foods and healthy fats, would lead to better control of chronic noncommunicable diseases and thus strengthen the immune system during the COVID-19 pandemic.¹⁵ The objective of the present study was to evaluate the frequency of lifestyles, physical activity and sleep characteristics, as well as changes in eating habits in the Peruvian population during the period of social isolation due to the COVID-19 pandemic.

2. Materials and Methods

2.1. Design and participants

This is a cross-sectional descriptive study based on a self-administered questionnaire. This questionnaire was addressed to the Peruvian adult population (over 18 years of age), and was applied using a non-probabilistic convenience sampling. Responses from 1218 participants were obtained, of which 42 were eliminated due to incompleteness and certain inconsistencies related to age, height, weight, etc.

The study followed the international ethical recommendations contained in The Declaration of Helsinki. All participants were informed about the characteristics of the study as well as their anonymous, voluntary and confidential participation. This study was approved by the Ethics in Research Committee of the Universidad Peruana Unión (No. 2020-CEUPeU-00013).

2.2. Procedure

The questionnaire was applied from July 16 to August 31, 2020 (period in which the Peruvian population was in partial confinement) using the Google form platform and disseminated through social networks (Facebook and Whastapp) and institutional

mailing lists. The questionnaire was completed through any electronic device with an internet connection. On the initial page, the participants were explained the voluntary nature of their participation with informed consent, as well as the justification, objectives, possible risks and benefits of the study before continuing with the instrument itself.

2.3. Questionnaire and variables

The questionnaire was developed by the authors to specifically address the particularities of the current pandemic context and its impact on lifestyle. The questionnaire included 42 questions divided into four sections: Sociodemographic data (10 questions: gender, marital status, age, level of education, occupational status, employment status, if the social isolation was carried out, duration of social isolation, people in the home before and during social isolation); anthropometric data and COVID-19 diagnosis reported (5 questions: weight, height, weight change during social isolation, number of kilograms, COVID-19 diagnosis); information on lifestyle habits (11 questions: physical activity, change and hours of sleep, meals eaten per day during confinement); information on the frequency of consumption of the main food groups (16 questions: vegetables, fruits, legumes, dried fruits/nuts, meat, fish, eggs, milk, yogurt, bakery products, snacks, beverages, beer, wine, other alcohol, fast food) before and during social isolation, based on five response categories: Participants who reported having consumed food from 1 to 4 times/week were considered as low consumption, while high consumption was considered when they reported having consumed from 5 times/week to 2 or more times/day.¹⁶

2.4. Statistical analysis

The data analysis was performed in RStudio v4.0.2 (R Foundation for Statistical Computing, Vienna, Austria; <http://www.R-project.org>). The variables were encoded and analyzed such as categorical. To descriptive analysis categorical variables were cluster in to tables with absolute and relative frequency (%). To comparative analysis χ^2 or Fisher's exact test were used to categorical variables depending of distribution of variable. Subsequently, to determinate differences before and during COVID-19 pandemic McNemar test was performed. A $p < 0.05$ was considered statistically significant.

3. Results

A summary of the sample's sociodemographic characteristics can be found in Table 1. In total 1176 participants were studied, 571 men and 605 women. The largest age groups were 18 to 25 years old (44.4%) with predominantly women and 26 to 40 years old (33.7%) with predominantly men. Overall, 67.7% were single and 38% had completed university studies. During social isolation, 39% of those sampled were students, 37.5% were workers, 30.8% worked with a formal contract, 46% were assert not to work and 67.6% were under social isolation for more than 61 days as of the date of the study.

A total of 35.7% of those studied were overweight and 8.4 % were obese, with a majority of men in both groups. A large portion reported having gained weight, 20.8% and 13.7% between 1 to 3 kg and 4 to 7 kg respectively. 8.7% of those surveyed had a confirmed diagnosis of COVID-19.

Some aspects of lifestyles showed patterns and changes when grouped by sex during the social isolation (Table 2). More than half of those studied affirmed to doing physical activity (54.8%) between 1 to 2 (14.7%) and 3 to 4 (26.4%) days per week, for at least 30 min. The reported sleep routine presented significate changes wherein

37.2% responded that they slept less and 36.9% slept more hours at night compared to before the pandemic. The hours of sleep per night were largely between 4 to 6 (35.6%) and 7 to 9 (55.4%) hours.

Eating patterns during the social isolation showed that approximately 90 to 95% ate breakfast, lunch and dinner. Additionally, some extra food consumption was reported for brunch (42.9 to 46.8%), afternoon meal (38.2 to 39.8%) and snacks (35.2 to 40.8%), these results presented no differences by sex (Figure 1).

The eating habits showed some changes before and during social isolation (Table 3). Fruit and vegetables presented elevated frequency of consumption; while, legumes and dried fruits/nuts showed low frequency of consumption (Figure 2). On the other hand, just vegetables (OR:1.56, CI95% 1.21 - 2.00), fruit (OR: 1.42, CI95% 1.10 – 1.81) and legumes (OR:1.67, CI95% 1.23 – 2.28) presented significantly consumption increase during social isolation. Approximately a quarter of the population presented high consumption of meat and eggs (Figure 3); while, dairy products and fish showed low frequency of consumption. In the animal-based foods only eggs (OR: 2.00, CI95% 1.52 – 2.65) presented significantly increasing during social isolation. The consumption of alcoholic drinks and ultra-processed food were the lowest; while, approximately 20% and 40% of population presented high consumption of beverages and bakery products, respectively (Figure 4). In the processed food, just bakery products (OR: 0.74, CI95% 0.56 – 0.97) presented change before and during social isolation.

4. Discussion

The objective of this study was to evaluate the frequency of lifestyles, physical activity and sleep characteristics, as well as changes in eating habits before and during social isolation due to the COVID-19 pandemic in a Peruvian population.

The findings of this study show that more than 60% of those surveyed remained at home, as a result of the imposition of social isolation due to outbreaks of COVID-19, and thus despite the more than 61 days since the start of the pandemic in Peru, only 8.7% had a confirmed diagnosis of COVID-19. Studies indicate that sustained transmission in the early stage of the epidemic in Peru (March 2020) showed an exponential growth dynamic despite non-pharmaceutical interventions by the authorities such as compulsory social isolation.¹⁷ In addition, Peru had some of the highest infection and mortality rates in the world, along with countries such as Italy, Spain, Brazil, the United Kingdom and France.¹⁸

This study highlights the potential impact of social isolation due to COVID-19 by including a variety of weight-related behaviors in Peruvian adults. The results of this study indicate that of the total number of participants subjected to social distancing, 35.7% were overweight and 8.4% were obese. Likewise, the findings show that the imposition of social isolation influenced lifestyles causing negative effects and weight changes, the population reported gaining weight with 20.8% between 1 to 3 kg and 13.7% from 4 to 7 kg respectively. Similar studies in 765 patients in Kurdistan, Iraq indicated a weight gain during social isolation of less than 2 kg in almost all patients (98.05%) and those who gained more than 3 kg were mostly women or from the center of large cities.¹⁹ Another study on the BMI of a Lebanese population showed 30.8% overweight and 12.9% obese.²⁰ On the other hand, 32% of the sample population (2364 participants) in a study in the United Kingdom was classified as obese during social isolation.²¹ In addition, a perceived increase in body weight was reported in the United States with an average body mass index (BMI) of 25.8.²²

Weight gain is associated with BMI, as those with a higher BMI gain weight more often compared to those with a normal BMI. Studies prior to the pandemic showed a

prevalence of overweight and obesity in Peruvians ≥ 20 years old and it was women who had a higher prevalence of obesity and abdominal obesity than men.²³ The results of this study indicate a weight gain of between 1 and 7 kg, it is likely that high rates of stress and depression will push people to consume foods with high sugar content and generate lifestyle changes and negative psychological impacts (increased prolonged naps, fears of infection, frustration, boredom, inadequate information, financial losses, negative psychological effects, etc.), causing long-term effects on obesity and the presence of co-morbidities, making obesity a higher risk of death related to the COVID-19 pandemic.²⁴

The association of severe COVID-19 cases with higher than normal BMI values has been demonstrated.²⁵ For example, 88.2% of patients admitted to Wuhan Union Hospital between January and February 2020 who were not survivors of this illness had a BMI > 25 kg/m².²⁶ Some explanations state that a higher than normal BMI increases susceptibility to respiratory infections due to the deterioration of lung mechanics and higher concentrations of pro-inflammatory molecules.²⁷ On the other hand, the excessive amount of adipose tissue in the abdominal area decreases diaphragmatic excursion and affects the ventilation of the base of the lung, which can lead to hypoxemia, clinical characteristics observed in this illness.²⁸

Pre-pandemic studies showed that the Peruvian population had a high prevalence of physical inactivity and sedentary behaviors.²⁹ This study found substantial changes in behavior during social isolation, the majority of participants claimed to be physically active (54.8%) for 1-2 (14.7%) or 3-4 (26.4%) days per week, for at least 30 minutes. Similar studies showed increased regular participation in moderate and vigorous physical activity in older adults during quarantine.³⁰ On the other hand, other studies show that total physical activity decreased significantly during social isolation, due to

greater sedentary behavior, more prolonged during the weekend.³¹ While some individuals report a decrease in physical activity due to mobility restrictions in the context of COVID-19, others report increases in physical exercise and other efforts to "stay fit".³²

In most studies, social isolation induced a reduction in physical activity and an increase in sitting time, or only low-intensity activities such as housework.³¹ This sudden cessation of physical exercise is detrimental in the context of this disease, since this decrease is associated with rapid muscle atrophy and reduced energy consumption by the muscles, promoting obesity and lipid accumulation, decreased ability of organ systems to resist viral infection, increased risk of damage to the immune system, respiratory system, cardiovascular system, musculoskeletal system and brain.³³ Therefore, the WHO recommends at least 150 minutes per week of moderate to vigorous intensity physical activity or 75 minutes of high-intensity physical activity per week, or a combination of both.³⁴

More than 70% of the participants in this study report substantial changes in sleep patterns during the period of isolation due to COVID-19. Similar studies revealed a high prevalence of anxiety sleep disorders and depressive symptoms in the population during the COVID-19 social isolation period, with women reporting a higher risk of sleep difficulties,³⁵ which was also observed in our study. It is likely that women were more receptive to expressing psychological distress and somatic symptoms.³⁶ On the other hand, the uncertainty of being infected by the coronavirus would cause greater pressure in the population, relating to hypochondriacal concerns, sleep apnea, and childhood trauma that can be reflected in sleep mentality, affecting its quality.³⁷ Sleep plays a critical role in physical and mental health, and adequate sleep duration and quality are essential to cope with major life events such as the COVID-19 pandemic.³⁸

This study also showed significant changes in the eating habits that this Peruvian population adopted during social isolation, among which the increase of eggs, legumes, fruit and vegetables, as well as the decrease of bakery products. Some studies observed an improvement in the pattern of consumption of healthy foods and a restriction of unhealthy foods, especially in the younger population (<30 years).³⁹ In contrast, studies in Italy indicated changes in eating habits in almost half the population, so that women were more likely than men to increase their food intake to feel better.⁸ In other countries, after-dinner snacks, carbohydrate consumption, consumption of sweets, cookies, and cakes, and consumption of fresh food products increased.⁴⁰ The lack of dietary and eating restriction may be due to a response to stress and reduced physical activity, which could lead to decreased nutritional quality.⁴⁰ This unhealthy behavior is observed in other studies primarily in adult populations with obesity and in settings with a lower eating environment index.²¹

In addition to changes in eating habits, alcoholic drinks and ultra-processed food consumption was the lowest and showed no changes before and during social isolation. Some studies suggest that increased alcohol consumption during the COVID-19 pandemic may be a response to boredom, inactivity, isolation, and as a means of combating anxiety, stress, and/or loss of sleep as a result of containment measures introduced to minimize the spread of the virus.⁴¹

Healthy lifestyles are relevant to various diseases as well as to maintaining people's mental health. Among these, eating habits may be a factor contributing to the geographical variation of COVID-19 infection rates, as the consumption of certain foods may be associated with lower mortality. A balanced diet is vital for the prevention of COVID-19, related chronic diseases and has a favorable impact on COVID-19 mortality.⁴²

The main limitation of this work is the lack of representativeness of the population sample. Therefore, it is not possible to expand the results to the population of the country. There is also an over-representation of younger individuals (< 25 years). Moreover, all data collected are self-reported and this could make them not completely reliable, especially when it comes to reporting behaviors for which there may be a social stigma (e.g., alcohol and tobacco consumption). Another limitation could be the lack of information about religious practices which could have influenced diet and lifestyle changes.

In conclusion, this study in a Peruvian population showed an important frequency of overweight and sleep disorders. On the other hand, efforts to included physical activity despite the social isolation measures, an increase in health eating habits, but nevertheless a majority reported gaining weight. Future studies should consider additionally potential factors, starting with the observation in this study that most participants had a changed sleep pattern during social isolation, and potentially including other factors like stress, type of exercise, pedometer step counts, and portion sizes.

Author statements

Funding: This research received no external funding.

Data Availability Statement: The data presented in this study are available on request from the corresponding author.

Competing interests: The authors have no conflict of interest

References

1. Velavan TP, Meyer CG. The COVID-19 epidemic. *Trop Med Int Health*. 2020;25(3):278-80.
2. Rosenkrantz L, Schuurman N, Bell N, Amram O. The need for GIScience in mapping COVID-19. *Health & Place*. 2020:102389.
3. Di Renzo L, Gualtieri P, Pivari F, Soldati L, Attina A, Cinelli G, et al. Eating habits and lifestyle changes during COVID-19 lockdown: an Italian survey. *J Transl Med*. 2020;18(1):229.
4. Gobierno del Perú. Decreto Supremo N 008-2020-SA que declara en emergencia sanitaria a nivel nacional por el plazo de noventa (90) días calendario y dicta medidas prevención y control del Covid-19. *Diario Oficial El Peruano*, 11 de marzo de 2020.
5. Gobierno del Perú. Decreto Supremo N° 027-2020-SA que Prorroga la Emergencia Sanitaria declarada por Decreto Supremo N° 008-2020-SA, prorrogada por Decreto Supremo N° 020-2020-SA. *Diario Oficial El Peruano*, 28 de agosto de 2020.
6. Pellegrini M, Ponzio V, Rosato R, Scumaci E, Goitre I, Benso A, et al. Changes in Weight and Nutritional Habits in Adults with Obesity during the "Lockdown" Period Caused by the COVID-19 Virus Emergency. *Nutrients*. 2020;12(7).
7. Najera H, Nandy S, Carrillo-Larco RM, Miranda JJ. Within-country migration and obesity dynamics: analysis of 94,783 women from the Peruvian demographic and health surveys. *BMC Public Health*. 2019;19(1):263.
8. Di Renzo L, Gualtieri P, Cinelli G, Bigioni G, Soldati L, Attinà A, et al. Psychological Aspects and Eating Habits during COVID-19 Home Confinement: Results of EHLC-COVID-19 Italian Online Survey. *Nutrients*. 2020;12(7).

9. Chaput JP, Dutil C. Lack of sleep as a contributor to obesity in adolescents: impacts on eating and activity behaviors. *Int J Behav Nutr Phys Act.* 2016;13(1):103.
10. Wang C, Pan R, Wan X, Tan Y, Xu L, Ho CS, et al. Immediate Psychological Responses and Associated Factors during the Initial Stage of the 2019 Coronavirus Disease (COVID-19) Epidemic among the General Population in China. *Int J Environ Res Public Health.* 2020;17(5).
11. Evers C, Dingemans A, Junghans AF, Boevé A. Feeling bad or feeling good, does emotion affect your consumption of food? A meta-analysis of the experimental evidence. *Neurosci Biobehav Rev.* 2018;92:195-208.
12. Butler MJ, Barrientos RM. The impact of nutrition on COVID-19 susceptibility and long-term consequences. *Brain Behav Immun.* 2020;87:53-4.
13. Garfield V, Llewellyn CH, Kumari M. The relationship between physical activity, sleep duration and depressive symptoms in older adults: The English Longitudinal Study of Ageing (ELSA). *Preventive medicine reports.* 2016;4:512-6.
14. Feng Q, Zhang QL, Du Y, Ye YL, He QQ. Associations of physical activity, screen time with depression, anxiety and sleep quality among Chinese college freshmen. *PLoS One.* 2014;9(6):e100914.
15. Reyes-Olavarría D, Latorre-Román P, Guzmán-Guzmán IP, Jerez-Mayorga D, Caamaño-Navarrete F, Delgado-Floody P. Positive and Negative Changes in Food Habits, Physical Activity Patterns, and Weight Status during COVID-19 Confinement: Associated Factors in the Chilean Population. *Int J Environ Res Public Health.* 2020;17(15).
16. Lazarevich I, Irigoyen Camacho ME, Velázquez-Alva MC, Flores NL, Nájera Medina O, Zepeda Zepeda MA. Depression and food consumption in Mexican college students. *Nutr Hosp.* 2018;35(3):620-6.

17. Munayco CV, Tariq A, Rothenberg R, Soto-Cabezas GG, Reyes MF, Valle A, et al. Early transmission dynamics of COVID-19 in a southern hemisphere setting: Lima-Peru: February 29(th)-March 30(th), 2020. medRxiv. 2020.
18. Vázquez-Rowe I, Gandolfi A. Peruvian efforts to contain COVID-19 fail to protect vulnerable population groups. *Public Health in Practice*. 2020:100020.
19. Ahmed HO. The impact of social distancing and self-isolation in the last corona COVID-19 outbreak on the body weight in Sulaimani governorate- Kurdistan/Iraq, a prospective case series study. *Ann Med Surg (Lond)*. 2020;59:110-7.
20. Haddad C, Zakhour M, Bou Kheir M, Haddad R, Al Hachach M, Sacre H, et al. Association between eating behavior and quarantine/confinement stressors during the coronavirus disease 2019 outbreak. *J Eat Disord*. 2020;8:40.
21. Robinson E, Boyland E, Chisholm A, Harrold J, Maloney NG, Marty L, et al. Obesity, eating behavior and physical activity during COVID-19 lockdown: A study of UK adults. *Appetite*. 2021;156:104853.
22. Keel PK, Gomez MM, Harris L, Kennedy GA, Ribeiro J, Joiner TE. Gaining "The Quarantine 15:" Perceived versus observed weight changes in college students in the wake of COVID-19. *Int J Eat Disord*. 2020;53(11):1801-8.
23. Woolcott OO, Gutierrez C, Castillo OA, Elashoff RM, Stefanovski D, Bergman RN. Inverse association between altitude and obesity: A prevalence study among andean and low-altitude adult individuals of Peru. *Obesity (Silver Spring)*. 2016;24(4):929-37.
24. Brooks SK, Webster RK, Smith LE, Woodland L, Wessely S, Greenberg N, et al. The psychological impact of quarantine and how to reduce it: rapid review of the evidence. *Lancet*. 2020;395(10227):912-20.

25. Huang R, Zhu L, Xue L, Liu L, Yan X, Wang J, et al. Clinical findings of patients with coronavirus disease 2019 in Jiangsu province, China: A retrospective, multi-center study. *PLoS Negl Trop Dis*. 2020;14(5):e0008280.
26. Peng YD, Meng K, Guan HQ, Leng L, Zhu RR, Wang BY, et al. [Clinical characteristics and outcomes of 112 cardiovascular disease patients infected by 2019-nCoV]. *Zhonghua Xin Xue Guan Bing Za Zhi*. 2020;48(6):450-5.
27. Pettit NN, MacKenzie EL, Ridgway JP, Pursell K, Ash D, Patel B, et al. Obesity is Associated with Increased Risk for Mortality Among Hospitalized Patients with COVID-19. *Obesity (Silver Spring)*. 2020;28(10):1806-10.
28. Simonnet A, Chetboun M, Poissy J, Raverdy V, Noulette J, Duhamel A, et al. High Prevalence of Obesity in Severe Acute Respiratory Syndrome Coronavirus-2 (SARS-CoV-2) Requiring Invasive Mechanical Ventilation. *Obesity (Silver Spring)*. 2020;28(7):1195-9.
29. Ferrari GLdM, Kovalskys I, Fisberg M, Gómez G, Rigotti A, Sanabria LYC, et al. Socio-demographic patterning of objectively measured physical activity and sedentary behaviours in eight Latin American countries: findings from the ELANS study. *European journal of sport science*. 2020;20(5):670-81.
30. Carriedo A, Cecchini JA, Fernández-Río J, Méndez-Giménez A. Resilience and physical activity in people under home isolation due to COVID-19: A preliminary evaluation. *Ment Health Phys Act*. 2020;19:100361.
31. Maugeri G, Castrogiovanni P, Battaglia G, Pippi R, D'Agata V, Palma A, et al. The impact of physical activity on psychological health during Covid-19 pandemic in Italy. *Heliyon*. 2020;6(6):e04315.
32. Rajkumar RP. Sleep, physical activity and mental health during the COVID-19 pandemic: complexities and opportunities for intervention. *Sleep Medicine*. 2020.

33. Woods J, Hutchinson NT, Powers SK, Roberts WO, Gomez-Cabrera MC, Radak Z, et al. The COVID-19 pandemic and physical activity. Elsevier; 2020.
34. Arnett DK, Blumenthal RS, Albert MA, Buroker AB, Goldberger ZD, Hahn EJ, et al. 2019 ACC/AHA Guideline on the Primary Prevention of Cardiovascular Disease: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *J Am Coll Cardiol.* 2019;74(10):e177-e232.
35. Pinto J, van Zeller M, Amorim P, Pimentel A, Dantas P, Eusébio E, et al. Sleep quality in times of Covid-19 pandemic. *Sleep Med.* 2020;74:81-5.
36. Jahrami H, BaHammam AS, AlGahtani H, Ebrahim A, Faris M, AlEid K, et al. The examination of sleep quality for frontline healthcare workers during the outbreak of COVID-19. *Sleep Breath.* 2020:1-9.
37. Huang Y, Zhao N. Generalized anxiety disorder, depressive symptoms and sleep quality during COVID-19 outbreak in China: a web-based cross-sectional survey. *Psychiatry Res.* 2020;288:112954.
38. Morin CM, Carrier J, Bastien C, Godbout R. Sleep and circadian rhythm in response to the COVID-19 pandemic. *Can J Public Health.* 2020;111(5):654-7.
39. Chopra S, Ranjan P, Singh V, Kumar S, Arora M, Hasan MS, et al. Impact of COVID-19 on lifestyle-related behaviours-a cross-sectional audit of responses from nine hundred and ninety-five participants from India. *Diabetes & Metabolic Syndrome: Clinical Research & Reviews.* 2020;14(6):2021-30.
40. Zachary Z, Brianna F, Brianna L, Garrett P, Jade W, Alyssa D, et al. Self-quarantine and weight gain related risk factors during the COVID-19 pandemic. *Obes Res Clin Pract.* 2020;14(3):210-6.

41. Yan AF, Sun X, Zheng J, Mi B, Zuo H, Ruan G, et al. Perceived risk, behavior changes and Health-related outcomes during COVID-19 pandemic: Findings among adults with and without diabetes in China. *Diabetes Res Clin Pract.* 2020;167:108350.
42. Jayawardena R, Misra A. Balanced diet is a major casualty in COVID-19. *Diabetes Metab Syndr.* 2020;14(5):1085-6.

Table 1. Sociodemographic and anthropometric data.

Variables	Total (n =1176)	Men (n =571)	Women (n =605)	p-value	
Age					
18-25 yrs	522 (44.4)	197 (34.5)	325 (53.7)	<0.001*	
26-40 yrs	396 (33.7)	213 (37.3)	183 (30.2)		
41-55 yrs	200 (17.0)	115 (20.1)	85 (14.0)		
56-70 yrs	58 (4.9)	46 (8.1)	12 (2.0)		
Marital status					
Single	796 (67.7)	339 (59.4)	457 (75.5)	<0.001*	
Married	333 (28.3)	205 (35.9)	128 (21.2)		
Separated or divorced	42 (3.6)	25 (4.4)	17 (2.8)		
Widow(er)	5 (0.4)	2 (0.4)	3 (0.5)		
Level of education					
Primary incomplete	4 (0.3)	4 (0.7)	0 (0.0)	0.004*	
Primary completed	9 (0.8)	4 (0.7)	5 (0.8)		
Secondary completed	342 (29.1)	143 (25.0)	199 (32.9)		
Preparatory or tertiary completed	199 (16.9)	89 (15.6)	110 (18.2)		
University degree completed	447 (38.0)	237 (41.5)	210 (34.7)		
Graduate degree completed	175 (14.9)	94 (16.5)	81 (13.4)		
Occupational status					
Student	459 (39.0)	159 (27.8)	300 (49.6)		<0.001*
Worker	441 (37.5)	272 (47.6)	169 (27.9)		
Worker and student	194 (16.5)	110 (19.3)	84 (13.9)		
Housewife	19 (1.6)	1 (0.2)	18 (3.0)		
Unemployed	58 (4.9)	25 (4.4)	33 (5.5)		
Retired	5 (0.4)	4 (0.7)	1 (0.2)		
Employment status					
I am not working	541 (46.0)	189 (33.1)	352 (58.2)	<0.001*	
Formal job with a contract	362 (30.8)	201 (35.2)	161 (26.6)		
Self-employed	159 (13.5)	114 (20.0)	45 (7.4)		
Public sector	114 (9.7)	67 (11.7)	47 (7.8)		
Days under social isolation					
Have not socially isolated	143 (12.2)	68 (11.9)	75 (12.4)	0.019*	
Less than 15 days	37 (3.1)	16 (2.8)	21 (3.5)		
15 to 30 days	59 (5.0)	36 (6.3)	23 (3.8)		
31 to 45 days	41 (3.5)	25 (4.4)	16 (2.6)		
45 to 60 days	101 (8.6)	60 (10.5)	41 (6.8)		
More than 61 days	795 (67.6)	366 (64.1)	429 (70.9)		
People in the home					
Before	4.24 (2.45)	4.20 (2.51)	4.28 (2.39)	0.578	
During	4.35 (2.35)	4.26 (2.27)	4.45 (2.42)	0.165	
Body Mass Index					
Malnourished	18 (1.5)	5 (0.9)	13 (2.1)	<0.001*	
Normal	639 (54.3)	245 (42.9)	394 (65.1)		
Overweight	420 (35.7)	258 (45.2)	162 (26.8)		
Obesity I	84 (7.1)	52 (9.1)	32 (5.3)		

Obesity II	15 (1.3)	11 (1.9)	4 (0.7)	
Weight change during social isolation (gained)				
1 - 3 kg	245 (20.8)	105 (18.4)	140 (23.1)	0.001*
4 - 7 kg	161 (13.7)	87 (15.2)	74 (12.2)	
8 - 11 kg	41 (3.5)	28 (4.9)	13 (2.1)	
>12 kg	14 (1.2)	6 (1.1)	8 (1.3)	
Yes, but unsure how much	185 (15.7)	72 (12.6)	113 (18.7)	
No	530 (45.1)	273 (47.8)	257 (42.5)	
COVID-19 diagnosis confirmed				
No	1074 (91.3)	521 (91.2)	553 (91.4)	1
Yes	102 (8.7)	50 (8.8)	52 (8.6)	

Data expressed as absolute and relative frequency (%). COVID-19, Coronavirus disease; yrs, years.

*Statistically significant, $p < 0.05$.

Table 2. Lifestyles during social isolation.

Variables	Total (n =1176)	Men (n =571)	Women (n =605)	p-value
Physical activity				
No	532 (45.2)	258 (45.2)	274 (45.3)	1
Yes	644 (54.8)	313 (54.8)	331 (54.7)	
Days per week				
None	532 (45.2)	258 (45.2)	274 (45.3)	0.204
1 to 2 days	173 (14.7)	76 (13.3)	97 (16.0)	
3 to 4 days	310 (26.4)	147 (25.7)	163 (26.9)	
5 to 6 days	131 (11.1)	71 (12.4)	60 (9.9)	
7 days	30 (2.6)	19 (3.3)	11 (1.8)	
Duration				
No physical activity	532 (45.2)	258 (45.2)	274 (45.3)	0.425
Less than 30 minutes	257 (21.9)	122 (21.4)	135 (22.3)	
30 minutes to 1 hour	335 (28.5)	159 (27.8)	176 (29.1)	
1 to 2 hours	46 (3.9)	28 (4.9)	18 (3.0)	
Other	6 (0.5)	4 (0.7)	2 (0.3)	
Sleep routine				
No change	304 (25.9)	171 (29.9)	133 (22.0)	0.006*
Sleep more	434 (36.9)	204 (35.7)	230 (38.0)	
Sleep less	438 (37.2)	196 (34.3)	242 (40.0)	
Hours of sleep				
0 to 3 hours	25 (2.1)	12 (2.1)	13 (2.1)	0.170
4 to 6 hours	419 (35.6)	188 (32.9)	231 (38.2)	
7 to 9 hours	651 (55.4)	335 (58.7)	316 (52.2)	
>10 hours	81 (6.9)	36 (6.3)	45 (7.4)	

Data expressed as absolute and relative frequency (%). *Statistically significant, $p < 0.05$




Figure 1. Frequency of eating patterns during social isolation.




Figure 2. Changes in the frequency of low and high consumption of plant-based foods before and during social isolation. *Before social isolation, **During social isolation.