

1 Respiratory cryptosporidiosis in Malawian children with diarrheal disease

2

3 Pui-Ying Iroh Tam,^{1,2,3a} Mphatso Chisala,¹ Wongani Nyangulu,^{1b} Herbert Thole,¹ James

4 Nyirenda¹

5

6 ¹ Paediatrics and Child Health Research Group, Malawi-Liverpool Wellcome Trust Clinical

7 Research Programme, Blantyre, Malawi

8 ² Department of Paediatrics, University of Malawi College of Medicine, Blantyre, Malawi

9 ³ Department of Clinical Sciences, Liverpool School of Tropical Medicine, Liverpool, United

10 Kingdom

11 ^a Corresponding author: Pui-Ying Iroh Tam, irohtam@mlw.mw

12 ^b Current affiliation: University of Malawi College of Medicine, Blantyre, Malawi

13

14 Running title: Respiratory cryptosporidium in diarrhea

15

16 Keywords: cryptosporidiosis, respiratory tract, gastrointestinal, children, longitudinal

17

18

19

20

21

22

23

24

25

26 **Abstract**

27 Background: Respiratory cryptosporidiosis has been documented in children with diarrhea.

28 We sought to describe the dynamics of respiratory involvement in children hospitalized with
29 gastrointestinal (GI) diarrheal disease.

30

31 Methods: We conducted a prospective, observational longitudinal study of Malawian children
32 2-24 months hospitalized with diarrhea. Nasopharyngeal (NP) swabs, induced sputum and
33 stool specimens were collected. Participants that were positive by cryptosporidium PCR in
34 any of the three compartments were followed up with fortnightly visits up to 8 weeks post-
35 enrollment.

36

37 Results: Of the 162 children recruited, participants had mild-moderate malnutrition, 36 (22%)
38 were PCR-positive for cryptosporidium at enrollment (34 stool, 11 sputum, and 4 NP) and 27
39 completed the majority of follow-up visits (75%). Cryptosporidium was detected in all
40 compartments over the 4 post-enrollment visits, most commonly in stool (94% at enrollment
41 with mean cycle thresholds (Ct) of 28.8 ± 4.3 to 44% at 8 weeks with Ct 29.9 ± 4.1), followed
42 by sputum (31% at enrollment with mean Ct 31.1 ± 4.4 to 20% at 8 weeks with Ct 35.7 ± 2.6),
43 then NP (11% with mean Ct 33.5 ± 1.0 to 8% with Ct 36.6 ± 0.7). Participants with
44 cryptosporidium detection in both the respiratory and GI tract over the study period reported
45 respiratory and GI symptoms in 81% and 62% of study visits, respectively, compared to 68%
46 and 27%, respectively, for those with only GI detection, and had longer GI shedding
47 (17.5 ± 6.6 v. 15.9 ± 2.9 days).

48

49 Conclusion: Cryptosporidium was detected in both respiratory and GI tracts throughout the 8
50 weeks post-enrollment. The development of therapeutics for cryptosporidium in children
51 should target the respiratory as well as GI tract.

52 (255)

53

54

55

56 **Author summary**

57 We conducted a prospective, observational longitudinal study of Malawian children 2-24
58 months hospitalized with diarrhea. Nasopharyngeal (NP) swabs, induced sputum and stool
59 specimens were collected. Participants that were positive by cryptosporidium PCR in any of
60 the three compartments were followed up with fortnightly visits up to 8 weeks post-
61 enrollment. Cryptosporidium was detected by PCR in 21%, 7% and 3% in stool, sputum and
62 nasopharynx of children hospitalized with diarrhea. Of those positive at enrollment, detection
63 was noted in 44%, 20%, and 8%, respectively, by 8 weeks post-enrollment.

64 (87)

65

66

67

68

69

70

71

72

73

74 Introduction

75 Cryptosporidiosis is a cause of diarrhea [1-3], excess mortality [4, 5], stunting [1, 6], and is
 76 associated with malnutrition [6, 7]. The Global Enteric Multicenter Study (GEMS) identified
 77 cryptosporidium as second most common cause of diarrhea among infants (0-11 months) in
 78 all four African countries studied (The Gambia, Mali, Mozambique, Kenya) regardless of
 79 HIV prevalence, and among the top five causes for older children (12-23 months) [4].
 80 Respiratory cryptosporidiosis has been documented in up to a third of children presenting
 81 with diarrhea [8]; furthermore, respiratory detection without intestinal involvement has been
 82 reported [9, 10], raising the possibility of primary respiratory infection with cryptosporidium,
 83 either by inhalation or by contact with fomites [11].

84
 85 Studies have identified cryptosporidium in sputum but have not looked at whether respiratory
 86 involvement is a transient phenomenon or a reservoir for gastrointestinal (GI) disease. This
 87 has implications for therapeutic development. Our primary objective was to evaluate whether
 88 respiratory involvement of cryptosporidium is a transient phenomenon in diarrheal disease,
 89 and to assess for respiratory and GI cryptosporidiosis concurrently and longitudinally in
 90 children hospitalized with diarrheal disease.

91

92 Methods

93 *Study design, setting, and participants.* We conducted a prospective, observational-
 94 longitudinal study of Malawian children hospitalized with diarrhea [12]. Children aged 2-24
 95 months and presenting with primary GI symptoms to Queen Elizabeth Central Hospital in
 96 Blantyre, Malawi, were screened. Eligible patients were those with at least three or more
 97 loose stools within the past 24 hours. Those with dysentery, or visible blood in loose stools,
 98 were excluded.

99

Clinical procedures. Once written consent was obtained, a detailed history and physical exam was conducted and all then enrolled subjects had a NP swab, induced sputum, and stool sample collected. For induced sputum collection, subjects were given a nebulized 3% sodium chloride mist to inhale for 5-15 minutes. For infants, suctioning of the oropharynx was done after nebulization to collect the specimen. To ensure safety of the subject, induced sputum was only collected if there were no contraindications (based on PERCH criteria for induced sputum collection) [13]: severe hypoxia <92% on supplemental oxygen; inability to protect airways; severe bronchospasm at admission (defined as continued hypoxia <92% after appropriate bronchodilator therapy, with other markers of respiratory distress); seizure within the past 24 hours; or deemed inappropriate by the clinician for another reason (e.g. midface trauma, inhalational injury, pulmonary effusion, congestive heart failure, congenital heart disease, etc). If the above symptoms/conditions resolved within 48 hours, induced sputum collection was reconsidered at that point.

HIV rapid testing was conducted on infants and, if positive, infants were referred to the pediatric HIV clinic for further care. Any relevant lab investigations (including full blood count, urea and electrolytes, liver function tests, malaria parasite screen, blood and/or CSF culture, TB GeneXpert and AFB testing) performed as part of routine care were recorded.

All participating children had enrollment NP, sputum and stool specimens evaluated for cryptosporidium by PCR. Sputum quality was evaluated by microscopy (good quality: ≤ 10 squamous epithelial cells/high powered field). Only subjects who were cryptosporidium PCR-positive on any one of NP/sputum/stool were followed up in the post-enrollment phase

up to 8 weeks post-enrollment, with a follow-up visit every two weeks for evaluation of symptoms, physical assessment, as well as NP/sputum/stool sampling.

Laboratory procedures. DNA was extracted in stool using QIAamp Fast Stool Mini Kit (Qiagen, Hilden, Germany) and for respiratory tract using the QIAamp DNA mini kit (Qiagen) for cryptosporidium, and UCP Mini kit (Qiagen) for respiratory co-pathogens. Cryptosporidium detection of specimens at enrollment were measured using quantitative polymerase chain reaction (qPCR), with appropriate positive and negative controls and a positive result being a cycle threshold (Ct) <35. In subsequent follow-up samples in stool, cryptosporidium was detected using qPCR in a TaqMan Array Card (Thermo Fisher, Waltham, MA) using a custom design developed at the Hout Laboratory (Charlottesville, VA) [14] and tested for 28 enteropathogens. In sputum, for multiplex PCR we used the respiratory pathogens 33 kit (Fast Track Diagnostics, Luxembourg, Luxembourg).

Statistical analysis. Frequencies and proportions of observed levels were reported for binary and categorical variables, with exact binomial 95% confidence intervals given where appropriate. Comparisons were performed using the Fisher's exact test for binary and categorical variables, t-test (two groups) or ANOVA (three or more groups) for approximately normally distributed variables and Wilcoxon rank sum (2 groups) or Kruskal-Wallis (3 or more groups) tests for variables with severely nonparametric distributions. Statistical analysis was performed using Stata software, version 16, and statistical significance was set at 0.05.

Ethical approval. The study was approved by the University of Malawi College of Medicine Research Ethics Committee (P.07/18/2438) and the Liverpool School of Tropical Medicine

Research Ethics Committee (18-066). Written consent was obtained from parents or guardians prior to enrollment.

Results

From 1 March 2019 to 3 April 2020, 162 children were recruited into the study. Two children did not submit specimens. Of the remaining 160 children, 36 (23%) were positive in any one of the three samples collected (34 stool (21%), 11 sputum (7%), and 4 NP (3%)) and were entered into follow-up. The study was discontinued early due to COVID-19, by which time 27 children had completed at least three of the four follow-up visits (75%).

Among the 162 children who were hospitalized with diarrhea, the mean age was 11.6 ± 5.0 months, and 59% were male (Table 1). Mean height-for-age, weight-for-age, and weight-for-height z scores were -1.41 ± 1.5 , -1.02 ± 1.2 , and -0.08 ± 1.6 , respectively. Nine percent of mothers were HIV-infected, and one-third of those children tested positive by HIV rapid test (3% of total). The participants had a mean of 4.5 ± 1.4 persons in a household, and the majority of children had access to a pit toilet/latrine (95%), piped drinking water (78%), and harbored a residential animal in the compound (66%). Among the 36 children who were cryptosporidium PCR-positive, the mean age was 12 ± 5.0 months, 63% were male and two (6%) were positive by HIV rapid test. No significant differences in characteristics between children who were cryptosporidium-positive versus -negative were noted (Supplementary Table 1).

Table 1. Characteristics of children hospitalized with diarrhea at enrollment

Characteristic	Children with diarrhea (n=162)	Cryptosporidium-positive (n=36)	Cryptosporidium-negative (n=124)	P value
----------------	--------------------------------	---------------------------------	----------------------------------	---------

Demographics				
Age, ^a months, mean (SD)	11.6 (5.0)	12 (5.0)	11.6 (5.1)	0.897
2-11 months	90 (56%)	21 (60%)	65 (53%)	0.565
12-24 months	72 (44%)	14 (40%)	58 (47%)	
Male sex	95 (59%)	22 (63%)	72 (59%)	0.649
Highest level of maternal education				
Did not complete any schooling	62 (38%)	14 (40%)	46 (37%)	0.829
Completed primary school	74 (46%)	15 (43%)	57 (46%)	
Completed secondary school and above	18 (11%)	5 (14%)	13 (11%)	
Mother is employed (%)	72 (44%)	19 (54%)	52 (42%)	0.208
Father is employed (%)	150 (93%)	33 (94%)	114 (92%)	0.742
Number of household members (SD)	4.5 (1.4)	4.1 (1.2)	4.6 (1.4)	0.061
Children <5 years (SD)	1.2 (0.5)	1.2 (0.4)	1.2 (0.6)	0.107
Other household members with respiratory symptoms	46 (28%)	13 (37%)	33 (27%)	0.236
Other household members with GI symptoms	13 (8%)	4 (11%)	9 (7%)	0.487
Residential animals in the compound (%)	107 (66%)	21 (20%)	84 (68%)	0.359
Facility for disposing feces in households (%)				
Pit toilet/latrine	154 (95%)	34 (97%)	116 (94%)	0.686

Pour/flush toilet	8 (5%)	1 (2%)	7 (60%)	
Child health indicators				
Mother HIV status (%)				
Positive	15 (9%)	8 (23%)	17 (14%)	0.391
Unknown	55 (34%)	10 (29%)	44 (36%)	
Child HIV rapid test (%)				
Positive	5 (3%)	2 (6%)	3 (2%)	0.158
Unknown/not done	120 (74%)	22 (63%)	95 (77%)	
Breastfeeding status (%)				
No breastfeeding	22 (14%)	5 (14%)	17 (14%)	1.000
Partial breastfeeding	134 (83%)	29 (82%)	101 (82%)	
Exclusive breastfeeding	6 (4%)	1 (3%)	5 (4%)	
Used antibiotics in 7 days before admission (%)	99 (61%)	20 (57%)	75 (61%)	0.699
Given intravenous fluids in 7 days before admission (%)	63 (61%)	15 (43%)	47 (38%)	0.696
Referred to a health facility for a diarrhea episode in the past 7 days (%)	136 (84%)	32 (91%)	101 (82%)	0.292
Z score (SD)				
Height-for-age	-1.45 (1.5)	-1.41 (1.5)	-1.46 (1.5)	0.807
Weight-for-age	-1.0 (1.3)	-1.02 (1.2)	-1.02 (1.3)	0.997
Weight-for-height	-0.08 (1.6)	-0.1 (1.6)	-0.12 (1.6)	0.952
MUAC, cm (SD)	13.7 (1.4)	13.8 (1.3)	13.7 (1.4)	0.692
Bipedal edema (%)	5 (3%)	1 (3%)	4 (3%)	1.000

Enrollment vitals and symptoms				
Temperature, °C (SD)	36.7 (0.7)	36.7 (0.6)	36.8 (0.7)	0.422
Respiratory rate, breaths/min (SD)	36 (6.1)	36.1 (8.5)	36 (5.4)	0.902
Oxygen saturations, % (SD)	98.6 (1.4)	98.6 (1.4)	98.7 (1.4)	0.719
GI symptoms (%)				
Vomiting	70 (43%)	16 (46%)	52 (42%)	0.847
Abdominal pain/ tenderness	65 (40%)	11 (31%)	51 (42%)	0.330
Poor feeding	66 (40%)	14 (40%)	50 (40%)	0.945
Dehydration	90 (56%)	20 (57%)	68 (55%)	0.845
Respiratory symptoms (%)				
Cough	98 (61%)	24 (69%)	72 (59%)	0.283
Runny nose	58 (36%)	16 (46%)	40 (33%)	0.150
Difficulty in breathing	3 (2%)	1 (3%)	2 (2%)	0.531
Wheezing	14 (9%)	4 (11%)	9 (7%)	0.487
Chest indrawings/retractions	2 (1%)	0	2 (2%)	1.000
Crackles	17 (11%)	6 (17%)	10 (8%)	0.125
Cryptosporidium-PCR positive				
Stool (%)	34 (21%)	34 (94%)	0	<0.001
Induced sputum (%)	11 (7%)	11 (31%)	0	<0.001
NP (%)	4 (2%)	4 (11%)	0	<0.001
Sputum quality (%)				
Good (≤10 sq epis/hpf)	131/160 (82%)	29 (81%)	100 (82%)	0.013
Poor (>10 sq epis/hpf)	15/160 (9%)	1 (3%)	14 (12%)	
Unreadable ^c	14/160 (9%)	6 (17%)	8 (7%)	

Outcomes

Mortality, inpatient (%)	NA	0	NA
Mortality, post-discharge (%)	NA	1 (3%)	NA

GI, gastrointestinal; HPF, high powered field; MUAC, mid-upper arm circumference; NA, not applicable; SD, standard deviation

^aValues are mean (SD).

Induced sputum quality at enrollment was good in 81%, poor in 9% and unreadable in 10%, with similar proportions of good quality sputum among cryptosporidium-positive and -negative participants (81% and 82%, respectively). Over the 8-week period, cryptosporidium was detected in both respiratory and GI tracts (Table 2): most commonly in stool (94% at enrollment to 44% at 8 weeks), followed by sputum (31% at enrollment to 20% at 8 weeks), then NP (11% to 8%). Lowest Ct counts were generally noted in the stool (mean Ct 28.8±4.3 at enrollment to 29.9±4.1 at 8 weeks), followed by sputum (mean Ct 31.1± 4.4 at enrollment to 35.7±2.6 at 8 weeks), and NP ((mean Ct 33.5±1.0 to 36.6±0.7); Figure 1). We detected cryptosporidium in respiratory but not GI tract in 7 study visits, including 2 at enrollment (stool Ct were above the 35 cutoff and therefore considered negative). Detection in the respiratory tract did not always correlate with symptoms (Figures 2A-O). For individual participants in whom cryptosporidium was detected at enrollment, detection was noted most consistently in the GI tract throughout the 8 weeks. Respiratory co-pathogens were detected in 87/104 (84%) visits over the follow-up period.

Table 2. Longitudinal detection of cryptosporidium in stool and respiratory tract and associated symptoms

Characteristics	Study period				
	Enrollment (n=36)	2 weeks ^a (n=27)	4 weeks ^b (n=24)	6 weeks ^c (n=26)	8 weeks ^d (n=25)
Cryptosporidium detection					
Detection in NP	4 (11%)	5 (19%)	2 (8%)	1 (4%)	2 (8%)
NP Ct values ^e (SD)	33.5 (1.0)	33.7 (2.0)	35.6 (0.6)	35.7	36.6 (0.7)
Detection in sputum	11 (31%)	10 (37%)	10 (42%)	8 (31%)	5 (20%)
Sputum Ct values (SD)	31.1 (4.4)	29.9(3.6)	33.0 (2.6)	34.3 (2.9)	35.7 (2.6)
Detection in GI tract	34 (94%)	22 (81%)	16 (67%)	11 (42%)	11 (44%)
Stool Ct values (SD)	28.8 (4.3)	26.0 (4.5)	30.3 (2.8)	31.3 (3.5)	29.9 (4.1)
Detection in respiratory tract only	2 (6%)	0	1 (4%)	2 (8%)	2 (8%)
Detection in GI tract only	26 (72%)	13 (48%)	6 (25%)	3 (12%)	6 (24%)
Detection in respiratory and GI tract	9 (25%)	9 (33%)	9 (26%)	6 (23%)	3 (12%)
Presentation					
Referral to health facility for an illness in the past 7 days	36 (100%)	10 (37%)	6 (25%)	5 (19%)	7 (28%)
Diarrhea in the past 7 days	36 (100%)	0	2 (8%)	4 (15%)	3 (12%)
Respiratory symptoms in the past 7 days ^f	26 (72%)	9/10 (90%)	4/6 (67%)	2/5(40%)	5/7 (71%)
Cough	24 (69%)	5/10 (50%)	2/6 (33%)	2/5 (40%)	5/7 (71%)

Runny nose	16 (46%)	7/10 (70%)	4/6 (67%)	1/5 (20%)	3/7 (43%)
Difficulty in breathing	1 (3%)	0	0	0	0
Wheezing	4 (11%)	3/10 (30%)	1/6 (17%)	1/5 (20%)	0
Chest indrawings/ retractions	0	0	0	0	0
Crackles	6 (17%)	5/10 (50%)	1/6 (17%)	1/5 (20%)	0/7
Vomiting	16 (46%)	0/10	0/6	1/5 (20%)	0/7
Abdominal pain/ tenderness	11 (31%)	1/10 (10%)	1/6 (17%)	1/5 (20%)	0/7
Poor feeding	14 (40%)	0	0	0	0
Other household members with respiratory symptoms	13 (37%)	1/6 (17%)	3/5 (60%)	0	0
Other household members with GI symptoms	4 (11%)	0	3 (13%)	1 (4%)	2 (8%)
Vital signs					
Respiratory rate, breaths/min (SD)	36 (6.1)	34.7 (3.4)	32.8 (3.2)	34.4 (5.7)	32.3 (3.7)
Oxygen saturations, % (SD)	98.6 (1.4)	99.1 (1.2)	98.5 (1.1)	99.4 (0.8)	98.8 (1.4)
Outcome					
Mortality	1 ^g (3%)	0	0	0	0

194 Ct, cycle threshold; GI, gastrointestinal; NP, nasopharynx; SD, standard deviation

195 ^aOne child withdrew after enrollment, five did not enter follow-up due to early stopping of
196 study, and three were lost to follow-up.

197 ^bThree children missed week 4 visit.

^cOne child missed week 6 visit.

^dTwo children did not complete week 8 visit due to early stopping of study.

^eValues are mean (SD).

^fIn follow-up study visits, symptoms were recorded only if these led to health facility referral.

^gPatient died after discharge before the week 2 visit.

Respiratory symptoms were noted in 72% of cryptosporidium-positive participants at enrollment, the most common symptom being cough (69%), but also included runny nose (46%), crackles (17%) and wheeze (11%). For those entered into the 8-week follow-up period, participants with cryptosporidium positivity in the respiratory tract had respiratory symptoms in 23/43 (53%) of visits.

Among participants with cryptosporidium detection in both the respiratory and GI tract at enrollment (Table 3A) compared to those with cryptosporidium detection in the GI tract only, a larger proportion reported respiratory symptoms (90% v. 65%), and GI shedding of cryptosporidium was longer (14.3 ± 2.1 v. 14.1 ± 0.7 days), but these were not statistically significant. Among participants whom cryptosporidium was detected in both the respiratory and GI tract over the 8-week study period compared to GI tract only (Table 3B), a larger proportion reported respiratory (81% v. 68%) and GI symptoms (62% v. 27%) per study visit, and GI shedding of cryptosporidium was longer (17.5 ± 6.6 v. 15.9 ± 2.9 days), but again these were not statistically significant.

Table 3. Characteristics and associated symptoms in participants with cryptosporidium detection in GI tract only v. GI and respiratory tract A) at enrollment; B) throughout study period

Characteristic	At enrollment			Throughout study period		
	GI detection only (n=26)	GI and respiratory detection (n=10)	P value	GI detection only (n=16)	GI and respiratory detection (n=11)	P value
Age, ^a months (SD)	11.3 (5.3)	13 (4.0)	0.370	12.2 (2.3)	12.0 (6.7)	0.922
Male sex	17 (65%)	6 (60%)	0.7664	9 (56%)	8 (72%)	0.448
Z score (SD)						
Height-for-age	-1.4 (2.4)	-1.3 (1.1)	0.896	-1.6 (2.5)	-1.4 (1.9)	0.837
Weight-for-age	-1.1 (1.4)	-0.6 (1.2)	0.319	-1.1 (1.2)	-0.8 (1.2)	0.474
Weight-for-height	-0.2 (1.6)	0.0 (1.7)	0.708	-0.1 (1.7)	0.0 (1.6)	0.945
Number of household members (SD)	4.2 (1.4)	4.3 (1.5)	0.771	4.4 (1.5)	4.1 (0.9)	0.504
Household children <5 years	1.3 (0.7)	1.1 (0.3)	0.449	1.1 (0.1)	1.2 (0.1)	0.696
Pit latrine	25 (96%)	10 (100%)	0.529	16 (100%)	10 (90%)	0.407
Piped water for drinking	4 (15%)	1 (10%)	0.916	2 (13%)	3 (27%)	0.370
Piped water for utensils	18 (69%)	6 (60%)	0.448	13 (81%)	4 (45%)	0.124

Treated water (%)	13 (50%)	2 (20%)	0.142	6 (37%)	5 (45%)	0.710
Residential animals in the compound	14 (54%)	7 (70%)	0.467	11 (69%)	7 (63%)	1.000
Respiratory symptoms	17 (65%)	9 (90%)	0.223	11 (68%)	9 (81%)	0.385
Enrollment oxygen sats, % (SD)	98.5 (1.4)	98.8 (1.2)	0.561	98.5% (1.1)	98.3% (1.7)	0.653
GI symptoms	23 (89%)	8 (80%)	0.603	3 (27%)	10 (62%)	0.120
Duration of GI shedding, ^b days (SD)	14.1 (0.7)	14.3 (2.1)	0.819	15.9 (2.9)	17.5 (6.6)	0.445

224 GI, gastrointestinal; SD, standard deviation

225 ^aValues are mean (SD).

226 ^bN changes to 27 who followed up through study period.

227

228

229 Discussion

230 This is the first longitudinal study to evaluate the respiratory cryptosporidiosis in pediatric
231 diarrheal disease. In children hospitalized with diarrheal disease, cryptosporidium was
232 detected most frequently in stool, followed by sputum and NP. The most common respiratory
233 symptom was cough. Longitudinally, we detected cryptosporidium in both respiratory and GI
234 tracts through 8 weeks post-enrollment, usually at lowest Ct counts in stool, followed by
235 sputum and then NP. Longer GI shedding of cryptosporidium was noted among those where
236 we detected cryptosporidium in both the respiratory and GI tracts, compared to GI detection
237 only, although findings did not reach statistical significance.

238

239 Our cryptosporidium detection rate is far higher than has been reported in diarrhea studies in
240 sub-Saharan Africa, with rates of positivity ranging from 9% in Kenya to 14.7% in
241 Mozambique [4, 8]. Subclinical cryptosporidium infection was noted to be high (6%) in
242 Tanzania [15]. A previous study done at QECH found cryptosporidium to be the third most
243 common cause of diarrhea with a prevalence in stool of 28% among cases [16]. In our study,
244 the positivity rate was 21%, and this would be 12% if we used the GEMS diarrheagenic
245 cutoffs [17]. This thus supports the high prevalence of cryptosporidium infection among
246 young children hospitalized with diarrheal illness in sub-Saharan Africa.

247

248 Few studies, however, have evaluated the prevalence of respiratory cryptosporidiosis in
249 pediatric diarrheal disease. A study in Uganda detected cryptosporidium in 35.4% of induced
250 sputum samples from children presenting with diarrhea who were cryptosporidium-positive
251 in the stool [8]. In our study, cough was noted in almost two-thirds of children presenting
252 with diarrhea, and the frequency of respiratory signs and symptoms tended to be higher, but
253 were not significantly different between those with cryptosporidium-positive and -negative
254 stool samples. In addition to sampling the lower respiratory tract with induced sputum, we
255 also collected NP specimens throughout the follow-up period, and documented detection
256 throughout the 8-week period, although in fewer numbers, more intermittently, and at higher
257 Ct counts compared to sputum. Respiratory symptoms were more common among those
258 cryptosporidium-positive in respiratory and GI tract compared to positive in GI tract alone,
259 although again these findings were not statistically significant. In contrast to the Mor study,
260 which only collected sputum samples from children with cough, unexplained tachypnea or
261 hypoxia, we collected respiratory specimens in all participants at enrollment and noted that
262 participants with cryptosporidium positivity in the respiratory tract had respiratory symptoms

in just over half of visits. These data support that almost half the participants with respiratory cryptosporidium detection are asymptomatic, and is a lower, rather than upper, respiratory tract pathogen or colonizer [18-21].

The continued detection of cryptosporidium in both GI and respiratory tract over 8 weeks may reflect the young age of this population. A study in Bangladesh noted that children ≤ 2 years positive for cryptosporidium shed it for a mean of 4.1 weeks, which was significantly longer than those > 2 years (mean 1.7 weeks) [22]. However, that was a household transmission study and did not evaluate children hospitalized with diarrhea. For respiratory pathogens, prolonged detection of respiratory viruses has been noted in a community surveillance study in Utah, with significantly longer detection noted in children < 5 years (mean 1.9 weeks) compared to other age groups (mean 1.6-1.7 weeks) [23]. To our knowledge, prolonged detection in the respiratory tract has not before been described for a protozoan parasite, which was not associated with higher prevalence of respiratory symptoms.

Prevalence of cryptosporidiosis is higher in cases of persistent v. acute diarrhea (15% v. 6.1% in a study in Guinea Bissau [5]). Persistent diarrhea has been described for cryptosporidiosis with duration of up to 5 months among severely immunocompromised infants [24]. In the Etiology, Risk Factors and Interactions of Enteric Infections and Malnutrition and the Consequences for Child Health and Development Program (MAL-ED) study, cryptosporidiosis was the fifth highest attributable pathogen in all pediatric diarrhea in the community setting, and had increased frequency among those with prolonged and severe diarrhea [3]. Among our participants, GI symptoms resolved within 2 weeks of hospitalization for all participants, but cryptosporidium was detected in stool for eight

participants throughout the 8-week period. Shedding duration for cryptosporidiosis among children hospitalized with diarrhea in sub-Saharan Africa has not been described previously using molecular techniques, although in Malawian HIV-infected adults we documented consistent shedding for up to 8 weeks [25]. Nutritional status may also account for the prolonged detection of up to 8 weeks that we documented in our study population. Even though the mean weight-for-height z score was within normal, the mean height-for-age and weight-for-age z scores of recruited participants both met WHO criteria for stunting and wasting, and 41%, 22%, and 10% had height-for-age, weight-for-age, and weight-for-height z scores <-2 , respectively, indicating mild-moderate malnutrition in this population. Reduced nutritional status could impact host immunity and predisposition to colonization/infection [26-28].

Cryptosporidiosis has been associated with excess mortality in children who had the infection in infancy, and this excess mortality persists into the second year of life [4, 5]. In our study we had one death, which occurred in a 7 month-old breastfeeding, HIV unknown male. This child had anthropometric parameters consistent with severe acute malnutrition (mid-upper arm circumference 11 cm, height-for-age z score -1.85, weight-for-age z score -3.01, weight-for-height z score -2.47, no nutritional edema) at time of enrollment, and an enrollment cryptosporidium Ct of 23 in the stool but negative in sputum and NP. At time of enrollment he was afebrile with normal vitals and had mild dehydration, diarrhea associated with abdominal tenderness and vomiting, as well as runny nose, cough and wheeze, but no shortness of breath. Death occurred two weeks after hospitalization after the child developed a cough and fever and was treated at an outpatient health facility.

This study has several limitations. The sample size was small, and the study population was drawn from a single hospital, and therefore findings may not be generalizable to other settings. Secondly, we did not evaluate for respiratory co-pathogens in our enrollment specimens, thereby limiting our ability ascribe symptoms to a single pathogen, although we did document co-pathogen detection in 84% during the follow-up period. Thirdly, we did not genotype cryptosporidium to ascertain whether re-infections occurred. However, these limitations are balanced out by the longitudinal observational design, which allowed us to document for the first time the dynamics of respiratory cryptosporidium among children hospitalized with diarrheal disease.

Conclusion

In summary, in this study we demonstrated that over 20% of young children hospitalized with diarrheal disease in our setting are positive for cryptosporidium, and they can shed cryptosporidium in the stool for up to 2 months. Concurrent respiratory cryptosporidiosis can be detected in a substantial proportion of young children, and detection continues over a prolonged period, with detection more prominent in the lower rather than upper respiratory tract. This has implications for the development of therapeutics for cryptosporidiosis, which is limited currently to nitazoxanide, and which is poorly efficacious in malnourished children [29, 30]. Since malnourished children are in greatest need for cryptosporidium therapeutics, scientists have been actively trying to find new, more effective drugs to treat cryptosporidiosis in this population. Because safety is of paramount concern in this age group, those developing drugs have debated whether a drug that only resides in the GI tract would be adequate for treating cryptosporidiosis in malnourished children, compared to one with broader systemic distribution [31]. Our findings suggest that the respiratory tract is a significant reservoir for cryptosporidium infection, and that to cure children, may require a

drug that distributes to the lungs as well as the GI tract. Thus, future research on therapeutic development should focus on drugs that target not only the GI but also the respiratory tract.

(2,628 words)

≡

Acknowledgements

We thank the patients and parents for participating in this study; research clinical and laboratory staff for conducting the study; David Moore and Tanja Adams with training the CryptoResp clinical and laboratory teams; Eric Houpt and Darwin Operario with providing TAC cards and technical assistance; Neema Toto for interim study support; Thokozani Ganiza for assistance with data management; and Wes Van Voorhis for reviewing drafts, providing feedback, and supporting us throughout the whole project.

Funding

The work was supported by the Bill & Melinda Gates Foundation (OPP1191165).

Author contributions

Conceived and designed the study: PI; Performed the study: MC, WN, HT, JN; Analyzed the data: MC, JN; Supervised the study and wrote first draft of the manuscript: PI; Reviewed, provided critical feedback, and approved the final draft: all authors.

Potential conflicts of interest

PI has received grants from BMGF outside of the submitted work. All other authors report no potential conflicts. All authors have submitted the ICMJE Form for Disclosure of Potential Conflicts of Interest.

References

1. Berkman DS, Lescano AG, Gilman RH, Lopez SL, Black MM. Effects of stunting, diarrhoeal disease, and parasitic infection during infancy on cognition in late childhood: a follow-up study. *Lancet* **2002**; 359(9306): 564-71.
2. Lima AA, Moore SR, Barboza MS, Jr., et al. Persistent diarrhea signals a critical period of increased diarrhea burdens and nutritional shortfalls: a prospective cohort study among children in northeastern Brazil. *J Infect Dis* **2000**; 181(5): 1643-51.
3. Platts-Mills JA, Babji S, Bodhidatta L, et al. Pathogen-specific burdens of community diarrhoea in developing countries: a multisite birth cohort study (MAL-ED). *Lancet Glob Health* **2015**; 3(9): e564-75.
4. Kotloff KL, Nataro JP, Blackwelder WC, et al. Burden and aetiology of diarrhoeal disease in infants and young children in developing countries (the Global Enteric Multicenter Study, GEMS): a prospective, case-control study. *Lancet* **2013**; 382(9888): 209-22.
5. Molbak K, Hojlyng N, Gottschau A, et al. Cryptosporidiosis in infancy and childhood mortality in Guinea Bissau, west Africa. *BMJ* **1993**; 307(6901): 417-20.
6. Korpe PS, Haque R, Gilchrist C, et al. Natural History of Cryptosporidiosis in a Longitudinal Study of Slum-Dwelling Bangladeshi Children: Association with Severe Malnutrition. *PLoS Negl Trop Dis* **2016**; 10(5): e0004564.
7. Costa LB, JohnBull EA, Reeves JT, et al. Cryptosporidium-malnutrition interactions: mucosal disruption, cytokines, and TLR signaling in a weaned murine model. *J Parasitol* **2011**; 97(6): 1113-20.
8. Mor SM, Tumwine JK, Ndeezi G, et al. Respiratory cryptosporidiosis in HIV-seronegative children in Uganda: potential for respiratory transmission. *Clin Infect Dis* **2010**; 50(10): 1366-72.

- 412 9. Hojlyng N, Jensen BN. Respiratory cryptosporidiosis in HIV-positive patients. *Lancet*
413 **1988**; 1(8585): 590-1.
- 414 10. Mercado R, Buck GA, Manque PA, Ozaki LS. *Cryptosporidium hominis* infection of
415 the human respiratory tract. *Emerg Infect Dis* **2007**; 13(3): 462-4.
- 416 11. Sponseller JK, Griffiths JK, Tzipori S. The evolution of respiratory Cryptosporidiosis:
417 evidence for transmission by inhalation. *Clin Microbiol Rev* **2014**; 27(3): 575-86.
- 418 12. Nyangulu W, Van Voorhis W, Iroh Tam PY. Evaluating respiratory cryptosporidiosis
419 in pediatric diarrheal disease: protocol for a prospective, observational study in
420 Malawi. *BMC Infect Dis* **2019**; 19(1): 728.
- 421 13. Grant LR, Hammitt LL, Murdoch DR, O'Brien KL, Scott JA. Procedures for
422 collection of induced sputum specimens from children. *Clin Infect Dis* **2012**; 54 Suppl
423 2: S140-5.
- 424 14. Liu J, Kabir F, Manneh J, et al. Development and assessment of molecular diagnostic
425 tests for 15 enteropathogens causing childhood diarrhoea: a multicentre study. *Lancet*
426 *Infect Dis* **2014**; 14(8): 716-24.
- 427 15. Korpe PS, Valencia C, Haque R, et al. Epidemiology and Risk Factors for
428 Cryptosporidiosis in Children From 8 Low-income Sites: Results From the MAL-ED
429 Study. *Clin Infect Dis* **2018**; 67(11): 1660-9.
- 430 16. Iturriza-Gomara M, Jere KC, Hungerford D, et al. Etiology of Diarrhea Among
431 Hospitalized Children in Blantyre, Malawi, Following Rotavirus Vaccine
432 Introduction: A Case-Control Study. *J Infect Dis* **2019**; 220(2): 213-8.
- 433 17. Liu J, Platts-Mills JA, Juma J, et al. Use of quantitative molecular diagnostic methods
434 to identify causes of diarrhoea in children: a reanalysis of the GEMS case-control
435 study. *Lancet* **2016**; 388(10051): 1291-301.

- 436 18. Clavel A, Arnal AC, Sanchez EC, et al. Respiratory cryptosporidiosis: case series and
437 review of the literature. *Infection* **1996**; 24(5): 341-6.
- 438 19. Dupont C, Bougnoux ME, Turner L, Rouveix E, Dorra M. Microbiological findings
439 about pulmonary cryptosporidiosis in two AIDS patients. *J Clin Microbiol* **1996**;
440 34(1): 227-9.
- 441 20. Lopez-Velez R, Tarazona R, Garcia Camacho A, et al. Intestinal and extraintestinal
442 cryptosporidiosis in AIDS patients. *Eur J Clin Microbiol Infect Dis* **1995**; 14(8): 677-
443 81.
- 444 21. Poirot JL, Deluol AM, Antoine M, et al. Broncho-pulmonary cryptosporidiosis in four
445 HIV-infected patients. *J Eukaryot Microbiol* **1996**; 43(5): 78S-9S.
- 446 22. Korpe PS, Gilchrist C, Burkey C, et al. Case-Control Study of *Cryptosporidium*
447 Transmission in Bangladeshi Households. *Clin Infect Dis* **2019**; 68(7): 1073-9.
- 448 23. Byington CL, Ampofo K, Stockmann C, et al. Community Surveillance of
449 Respiratory Viruses Among Families in the Utah Better Identification of Germs-
450 Longitudinal Viral Epidemiology (BIG-LoVE) Study. *Clin Infect Dis* **2015**; 61(8):
451 1217-24.
- 452 24. Kocoshis SA, Cibull ML, Davis TE, Hinton JT, Seip M, Banwell JG. Intestinal and
453 pulmonary cryptosporidiosis in an infant with severe combined immune deficiency. *J*
454 *Pediatr Gastroenterol Nutr* **1984**; 3(1): 149-57.
- 455 25. Iroh Tam PY, Arnold SLM, Barrett LK, et al. Clofazimine for treatment of
456 cryptosporidiosis in HIV-infected adults (CRYPTOFAZ): an experimental medicine,
457 randomized, double-blind, placebo-controlled phase 2a trial. *Clin Infect Dis* **2020**.
- 458 26. Bourke CD, Jones KDJ, Prendergast AJ. Current Understanding of Innate Immune
459 Cell Dysfunction in Childhood Undernutrition. *Front Immunol* **2019**; 10: 1728.

27. Gwela A, Mupere E, Berkley JA, Lancioni C. Undernutrition, Host Immunity and Vulnerability to Infection Among Young Children. *Pediatr Infect Dis J* **2019**; 38(8): e175-e7.
28. Ibrahim MK, Zambruni M, Melby CL, Melby PC. Impact of Childhood Malnutrition on Host Defense and Infection. *Clin Microbiol Rev* **2017**; 30(4): 919-71.
29. Checkley W, White AC, Jr., Jaganath D, et al. A review of the global burden, novel diagnostics, therapeutics, and vaccine targets for cryptosporidium. *Lancet Infect Dis* **2015**; 15(1): 85-94.
30. Striepen B. Parasitic infections: Time to tackle cryptosporidiosis. *Nature* **2013**; 503(7475): 189-91.
31. Huston CD, Spangenberg T, Burrows J, Willis P, Wells TN, van Voorhis W. A Proposed Target Product Profile and Developmental Cascade for New Cryptosporidiosis Treatments. *PLoS Negl Trop Dis* **2015**; 9(10): e0003987.

Table 1. Characteristics of the study population at enrollment

Table 2. Longitudinal detection of cryptosporidium in GI and respiratory tract and associated symptoms


Table 3. Characteristics and associated symptoms in participants with cryptosporidium detection in GI tract only v. GI and respiratory tract A) at enrollment; B) throughout study period

Supplementary Table 1. Full characteristics of study population at enrollment

Figure legend

Figure 1. Comparison of cryptosporidium detection in GI and respiratory tracts (mean cycle threshold with 95% confidence intervals) over 8 weeks in children hospitalized with diarrheal disease

Figure 2. Patient-level dynamics of cryptosporidium detection in GI and respiratory tracts over 8 weeks in a: A) 10 month-old; B) 14 month-old; C) 9 month-old; D) 12 month-old; E) 14 month-old; F) 7 month-old; G) 7 month-old; H) 11 month-old; I) 14 month-old; J) 11 month-old; K) 17 month-old; L) 10 month-old; M) 13 month-old; N) 20 month-old; and O) 15 month-old.



Cycle threshold (Ct)

Study visit

—◆— stool - - -■- - - sputum —○— nasopharyngeal ↑ respiratory symptoms

