

1 Skill-level based examination of forearm muscle activation
2 associated with efficient wrist and finger movements during
3 typing

4 **Keywords**

5 Typing; Upper extremity musculoskeletal disorders (UE-
6 MSDs); Motion capture system; Electromyography;
7 Wrist/finger joint movements; Muscle activation

8 **Abstract**

9 This study aims to elucidate the relationship between the
10 wrist and finger movements and forearm muscle activation of
11 people with different typing skills. It hypothesizes that skilled
12 typists (STs) can move their wrist and finger joints faster than
13 unskilled typists (UTs) can because they can efficiently use
14 their muscles according to the activity characteristics of the
15 flexors and extensors of the wrist joint.

16 During the typing task, we measured wrist and finger
17 movements using a 3D motion capture system and forearm
18 muscle activation using surface electromyography.

19 The angular velocity of the wrist and finger

20 flexion/extension and the muscle activation of the wrist flexors
 21 was higher in the STs than that in the UTs while the muscle
 22 activation of the wrist extensors was higher in the latter than
 23 that in the former.

24 Our results showed that STs may have used their forearm
 25 muscles to take advantage of the physical characteristics of the
 26 keys and the spring characteristics of their muscles and tendons.
 27 It was suggested that they placed less mechanical stress on their
 28 finger muscles and tendons when pressing and releasing the
 29 keys.

30

31 **1. Introduction**

32 Computers are essential tools in modern society that we
33 frequently use for long periods when working or studying.
34 Most people, regardless of age, gender, or occupation, are
35 required to use computers, with the continuing expansion of
36 digital transformation. Typing is a particularly complex task
37 that relies on many joints and muscles. During typing, finger
38 movements are usually rapid and repetitive and people hold
39 their hands continuously above the keyboard, exposing the
40 tendons and sheaths of their wrist and fingers to mechanical
41 stress (Keller et al. , 1998).

42 Bergqvist et al. (1992) first reported a prospective study
43 wherein typing work induced the risk of upper extremity
44 musculoskeletal disorders (UE-MSDs) of the wrist and finger
45 joints (Bergqvist et al. , 1992). In the next largest prospective
46 study, 21% of workers who used computers for more than 15 h/
47 week developed wrist or finger-related disorders in 1 year, most
48 of which were tendon disorders (Gerr et al. , 2002). In a recent
49 review, a dose–response relationship was plotted between the

50 exposure of people to mechanical factors and the risk of UE-
51 MSDs associated with their wrists and finger tendons (Keir et
52 al. , 2021). Therefore, it is necessary to investigate how the
53 overuse of wrists and fingers may contribute to the
54 development of UE-MSDs.

55 In this context, typing is required in bodily kinesthetic
56 learning to avoid overloads on wrist and finger joints and UE-
57 MSDs, such as sports injuries. Previous research on typing
58 proficiency and physical movements has mainly focused on
59 skilled typists (STs) of high skill. Additionally, few studies
60 have assessed the kinematic properties of the wrist and finger
61 joints, even though typing primarily comprises wrist and finger
62 movements (Alizadehkhayat and Frostick, 2015, Eygendaal et
63 al. , 2007).

64 To quantify the complex kinematic properties of wrists
65 and fingers during typing, it is necessary to understand their
66 structural characteristics and properly evaluate the coordination
67 between them. The relationship between the synergistic motion
68 of the wrist and finger joints in the sagittal plane is called the

69 tenodesis function (Cabri et al. , 2009, Marta et al. , 2012). It
70 refers to when passive finger joint flexion is induced during
71 wrist joint extension; and conversely, finger joint extension is
72 induced during wrist joint flexion (Cooney et al. , 1989, Horii
73 et al. , 1992). Particularly, the muscle tension of the finger joint
74 does not increase excessively and the amount of tendon
75 extension can be increased during wrist joint flexion. In this
76 context, evaluating only the movements of the wrist or finger
77 joint and related muscle activities may lead to a
78 misinterpretation of the mechanical activation of the body.
79 Therefore, the relationship between hand kinematic
80 coordination and muscle activities needs to be properly
81 assessed and the biomechanical risk factors that arise need to
82 be determined. Previous studies have discussed hand and finger
83 kinematics by analyzing the simple tapping of an index finger.

84 In this study, we evaluated repeated keystroke motions
85 during typing to analyze wrist and finger kinematics during
86 more complex repetitive motions. Movement styles may vary
87 with proficiency because typing is a complex task controlled by

many joints and muscles. This study aims to elucidate the relationship between wrist and finger movements and forearm muscle activation for different typing skills. We hypothesized that STs could move their wrists and fingers more efficiently than unskilled typists (UTs) could. The movement of their muscles depends on the characteristics of the flexor and extensor activities at the wrist joint.

2. Methods

2.1 Subjects

Twelve healthy subjects (nine men and three women) participated in this experiment; they were all right-handed. All the subjects provided written informed consent, according to the Declaration of Helsinki, after receiving a detailed explanation of the purpose of the study and the risks involved. This study was approved by the Ethics Review Committee (approval number: 20,508).

2.2 Measurement settings

A PC monitor and QWERTY keyboard (Microsoft, Wired Keyboard 600 ANB-00040, Washington, USA) were

107 placed at a height of 70 cm on a desk. Before beginning the
108 tests, each subject adjusted the chair according to their
109 preferences. We used a 3D motion capture system (Vicon
110 Motion Systems Ltd., Vicon, Oxford Metrics PLC, London,
111 UK; sampling frequency: 100 Hz) and wireless multipoint
112 surface electromyography (EMG) (DELSYS, Delsys Trigno
113 Wireless System, Massachusetts, USA; sampling frequency:
114 1,000 Hz).

115 ***2.3 Procedure***

116 We measured the height and weight of each subject and
117 collected the other necessary physical information for the
118 experiment. Infrared-reflecting markers (diameters 4 and 14
119 mm) were attached to points on the subjects (Fig. 1)—
120 according to the Vicon plug-in-gait full-body (PIG) and wrist
121 and finger joint models (Baker et al. , 2007, Cook et al. , 2007)
122 —to calculate the sagittal plane of the wrist and the angular
123 velocity of the metacarpophalangeal joint (MCP) of the index
124 finger.

125 Surface EMG sensors were attached to the right side of

126 the flexor carpi radialis (FCR), flexor carpi ulnaris (FCU), both
127 as the term extensor carpi radialis (ECR), and extensor carpi
128 ulnaris (ECU). The strength of each muscle was measured
129 using maximum voluntary isometric contraction (MVC).

130 Each subject practiced typing for 5 min before the
131 experiment to become accustomed to the test environment.
132 During the practice trial, each subject adjusted the height of the
133 chair and the angle of the monitor according to their preference.
134 The typing skills of the subjects were pretested on an electronic
135 keyboarding program (e-typing, e-typing Co., Ltd., Aichi,
136 Japan; <http://www.e-typing.ne.jp>), which had a text display with
137 text that moved as each subject typed.

138 In the typing test, we instructed the subjects to begin the
139 task after confirming that their hands were placed on the
140 keyboard. All the subjects typed the same sentences—each
141 comprising approximately 5,000 Japanese characters—that
142 were displayed on the monitor. No instructions were provided
143 regarding the typing speed, rhythm, or time limit; all the
144 subjects typed according to their regular style and pace. Each

145 subject performed the task thrice, with short pauses in between.

146

147 ***2.4 Data processing***

148 The wrist joint angle was calculated from markers
149 attached to the metacarpal head of the index finger, radial
150 styloid process, ulnar styloid process, and forearm in the model
151 (Fellinger et al. , 2010). We expressed that the positive value
152 was the extension and that the negative value was the flexion
153 angle of the wrist joint. The MCP joints were used to define the
154 hand as a rigid body, and the angle of the proximal phalanx
155 segment of the hand on the sagittal plane was calculated and
156 used as the flexion/extension angle (Fig. 1). We expressed that
157 the positive value was the flexion angle and that the negative
158 value was the extension angle for the MCP angle. The 4 mm
159 marker was attached to the enter key on the keyboard to record
160 the typing time and calculate the analysis interval.

161 The kinematic data were filtered using a fourth-order
162 Butterworth low-pass filter with a cut-off frequency of 6 Hz.
163 After filtering the data, the mean joint angle, changes in the

164 joint angle based on the root mean square (RMS), and joint
165 velocity of the wrist and MCP joint were calculated for each
166 trial.

167 The EMG data were passed through a fourth-order
168 Butterworth bandpass filter with a cutoff frequency of 20–480
169 Hz; the filter was full-wave rectified and smoothed at 10 Hz.
170 The maximum voluntary contraction (MVC) normalization
171 technique was applied to the EMG data obtained during the
172 typing sessions.

173 The 3D motion capture system was used to capture the
174 timing of the first end of the descent of the marker, and the
175 timings of the key input signal and start of the EMG data
176 recording were synchronized in time. The analysis interval was
177 sampled between the first and last entries of the task sentences.

178 ***2.5 Data analyses***

179 ***2.5.1 Grouping by the pretest results***

180 All data were obtained using Nexus 2.9 (Vicon Motion
181 Systems Ltd., Vicon, Oxford Metrics PLC, London, UK) and
182 Python 3.7 (Python Software Foundation, Beaverton, Orlando,

183 USA). The accuracy (%) and typing speed (words per minute,
184 WPM) in Japanese for all the subjects in the pretest were
185 obtained, where accuracy is the percentage of number of keys
186 that can be inputted without any mistakes. In a study
187 investigating the keystroke characteristics of both typing
188 novices and experts (Takaoka et al. , 2014), both were
189 classified as having a correct response rate of 95% or higher or
190 rates close to it, reaching the required condition for
191 employment.

192 *2.5.2 Kinematic data of wrist and metacarpophalangeal joints*

193 The mean and SD values of the flexion/extension angles
194 of the wrist and MCP flexion/extension angles of the index
195 finger were calculated to analyze the average posture of the
196 wrist and MCP joints during the analysis interval. Similarly, the
197 mean and standard deviation (SD) of the RMS of the data were
198 calculated. The angular velocities of the wrist and index finger
199 MCP joints were investigated by calculating the mean and SD
200 values of the changes in the angles/s .

201 *2.5.3 Analysis of muscle activation in the forearm*

202 To investigate the muscle activation in each forearm
203 muscle, we used amplitude probability distribution function
204 (APDF) analysis for the time series data of %MVC. APDF
205 analysis is a method that expresses the percentage of the total
206 time that the output is below a certain level as the probability of
207 occurrence (P) for that output when the EMG is being
208 measured (Jonsson, 1982). Based on these probability values,
209 the muscle activities can be classified into the static activity
210 level, which is equivalent to the activity involved in static
211 posture holding or pausing movement ($P = 0.1$); the average
212 activity level, which is exhibited throughout the movement (P
213 $= 0.5$); and the maximum activity level, which is equivalent to
214 the maximum activity exhibited during the movement ($P = 0.9$).
215 After plotting the function of the power output and appearance
216 probability of each muscle of each subject in each trial, we
217 calculated the mean and SD values of the muscle activation at
218 $P = 0.1, 0.5$, and 0.9 .

219 2.5.4 Identification of press and release times

220 From analyzing the key input signals of the subjects, we

221 obtained the time at which the input of each key started and
 222 ended. The time at which the “U” key was pressed, which is the
 223 start of the input, was defined as the press time; and the time at
 224 which the fingertip left the “U” key, indicating the end of the
 225 inputs, was defined as the release time. The mean and SD
 226 values of each data point were calculated and compared
 227 between the two groups.

228 **2.6 Statistics**

229 After confirming the normality of all the kinematic and
 230 muscle activity data using the Shapiro–Wilk test, we
 231 investigated the homoscedasticity of the data using the F-test.
 232 Since there was no difference in the EMG and kinematic data
 233 in each trial (confirmed by one-way ANOVA; $P < 0.05$), the
 234 average values of the three trials were used as representative
 235 values. The data of the two groups were compared using the
 236 unpaired t-test ($P < 0.05$) or Wilcoxon signed-rank test.

237 **3. Results**

238 **3.1 Subjects and environments**

239 The mean accuracy (SD) values in the typing test are

240 shown in Table 1. The mean value of the WPM typed by all the
241 subjects was used as the benchmark for measuring the skill
242 level; subjects who achieved scores higher and lower than the
243 mean were classified into the ST and UT groups, respectively.

244 The mean (SD) age, height, weight, hand width, finger
245 length, male-to-female ratio, chair height, and screen angle
246 values are presented in Table 1. The ratio of subjects that used
247 desktop to notebook PCs daily was 1:11. The mean (SD) values
248 for the frequency and duration of computer use were 5.2 (1.2)
249 times per week and 3.3 (1.8) h per day, respectively. There
250 were no significant differences between the two groups in any
251 of the data.

252 ***3.2 Wrist and metacarpophalangeal joint motion***

253 There was no significant difference in the mean
254 extension angle of the wrist joint at the beginning of the typing
255 motion between the two groups. The mean values (SD) of the
256 wrist joint extension angle, RMS, and angular velocity are
257 shown in Fig. 2. The angular velocity of the flexion/extension
258 was predominantly higher in the ST group ($P = 0.003$) than that

259 in the UT group.

260 The mean (SD) values of the flexion angle, RMS, and
261 angular velocity of the MCP joint of the index finger are shown
262 in Fig. 2. The mean angular velocity of the MCP joint in the ST
263 group was significantly higher than that in the UT group ($P =$
264 0.002). The mean values (SD) of %MVC of FCR, FCU, ECR,
265 and ECU at the static activity level ($P = 0.1$), mean activity
266 level ($P = 0.5$), and maximum activity level ($P = 0.9$) are
267 shown in Fig. 3. The mean %MVC of FCR at $P = 0.9$ was
268 significantly higher than that in the ST group ($P = 0.023$). The
269 mean %MVC of ECR at $P = 0.1, 0.5, 0.9$ were significantly
270 higher than that in the UT group at $P = 0.040, 0.020$, and 0.020 ,
271 respectively.

272 ***3.3 Data on press and release time***

273 The means (SD) of the press and release times were 0.12
274 (0.02) and 0.20 s (0.14), respectively. The press and release
275 times of the ST group were 0.13 (0.01) and 0.11 s (0.01),
276 respectively. The press and release times of the UT group were
277 0.11 (0.02) and 0.26 (0.15), respectively. There were no

278 significant differences between the two groups. The mean
279 values (SD) of the wrist joint extension angle, angular velocity,
280 flexion angle of the MCP joint of the index finger, and angular
281 velocity during the press and release times are shown in Figs. 4
282 and 5. The mean angular velocity of the wrist joint extension
283 during the release time in the ST group was significantly higher
284 than that in the UT group ($P = 0.005$). The mean value of the
285 angular velocity of the MCP joint flexion/extension of the
286 index finger during the press and release times in the ST group
287 was significantly higher than that in the UT group ($P = 0.002$,
288 0.006).

289 The mean values (SD) of the %MVC of the FCR, FCU,
290 ECR, and ECU during press and release are shown in Figs. 4
291 and 5. The mean values (SD) of the %MVC of the ECR during
292 the press time of the ST group were significantly lower than
293 those of the UT group ($P = 0.020$).

294 **4. Discussion**

295 We hypothesized that the subjects of the ST group could
296 move their wrist and finger joints faster than those of the UT

group could because they could use their muscles according to the activity characteristics of the flexors and extensors of their wrist joint. The ST group had a higher muscle activation at the maximum activity level of the FCU, whereas the UT group had a higher muscle activation at all the ECR activity levels. The subjects of the ST group also moved the MCP joints of their wrists and index fingers faster than those of the UT group did. The press and release time results clearly explained the muscle activation and kinematics of the subjects during the experiment. This study revealed that repetitive keystrokes with muscle activity controls for efficient wrist and finger movements reduced muscle activation during typing. This study emphasizes the significance of assessing not only muscle activity but also the complementary movements of the wrists and fingers simultaneously when considering muscle activation and the risks associated with UE-MSDs during typing.

4.1 Muscle activation and movement during experiment

The maximum activity level of the FCR ($P = 0.9$) was higher in the ST group. This value is an index of the peak level

316 of muscle activity involved in the task performance (Szeto et
317 al. , 2009); it refers to the muscle activity acting on the pressed
318 keys during typing.

319 Key presses are coordinated by the muscle activities of
320 the wrist and finger flexors (Dennerlein et al. , 2007).
321 Additionally, because the finger flexors were stretched in the
322 wrist extension posture, the mechanical stress on them and the
323 tendons may have increased if active tension was generated in
324 the stretched state when pressing the keys. Therefore, the
325 controlled muscle activity of the ST group may be attributed to
326 the increase in the peak level of muscle activity in the wrist
327 flexors, thereby decreasing the muscle load and mechanical
328 stress on the finger flexors.

329 All the ECR activity levels ($P = 0.1, 0.5, \text{ and } 0.9$) were
330 higher in the UT group. Static activity level is an index of
331 muscle activity related to maintaining a static posture
332 (Simoneau et al. , 2003); it was inferred that the muscle activity
333 was inefficiently controlled due to static posture and the highly
334 antagonistic action of the finger flexor when pressing the keys.

335 ***4.2 Kinematics of wrist and finger joints during experiment***

336 No significant differences were observed in the extension
337 angle and RMS of the wrist between the ST and UT groups.
338 However, the angular velocity of flexion/extension was higher
339 in the ST group than that in the UT group. The kinematic
340 parameters of the MCP joint of the index finger showed no
341 significant differences in the joint angle and RMS. The higher
342 typing speed of the ST group was due to the fast movements of
343 the wrist and finger joints and coordinated fast actions of the
344 joints.

345 To move the wrist and finger joints efficiently, it is
346 important to convert variables, such as inertia and gravity, into
347 joint movements (Dennerlein, Kingma, 2007, Qin et al. , 2014).

348 As mentioned in 4.1, the ST group had muscle
349 coordination that increased and decreased the FCR and ECR
350 muscle activities, respectively. This may have allowed the ST
351 group to maintain the downward inertial force generated in the
352 fingertip immediately before pressing the keys and reduced the
353 braking action. As a result, the joint angular velocity of the

354 wrist and finger when pressing the keys was higher in the ST
355 group.

356 Additionally, the ST group performed muscle activity
357 control to reduce the ECR muscle activity during the
358 experiment. This indicated that the ST group effectively used
359 the spring action of the finger flexors to support the movement
360 of the fingertips in resisting gravity during the transition from
361 pressing the keys to releasing them. This inference also
362 indicated the existence of a relationship between the fast
363 movements of the wrist and finger joints.

364 *4.3 Muscle activation and movement when pressing and* 365 *releasing keys*

366 We evaluated the muscle activation of the wrist joint
367 during press and release times. The types and combinations of
368 fingers that a person uses while typing vary by skill; however,
369 the number of typists that do not use the index finger is small
370 (Feit et al. , 2016). Additionally, the "U" key was most
371 frequently entered using the right index finger, because vowels
372 (a, e, i, o, u) in Japanese are always entered after the

373 consonants.

374 The two groups showed similar activations of the wrist
375 flexors (FCR and FCU) during the task and muscle activities
376 during the press and release times. The ST group was also able
377 to perform finger and wrist joint movements faster than the UT
378 group could. Since the flexion movements of the MCP joint
379 during the typing movements were synergistic with those of the
380 wrist joint (Dennerlein, Kingma, 2007), it can be assumed that
381 the ST group took advantage of the excessive muscle activity
382 of the wrist flexors during keystrokes. Additionally, because the
383 fingertips were in high-speed contact with the keys when
384 pressing them, the wrist flexors may have acted excessively to
385 maintain the posture of the hand against the reaction force from
386 the keys. Finger flexor activity is activated after the onset of
387 downward finger movement (Kuo et al. , 2006); the ST group
388 may have activated their wrist flexors in a coordinated manner.
389 The ST group used this strategy because it increased the kinetic
390 energy of their fingertips immediately after pressing the keys,
391 producing a key release, which facilitated keystroke switching.

392 Wrist posture affects muscle strength and activation
393 patterns (Visser et al. , 2000), and sustained low-intensity
394 muscle activity has been associated with wrist extension
395 postures during typing (Dennerlein and Johnson, 2006) with the
396 wrist extensor muscle activity decreasing as the extension angle
397 of the wrist decreases (Simoneau, Marklin, 2003), possibly
398 because of changes in the muscle length and moment arm.

399 In this study, there was no difference in the angle of the
400 wrist joint or the degree of change in the angle between the two
401 groups; however, the muscle activation of the static activity
402 level of the ECR, an extensor muscle of the wrist joint, was
403 significantly higher in the UT group than that in the ST group.
404 This may be due to the over-activation of the ECR in the UT
405 group due to the lack of minimal muscle activity control
406 despite the longer moment arm of the muscle. This action is
407 antagonistic to the flexion movement of the finger and,
408 therefore, may increase the muscle load on the finger flexors
409 when pressing the keys.

410 Furthermore, the ECU muscles cause the hand to be

411 raised after pressing the keys (Dennerlein, Kingma, 2007). The
 412 UT group could not efficiently utilize the spring properties of
 413 the finger flexor tendons, resulting in the overuse of the
 414 muscles acting on the extensor muscles of their wrist joint
 415 when lifting their fingertips. This may have increased the EMG
 416 load on the ECU muscles when releasing the keys. Additionally,
 417 there is a proportional relationship between the distance of the
 418 spatial movement of the hand after key release and the time
 419 interval before the next keystroke (Feit, Weir, 2016). Therefore,
 420 the UT group, which tended to have longer key release times,
 421 may have increased the muscle load on the ECU muscles
 422 during key release because of excessive attempts to lift and
 423 move the hands.

424 ***4.4 Limitations***

425 The measurement conditions in this study were not strict.
 426 The postures of the subjects were not constrained during the
 427 test; they could place their forearms on the desk or keep them
 428 floating. Since placing the forearm on the desk may decrease
 429 the activation of the extensor muscles of the wrist joint, the

430 distance between the upper body and the keyboard and the
431 space on the desk in front of the keyboard needs to be checked
432 at the beginning of the measurement; and the effects of
433 differences in the contact time and contact area between the
434 desk and the forearm on the body during the task should be
435 determined.

436 In this study, we examined the differences in the physical
437 parameters of skill levels among healthy adult students.
438 Therefore, the findings of this study cannot be generalized
439 completely.

440 The press time and release time were used as the analysis
441 intervals, and the index finger movements were analyzed
442 thoroughly. However, we could not confirm which finger each
443 subject used to input the “U” key. Thus, we may not have been
444 able to exclusively evaluate the movement of the index finger.

445 Furthermore, because we did not measure the force
446 exerted by the fingertip during keystrokes, we could not
447 demonstrate a direct relationship between the angular velocity
448 of the MCP joint and the actual force generated during

449 keystrokes. In future investigations, we intend to evaluate the
450 movements of fingers other than that of the index finger and
451 examine the relationships between changes in the posture,
452 muscle activation, and the force exerted by fingertips during
453 specific keystrokes.

454 **5. Conclusions**

455 The velocity of the wrist and finger joints and muscle
456 activation of the forearm during typing differed according to
457 the skill level of the subjects, and it was inferred that these
458 differences were caused by using muscles during keystrokes.

459 This study emphasizes the significance of assessing not
460 only muscle activity but also the complementary movements of
461 wrists and fingers simultaneously when considering muscle
462 activation and the risks associated with UE-MSDs during
463 typing.

464 **Conflicts of Interest**

465 The authors declare that there are no conflicts of interest.

466 **Acknowledgments**

467 **References**

468 Alizadehkhayat O, Frostick SP. Electromyographic assessment
469 of forearm muscle function in tennis players with and without
470 Lateral Epicondylitis. *Journal of Electromyography and*
471 *Kinesiology*. 2015;25:876-86.

472 Baker NA, Cham R, Cidboy EH, Cook J, Redfern MS.
473 Kinematics of the fingers and hands during computer keyboard
474 use. *Clin Biomech (Bristol, Avon)*. 2007;22:34-43.

475 Bergqvist U, Knave B, Voss M, Wibom R. A longitudinal study
476 of VDT work and health. *International Journal of Human-
477 Computer Interaction*. 1992;4:197-219.

478 Cabri J, Sousa JP, Kots M, Barreiros J. Golf-related injuries: A
479 systematic review. *European Journal of Sport Science*.
480 2009;9:353-66.

481 Cook JR, Baker NA, Cham R, Hale E, Redfern MS.
482 Measurements of wrist and finger postures: A comparison of
483 goniometric and motion capture techniques. *Journal of Applied*
484 *Biomechanics*. 2007;23:70-8.

485 Cooney WP, Lin GT, An K-N. Improved tendon excursion
486 following flexor tendon repair. *Journal of Hand Therapy*.

1989;2:102-6.

Dennerlein JT, Johnson PW. Different computer tasks affect the exposure of the upper extremity to biomechanical risk factors. *Ergonomics*. 2006;49:45-61.

Dennerlein JT, Kingma I, Visser B, van Dieen JH. The contribution of the wrist, elbow and shoulder joints to single-finger tapping. *J Biomech*. 2007;40:3013-22.

Eyendaal D, Rahussen FTG, Diercks RL. Biomechanics of the elbow joint in tennis players and relation to pathology. *Br J Sports Med*. 2007;41:820-3.

Feit AM, Weir D, Oulasvirta A. How We Type. *Proceedings of the 2016 CHI Conference on Human Factors in Computing Systems* 2016. p. 4262-73.

Fellinger M, Passler J, Seggl W. Plug-in Gait Reference Guide. *Human and Nonhuman Bone Identification*. 2010:227-46.

Gerr F, Marcus M, Ensor C, Kleinbaum D, Cohen S, Edwards A, et al. A prospective study of computer users: I. Study design and incidence of musculoskeletal symptoms and disorders. *Am J Ind Med*. 2002;41:221-35.

506 Horii E, Lin GT, Cooney WP, Linscheid RL, An KN.
507 Comparative flexor tendon excursion after passive
508 mobilization: An in vitro study. Journal of Hand Surgery.
509 1992;17:559-66.

510 Jonsson B. Measurement and evaluation of local muscular
511 strain in the shoulder during constrained work. Journal of
512 human ergology. 1982;11:73-88.

513 Keir PJ, Farias Zuniga A, Mulla DM, Somasundram KG.
514 Relationships and Mechanisms Between Occupational Risk
515 Factors and Distal Upper Extremity Disorders. Hum Factors.
516 2021;63:5-31.

517 Keller K, Corbett J, Nichols D. Repetitive strain injury in
518 computer keyboard users: Pathomechanics and treatment
519 principles in individual and group intervention. Journal of Hand
520 Therapy. 1998;11:9-26.

521 Kuo PL, Lee DL, Jindrich DL, Dennerlein JT. Finger joint
522 coordination during tapping. J Biomech. 2006;39:2934-42.

523 Marta S, Silva L, Castro MA, Pezarat-Correia P, Cabri J.
524 Electromyography variables during the golf swing: A literature

525 review. Journal of Electromyography and Kinesiology.
526 2012;22:803-13.

527 Qin J, Trudeau M, Buchholz B, Katz JN, Xu X, Dennerlein JT.
528 Joint contribution to fingertip movement during a number entry
529 task: an application of Jacobian matrix. J Appl Biomech.
530 2014;30:338-42.

531 Simoneau GG, Marklin RW, Berman JE. Effect of Computer
532 Keyboard Slope on Wrist Position and Forearm
533 Electromyography of Typists Without Musculoskeletal
534 Disorders. Physical Therapy. 2003;83:816-30.

535 Szeto GP, Straker LM, O'Sullivan PB. Examining the low, high
536 and range measures of muscle activity amplitudes in
537 symptomatic and asymptomatic computer users performing
538 typing and mousing tasks. Eur J Appl Physiol. 2009;106:243-
539 51.

540 Takaoka E, Sugiura M, Komiya H. A study on typist
541 characterization. IPSJ SIG Technical Report. 2014;No.9:1-14.

542 Visser B, de Korte E, van der Kraan I, Kuijer P. The effect of
543 arm and wrist supports on the load of the upper extremity

544 during VDU work. Clinical Biomechanics. 2000;15:S34-S8.

545

Table.1 Subjects and Environments

	Skilled Typist (ST)	Unskilled Typist (UT)	<i>p</i>
Number	5	7	—
Correct answers(%)	94.0 (2.3)	95.1 (2.6)	0.513
WPM ^a	219.5 (21.3)	170.6 (36.2)	*0.033
Age	21.6 (0.8)	20.7 (0.5)	0.050
Height (cm)	167.3 (5.0)	166.1 (4.2)	0.700
Weight (kg)	63.2 (8.9)	56.6 (6.4)	0.202
Hand width (cm)	7.0 (0.4)	6.8 (0.1)	0.466
Finger length (cm)	10.2 (0.4)	9.9 (0.3)	0.305
Chair height (cm)	42.4 (3.9)	45.7 (2.8)	0.127
Screen angle (°)	129.0(12.4)	122.9(17.5)	0.552
Male-to-female ratio	3:2	6:1	—
Right-left handed ratio	5:0	7:0	—
Desktop - notebook PC ratio	1:4	0:7	—
Frequency; PC use times per week	5.6 (1.0)	4.86 (1.3)	0.340
Frequency; PC use times hours per days	3.8 (1.5)	2.93 (1.9)	0.445

* $p < 0.05$, ^a WPM = Words per minutes

546

547 **Figure legends**

548 **Fig. 1** Experimental equipment.

549 (A) Positions of the markers and EMGs

550 (B) Calculation of MCP joint angle

551 **Fig. 2** Wrist and metacarpophalangeal (MCP) joint motion

552 The error bar represents the standard deviation between the
553 subjects.

554 $p < 0.05$, ST = skilled typist, UT = unskilled typist, RMS = root
555 mean square

556 **Fig. 3** Muscle activation

557 From left to right, the static activity level ($P = 0.1$), mean
558 activity level ($P = 0.5$), and maximum activity level ($P = 0.9$)
559 are shown.

560 The error bar represents the standard deviation between the
561 subjects.

562 * $p < 0.05$, FCR (flexor carpi radialis); FCU (flexor carpi
563 ulnaris); ECR (extensor carpi radialis); ECU (extensor carpi
564 ulnaris).

565 **Fig. 4** Press time data

566 Data on when the “U” key is pressed. From left to right, the
 567 wrist joint extension angle, angular velocity, flexion angle of
 568 the MCP joint of the index finger, angular velocity, and %MVC
 569 of the FCR, FCU, ECR, and ECU are listed.

570 The error bar represents the standard deviation between the
 571 subjects.

572 * $p < 0.05$

573 **Fig. 5** Data on release time


574 Data on the time between releasing the “U” key and the start of
 575 the next keystroke. From left to right, the wrist joint extension
 576 angle, angular velocity, flexion angle of the MCP joint of the
 577 index finger, angular velocity, and %MVC of FCR, FCU, ECR,
 578 and ECU are listed.

579 The error bar represents the standard deviation between
 580 subjects.

581 * $p < 0.05$


582

583 **Fig.1**



584


585 **Fig.2**




586

587 **Fig.3**

588




589 **Fig.4**



590

591 **Fig.5**



592





Fig. 1 Experimental equipment.
 (A) Positions of the markers and EMGs
 (B) Calculation of MCP joint angle

(A) Wrist Joint



(B) MCP joint




Fig. 2 Wrist and metacarpophalangeal (MCP) joint motion

The error bar represents the standard deviation between the subjects.

$p < 0.05$, ST = skilled typist, UT = unskilled typist, RMS = root mean square




Fig. 3 Muscle activation

From left to right, the static activity level ($P = 0.1$), mean activity level ($P = 0.5$), and maximum activity level ($P = 0.9$) are shown.

The error bar represents the standard deviation between the subjects.

* $p < 0.05$, FCR (flexor carpi radialis); FCU (flexor carpi ulnaris); ECR (extensor carpi radialis); ECU (extensor carpi ulnaris).

Press the "U" key

Release the "U" key




Fig. 4 Press time data

Data on when the "U" key is pressed. From left to right, the wrist joint extension angle, angular velocity, flexion angle of the MCP joint of the index finger, angular velocity, and %MVC of the FCR, FCU, ECR, and ECU are listed.

The error bar represents the standard deviation between the subjects.

* $p < 0.05$

Press the "U" key

Release the "U" key





Fig. 5 Data on release time

Data on the time between releasing the "U" key and the start of the next keystroke. From left to right, the wrist joint extension angle, angular velocity, flexion angle of the MCP joint of the index finger, angular velocity, and %MVC of FCR, FCU, ECR, and ECU are listed. The error bar represents the standard deviation between subjects.

* $p < 0.05$

Table.1 Subjects and Environments

	Skilled Typist (ST)	Unskilled Typist (UT)	<i>p</i>
Number	5	7	—
Correct answers(%)	94.0 (2.3)	95.1 (2.6)	0.513
WPM ^a	219.5 (21.3)	170.6 (36.2)	*0.033
Age	21.6 (0.8)	20.7 (0.5)	0.050
Height (cm)	167.3 (5.0)	166.1 (4.2)	0.700
Weight (kg)	63.2 (8.9)	56.6 (6.4)	0.202
Hand width (cm)	7.0 (0.4)	6.8 (0.1)	0.466
Finger length (cm)	10.2 (0.4)	9.9 (0.3)	0.305
Chair height (cm)	42.4 (3.9)	45.7 (2.8)	0.127
Screen angle (°)	129.0(12.4)	122.9(17.5)	0.552
Male-to-female ratio	3:2	6:1	—
Right-left handed ratio	5:0	7:0	—
Desktop - notebook PC ratio	1:4	0:7	—
Frequency; PC use times per week	5.6 (1.0)	4.86 (1.3)	0.340
Frequency; PC use times hours per days	3.8 (1.5)	2.93 (1.9)	0.445

* $p < 0.05$, ^a WPM = Words per minutes