

1 **Running Title: Shared brain and genetic architectures between mental health and physical**
2 **activity**

3
4 Wei Zhang¹, Sarah E. Paul², Anderson Winkler³, Ryan Bogdan², Janine D. Bijsterbosch¹

5 ¹Radiology Department, Washington University School of Medicine, St. Louis

6 ²Department of Psychological & Brain Sciences, Washington University in St. Louis

7 ³National Institute of Mental Health / National Institutes of Health

8

9

10

11 **Abstract**

12 Physical activity is correlated with, and effectively treats various forms of psychopathology.
13 However, whether biological correlates of physical activity and psychopathology are shared
14 remains unclear. Here, we examined the extent to which the neural and genetic architecture of
15 physical activity and mental health are shared. Using data from the UK Biobank (N=6,389),
16 canonical correlation analysis was applied to estimate associations between the amplitude and
17 connectivity strength of sub-networks of three major neurocognitive networks (default mode,
18 DMN; salience, SN; central executive networks, CEN) with accelerometer-derived measures of
19 physical activity and self-reported mental health measures (primarily of depression, anxiety
20 disorders, neuroticism, subjective well-being, and risk-taking behaviors). We estimated the
21 genetic correlation between mental health and physical activity measures, as well as putative
22 causal relationships by applying linkage disequilibrium score regression, genomic structural
23 equational modeling, and latent causal variable analysis to genome-wide association summary
24 statistics (GWAS N=91,105-500,199). Physical activity and mental health were associated with
25 connectivity strength and amplitude of the DMN, SN, and CEN ($r's \geq 0.12$, $p's < 0.048$). These
26 neural correlates exhibited highly similar loading patterns across mental health and physical
27 activity models even when accounting for their shared variance. This suggests a largely shared
28 brain network architecture between mental health and physical activity. Mental health and
29 physical activity (including sleep) were also genetically correlated ($|r_g| = 0.085-0.121$), but we
30 found no evidence for causal relationships between them. Collectively, our findings provide
31 empirical evidence that mental health and physical activity have shared brain and genetic
32 architectures and suggest potential candidate sub-networks for future studies on brain
33 mechanisms underlying beneficial effects of physical activity on mental health.

34

35

36

37

38

39 **Keywords:** mental health, physical exercise, sleep, canonical correlation, multivariate analysis,

40

genetic correlation

41

42 **1. Introduction**

43 Mental health and physical activity have both been linked to emotion, cognition, and brain
44 correlates¹⁻⁴. Furthermore, physical activity is known to improve various psychiatric conditions⁵⁻
45 ⁷. Yet, it remains elusive whether mental health and physical activity overlap in neural and
46 genetic architectures. Here, in a large population sample (n=6,389 from the UK Biobank⁸), we
47 estimate the extent to which physical activity and general mental health share patterns of resting-
48 state functional MRI network measures and genetic architectures.

49

50 *1.1 Potential overlap between mental health and physical activity in neural networks*

51 Functional alterations in large-scale brain networks have been consistently implicated in a wide
52 range of psychiatric disorders⁹. Dysfunctional configuration of neurocognitive networks such as
53 the default mode, salience, and central executive networks has been hypothesized to characterize
54 major psychiatric disorders including depression and anxiety (DMN, SN, CEN)¹⁰. In line with
55 this hypothesis, findings from meta-analyses have shown that core cognitive and affective
56 abnormalities in major depression can be accounted for by hypo-connectivity within the CEN
57 and hyper-connectivity within the DMN¹¹, together with hypo-connectivity between the control
58 systems (i.e., CEN) and salience, emotion processing systems (i.e., SN)¹². Similarly, a recent
59 meta-analysis suggests that anxiety disorders are characterized by hypo-connectivity between
60 subcortical limbic circuits that partially overlap with the SN, CEN, and DMN, as well as
61 decoupling between the CEN and DMN¹³. Furthermore, the personality trait neuroticism, which
62 is considered a risk marker for psychopathology, has also been linked to alterations in functional
63 brain networks¹⁴. These same networks have also been linked to physical activity. Evidence from
64 fMRI studies on physical activity demonstrated the changes in activity of and functional
65 connectivity between these network hub regions including the hippocampus, parahippocampus,
66 dorsal anterior cingulate cortex, and ventromedial prefrontal cortex that primarily subserve
67 executive functions such as working memory, attention, and inhibition¹⁵⁻¹⁹. At the more system
68 level, a 12-month aerobic walking intervention was found to increase resting-state functional
69 connectivity between subnetworks of DMN, and between subnetworks of SN²⁰, whereas
70 connectivity in the CEN was found to increase after multiple sessions of high intensity interval
71 training²¹. Interestingly, the intensity of physical exercise appeared to modulate functional
72 connectivity changes in the hub regions of the CEN²², as well as in the DMN subsystems acutely

73 and after 3 months of training²³. Additionally, although sleep might not be considered a type of
74 physical activity for its inactive nature, it is closely connected to mental health such that sleep
75 problems have been considered as a risk factor for subsequent development of depressive
76 symptoms^{24,25}, and disrupted sleep is often seen in patients with various types of mood and
77 anxiety disorders²⁶⁻²⁸.

78 Additionally, sleep disturbance can reduce physical activity levels and increase the risk of
79 exercise-related injuries²⁹. Although sleep has been largely overlooked in the literature for
80 physical activity in relation to mental health, we included it in the current study for its close
81 relationships with both mental health and physical activity. Sleep duration can be accurately
82 measured with ecological sampling approaches such as wrist-worn accelerometers, which have
83 been adopted in medical and mental health research^{30,31}. At the neural network level, sleep
84 deprivation has been linked to reduction in functional connectivity within the DMN³²⁻³⁴,
85 whereas increased sleep duration has been linked to strengthened connectivity within the DMN
86 but reduced connectivity between DMN and SN³⁵. Fluctuations in arousal (i.e., indication of
87 drowsiness) during resting-state scan acquisitions have also been linked to the fMRI signal
88 amplitude in sensorimotor networks³⁶. These studies together highlight the engagement of the
89 DMN, SN and CEN in both mental health and physical activity and demonstrate that the
90 interplays between these large-scale intrinsic networks and subnetworks are coupled with
91 changes in mental health symptoms and physical exercises. Yet, it remains unclear whether
92 similar connectivity patterns or signal changes of these networks are shared by mental health and
93 physical activity.

94

95 *1.2 Potential overlap between mental health and physical activity in genetic variance*

96 Physical activity is known to promote resilience to various psychiatric conditions, alleviating
97 symptoms of depression, anxiety, and negative mood^{37,38}. This buffering effect may be rooted in
98 the shared genetic variance between mental health and physical activity. For instance, recent
99 research employing summary statistics from independent genome-wide associate studies (GWAS)
100 showed that higher polygenic risk scores for depression are associated with increased odds of
101 incident depression, whereas self-reported physical activities such as walking, jogging, running,
102 dancing and yoga appeared to reduce the odds with similar magnitude³⁹. This effect has also
103 been observed using more objective measurement of physical activity such that reduced activity

104 levels measured by accelerometer were found to associate with diagnoses of schizophrenia,
105 bipolar disorder, depression, or autism spectrum disorders (ASD) and that healthy participants
106 without disorder diagnoses were observed to perform less physical activity if they had a higher
107 polygenic risk score for schizophrenia, depression, and ASD⁴⁰.

108 Additionally, overlapping genetic architectures may exist across various psychiatric disorders
109 including anxiety and depression^{41–43} as psychiatric phenotypes are highly polygenic⁴⁴. Thus,
110 latent genetic factors capturing shared variance across clusters of psychiatric symptoms may
111 improve identification of associations between mental health and physical activity. For instance,
112 genomic contributions to disorders such as depression and anxiety are captured by a genetic
113 factor for internalizing disorders that are primarily characterized by these two disorders as
114 indicated by a confirmatory factor analysis, and this genetic factor is positively genetically
115 associated with various adverse health outcomes and negatively genetically associated with
116 physical movement patterns⁴⁵. These findings point to the possibility that the genetic
117 architectures of mental health and physical activity may overlap. The degree and phenotypic
118 specificity of overlap remains to be tested.

119

120 *1.3 Shared brain and genetic architectures?*

121 Together, the studies reviewed above show that mental health and physical activity both involve
122 large-scale brain networks such as the DMN, SN and CEN. Additionally, mental health and
123 physical activity may have partially overlapping genetic architectures, with evidence showing
124 associations between genetic liabilities of psychiatric disorders and physical activity, as well as
125 genetic associations between latent factors of psychopathology and physical activity. As mental
126 health and physical activity are also tightly related at the behavioral level (e.g., emotion,
127 cognition), it is reasonable to speculate that these two constructs may partially overlap in the
128 underlying neurobiological mechanisms. In this study, we aim to determine whether mental
129 health and physical activity have shared variance in brain and genetic architectures, using brain
130 network measures and genomic summary statistics.

131

132

133 **2. Methods and Materials**

134 *2.1 Participants*

135 The UK Biobank is an openly accessible population dataset with neuroimaging data collection,
136 in addition to extensive demographic, behavioral, lifestyle, and cognitive measures^{8,46}. An initial
137 sample of N=8,378 participants from the UK Biobank (UKB) was considered for this study.
138 These participants had participated in accelerometer-based physical activity evaluations and
139 visited the assessment center, where the resting-state fMRI and mental health data were acquired.
140 Data quality assurance resulted in exclusion of N=64 participants for insufficient accelerometer
141 data (see details below in section 2.2.2), and N=1,925 participants with considerable missing
142 data in the mental health questionnaire (see details below in section 2.2.1). The final sample had
143 N=6,389 participants with 2,994 (46.9%) females (sample mean age=63.74 \pm SD 7.53). All
144 participants provided informed consent. UK Biobank has ethical approval from the North West
145 Multi-Centre Research Ethics Committee (MREC). Data access was obtained under UK Biobank
146 application ID 47267.

147

148 *2.2 Data Acquisition and Preprocessing*

149 *2.2.1 General mental health measures*

150 The UKB general mental health questionnaire consisting of 41 items was conducted on the same
151 day as fMRI data acquisition (<https://biobank.ndph.ox.ac.uk/showcase/label.cgi?id=100060>).
152 This self-reported questionnaire primarily measured depression, anxiety, and neuroticism, as
153 well as subjective well-being. These variables had varying degrees of missing data points partly
154 due to responses such as “do not know”, “prefer not to answer”, or “none of above,” and partly
155 due to question dependencies. To ensure robust model estimation while maximizing statistical
156 power, individual variables or questions that had more than 30% missing values were excluded
157 (N=10; see full descriptions for each included individual question in *Table S1*). A multivariate
158 imputation procedure was then leveraged to handle the missingness in the remaining data. By
159 default, this procedure implements multiple imputations with separate imputation models for
160 each incomplete variable⁴⁷. Predictive mean matching (PMM) approach was employed for
161 imputing continuous variables, which first estimates a linear regression model for the target
162 variable (e.g., Y) from all other variables in the data (e.g., non-Y variables) with complete
163 observations. New coefficients are then drawn from the posterior predictive distribution of the

164 estimated regression coefficients and used to calculate the predicted values for the missing
165 entries in Y. The predictive values for the observed Y are also calculated using the estimated
166 regression coefficients. Finally, a small set of candidate donors is formed from the observed Y
167 (i.e., usually 3 or 5 donors) that have the closest predicted values to the missing Y, and the
168 observed value from one donor will be randomly selected to replace the missing value⁴⁷. Using
169 PMM, 20 iterations was performed for each incomplete variable of mental health, and the final
170 imputed value for any given missing entry was averaged across all iterations.

171 As the current mental health questionnaire covers a broad range of multiple constructs, including
172 depression and anxiety symptoms, neuroticism, and subjective well-being, we performed data
173 decomposition using principal component analysis (PCA) on the imputed data to extract the most
174 relevant information about general psychopathology, using a R package (see details below in
175 section 2.4). We retained the top principal components that collectively explained more than 50%
176 variance of the data in the subsequent statistical analyses. PCA loadings of each individual
177 question per component can be found in *Table 1*.

178

179

180 2.2.2 Physical activity measures

181 Accelerometer data were acquired for a subset of UKB participants during a seven-day
182 monitoring period (<https://biobank.ndph.ox.ac.uk/ukb/label.cgi?id=1008>). This enabled real-time
183 measuring of physical activity for the participants throughout the entire week. Following a
184 recommendation for quality control⁴⁸, data from participants who had less than 72h device
185 wearing time or had no data in each one-hour period of the 24h cycle were excluded (N=64).

186 Using a publicly available machine learning algorithm, we extracted measures of five types of
187 physical activity including sleep, sedentary, walking, light task, and moderate activities⁴⁸. This
188 algorithm applied random forest and hidden Markov models to a 126-dimensional vector that
189 represented a range of time and frequency domain features for every non-overlapping 30-sec
190 epoch. The resulting probability of each physical activity was then defined as the count of
191 predicted activity type per 30-sec epoch divided by the number of epochs⁴⁸. In addition to these
192 probability measures, the average acceleration magnitude, and metabolic equivalents of task
193 (MET) were included to indicate overall activity intensity. The mean values of these features
194 were calculated across weekdays and weekends, respectively, as well as across the entire
195 monitoring period (i.e., average over weekdays and weekends). To account for variation in each

196 individual physical activity measure at different time points, all mean values were standardized
197 by standard deviations for each participant. In total, 21 standardized physical activity measures
198 were included in our analyses.

199

200 2.2.3 Resting-state fMRI data and brain network measure

201 Resting-state fMRI was acquired using a multiband sequence with an acceleration factor of 8
202 (TR=0.735; voxel size=2.4x2.4x2.4mm³). Preprocessing steps included motion correction,
203 grand-mean intensity normalization, high-pass temporal filtering, unwarping and ICA-FIX
204 denoising (Alfaro-Almagro et al., 2018). Full details can be found in UK Biobank Brain Imaging
205 Documentation (https://biobank.ctsu.ox.ac.uk/crystal/crystal/docs/brain_mri.pdf).

206 For this study, we used IDPs (imaging-derived phenotypes) that were generated and released by
207 the UKB⁴⁶. Specifically, partial connectivity matrices and network amplitudes from ICA with
208 dimensionalities of 100 (ICA100) were considered (see *Supplementary Results* for the
209 comparison with ICA25). ICA is a data-driven approach that can estimate the resting-state brain
210 networks reliably and reproducibly, including the three networks of interest. It further helps
211 eliminate noise in the data by separating noise and signal components⁴⁹. These advantages of
212 ICA make it an unbiased and powerful technique to study the resting-state networks. The
213 mapping of the DMN, SN and CEN onto the ICA components and the calculation of partial
214 connectivity matrices and network amplitudes are described below.

215

216 2.2.3.1 Mapping the networks of interest

217 We used the Stanford FIND atlas⁵⁰ to construct canonical spatial maps as the reference to
218 identify the three brain networks of interest, namely the DMN, SN and CEN in our sample.
219 Notably, the FIND atlas is a functional connectivity-based parcellation atlas, from which we
220 selected seven functional parcels that are all part of these three intrinsic, including dorsal (i.e.,
221 posterior cingulate cortex and medial prefrontal cortex) and ventral default mode networks (i.e.,
222 retrosplenial cortex and medial temporal lobe), precuneus network, anterior (i.e., anterior insula
223 and dorsal anterior cingulate cortex) and posterior salience networks (i.e., posterior insula), as
224 well as left and right executive control networks (i.e., dorsolateral prefrontal cortex and parietal
225 cortex in the left and right hemispheres respectively). These selected parcels well represent the
226 subsystems of the three intrinsic networks that have been associated with physical exercises (e.g.,

227 via cognitive and interoceptive processing)^{5,6} and various psychiatric disorders including
228 depression and anxiety⁷⁻⁹. We therefore selected them as our reference networks. To identify
229 ICA components that can be mapped onto these networks of interest, we examined fifty-five
230 signal components that were generated from ICA100 and detected the best matching components
231 corresponding to each of seven selected network parcels, based on spatial correlations. Seven
232 ICA components showing the highest spatial correlations with the FIND atlas were identified
233 (mean $r = 0.37$). We refer to these ICA components as ‘subnetworks’ from the large-scale DMN,
234 SN and CEN as they represented the subsystems of the three intrinsic networks.

235

236 2.2.3.2 Extracting subnetwork edges

237 Dual regression was performed to obtain timeseries for each ICA component or subnetwork at
238 the individual level⁵⁶. These extracted subnetwork timeseries were then used to calculate the
239 partial connectivity matrix and amplitude measures of interest. As described in Miller et al.⁴⁶,
240 pairwise partial correlation coefficient were estimated using L2-regularized partial correlations
241 between all 55 signal components from ICA100, which are deemed to be non-artifactual. This
242 analysis resulted in a 55x55 partial correlation matrix, from which we selected the correlation
243 coefficients corresponding to our seven components of interest (i.e., subnetwork edges). In total,
244 21 subnetwork edges were extracted and included in the subsequent analyses.

245

246 2.2.3.3 Calculating subnetwork amplitudes

247 In addition to pairwise partial correlations between these 7 subnetworks, we further considered
248 the signal amplitude of each subnetwork as the brain network variables. The amplitude is defined
249 as the standard deviation of the ICA component timeseries³⁶. Previous work has shown that
250 amplitudes capture the overall signal fluctuations within each subnetwork and can offer
251 complementary information in relation to behavioral measures. For example, a recent study using
252 the UK Biobank dataset demonstrated independent associations between network amplitudes and
253 behavioral measures in addition to connectivity strength⁴⁶. Thus, we also included seven
254 subnetwork amplitudes in the subsequent analyses.

255

256 2.2.3.4 Overview of resting state measures

257 In summary, subnetwork edges from the partial connectivity matrix indicate the connection
258 strength between each pair of 7 subnetworks while controlling for all other ICA components,
259 whereas subnetwork amplitudes capture the variance of signal changes within each subnetwork.
260 In total, 28 brain measures including 21 partial connectivity strength measures (i.e., subnetwork
261 edges) and 7 amplitudes were included (see the overlaid subnetworks in *Figure 1*; also see
262 subnetwork selection in *Figure SI*). All of these resting-state imaging measures are available
263 from the UK Biobank showcase (bulk field IDs 25753 and 25755).

264

265

266 2.3 *Statistical Analysis*

267 Separate statistical analyses were performed to examine the shared brain network architecture
268 and genetic architecture between mental health and physical activity (see *Figure 1* for an
269 overview for the relevant variables and analyses).

270

271 2.3.1 Shared variance between mental health and physical activity

272 Pearson's correlation was used to identify the shared variance between phenotypes of mental
273 health and physical activity. As we decomposed the data of mental health into principal
274 components, all correlations were performed using the individual-specific component scores. We
275 further calculated the false discovery rate (FDR) to account for multiple testing on all pairwise
276 correlations between mental health and physical activity phenotypes.

277

278 2.3.2 Brain Associations with mental health and physical activity

279 Canonical Correlation Analysis (CCA) has been recognized as a key tool for population
280 neuroimaging that allows for investigating associations between imaging and non-imaging
281 variables⁵⁷. Here in this study, CCA was performed to investigate the associations of brain
282 measures with physical activity and with mental health separately (i.e., simple CCA models).
283 Specifically, CCA finds a linear combination of brain measures that is maximally correlated with
284 a linear combination of mental health or physical activity variables respectively, as defined in Y^*
285 $A = U \sim V = X * B^{58}$ where Y is the set of brain measures, X the set of mental health or physical
286 activity measures, A and B are the linear weights, and U and V the canonical variables or
287 canonical variate pair. The canonical correlation for each pair of canonical variates is defined as

288 the correlation between U and V . Canonical loadings that indicate the shared variance between
289 the original observations and canonical variables are calculated as the correlations between Y and
290 U , or between X and V .

291 To further identify unique brain associations with physical activity and with mental health
292 respectively, variance in brain network data explained by one set of variables was partialled out
293 in the CCA model for the other set of variables (i.e., the unique CCA model included physical
294 activity measures as covariates in the model for assessing brain-mental health associations and
295 vice versa). Confounding variables were included in all four CCA models and statistical
296 inference for CCA results was made via 1,000 permutations (i.e., breaking correspondence of
297 participant identity with brain measures and mental health/physical activity measures), as
298 implemented in the permCCA package⁵⁷. Notably, the CCA model with the largest number of
299 non-confounding variables included 28 brain measures and 21 physical activity measures,
300 resulting in a ratio of approximately 130 observations (i.e., individuals) per feature. This is
301 expected to ensure sufficient stability for our study⁵⁹.

302 To further investigate whether the patterns of brain measures in relation to mental health and
303 physical activity overlap, post-hoc analyses were carried out to test the significance of canonical
304 loadings for each individual brain variable. Specifically, we aimed to determine whether the
305 same brain measures contributed significantly to the canonical associations both with mental
306 health and physical activity, and thus could indicate a shared brain basis. These analyses were
307 conducted only for significant canonical variates within each individual CCA model using
308 permutation testing, where correspondence between brain measures and mental health/physical
309 activity measures for each individual participant is shuffled. Canonical loading for each brain
310 variable was recorded per permutation, which resulted in separate null distributions of loadings
311 for each brain variable. The loadings from the true (unpermuted) CCA were then compared
312 against the matching null distributions for each individual brain variable. Statistical significance
313 was determined as the proportion of permuted loadings equal or higher than the observed
314 loadings from the unpermuted analysis, divided by the total number of 1,000 permutations.
315 These permutation-derived p values were further corrected for the number of significant
316 canonical variates within each model (i.e., record the permuted loadings across canonical
317 variates). To compare brain variable patterns across different CCA models, we matched the first
318 significant canonical variates from each model based on the correlations between the canonical

319 variate for the brain measures (i.e., correlating the vector U obtained from the mental health
320 CCA models with the vector U obtained from the physical activity CCA models).

321 To characterize the individual mental health questions and physical activity types in relation to
322 the tested brain associations, we further examined the loading patterns of each individual
323 question and physical activity type for the first canonical variate from all models, without testing
324 for statistical significance.

325

326 2.3.3 Shared genetic architecture between mental health and physical activity

327 Genetic correlations between mental health and physical activity were examined by leveraging
328 GWAS summary statistics for the relevant phenotypes.

329

330 2.3.3.1 Summary statistics

331 The mental health questionnaire used in this study includes items that measure neuroticism,
332 anxiety, subjective well-being, depression, and risk taking. We therefore sought to obtain
333 summary statistics for these psychopathological phenotypes. First, we extracted summary
334 statistics for *Neuroticism* from a GWAS meta-analysis of self-reported neuroticism in the UKB
335 (using the same questions as in our study) and Psychiatric Genetics Consortium (using the NEO-
336 FFI personality inventory)⁶⁰. For *Generalized Anxiety Disorder*, we leveraged summary statistics
337 from a GWAS of self-reported Generalized Anxiety Disorder 2-item scale scores in the Million
338 Veteran Program⁶¹. We further obtained summary statistics for *Subjective Well-Being* from a
339 GWAS meta-analysis of life satisfaction, positive affect, or both life satisfaction and positive
340 affect across 59 cohorts⁶². For *Major Depressive Disorder*, we meta-analyzed summary statistics
341 from case-control GWAS in the UK Biobank and Psychiatric Genomics Consortium⁶³ and the
342 Million Veteran Program⁶⁴ (see *Supplementary Results* for further details). Lastly, we obtained
343 summary statistics for *Risk Taking* from a GWAS study using the UKB data, which included the
344 same question of risk taking as in our study⁶⁵. Although these psychopathological phenotypes
345 obtained from the independent GWAS studies were not directly equivalent to the phenotypes
346 derived from mental health questionnaire in our brain association analyses due to different
347 measurements, the underlying constructs of depression, anxiety disorders, and neuroticism,
348 subjective well-being and risk-taking are identical. The phenotypes obtained to conduct genetic
349 association analyses are therefore similar to those in the brain association analyses.

350 Summary statistics for accelerometer data-derived physical activity phenotypes, including
351 moderate activity, overall activity, sedentary activity, walking, and sleep duration were derived
352 from a GWAS of N=91,105 participants of European ancestry in the UK Biobank⁴⁸. The exact
353 phenotypes were also included in our brain association analyses.

354 Please refer to *Table 2* for an overview for all the summary statistics used in this study, including
355 sample size and SNP heritability.

356

357 2.3.3.2 Genetic correlations

358 We used linkage disequilibrium score regression (LDSR) and genomic structural equation
359 modeling (gSEM) to test whether the genomic architecture associated with general mental
360 health is shared with physical activity. LDSR leverages GWAS summary statistics to estimate
361 genetic correlations by regressing the SNP statistics on the SNP linkage disequilibrium (LD)
362 scores, or correlations between nearby genomic loci due to population stratification (i.e.,
363 systematic differences in allele frequencies due to differences in ancestry). gSEM characterizes
364 the latent genetic architecture across phenotypes based on the LDSR-derived genetic correlation
365 matrices⁶⁶. To this end, we first applied LDSR to existing GWAS summary statistics of
366 psychopathological phenotypes (i.e., neuroticism, generalized anxiety disorder, subjective well-
367 being, major depressive disorder, and risk taking) and physical activity phenotypes (i.e., overall
368 activity, moderate activity, sedentary activity, sleep duration, and walking), respectively, to
369 estimate pairwise genetic correlations within each construct (i.e., within mental health and within
370 physical activity respectively). We also examined genetic correlations between mental health
371 phenotypes and physical activity phenotypes adjusted for sex and BMI. We then applied gSEM
372 to the covariance matrix of psychopathology and that of physical activity separately, allowing
373 one single latent factor to load freely within each model. Metrics indicating model fit (i.e., CFI,
374 comparative fit index; SRMR, standardized root mean squared residual) and factor loadings from
375 each of these models were used to determine whether one common genetic factor fit the physical
376 activity and mental health data well, respectively.

377 Because our results indicated poor model fit for some gSEM analyses (see details below in
378 Results section 3.3), we focused on the model of mental health, where “risk-taking” was
379 excluded to generate a latent factor of “negative affect” across other phenotypes. Specifically, we
380 explored genetic correlations between the latent factor of “negative affect” (i.e., without risk-

381 taking) and each of the five physical activity phenotypes. In addition, we examined genetic
382 correlations between “risk-taking” alone and each individual physical activity phenotype, using
383 LDSR. False discovery rate correction was used to correct for multiple testing (N=10 tests). We
384 also repeated analyses that returned significant results, with adjustment for sex and BMI.
385 Adjusted summary statistics from Doherty and colleagues (2018) were used in these analyses.

386

387 2.3.3.3 Causal relationships

388 To examine plausible causal associations between physical activity phenotypes and negative
389 affect, we conducted Latent Causal Variable Analysis (LCV). This approach finds a latent
390 variable that mediates the genetic correlation between two traits, such as negative affect and
391 sleep duration. Generally, if the latent variable has a stronger genetic correlation with trait 1 (e.g.,
392 sleep duration) than with trait 2 (e.g., negative affect), part of the genetic component of trait 1 is
393 thought to be causal for trait 2. This partial causality can be quantified using the genetic causality
394 proportion (GCP) of trait 1 (sleep duration) on trait 2 (negative affect), which can range between
395 0 (no partial genetic causality) and 1 (full genetic causality)⁶⁷. The advantages of using LCV
396 over the more traditional Mendelian Randomization for causal inference include increased power
397 by leveraging SNPs across the genome and less susceptibility to confounding by horizontal
398 pleiotropy⁶⁷.

399

400 2.4 *Confounding variables*

401 Based on the literature, BMI, smoking and drinking status were included as confounding factors
402 in all statistical analyses for brain associations^{68,69}. Age, sex, head motion during rs-fMRI
403 acquisition (i.e., mean frame-wise displacement), time difference in days between accelerometer
404 recording (i.e., start date) and assessment center visit date (i.e., acquisition date for both mental
405 health questionnaire and rs-fMRI), as well as the scanning site were further included. Due to
406 varying degrees of missingness in the confound variables (i.e., up to 22%), the same imputation
407 procedure, as described in *Section 2.2.2*, was performed except that additional usage of
408 multinomial logistic regression was employed to impute categorical data with more than two
409 levels (i.e., smoking and drinking status). Complete observations from all variables including the
410 confounding variables were used in the imputation procedure. When calculating the final
411 predicted values to replace missing data, the predicted values for continuous variables were

412 averaged across 20 iterations and the level with the highest count across iterations was selected.
413 All categorical variables after imputation were dummy coded for subsequent statistical analyses.
414 In the genetic correlation analyses, to ensure robust test effects, we repeated models with
415 significant results including sex and BMI as covariates.

416

417

418 2.5 *Data and Code Availability*

419 The UK Biobank data used in this study can be accessed by researchers upon application
420 (<https://www.ukbiobank.ac.uk/register-apply>). A machine learning algorithm shared on github
421 was used to extract physical activity measures
422 (<https://github.com/activityMonitoring/biobankAccelerometerAnalysis>). The Matlab code for
423 running permutation inference for CCA are also available on github, including the new extension
424 for testing canonical variable loadings (<https://github.com/andersonwinkler/PermCCA>). We
425 performed Pearson's correlations and imputations in R (version 4.1.0)⁷⁰, using basic *stat*
426 function and *MICE* package⁴⁷, respectively. Genetic analyses including LDSR and gSEM were
427 performed using the GenomicSEM package also in R⁶⁶, with code available on github
428 (<https://github.com/GenomicSEM/GenomicSEM>). LCV was performed using the open-source
429 LCV software (<https://github.com/lukejconnor/LCV>). We also shared the derived data from
430 statistical analyses and code for producing the figures on the Open Science Framework
431 (<https://osf.io/p3fzv/>). To note, the GWAS summary statistics used in the current study are hosted
432 elsewhere, for which we only shared the downloading links.

433

434

435 **3. Results (1,054 words)**

436 *3.1 Correlations between mental health and physical activity measures*

437 In this sample, phenotypic measures of mental health and physical activity demonstrated overall
438 small but significant correlations, with coefficients ranging between -0.08 and 0.11 (FDR-
439 corrected p -values ≤ 0.045). As shown in *Table 1*, the mental health measures (i.e., principal
440 component scores of the seven principal components) were broadly associated with all physical
441 activity measures. In particular, sleep showed the strongest association with the first principal
442 component (PC1) of mental health such that greater sleep duration was associated with poorer
443 overall mental health (i.e., loadings of PC1 were mostly negative as shown in Table S1) and
444 walking had the greatest association with the fifth principal component (PC5) of mental health
445 such that higher depression scores was linked to reduced walking time. The overall coefficient
446 patterns were largely consistent for correlations between physical activity and mental health
447 measures despite of minor differences across different time windows (i.e., overall, weekdays,
448 weekend; see full correlations in *Table 1*).

449

450

451 *3.2 Shared neural correlates of mental health and physical activity*

452 *3.2.1 Brain associations with mental health and physical activity*

453 In simple CCA models (i.e., without accounting for the potentially shared variance in brain
454 measures between physical activity and mental health), 2 and 3 significant canonical variate pairs
455 were observed respectively for brain-mental health ($r_1=0.16$, $p_1=0.001$; $r_2=0.12$, $p_2=0.001$) and
456 brain-physical activity associations ($r_1=0.23$, $p_1=0.001$; $r_2=0.15$, $p_2=0.001$; $r_3=0.13$, $p_3=0.047$).
457 When controlling for the shared variance in brain measures between mental health and physical
458 activity (i.e., unique models), we found 2 significant canonical variates for both brain-mental
459 health ($r_1=0.14$, $p_1=0.001$; $r_2=0.12$, $p_2=0.002$) and brain-physical activity associations ($r_1=0.21$,
460 $p_1=0.001$; $r_2=0.15$, $p_2=0.001$), with slightly decreased canonical correlation coefficients
461 comparing to those from the simple models. The canonical variates for brain measures after
462 accounting for the shared variance (i.e., from the unique models) mapped well with those from
463 the simple models, as indicated by correlation coefficients between brain canonical variables (i.e.,
464 \square ; all r 's > 0.96 ; see full results in *Table S2*). These results are in line with our expectations that
465 both mental health and physical activity are closely associated with the functional networks
466 under investigation and that mental health and physical activity have shared variance in the

467 functioning of these networks, as indicated by the reduced canonical correlation coefficients
468 from the unique models.

469

470 3.2.2 Canonical loadings of brain measures

471 Findings from the post-hoc analyses on the significant canonical variates showed considerable
472 overlap between the brain measures with significant loadings associated with mental health and
473 with physical activity. Overall, the amplitude of subnetworks that indicates the overall signal
474 fluctuations in each subnetwork over time loaded higher than the edges, with the highest
475 loadings on the amplitude of dorsal DMN or left CEN for the first canonical variates in both the
476 mental health and physical activity models (*permuted p-values* ≤ 0.001 ; *Figure 2*). This
477 suggested that the magnitude of fluctuations of intra-network signal (i.e., amplitude) had higher
478 contributions to the observed canonical associations with both mental health and physical
479 activity than the inter-network connectivity strength. The similar loading pattern of all brain
480 measures were largely retained even when the shared variance was partialled out for mental
481 health and for physical activity, respectively, in the unique models (*permuted p-values* ≤ 0.035).
482 In addition to amplitude, connectivity between the dorsal DMN and left CEN (i.e., subnetwork
483 edge) also exhibited statistically significant loadings in the first canonical variates of both the
484 mental health and the physical activity models (simple and unique models; *permuted p-values* \leq
485 0.001 ; *Figure 2*). The brain variable loadings for the second canonical variate were highly
486 similar between simple and unique models for either the mental health or physical activity
487 models, with the amplitude and subnetwork edges showing evenly important involvement. Yet,
488 the patterns of these loadings differed between the mental health and physical activity model
489 (*Figure S2*).

490

491 3.2.3 Canonical loadings of mental health questions and physical activities

492 Canonical loadings of both individual mental health questions and physical activity types for the
493 first canonical variate also exhibited similar patterns between the simple and unique models
494 (*Figure S3*). In the models for brain-mental health associations, “risk taking” and “ever irritable/
495 argumentative for 2 days” had the highest loadings in both simple and unique models, whereas in
496 the models for brain-physical activity associations, “walking” in all time windows (i.e., overall,
497 weekdays, weekend) showed the greatest importance across models. Interestingly, among all

498 physical activity types, only “sleep” showed the opposite direction in canonical loadings (*Figure*
499 *S3*).

500

501 3.3 Genetic correlations

502 3.3.1 Genetic correlations for individual phenotypes

503 Pairwise LDSR was performed separately for psychopathological phenotypes and physical
504 activity phenotypes. All results were significant after controlling for multiple comparisons,
505 except for the genetic correlations between neuroticism and risk-taking ($rg = 0.039$, $p = 0.075$),
506 and between subjective well-being and risk-taking ($rg = 0.054$, $p = 0.146$; *Figure 3*). Specifically,
507 among psychopathological phenotypes, the smallest significant genetic correlation was observed
508 for generalized anxiety disorder and risk-taking ($rg = 0.151$, $p = 2.69e-05$, *FDR-corrected p-*
509 *value = 3.36e-05*), and the largest effect was observed for MDD and generalized anxiety disorder
510 ($rg = 0.768$, $p = 2.62e-90$, *FDR-corrected p-value = 1.31e-89*). For physical activity, genetic
511 correlations ranged from -0.217 ($p = 1.03e-04$, *FDR-corrected p-value = 1.03e-04*) for sleep
512 duration and walking to 0.796 ($p = 1.77e-31$, *FDR-corrected p-value = 8.85e-31*) for moderate
513 and overall activity (*Figure 3*). Physical activity phenotypes adjusted for sex and BMI largely
514 recapitulated these results, with genetic correlations ranging from -0.209 ($p = 2.49e-04$, *FDR-*
515 *corrected p-value = 2.49e-04*) for sleep duration and walking, to 0.780 ($p = 2.12e-23$, *FDR-*
516 *corrected p-value = 1.06e-22*) for moderate and overall activities.

517

518 3.3.2 Genetic correlations for latent factors

519 Analyses using gSEM across all mental health phenotypes returned results indicating suboptimal
520 estimations with a low factor loading on risk taking from the model for a general
521 psychopathology latent factor (standardized loading = 0.065), which likely reflected its
522 conceptually distinct construct from all other phenotypes (i.e., depression, anxiety, neuroticism,
523 and subjective well-being). Additionally, initial gSEM results for physical activity phenotypes
524 indicated overall poor model fit (CFI = 0.569, SRMR = 0.160; *Figure S4*). Thus, we performed
525 gSEM for psychopathological phenotypes grouped as one “negative affect” latent factor after
526 excluding “risk-taking”, with an effective sample size of 571,170 and good model fit (CFI =
527 0.987, SRMR = 0.0535; *Figure S5*). After correction for multiple testing, genetic variance in
528 negative affect was significantly and positively correlated with sleep duration ($rg = 0.121$, $p =$

529 1.25e-05, *FDR-corrected* $p = 1.25e-04$), and negatively correlated with moderate ($rg = -0.117$, p
530 $= 1.33e-03$, *FDR-corrected* $p = 4.43e-03$) and overall activity ($rg = -0.085$, $p = 7.70e-04$, *FDR-*
531 *corrected* $p = 3.85e-03$; *Figure 4*; also see the results for a latent negative affect factor including
532 “risk-taking” [in the *Supplementary Results*). No significant genetic correlation was observed
533 between negative affect and sedentary activity or walking after multiple comparison corrections
534 ($|rgs| \leq 0.062$, p -values ≥ 0.036 , *FDR-corrected* p -values ≥ 0.060 ; *Table 3*). Post-hoc analyses
535 revealed that after adjustment for sex and BMI, only the genetic correlation between negative
536 affect and sleep duration remained significant ($rg = 0.122$, $p = 9.38e-06$). Using LDSR, risk-
537 taking was genetically correlated with both overall activity and walking ($|rgs| \geq 0.074$, $ps \leq 0.020$,
538 $psFDR \leq 0.040$) but not moderate or sedentary activity or sleep duration ($|rgs| \leq 0.081$, $ps \geq$
539 0.092).

540

541 3.3.3 Latent Causal Variable Analysis.

542 Results from the LCV analyses did not indicate causal relationships between any physical
543 activity phenotypes and negative affect, in either direction ($|GCPs| < 0.425$; $ps > 0.163$).

544

545

546

547

548 **4. Discussion (1,107 words)**

549 In this study, we investigated whether mental health and physical activity have shared brain and
550 genetic architectures using the UK Biobank cohort. Our findings showed significant associations
551 of mental health and physical activity separately with a set of brain measures that represent the
552 connectivity strength and amplitude of subnetworks from the DMN, SN and CEN. Critically,
553 these significant associations exhibited highly similar patterns of brain variable loadings across
554 mental health and physical activity models even when the shared variance between these two
555 constructs was accounted for, suggesting a potential overlap in brain network architecture
556 between these two constructs. Further analyses examining genetic correlations for mental health
557 and physical activity showed that negative affect exhibited significant genetic correlations with
558 several physical activity types, of which sleep duration demonstrated the strongest genetic
559 correlation that remained significant after controlling for BMI and sex effects. Together, these
560 results support the presence of shared multivariate brain and genetic architectures between
561 mental health and physical activity.

562 The three intrinsic brain networks, namely the DMN, SN and CEN, have been consistently
563 implicated in a wide range of psychiatric disorders including major depression and anxiety¹⁰.
564 Interestingly, connections between or the configurations of these networks have also been
565 associated with physical exercises^{71,72}. The current study therefore focused specifically on the
566 subnetworks from these large-scale networks and used the amplitude and connectivity strength
567 (i.e., subnetwork edge) to examine the associations of these networks with mental health and
568 with physical activity, respectively. In line with the literature, we observed significant
569 multivariate associations for all three networks with either mental health or physical activity, and
570 significant loadings on most of the subnetworks. In particular, the dorsal DMN and left CEN
571 showed the greatest involvement in the observed brain associations with both mental health and
572 physical activity (*Figure 2*). In this study, the dorsal DMN subnetwork primarily consisted of the
573 posterior cingulate cortex (PCC) and the ventromedial prefrontal cortex (vmPFC), the two brain
574 areas that are commonly considered as the core subsystem of the DMN⁷³. Similarly, the
575 subnetwork of the CEN here included two most typical hub regions: the dorsolateral prefrontal
576 cortex (dlPFC) and posterior parietal cortex (PPC) for each individual hemisphere (*Figure S1*).
577 These major hubs of the DMN and CEN have been implicated in various mental illnesses
578 including depression and anxiety. For example, the PCC and mPFC have been suggested to

579 collectively support multiple cognitive functions including decision making and memory⁷⁴, the
580 impairment of which has often been reported in patients with major depression and anxiety
581 disorders⁷⁵⁻⁷⁷. Additionally, the dlPFC is known to be involved in emotion regulation⁷⁸⁻⁸⁰ and
582 dysfunction of this region is often seen in abnormal processing of emotional experiences in
583 patients with depressive and anxiety symptoms^{81,82}. As for physical activity, increased dlPFC
584 activity has been observed after acute physical exercises in participants with higher scores in the
585 Stroop test⁸³, whereas the involvement of the DMN subsystems in the medial temporal lobe
586 (MTL) including the hippocampus and its connection with the medial PFC are often observed in
587 relation to enhanced memory after physical exercises^{20,84,85}. In our findings, the amplitude of
588 dorsal DMN and left CEN that indicated the magnitude of fluctuations of intra-network signal
589 (i.e., variance in the connections between PCC and vmPFC or between dlPFC and PCC), as well
590 as the connectivity strength between these two subnetworks, showed significantly high loadings
591 for the most critical association between brain measures and mental health measures (i.e., the
592 first canonical variate). These observations are in line with separate literature on mental health
593 and physical activity, and provide empirical evidence that mental health and physical activity
594 may share brain architecture involved in major cognitive functions.

595 Interestingly, mental health and physical activity also appear to have partially overlapping
596 genetic architectures. In line with previous reports that internalizing problems are negatively
597 genetically correlated with physical movement⁴⁵, we showed that a latent negative affect factor
598 capturing genetic covariance between subjective well-being, neuroticism, major depressive
599 disorder, and generalized anxiety disorder was negatively genetically correlated with overall
600 physical activity as well as a more fine-grained phenotype of moderate activity, and positively
601 genetically correlated with sleep duration. Protective effects of physical activity on mental health
602 have long been documented, as have negative health consequences of psychiatric disorders^{5-7,86}.

603 Here, we demonstrate that these relationships can be partially explained by shared genetic
604 predisposition, although results from our latent causal variable analysis indicate that these
605 associations do not reflect causal influences. Interestingly, with the adjustment of sex and BMI,
606 only the correlation with sleep duration remained significant. Sleep duration also showed the
607 lowest loading onto a latent physical activity genetic factor (*Figure S4*). These results suggest
608 that the sleep phenotype is somewhat distinct from the remaining physical activity phenotypes
609 and that the shared genetic architecture between negative affect and sleep duration is more

610 pronounced than that between negative affect and the degree of daily physical or sedentary
611 activity. This is in line with the frequent documentation of symptomatic sleep disturbances
612 across forms of psychopathology, including depression and anxiety^{87,88}, even in children⁸⁹.
613 Specifically, this aligns well with the literature showing reduced sleep duration in older adults
614 with depression and anxiety disorders⁹⁰⁻⁹².

615 Despite being the first to jointly investigate the shared brain network architecture and genetic
616 basis of mental health and physical activity in a large population cohort, our study has some
617 limitations. First, the brain measures in our study are derived from resting-state fMRI measures.
618 Although our choice reflected a rich literature that has implicated these measures in both mental
619 health and physical activity, addressing smaller-scale brain structures (e.g., specific regions) with
620 relevant hypotheses and inclusion of multimodal brain measures such as structural gray matter
621 volume, cortical thickness, and white matter integrity can provide complementary insights into
622 brain architecture in relation to mental health and physical activity, and thus may be of interest
623 for future investigations. Second, although the accelerometer recording took place prior to the
624 acquisition of resting-state fMRI and mental health assessment for most of the participants in our
625 study (i.e., 96%), the degrees of time difference between these measurements varied greatly at
626 the individual level (i.e., ranging between -473 to 2281 days). This time discrepancy was
627 accounted for in all CCA models as a covariate and cautions should be taken when interpreting
628 the observed brain associations with reference to time effects. It should also be noted that the
629 mean age of the current sample is relatively high as the UKB cohort comprises predominantly
630 middle-to-late aged individuals. Our findings therefore should be interpreted carefully in the
631 relevant context. Lastly, socioeconomic variables including education attainment can be relevant
632 to mental health and physical activity phenotypes and inclusion of these variables as
633 confounding factors may be considered in future investigations.

634

635 In conclusion, our study jointly analyzed resting-state network measures and genetic correlations
636 in a large cohort to test the hypothesis of a shared neurobiological basis of mental health and
637 physical activity. Our findings revealed that multivariate patterns of brain correlates were highly
638 similar between mental health and physical activity and highlighted genetic correlations between
639 mental health (negative affect) and overall physical activity, moderate activity levels, and sleep

640 duration. Taken together, these findings point towards neural and genetic mechanisms that may
641 subserve the protective influence of physical exercise and sleep on mental health.

642

643 **Acknowledgements**

644 JDB is supported by the NIH (1 R34 NS118618-01) and the McDonnell Center for Systems

645 Neuroscience. SEP is supported by the NIH (1 F31 AA029934-01).

646

647 **Conflict of Interest**

648 The authors report no biomedical financial interests or potential conflicts of interest.

649

650

651

652

653

654

655

656

657

658

659

660

661 **Reference**

- 662 1 Zhang Z, Chen W. A Systematic Review of the Relationship Between Physical Activity and
663 Happiness. *J. Happiness Stud.* 2019. doi:10.1007/s10902-018-9976-0.
- 664 2 Fernandes J, Arida RM, Gomez-Pinilla F. Physical exercise as an epigenetic modulator of brain
665 plasticity and cognition. *Neurosci. Biobehav. Rev.* 2017. doi:10.1016/j.neubiorev.2017.06.012.
- 666 3 Kays JL, Hurley RA, Taber KH. The dynamic brain: Neuroplasticity and mental health. *J*
667 *Neuropsychiatry Clin Neurosci* 2012. doi:10.1176/appi.neuropsych.12050109.
- 668 4 Millan MJ, Agid Y, Brüne M, Bullmore ET, Carter CS, Clayton NS *et al.* Cognitive dysfunction in
669 psychiatric disorders: Characteristics, causes and the quest for improved therapy. *Nat. Rev. Drug*
670 *Discov.* 2012. doi:10.1038/nrd3628.
- 671 5 Biddle S. Physical activity and mental health: Evidence is growing. *World Psychiatry.* 2016.
672 doi:10.1002/wps.20331.
- 673 6 Deslandes A, Moraes H, Ferreira C, Veiga H, Silveira H, Mouta R *et al.* Exercise and mental
674 health: Many reasons to move. *Neuropsychobiology.* 2009. doi:10.1159/000223730.
- 675 7 Heinze K, Cumming J, Dosanjh A, Palin S, Poulton S, Bagshaw AP *et al.* Neurobiological
676 evidence of longer-term physical activity interventions on mental health outcomes and cognition
677 in young people: A systematic review of randomised controlled trials. *Neurosci. Biobehav. Rev.*
678 2021. doi:10.1016/j.neubiorev.2020.10.014.
- 679 8 Sudlow C, Gallacher J, Allen N, Beral V, Burton P, Danesh J *et al.* UK Biobank: An Open Access
680 Resource for Identifying the Causes of a Wide Range of Complex Diseases of Middle and Old
681 Age. *PLoS Med* 2015. doi:10.1371/journal.pmed.1001779.
- 682 9 Woodward ND, Cascio CJ. Resting-state functional connectivity in psychiatric disorders. *JAMA*
683 *Psychiatry* 2015; **72**: 743–744.
- 684 10 Menon V. Large-scale brain networks and psychopathology: A unifying triple network model.
685 *Trends Cogn Sci* 2011; **15**: 483–506.
- 686 11 Tozzi L, Zhang X, Chesnut M, Holt-Gosselin B, Ramirez CA, Williams LM. Reduced functional
687 connectivity of default mode network subsystems in depression: Meta-analytic evidence and
688 relationship with trait rumination. *NeuroImage Clin* 2021. doi:10.1016/j.nicl.2021.102570.
- 689 12 Kaiser RH, Andrews-Hanna JR, Wager TD, Pizzagalli DA. Large-Scale Network Dysfunction in
690 Major Depressive Disorder. *JAMA Psychiatry* 2015. doi:10.1001/jamapsychiatry.2015.0071.
- 691 13 Xu J, Van Dam NT, Feng C, Luo Y, Ai H, Gu R *et al.* Anxious brain networks: A coordinate-
692 based activation likelihood estimation meta-analysis of resting-state functional connectivity
693 studies in anxiety. *Neurosci. Biobehav. Rev.* 2019. doi:10.1016/j.neubiorev.2018.11.005.
- 694 14 Servaas MN, Geerligs L, Renken RJ, Marsman JBC, Ormel J, Riese H *et al.* Connectomics and
695 neuroticism: An altered functional network organization. *Neuropsychopharmacology* 2015.
696 doi:10.1038/npp.2014.169.
- 697 15 Colcombe SJ, Kramer AF, Erickson KI, Scalf P, McAuley E, Cohen NJ *et al.* Cardiovascular
698 fitness, cortical plasticity, and aging. *Proc Natl Acad Sci U S A* 2004.
699 doi:10.1073/pnas.0400266101.
- 700 16 Herold F, Aye N, Lehmann N, Taubert M, Müller NG. The contribution of functional magnetic
701 resonance imaging to the understanding of the effects of acute physical exercise on cognition.
702 *Brain Sci* 2020. doi:10.3390/brainsci10030175.
- 703 17 Ikuta T, Loprinzi PD. Association of cardiorespiratory fitness on interhemispheric hippocampal
704 and parahippocampal functional connectivity. *Eur J Neurosci* 2019. doi:10.1111/ejn.14366.
- 705 18 Talukdar T, Nikolaidis A, Zwilling CE, Paul EJ, Hillman CH, Cohen NJ *et al.* Aerobic fitness
706 explains individual differences in the functional brain connectome of healthy young adults. *Cereb*
707 *Cortex* 2018. doi:10.1093/cercor/bhx232.
- 708 19 Weng TB, Pierce GL, Darling WG, Falk D, Magnotta VA, Voss MW. The Acute Effects of
709 Aerobic Exercise on the Functional Connectivity of Human Brain Networks. *Brain Plast* 2016.
710 doi:10.3233/bpl-160039.

- 711 20 Voss MW, Prakash RS, Erickson KI, Basak C, Chaddock L, Kim JS *et al.* Plasticity of brain
712 networks in a randomized intervention trial of exercise training in older adults. *Front Aging*
713 *Neurosci* 2010. doi:10.3389/fnagi.2010.00032.
- 714 21 Greeley B, Chau B, Jones CB, Neva JL, Kraeutner SN, Campbell KL *et al.* Multiple bouts of high-
715 intensity interval exercise reverse age-related functional connectivity disruptions without affecting
716 motor learning in older adults. *Sci Rep* 2021. doi:10.1038/s41598-021-96333-4.
- 717 22 Schmitt A, Upadhyay N, Martin JA, Rojas S, Strüder HK, Boecker H. Modulation of Distinct
718 Intrinsic Resting State Brain Networks by Acute Exercise Bouts of Differing Intensity. *Brain Plast*
719 2019. doi:10.3233/bpl-190081.
- 720 23 Voss MW, Weng TB, Narayana-Kumanan K, Cole RC, Wharff C, Reist L *et al.* Acute exercise
721 effects predict training change in cognition and connectivity. *Med Sci Sports Exerc* 2020.
722 doi:10.1249/MSS.0000000000002115.
- 723 24 Nutt DJ, Wilson S, Paterson L. Sleep disorders as core symptoms of depression. *Dialogues Clin*
724 *Neurosci* 2008. doi:10.31887/dcms.2008.10.3/dnutt.
- 725 25 Riemann D, Krone LB, Wulff K, Nissen C. Sleep, insomnia, and depression.
726 *Neuropsychopharmacology*. 2020. doi:10.1038/s41386-019-0411-y.
- 727 26 Cox RC, Olatunji BO. Sleep in the anxiety-related disorders: A meta-analysis of subjective and
728 objective research. *Sleep Med. Rev.* 2020. doi:10.1016/j.smrv.2020.101282.
- 729 27 Mellman TA. Sleep and Anxiety Disorders. *Psychiatr. Clin. North Am.* 2006.
730 doi:10.1016/j.psc.2006.08.005.
- 731 28 Rumble ME, White KH, Benca RM. Sleep Disturbances in Mood Disorders. *Psychiatr. Clin. North*
732 *Am.* 2015. doi:10.1016/j.psc.2015.07.006.
- 733 29 Chennaoui M, Arnal PJ, Sauvet F, Léger D. Sleep and exercise: A reciprocal issue? *Sleep Med.*
734 *Rev.* 2015. doi:10.1016/j.smrv.2014.06.008.
- 735 30 Wainberg M, Jones SE, Beaupre LM, Hill SL, Felsky D, Rivas MA *et al.* Association of
736 accelerometer-derived sleep measures with lifetime psychiatric diagnoses: A cross-sectional study
737 of 89,205 participants from the UK Biobank. *PLOS Med* 2021; **18**: e1003782.
- 738 31 Full KM, Malhotra A, Gallo LC, Kerr J, Arredondo EM, Natarajan L *et al.* Accelerometer-
739 Measured Sleep Duration and Clinical Cardiovascular Risk Factor Scores in Older Women.
740 *Journals Gerontol - Ser A Biol Sci Med Sci* 2020. doi:10.1093/gerona/glz201.
- 741 32 De Havas JA, Parimal S, Soon CS, Chee MWL. Sleep deprivation reduces default mode network
742 connectivity and anti-correlation during rest and task performance. *Neuroimage* 2012.
743 doi:10.1016/j.neuroimage.2011.08.026.
- 744 33 Sämann PG, Tully C, Spoormaker VI, Wetter TC, Holsboer F, Wehrle R *et al.* Increased sleep
745 pressure reduces resting state functional connectivity. *Magn Reson Mater Physics, Biol Med* 2010.
746 doi:10.1007/s10334-010-0213-z.
- 747 34 Yeo BTT, Tandi J, Chee MWL. Functional connectivity during rested wakefulness predicts
748 vulnerability to sleep deprivation. *Neuroimage* 2015. doi:10.1016/j.neuroimage.2015.02.018.
- 749 35 Khalsa S, Mayhew SD, Przewdzik I, Wilson R, Hale J, Goldstone A *et al.* Variability in cumulative
750 habitual sleep duration predicts waking functional connectivity. *Sleep* 2016.
751 doi:10.5665/sleep.5324.
- 752 36 Bijsterbosch J, Harrison S, Duff E, Alfaro-Almagro F, Woolrich M, Smith S. Investigations into
753 within- and between-subject resting-state amplitude variations. *Neuroimage* 2017.
754 doi:10.1016/j.neuroimage.2017.07.014.
- 755 37 Anderson E, Shivakumar G. Effects of exercise and physical activity on anxiety. *Front Psychiatry*
756 2013. doi:10.3389/fpsy.2013.00027.
- 757 38 Blumenthal JA, Babyak MA, Doraiswamy PM, Watkins L, Hoffman BM, Barbour KA *et al.*
758 Exercise and pharmacotherapy in the treatment of major depressive disorder. *Psychosom Med*
759 2007. doi:10.1097/PSY.0b013e318148c19a.
- 760 39 Choi KW, Zheutlin AB, Karlson RA, Wang MJ, Dunn EC, Stein MB *et al.* Physical activity
761 offsets genetic risk for incident depression assessed via electronic health records in a biobank

- 762 cohort study. *Depress Anxiety* 2020. doi:10.1002/da.22967.
- 763 40 Dennison CA, Legge SE, Bracher-Smith M, Menzies G, Escott-Price V, Smith DJ *et al.*
- 764 Association of genetic liability for psychiatric disorders with accelerometer-assessed physical
- 765 activity in the UK Biobank. *PLoS One* 2021. doi:10.1371/journal.pone.0249189.
- 766 41 Anttila V, Bulik-Sullivan B, Finucane HK, Walters RK, Bras J, Duncan L *et al.* Analysis of shared
- 767 heritability in common disorders of the brain. *Science (80-)* 2018. doi:10.1126/science.aap8757.
- 768 42 Lee SH, Ripke S, Neale BM, Faraone S V., Purcell SM, Perlis RH *et al.* Genetic relationship
- 769 between five psychiatric disorders estimated from genome-wide SNPs. *Nat Genet* 2013.
- 770 doi:10.1038/ng.2711.
- 771 43 Taylor MJ, Martin J, Lu Y, Brikell I, Lundström S, Larsson H *et al.* Association of Genetic Risk
- 772 Factors for Psychiatric Disorders and Traits of These Disorders in a Swedish Population Twin
- 773 Sample. *JAMA Psychiatry* 2019. doi:10.1001/jamapsychiatry.2018.3652.
- 774 44 Smoller JW, Andreassen OA, Edenberg HJ, Faraone S V., Glatt SJ, Kendler KS. Psychiatric
- 775 genetics and the structure of psychopathology. *Mol. Psychiatry*. 2019. doi:10.1038/s41380-017-
- 776 0010-4.
- 777 45 Grotzinger AD, Mallard TT, Akingbuwa WA, Ip HF, Adams MJ, Lewis CM *et al.* Genetic
- 778 architecture of 11 major psychiatric disorders at biobehavioral, functional genomic, and molecular
- 779 genetic levels of analysis. medRxiv. 2020. doi:10.1101/2020.09.22.20196089.
- 780 46 Miller KL, Alfaro-Almagro F, Bangerter NK, Thomas DL, Yacoub E, Xu J *et al.* Multimodal
- 781 population brain imaging in the UK Biobank prospective epidemiological study. *Nat Neurosci*
- 782 2016. doi:10.1038/nn.4393.
- 783 47 van Buuren S, Groothuis-Oudshoorn K. mice: Multivariate imputation by chained equations in R.
- 784 *J Stat Softw* 2011. doi:10.18637/jss.v045.i03.
- 785 48 Doherty A, Smith-Byrne K, Ferreira T, Holmes M V., Holmes C, Pulit SL *et al.* GWAS identifies
- 786 14 loci for device-measured physical activity and sleep duration. *Nat Commun* 2018; **9**.
- 787 doi:10.1038/s41467-018-07743-4.
- 788 49 Griffanti L, Douaud G, Bijsterbosch J, Evangelisti S, Alfaro-Almagro F, Glasser MF *et al.* Hand
- 789 classification of fMRI ICA noise components. *Neuroimage* 2017.
- 790 doi:10.1016/j.neuroimage.2016.12.036.
- 791 50 Shirer WR, Ryali S, Rykhlevskaia E, Menon V, Greicius MD. Decoding Subject-Driven Cognitive
- 792 States with Whole-Brain Connectivity Patterns. *Cereb Cortex* 2012; **22**: 158–165.
- 793 51 Voss MW, Nagamatsu LS, Liu-Ambrose T, Kramer AF. Exercise, brain, and cognition across the
- 794 life span. *J. Appl. Physiol.* 2011. doi:10.1152/jappphysiol.00210.2011.
- 795 52 Wallman-Jones A, Perakakis P, Tsakiris M, Schmidt M. Physical activity and interoceptive
- 796 processing: Theoretical considerations for future research. *Int. J. Psychophysiol.* 2021.
- 797 doi:10.1016/j.ijpsycho.2021.05.002.
- 798 53 Sha Z, Wager TD, Mechelli A, He Y. Common Dysfunction of Large-Scale Neurocognitive
- 799 Networks Across Psychiatric Disorders. *Biol Psychiatry* 2019.
- 800 doi:10.1016/j.biopsycho.2018.11.011.
- 801 54 Sha Z, Xia M, Lin Q, Cao M, Tang Y, Xu K *et al.* Meta-connectomic analysis reveals commonly
- 802 disrupted functional architectures in network modules and connectors across brain disorders.
- 803 *Cereb Cortex* 2018. doi:10.1093/cercor/bhx273.
- 804 55 Kolesar TA, Bilevicius E, Wilson AD, Kornelsen J. Systematic review and meta-analyses of
- 805 neural structural and functional differences in generalized anxiety disorder and healthy controls
- 806 using magnetic resonance imaging. *NeuroImage Clin.* 2019. doi:10.1016/j.nicl.2019.102016.
- 807 56 Nickerson LD, Smith SM, Öngür D, Beckmann CF. Using dual regression to investigate network
- 808 shape and amplitude in functional connectivity analyses. *Front Neurosci* 2017; **11**.
- 809 doi:10.3389/fnins.2017.00115.
- 810 57 Winkler AM, Renaud O, Smith SM, Nichols TE. Permutation inference for canonical correlation
- 811 analysis. *Neuroimage* 2020. doi:10.1016/j.neuroimage.2020.117065.
- 812 58 HOTELLING H. RELATIONS BETWEEN TWO SETS OF VARIATES. *Biometrika* 1936.

- 813 doi:10.1093/biomet/28.3-4.321.
- 814 59 Helmer M, Warrington S, Mohammadi-Nejad A-R, Ji LJ, Howell A, Rosand B *et al.* On stability
815 of Canonical Correlation Analysis and Partial Least Squares with application to brain-behavior
816 associations. *bioRxiv* 2020.
- 817 60 Nagel M, Jansen PR, Stringer S, Watanabe K, De Leeuw CA, Bryois J *et al.* Meta-analysis of
818 genome-wide association studies for neuroticism in 449,484 individuals identifies novel genetic
819 loci and pathways. *Nat Genet* 2018. doi:10.1038/s41588-018-0151-7.
- 820 61 Levey DF, Gelernter J, Polimanti R, Zhou H, Cheng Z, Aslan M *et al.* Reproducible Genetic Risk
821 Loci for Anxiety: Results From ~200,000 Participants in the Million Veteran Program. *Am J*
822 *Psychiatry* 2020. doi:10.1176/appi.ajp.2019.19030256.
- 823 62 Okbay A, Baselmans BML, De Neve JE, Turley P, Nivard MG, Fontana MA *et al.* Genetic
824 variants associated with subjective well-being, depressive symptoms, and neuroticism identified
825 through genome-wide analyses. *Nat Genet* 2016. doi:10.1038/ng.3552.
- 826 63 Howard DM, Adams MJ, Clarke TK, Hafferty JD, Gibson J, Shirali M *et al.* Genome-wide meta-
827 analysis of depression identifies 102 independent variants and highlights the importance of the
828 prefrontal brain regions. *Nat Neurosci* 2019. doi:10.1038/s41593-018-0326-7.
- 829 64 Levey DF, Stein MB, Wendt FR, Pathak GA, Zhou H, Aslan M *et al.* Bi-ancestral depression
830 GWAS in the Million Veteran Program and meta-analysis in >1.2 million individuals highlight
831 new therapeutic directions. *Nat Neurosci* 2021. doi:10.1038/s41593-021-00860-2.
- 832 65 Karlsson Linnér R, Biroli P, Kong E, Meddens SFW, Wedow R, Fontana MA *et al.* Genome-wide
833 association analyses of risk tolerance and risky behaviors in over 1 million individuals identify
834 hundreds of loci and shared genetic influences. *Nat Genet* 2019. doi:10.1038/s41588-018-0309-3.
- 835 66 Grotzinger AD, Rhemtulla M, de Vlaming R, Ritchie SJ, Mallard TT, Hill WD *et al.* Genomic
836 structural equation modelling provides insights into the multivariate genetic architecture of
837 complex traits. *Nat Hum Behav* 2019. doi:10.1038/s41562-019-0566-x.
- 838 67 O'Connor LJ, Price AL. Distinguishing genetic correlation from causation across 52 diseases and
839 complex traits. *Nat Genet* 2018. doi:10.1038/s41588-018-0255-0.
- 840 68 Jia H, Zack MM, Gottesman II, Thompson WW. Associations of Smoking, Physical Inactivity,
841 Heavy Drinking, and Obesity with Quality-Adjusted Life Expectancy among US Adults with
842 Depression. *Value Heal* 2018. doi:10.1016/j.jval.2017.08.002.
- 843 69 Metse AP, Clinton-Mcharg T, Skinner E, Yogaraj Y, Colyvas K, Bowman J. Associations between
844 suboptimal sleep and smoking, poor nutrition, harmful alcohol consumption and inadequate
845 physical activity ('snap risks'): A comparison of people with and without a mental health
846 condition in an australian community survey. *Int J Environ Res Public Health* 2021.
847 doi:10.3390/ijerph18115946.
- 848 70 R Core Team (2021). R: A Language and Environment for Statistical Computing. R Found. Stat.
849 Comput. 2021.
- 850 71 Baniqued PL, Gallen CL, Voss MW, Burzynska AZ, Wong CN, Cooke GE *et al.* Brain network
851 modularity predicts exercise-related executive function gains in older adults. *Front Aging*
852 *Neurosci* 2018. doi:10.3389/fnagi.2017.00426.
- 853 72 Chaddock-Heyman L, Weng TB, Kienzler C, Weissappel R, Drollette ES, Raine LB *et al.* Brain
854 Network Modularity Predicts Improvements in Cognitive and Scholastic Performance in Children
855 Involved in a Physical Activity Intervention. *Front Hum Neurosci* 2020.
856 doi:10.3389/fnhum.2020.00346.
- 857 73 Smallwood J, Bernhardt BC, Leech R, Bzdok D, Jefferies E, Margulies DS. The default mode
858 network in cognition: a topographical perspective. *Nat. Rev. Neurosci.* 2021. doi:10.1038/s41583-
859 021-00474-4.
- 860 74 Wang S, Tepfer LJ, Taren AA, Smith D V. Functional parcellation of the default mode network: a
861 large-scale meta-analysis. *Sci Rep* 2020. doi:10.1038/s41598-020-72317-8.
- 862 75 Mirza SS, Ikram MA, Bos D, Mihaescu R, Hofman A, Tiemeier H. Mild cognitive impairment and
863 risk of depression and anxiety: A population-based study. *Alzheimer's Dement* 2017.

- 864 doi:10.1016/j.jalz.2016.06.2361.
- 865 76 Perini G, Ramusino MC, Sinforiani E, Bernini S, Petrachi R, Costa A. Cognitive impairment in
866 depression: Recent advances and novel treatments. *Neuropsychiatr Dis Treat* 2019.
867 doi:10.2147/NDT.S199746.
- 868 77 Rock PL, Roiser JP, Riedel WJ, Blackwell AD. Cognitive impairment in depression: A systematic
869 review and meta-analysis. *Psychol. Med.* 2014. doi:10.1017/S0033291713002535.
- 870 78 Delgado MR, Nearing KI, LeDoux JE, Phelps EA. Neural Circuitry Underlying the Regulation of
871 Conditioned Fear and Its Relation to Extinction. *Neuron* 2008. doi:10.1016/j.neuron.2008.06.029.
- 872 79 Morawetz C, Bode S, Baudewig J, Kirilina E, Heekeren HR. Changes in Effective Connectivity
873 Between Dorsal and Ventral Prefrontal Regions Moderate Emotion Regulation. *Cereb Cortex*
874 2016. doi:10.1093/cercor/bhv005.
- 875 80 Ochsner KN, Gross JJ. The cognitive control of emotion. *Trends Cogn. Sci.* 2005.
876 doi:10.1016/j.tics.2005.03.010.
- 877 81 Koenigs M, Grafman J. The functional neuroanatomy of depression: Distinct roles for
878 ventromedial and dorsolateral prefrontal cortex. *Behav. Brain Res.* 2009.
879 doi:10.1016/j.bbr.2009.03.004.
- 880 82 Nejati V, Majdi R, Salehinejad MA, Nitsche MA. The role of dorsolateral and ventromedial
881 prefrontal cortex in the processing of emotional dimensions. *Sci Rep* 2021. doi:10.1038/s41598-
882 021-81454-7.
- 883 83 Yanagisawa H, Dan I, Tsuzuki D, Kato M, Okamoto M, Kyutoku Y *et al.* Acute moderate exercise
884 elicits increased dorsolateral prefrontal activation and improves cognitive performance with
885 Stroop test. *Neuroimage* 2010. doi:10.1016/j.neuroimage.2009.12.023.
- 886 84 Li MY, Huang MM, Li SZ, Tao J, Zheng GH, Chen LD. The effects of aerobic exercise on the
887 structure and function of DMN-related brain regions: a systematic review. *Int. J. Neurosci.* 2017.
888 doi:10.1080/00207454.2016.1212855.
- 889 85 Tao J, Liu J, Egorova N, Chen X, Sun S, Xue X *et al.* Increased hippocampus-medial prefrontal
890 cortex resting-state functional connectivity and memory function after Tai Chi Chuan practice in
891 elder adults. *Front Aging Neurosci* 2016. doi:10.3389/fnagi.2016.00025.
- 892 86 De Hert M, Correll CU, Bobes J, Cetkovich-Bakmas M, Cohen DAN, Asai I *et al.* Physical illness
893 in patients with severe mental disorders. I. Prevalence, impact of medications and disparities in
894 health care. *World Psychiatry.* 2011. doi:10.1002/j.2051-5545.2011.tb00014.x.
- 895 87 Fang H, Tu S, Sheng J, Shao A. Depression in sleep disturbance: A review on a bidirectional
896 relationship, mechanisms and treatment. *J. Cell. Mol. Med.* 2019. doi:10.1111/jcmm.14170.
- 897 88 Cox RC, Olatunji BO. A systematic review of sleep disturbance in anxiety and related disorders. *J.*
898 *Anxiety Disord.* 2016. doi:10.1016/j.janxdis.2015.12.001.
- 899 89 Cheng W, Rolls E, Gong W, Du J, Zhang J, Zhang XY *et al.* Sleep duration, brain structure, and
900 psychiatric and cognitive problems in children. *Mol Psychiatry* 2021. doi:10.1038/s41380-020-
901 0663-2.
- 902 90 Brenes GA, Miller ME, Stanley MA, Williamson JD, Knudson M, McCall WV. Insomnia in older
903 adults with generalized anxiety disorder. *Am J Geriatr Psychiatry* 2009.
904 doi:10.1097/JGP.0b013e3181987747.
- 905 91 Lippman S, Gardener H, Rundek T, Seixas A, Elkind MSV, Sacco RL *et al.* Short sleep is
906 associated with more depressive symptoms in a multi-ethnic cohort of older adults. *Sleep Med*
907 2017. doi:10.1016/j.sleep.2017.09.019.
- 908 92 Li Y, Wu Y, Zhai L, Wang T, Sun Y, Zhang D. Longitudinal Association of Sleep Duration with
909 Depressive Symptoms among Middle-aged and Older Chinese. *Sci Rep* 2017. doi:10.1038/s41598-
910 017-12182-0.
- 911
- 912

913 **Figure Legends**

914

915 **Figure 1. Overview of variables and analyses.** Separate analyses were conducted for brain (left panel) and
916 genetic associations (right panel). For brain associations, canonical correlation analysis (CCA) was employed for
917 mental health (MH) and physical activity (PA) separately. Simple and unique CCA models only differed in whether
918 the shared variance between MH and PA was accounted for. For genetic associations, GWAS summary statistics for
919 5 mental health phenotypes (MHP₁-MHP₅) and for 5 physical activity phenotypes (PAP₁-PAP₅) were leveraged into
920 pairwise linkage disequilibrium score regression (LDSR) analyses in MHP and PAP models separately. Genomic
921 structural equation modeling (gSEM) was also employed to identify genetic associations between a latent factor
922 from MH phenotypes and each of PA phenotype, followed by latent causal variable analysis (LCV) that allows for
923 inferring causal genetic relationships among MH and PA phenotypes.

924

925

926 **Figure 2. Canonical loadings of brain measures on the first canonical variates.** These loadings represent the
927 linear correlation between the original brain measures (Y) and the first canonical variate (U) per model. Color
928 coding was made for brain variable names along the Y axis (i.e., subnetwork edges in gray with “-” between
929 subnetwork names and amplitude in orange), and for the bars representing canonical loadings (i.e., significance in
930 cyan, insignificance in yellow). Simple and unique models differ in whether the model accounted for the shared
931 variance in brain measures between mental health and physical activity. vDMN = ventral default mode network; R/L
932 CEN = right/ left central executive network; PCu = precuneus; a/p SN = anterior/posterior salience network.

933

934

935 **Figure 3. Heatmap of bivariate genetic correlations.** Pairwise genetic correlations were calculated separately for
936 phenotypes of mental health (A) and those of physical activity (B). rg = genetic correlation coefficient, GAD =
937 Generalized Anxiety Disorder, MDD = Major Depressive Disorder, Risk = Risk Tolerance, SWB = Subjective Well-
938 Being.

939

940

941 **Figure 4. Genetic correlations between negative affect and physical activity phenotypes.** Significant genetic
942 correlations with negative affect were observed for moderate (A), overall activities (B), and sleep duration (C). The
943 latent negative affect factor loaded positively on neuroticism, major depression disorder (MDD) and general anxiety
944 disorder (GAD) phenotypes, but negatively on the subjective well-being (SWB) phenotype.

945

946

947

948

949 **Tables**

950

951 **Table 1.** Correlations between mental health and physical activity measures.

Physical Activity		Mental Health						
		PC1	PC2	PC3	PC4	PC5	PC6	PC7
Overall	Sleep Duration	-0.08*	0.04*	0.01	0.00	0.04*	-0.01	0.03
	Sedentary	-0.03*	0.07*	0.07*	-0.02	-0.02	-0.03	0.03*
	Light Tasks	0.02	-0.02	0.01	-0.01	-0.04*	-0.01	0.00
	Moderate	-0.02	-0.06*	0.02	0.01	0.03*	0.04*	0.01
	Walking	0.01	-0.01	0.02	-0.03*	-0.06*	0.00	0.04*
	Overall Activity	-0.06*	0.04*	0.03*	-0.02	0.04*	0.01	0.08*
	MET	-0.06*	0.10*	0.07*	-0.03	-0.01	-0.05*	0.02
Weekdays	Sleep Duration	-0.08*	0.04*	0.02	0.00	0.03*	-0.01	0.03*
	Sedentary	-0.05*	0.08*	0.08*	-0.01	-0.04*	-0.02	0.04*
	Light Tasks	0.01	-0.02	0.03	-0.03	-0.03*	-0.01	0.01
	Moderate	-0.02	-0.05*	0.03	0.01	0.02	0.03*	0.03
	Walking	0.02	0.00	0.04*	-0.04*	-0.09*	-0.01	0.04*
	Overall Activity	-0.05*	0.04*	0.06*	-0.04*	-0.01	0.00	0.07*
	MET	-0.06*	0.11*	0.08*	-0.03	-0.07*	-0.04*	0.02
Weekend	Sleep Duration	-0.06*	0.04*	0.00	0.00	0.02	-0.01	0.02
	Sedentary	-0.02	0.05*	0.06*	-0.02	-0.01	-0.02	0.02
	Light Tasks	0.02	-0.01	0.00	-0.01	-0.05*	-0.02	-0.02
	Moderate	-0.02	-0.04*	0.01	0.00	0.01	0.04*	0.00
	Walking	0.02	0.00	0.02	-0.03*	-0.06*	0.00	0.04*
	Overall Activity	-0.05*	0.04*	0.03	-0.02	0.03*	0.01	0.06*
	MET	-0.05*	0.08*	0.05*	-0.02	0.01	-0.04*	0.02

952 Phenotypes of mental health were represented by the first seven principal components (i.e., PC1-PC7)
 953 that altogether explained 51.05% variance. Significant correlations are indicated by * after FDR
 954 corrections (i.e., all corrected p 's ≤ 0.045) and negative correlation coefficients are highlighted with light
 955 grey shading

956

957

958 **Table 2. Summary statistics from GWAS studies.**

Reference	Phenotype	Dataset	Sample Size	SNP h^2
Nagel et al 2018 ⁶⁰	Neuroticism	UK Biobank & PGC	390,279	0.10
Levey et al 2020 ⁶¹	GAD	Million Veteran Program	199,611	0.056
Okbay et al 2016 ⁶²	SWB	Meta-analysis across 59 cohorts	298,420	0.040
Howard et al 2019 ⁶³	MDD	UK Biobank & PGC	500,199	0.11
Levey et al 2021 ⁶⁴		Million Veteran Program	250,215	
Linnér et al 2019 ⁶⁵	Risk-taking	UK Biobank	431,126	0.050

Doherty et al 2018 ⁴⁸	Physical activities	UK Biobank	91,105	0.10 – 0.21
----------------------------------	---------------------	------------	--------	-------------

959 GAD = general anxiety disorder; SWB = subjective well-being; MDD = major depressive disorder; PGC
 960 = Psychiatric Genetics Consortium.

961

962

Table 3. Genetic correlations between mental health and physical activity phenotypes.

Mental Health Phenotype	Physical Activity Phenotype	<i>rg</i>	<i>p</i>	<i>pFDR</i>
Risk Tolerance	Sleep Duration	-0.029	0.433	0.447
	Sedentary	-0.028	0.447	0.447
	Moderate	0.081	0.092	0.131
	Walking	0.115	4.95e-3	0.012
	Overall	0.074	0.020	0.040
Negative Affect	Sleep Duration	0.121	1.25e-5	1.25e-4
	Sedentary	-0.062	0.036	0.060
	Moderate	-0.117	1.33e-3	4.43e-3
	Walking	-0.029	0.383	0.447
	Overall	-0.085	7.70e-4	3.85e-3

963 *Note*, Significant results after FDR corrections were highlighted in bold. *rg* = genetic correlation,

964 *pFDR* = FDR-corrected p-value.

965

966

967