

Immunogenetics of Gastrointestinal Cancers: A Systematic Review of Inborn Errors of Immunity in Humans

Beishi Zheng^{1,2}, Howard Chung¹, Chen Bing¹, Siming Sun¹, Peter H.R. Green¹, Timothy C. Wang¹, Xiao-Fei Kong^{1*}

1. Division of Digestive and Liver Disease, Department of Medicine, Columbia University Irving Medical Center, New York, NY, 10032
2. Department of Internal Medicine, Woodhull Medical and Mental Health Center, Brooklyn, NY, 11206

*Corresponding author: Xiao-Fei Kong (xk2137@cumc.columbia.edu)

Address: 622 West 168th St, PH7W, Suite 318, New York, NY, 10032

Conflict Of Interest: Beishi Zheng, Howard Chung, Chen Bing, Siming Sun, Peter H.R. Green, Timothy C. Wang, Xiao-Fei Kong have no conflict of interest (financial, professional, or personal) to disclose.

Grant Support: Research reported in this publication was supported by the National Institute Of Diabetes And Digestive And Kidney Diseases of the National Institutes of Health under Award Number K08DK128631. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

Abstract:

Background/Aims: The immune system plays a key role in cancer surveillance and modulation of the tumor microenvironment. Humans with inborn errors of immunity (IEI), or primary immunodeficiencies, may be prone to recurrent mucosal bacterial and viral infections and chronic inflammation, associated with intrinsic or secondary epithelium dysfunction, a potential risk factor for early-onset gastrointestinal (GI) cancer.

Methods: We systematically reviewed all cases with clinical diagnoses of both an IEI and a GI cancer in three databases (MEDLINE, SCOPUS, EMBASE). In total, 76 publications satisfying our inclusion criteria were identified, and data for 149 cases were analyzed.

Results: Of the 149 patients with IEIs, 95 presented with gastric cancer, 13 with small bowel cancer, 35 with colorectal cancer, and six with unspecified cancer or cancer at another site. Gastric and colonic adenocarcinoma was the most common. For both gastric and colorectal cancer, age at onset was significantly earlier in patients with IEIs than in the general population, based on the SEER database. Common variable immune deficiency (CVID) was the most common IEI associated with gastrointestinal cancer. About 12% of patients had molecular genetic diagnoses, the three most frequently implicated genes being *ATM*, *CARMIL2*, *CTLA4*. Impaired humoral immunity and Epstein-Barr virus (EBV) infection were frequently reported as the factors potentially underlying early-onset GI malignancy in patients with IEIs.

Conclusion:

Patients with IEIs should be considered at risk of early-onset GI cancers, and should therefore undergo cancer screening at an earlier age. Surveillance guidance based on stratifications for genetic risk should be revised to take into account the immunogenetic contribution to GI cancers.

Keywords: inborn errors of immunity, gastrointestinal cancer, immunosurveillance, human genetics

Introduction

Cancer immunosurveillance, first proposed by Frank Macfarlane Burnet and Lewis Thomas in 1957, plays an indispensable role in the prevention of gastrointestinal cancers¹. Cancers were found to occur more frequently in immunocompromised individuals in studies of AIDS patients or organ recipients, particularly for non-melanoma skin cancer^{2,3}. Studies of transgenic mice with defects of crucial components of the immune system have improved our understanding of the effector molecules involved in immunosurveillance and led to the development of immune checkpoint inhibitors, a key breakthrough in immunotherapy for cancer treatment^{4,5}. Inborn errors of immunity (IEIs), also known as primary immunodeficiencies, manifest as recurrent infections, autoimmunity, autoinflammatory diseases, allergy, and cancer⁶. Humans with IEIs provide us with a unique angle of study, or an experiment in nature, for elucidating immunosurveillance for GI cancer and the interaction between environment and host. However, the rare cases of GI cancer in patients with IEIs are challenging for the physicians responsible for their care. For instance, IEIs can manifest as an early-onset GI cancer, but unlike high-risk hereditary risk factors for gastrointestinal cancer, IEIs are not yet generally considered as a risk factor for GI cancers, and there are currently no practice guidelines for screening for gastrointestinal cancers in patients with IEIs^{7,8}.

The diagnosis of an early-onset (before the age of 50 years) gastrointestinal cancer, whether gastric or colorectal cancer, is devastating for any patient⁹⁻¹¹. The field of primary immunodeficiencies has progressed considerably with recent advances in genetic sequencing technology, which are facilitating genetic diagnosis and discoveries. More than 400 genes have been shown to cause a wide array of immunodysregulation, with a broad spectrum of clinical presentations, some dominant and others recessive, and complete or incomplete penetrance⁶. Epidemiological data suggest that inborn errors of immunity are more prevalent than previously thought, affecting about 1:8,500 to 1:100,000 individuals in the general population¹². With earlier clinical recognition, appropriate treatments, and

hematopoietic stem cell transplantation, many patients with IEs now survive into adulthood, overcoming severe infections or inflammation. Given these improvements in the overall survival of patients with IEI, physicians may begin to see more delayed clinical phenotypes with an incomplete penetrance, including gastrointestinal cancers. It is often difficult to understand the human genetics and immunologic basis of IEs, due to phenotypic and allelic heterogeneity, differences in expressivity and incomplete penetrance. Given the complexity of human genetics and immunology, and the rareness of individuals with identified IEs and gastrointestinal cancer, we systematically reviewed all the reported cases with both an IEI and a gastrointestinal cancer. We summarized their genetic and immunological features and highlighted the elements of importance for the clinical surveillance and management of this special population.

Methods

Case identification and literature search

We followed a Preferred Reporting Items for Systemic Reviews and Meta-Analysis (PRISMA)¹³ flow chart for the literature search and data collection for this systematic review (Supplementary Figure 1). Briefly, we conducted a search of the MEDLINE, SCOPUS, EMBASE databases for all publications up to December 1, 2020, focusing on IEs and GI cancers. We focused on luminal GI cancers, and excluded pancreatic or hepatobiliary cancers. We followed the International Union of Immunological Societies (IUIS) guidelines⁶ to ensure the inclusion of all IEs. We searched three databases for the following terms: “gastrointestinal neoplasms”, “primary immunodeficiency”, “agammaglobulinemia”, “common variable immunodeficiency”, “dysgammaglobulinemia”, “lymphopenia”, “immune dysregulation” and 14 additional Medical Subject Headings (MeSH) terms relating to IEs in the title, abstract, or keywords of articles (Supplementary Table 1). The results of three database searches ($n=1146$) were then carefully reviewed, as illustrated in Supplementary Figure 1. Additional publications ($n=24$) were identified and included manually if missed by the search strategy described above search. Duplicated articles

were consolidated. All records ($n=1099$) were then reviewed independently by two authors (B.Z and H.C) by comprehensive reading, and any discrepancies were resolved by a third author (X-F.K). The inclusion criteria were as follows: 1) Clinical diagnosis of IEI for cases; 2) Clinical diagnosis of gastrointestinal cancers for cases. Exclusion criteria were: 1) Not written in English; 2). Lack of detailed clinical information, such as age, sex, and age-at-onset data for the patient. In total, 76 articles satisfied with our inclusion criteria and had no exclusion criteria.

Data curation, quality assessment and statistical analysis

We collected information by scrutinizing the publications identified to generate a dataset with the following information: author and year of publication of the article and, for the patients, sex, ethnicity, age at cancer diagnosis, genetic defects, immunological dysfunctions, IEI diagnosis, cancer histology and prognosis. Standardized forms were used to extract the required information from the articles. A quality assessment form was developed based on the CARE (CAse REports)¹³ tool for quality assessment (Supplementary Table 2). The expected distributions of age, sex, and survival interval of GI cancers were calculated from the data in the SEER (Surveillance, Epidemiology and End Results) database. Time-to-event and survival analyses (Log rank) were performed in GraphPad Prism 8.0 to calculate the hazard ratio. Categorical variables were analyzed with a Fisher's exact test or *Chi*-squared test. Continuous variables were analyzed with *t*-tests. All data were analyzed with the statistical packages in R software (4.0.0).

Results

1. A clinical summary of GI cancers in patients with IEIs

The 76 publications identified¹⁴⁻⁸⁹ included 149 cases in which both IEIs and gastrointestinal cancers have been diagnosed. A summary of the clinical characteristics of the patients concerned is provided in Figure 1. Briefly, 95 cases were diagnosed with gastric cancer, 13 cases presented with small bowel cancer, 35

cases with colorectal cancer, three cases with esophageal cancer, and three cases with a GI tract tumor unspecified location (Figure 1A). In total, 59 cases (40%) were reported after 2010 (Figure 1B). The dataset consisted of 84 (56.4%) male patients, 58(38.9%) female patients and seven patients for whom sex was not specified (Figure 1C). Age at onset for the cancer ranged from three to 82 years, with a median age at onset of 42 years (Figure 1D). Most of the reported cases (64.4%) were diagnosed before the age of 50 years and their cancers should therefore be considered early-onset. Age at onset for GI was normally distributed, contrasting with the left-skewed distribution in the general population. Fisher's exact test showed that the proportion of male patients was higher in the younger age groups (ages 0-40) than the older age groups (ages above 40, $p=0.034$) (Figure 1D). Age at onset of the GI cancer was 26 to 30 years lower for the individuals with IEIs than for the general population (42 vs 68-72 years), according to the data of the SEER database. Ethnicity data were available for only 18 patients: seven white, four non-Hispanic, three Asian, two African American, and two Hispanic or Latino individuals. The IEI diagnosis was common variable immunodeficiency (CVID) for 88 patients in the cohort. The other IEI diagnoses included X-linked agammaglobulinemia (XLA) in 11 patients, selective IgA deficiency (SIgAD) in 10 patients, and ataxia-telangiectasia mutated gene deficiency (*ATM*) in 10 patients. The rate of molecular genetic diagnosis for IEIs was low, with disease-causing genotypes reported for only 18/149 (12.1%).

2. Gastric cancers in patients with IEIs

Our literature search identified 95 patients with IEIs and gastric cancer, which was the most common GI cancer (Supplementary Table 3). Most (87/95, 91.5%) of the patients were diagnosed with gastric adenocarcinoma. The other cancer diagnosed were lymphoma (7/95, 7.4%), and neuroendocrine carcinoma (1/95, 1.1%) (Figure 2A). The mean age at gastric cancer diagnosis was 45.86 years, male patients outnumbered female patients in this group (55.8% vs 43.2%), and there were no significant differences with respect to the expected value (Figure 2B). Age at onset of gastric cancer in patients with IEIs was significantly different from that in the general

population, with most patients with IEIs diagnosed with gastric cancer before the age of 65 years (Figure 2B). Gastric cancer was diagnosed between the ages of 25 and 44 years in 32.6% of IEI cases, whereas such early diagnosis was uncommon in the general population (Figure 2B). A time-to-event data analysis based on the observed and expected age at onset for gastric cancer yielded a hazard ratio (HR) of 3.57 for the IEI group, indicating a significantly earlier than expected age at onset for this group (Figure 2C). Prognosis was as follows: 31 (32.6%) patients died within one year, and nine (9.5%) patients died within five years of a gastric cancer diagnosis. The five-year survival rate in IEI patients with gastric cancer was 27.3%, which is close to the value reported for the general population, as shown by the survival analysis (Figure 2D). CVID was the most common IEI and was found in 69 cases (72.6%) of gastric cancer. *ATM* defects and *XLA* were the next most frequent IEIs (Figure 2E). Four patients with *CTLA4* deficiency and one patient with *LRBA* deficiency were reported to have gastric cancer. However, most of the publications did not report the history of *Helicobacter pylori* infection. In summary, inborn errors resulting in B-cell defects, or dysfunctions of humoral immunity were the most common IEIs associated with gastric cancer.

3. Small bowel cancers reported in patients with IEIs

The small bowel is an uncommon site for cancers. To date, 13 patients with IEIs and small bowel cancer have been reported (Supplementary Table 4). Lymphoma was the most common type of cancer (11/13, 84.6%) affecting the small bowel, with 10 cases of B-cell lymphoma, one of T-cell lymphoma. Neuroendocrine carcinoma and reticular lymphosarcoma were reported in one case. Mean age at cancer diagnosis was 43.92 years and there were seven male patients and six female patients. CVID (41.6%, 6/13) was the most common IEIs associated with small bowel cancer, consistent with the recent meta-analysis of CVID patients showing that lymphoma was the cancer most frequently reported in these patients⁹⁰. One patient with Wiskott-Aldrich syndrome and another with familial Mediterranean fever were reported. Overall, 41.6% (6/13) of the patients survived for more than five years, 15.4% (2/13) survived for less

than three years, and 23.1% (3/13) died within one year. The genetic etiology of small-bowel lymphoma in CVID remains to be explored.

4. Colorectal cancers in patients with IEIs

There were 35 cases of colorectal cancer reported in patients with IEIs (Supplementary Table 5). The most common type of cancer was adenocarcinoma (68.1%, 24/35), followed by lymphoma (11.4%, 4/35), and smooth-muscle tumors (5.7%, 2/35) (Figure 3A). Other cellular types of cancer were reported in one case each, including neuroendocrine carcinoma, rhabdomyosarcoma, and squamous cell carcinoma (Figure 3A). Mean age at onset was 32.94 years, and this group comprised 60% (21/35) male and 28.6% (10/35) female patients. Most of the cases (71.4%, 25/35) were diagnosed before 45 years of age, which is significantly younger than expected (Figure 3B). Time-to-event analysis (Figure 3C) revealed a HR of 5.23 (95% CI 2.65- 10.30) for the IEI group. Overall, 42.9% (15/35) of the patients with IEIs died within one year of colon cancer diagnosis, and five-year survival rate for IEIs patients with colon cancer was 31.4% (11/35). A survival analysis showed prognosis to be worse for the IEI group than for the general population (HR = 2.55, $p = 0.002$) (Figure 3D). The most common immune dysregulations were B-cell defects (71.4%, 25/35), including CVID, XLA or selective IgA deficiency. However, DNA repair defects, including *BLM*, *NBS1* and *ATM* gene mutations, accounted for 20% (7/35) of IEI cases with colorectal cancer (Figure 3E). In summary, colorectal cancer in patients with IEIs appeared to be more heterogeneous in terms of immunological dysfunction and to lead to worse outcomes than such cancers in the general population.

Discussion

The Fearon-Vogelstein model has prevailed over the last few decades, and genetic practices for GI cancer have now emerged and can facilitate the management of about 3% to 5% of all colorectal cancer patients⁹¹. Pathogenic variants with various degrees of clinical penetrance have been identified in 9-26% of patients with early-onset colorectal cancer⁹². However, the detection of such mutations is limited by a lack of

knowledge of the full genetic landscape for GI cancers. We investigated the immunogenetic determinants of GI cancers by performing a systemic review of cases in patients with IEIs, who usually have a genetic predisposition for cancer and abnormal immunological dysregulation. Even though IEIs are usually diagnosed during childhood and mortality rates are high at a young age, 2-10% of patients have been found to develop cancers with their short lifespans⁷⁸. Our analysis indicates that patients with IEIs may be prone to early-onset GI cancers. Longitudinal data from a large IEI cohort are lacking, but our findings are consistent with previous CVID studies^{58, 93}. Further studies are required to delineate the clinical epidemiology of IEIs in patients with early-onset GI cancers, through immunological or genetic testing. A collaboration between clinical immunologists and pediatricians might facilitate GI cancer surveillance in individuals with IEIs. We need to address the question as to whether there is an association between clinical symptoms or biomarkers of IEIs and early-onset GI cancer? It would be intriguing to determine whether immunoglobulin level, lymphocytopenia, a clinical diagnosis of CVID, or a history of IVIG use is associated with early-onset GI cancer, ideally based on data from a large registry of GI cancer patients.

Several factors could, conceivably, account for the early-onset gastrointestinal cancers observed in patients with IEIs. First, primary immunodeficiency may lead to chronic mucocutaneous infection (with *Helicobacter. pylori* or EBV, for example,) in the gastrointestinal tract, which is a risk factor for gastric cancer. *CARMIL2* deficiency has been associated with EBV⁺ smooth muscle tumors in the GI tract⁶⁵. Previous studies on CVID have revealed a high prevalence of *H. pylori* infection and multifocal atrophic gastritis, but *H.pylori* infection rates in CVID patients vary between reports and may not, necessarily, explain the risk of GI cancer⁶⁷. Nevertheless, one recent large CVID cohort meta-analysis⁹⁰ on 8123 CVID patients revealed an estimated prevalence of gastric cancer in CVID patients of about 1.5%, lower than the prevalence of 4.1% for lymphoma, consistent with another report suggesting that gastric cancer may be the leading cause of death in CVID patients⁶⁷. Gastrointestinal cancers may also arise due to defects of immune cells in the mucosa. As most reported

cases have impaired humoral immunity, this suggests that B cells may play an important role in gastrointestinal mucosal surveillance. Only a few cases with genetic abnormalities have been reported, but several genetic disorders have well-defined B-cell abnormalities, including *CTLA4*, *LBRA*, and *CARMIL2* mutations.

A third plausible explanation is related to intrinsic epithelium abnormalities due to the pleiotropic effects of some genes. Three disorders, ataxia-telangiectasia due to *ATM* mutations, Nijmegen breakage syndrome due to *NBS1* defects, and Bloom syndrome due to *BLM* mutations, have been reported to cause heterogeneous clinical phenotypes and chromosomal instability. ATM is a serine/threonine kinase activated by DNA damage, it phosphorylates several tumor suppressors, including p53, NBS1, and H2AX. Eight patients (9.5%) with ataxia-telangiectasia due to biallelic *ATM* gene mutations have been reported to have gastric cancer and immunodeficiency^{33, 50, 76, 82, 83, 85}. Individuals with heterozygous germline *ATM* mutations might have an increased risk for early-onset gastric and pancreatic cancer⁹⁴. However, some IEs characterized by impaired DNA, such as MCM4 or GINS1 deficiencies, have not been found to lead to cancers^{95, 96}. It is possible that GI cancers in patients with DNA repair disorders are more strongly related to intrinsic epithelial cell defects than to compromised immunity. With advances in human genetics and immunological technologies, studies of humans with IEs might improve our understanding of immune surveillance in the gastrointestinal tract in general.

Figure and table legend:

Figure 1: A descriptive summary of patients with inborn errors of immunity and GI cancers. A. A schematic presentation of the numbers of esophageal, gastric, small bowel and colon cancers reported in patients with IEs. B. The table describes the number of cases published over the last six decades. C. A pie chart of the sex distribution for this cohort. D. A histogram presents the age and sex distributions of the cohort. NA: Not available.

Figure 2: Gastric cancers reported in patients with IEs. A. Cellular types of

gastric cancer reported in patients with IEIs. Adenoca: Adenocarcinoma, NET: Neuroendocrine tumor. B. The table shows the sex and age distributions of gastric cancer patients from the IEI cohort relative to the expected numbers in the general population imputed from the SEER database. OR: Odds ratio. C. A pie chart of the different types of IEI and numbers of cases with gastric cancer. CVID: Common variable immune deficiency; ATM: Ataxia-telangiectasia mutated, XLA: X-linked agammaglobulinemia, CTLA4: Cytotoxic T-lymphocyte associated Protein 4, SIgAD: Selective IgA deficiency, FMF: Familial Mediterranean fever, LRBA: LPS responsive beige-like anchor protein. D. Survival curve for gastric cancer comparing patients with and without IEIs. NA: Not available.

Figure 3: Colon cancer in patients with IEIs. A. Cellular types of colon cancer reported in patients with IEIs. Adenoca: Adenocarcinoma, NET: Neuroendocrine tumor, SMT: Smooth muscle tumor, RMS: Rhabdomyosarcoma, GS: Granulocytic sarcoma, SCC: Squamous cell carcinoma. B. The table shows the sex and age distribution of colon cancer patients from the IEI cohort relative to the expected numbers in the general population imputed from the SEER database. OR: Odds ratio. C. A pie chart of the different types of IEIs and numbers of cases with colon cancer. CVID: Common variable immune deficiency; ATM: Ataxia-telangiectasia mutated, XLA: X-linked agammaglobulinemia, BLM: Bloom syndrome, SIgAD: Selective IgA deficiency, *CARMIL2*: Capping Protein Regulator And Myosin 1 Linker 2, ATM: Ataxia-telangiectasia mutated, FMF: Familial Mediterranean fever, NBS: Nijmegen breakage syndrome. D. Survival curve for colon cancer comparing patients with and without IEIs. NA: Not available.

Acknowledgments:

We thank Dr. Jean-Laurent Casanova, Laurent Abel, Anil K Rustgi and Benjamin Lebwohl for their careful reading of the manuscript and suggestions. Figure 1 was created with BioRender.com.

Reference:

1. Burnet M. Cancer; a biological approach. I. The processes of control. Br Med J 1957;1:779-86.
2. Frisch M, Biggar RJ, Engels EA, et al. Association of cancer with AIDS-related immunosuppression in adults. JAMA 2001;285:1736-45.
3. Ramsay HM, Fryer AA, Hawley CM, et al. Non-melanoma skin cancer risk in the Queensland renal transplant population. Br J Dermatol 2002;147:950-6.
4. Kroemer G, Zitvogel L. Immune checkpoint inhibitors. J Exp Med 2021;218.
5. Kim W, Chu TH, Nienhuser H, et al. PD-1 Signaling Promotes Tumor-Infiltrating Myeloid-Derived Suppressor Cells and Gastric Tumorigenesis in Mice. Gastroenterology 2021;160:781-796.
6. Tangye SG, Al-Herz W, Bousfiha A, et al. Human Inborn Errors of Immunity: 2019 Update on the Classification from the International Union of Immunological Societies Expert Committee. J Clin Immunol 2020;40:24-64.
7. Shaukat A, Kahi CJ, Burke CA, et al. ACG Clinical Guidelines: Colorectal Cancer Screening 2021. Am J Gastroenterol 2021;116:458-479.
8. Milne AN, Offerhaus GJ. Early-onset gastric cancer: Learning lessons from the young. World J Gastrointest Oncol 2010;2:59-64.
9. Bergquist JR, Leiting JL, Habermann EB, et al. Early-onset gastric cancer is a distinct disease with worrisome trends and oncogenic features. Surgery 2019;166:547-555.
10. Kolb JM, Ahnen DJ, Samadder NJ. Evidenced-Based Screening Strategies for a Positive Family History. Gastrointest Endosc Clin N Am 2020;30:597-609.
11. Machlowska J, Baj J, Sitarz M, et al. Gastric Cancer: Epidemiology, Risk Factors, Classification, Genomic Characteristics and Treatment Strategies. Int J Mol Sci 2020;21.
12. Kobrynski L, Powell RW, Bowen S. Prevalence and morbidity of primary immunodeficiency diseases, United States 2001-2007. J Clin Immunol

2014;34:954-61.

13. Tawfik GM, Dila KAS, Mohamed MYF, et al. A step by step guide for conducting a systematic review and meta-analysis with simulation data. Tropical medicine and health. Volume 47, 2019:46.
14. Aguilar FP, Alfonso V, Rivas S, et al. Jejunal malignant lymphoma in a patient with adult-onset hypo-gamma-globulinemia and nodular lymphoid hyperplasia of the small bowel. The American journal of gastroenterology. Volume 82, 1987:472-475.
15. Cunningham-Rundles C, Siegal FP, Cunningham-Rundles S, et al. Incidence of cancer in 98 patients with common varied immunodeficiency. Journal of clinical immunology. Volume 7, 1987:294-9.
16. Durham JC, Stephens DS, Rimland D, et al. Common variable hypogammaglobulinemia complicated by an unusual T-suppressor/cytotoxic cell lymphoma. Cancer. Volume 59, 1987:271-6.
17. Takemiya M, Shiraishi S, Teramoto T, et al. Bloom's syndrome with porokeratosis of Mibelli and multiple cancers of the skin, lung and colon. Clinical genetics. Volume 31, 1987:35-44.
18. Conley ME, Ziegler MM, Borden St, et al. Multifocal adenocarcinoma of the stomach in a child with common variable immunodeficiency. Journal of pediatric gastroenterology and nutrition. Volume 7, 1988:456-460.
19. Cunningham-Rundles C. Clinical and immunologic analyses of 103 patients with common variable immunodeficiency. Journal of clinical immunology. Volume 9, 1989:22-33.
20. Vorechovsky I, Litzman J, Lokaj J, et al. Common variable immunodeficiency and malignancy: a report of two cases and possible explanation for the association. Cancer immunology, immunotherapy : CII. Volume 31, 1990:250-4.
21. Paraf F, Paraf A, Brousse N. Primary gastric lymphoma occurring in familial Mediterranean fever. Journal of clinical gastroenterology. Volume 13, 1991:587-588.

22. Adachi Y, Mori M, Kido A, et al. Multiple colorectal neoplasms in a young adult with hypogammaglobulinemia. Report of a case. *Diseases of the colon and rectum*. Volume 35, 1992:197-200.
23. Castellano G, Moreno D, Galvao O, et al. Malignant lymphoma of jejunum with common variable hypogammaglobulinemia and diffuse nodular hyperplasia of the small intestine. A case study and literature review. *Journal of clinical gastroenterology*. Volume 15, 1992:128-135.
24. Matz J, Routes JM, Borish LC. Diffuse pulmonary infiltrates in hypogammaglobulinemia. *Annals of allergy*. Volume 69, 1992:485-490.
25. de Bruin NC, de Groot R, den Hollander JC, et al. Small-cell undifferentiated (neuroendocrine) carcinoma of the cecum in a child with common variable immunodeficiency. *The American journal of pediatric hematology/oncology*. Volume 15, 1993:258-261.
26. Lavilla P, Gil A, Rodríguez MC, et al. X-linked agammaglobulinemia and gastric adenocarcinoma. *Cancer*. Volume 72, 1993:1528-1531.
27. Ohnoshi T, Hayashi K, Tagawa S, et al. Successful treatment of non-Hodgkin's lymphoma in a patient with common variable immunodeficiency. *Internal medicine (Tokyo, Japan)*. Volume 32, 1993:152-155.
28. van der Meer JW, Weening RS, Schellekens PT, et al. Colorectal cancer in patients with X-linked agammaglobulinaemia. *Lancet (London, England)*. Volume 341, 1993:1439-40.
29. Chiamonte C, Glick SN. Nodular lymphoid hyperplasia of the small bowel complicated by jejunal lymphoma in a patient with common variable immune deficiency syndrome. *AJR. American journal of roentgenology*. Volume 163, 1994:1118-1119.
30. Cox JE, Ott DJ. Gastric adenocarcinoma in a patient with common variable immunodeficiency. *Abdominal imaging*. Volume 19, 1994:501-502.
31. Zirkin HJ, Levy J, Katchko L. Small cell undifferentiated carcinoma of the colon associated with hepatocellular carcinoma in an immunodeficient patient. *Human pathology*. Volume 27, 1996:992-6.

32. Balci S, Aktas D. Mucinous carcinoma of the colon in a 16-year-old Turkish boy with Bloom syndrome: cytogenetic, histopathologic, TP53 gene and protein expression studies. *Cancer genetics and cytogenetics*. Volume 111, 1999:45-48.
33. Murphy RC, Berdon WE, Ruzal-Shapiro C, et al. Malignancies in pediatric patients with ataxia telangiectasia. *Pediatric radiology*. Volume 29, 1999:225-30.
34. Wang J, German J, Ashby K, et al. Ulcerative colitis complicated by dysplasia-adenoma-carcinoma in a man with Bloom's syndrome. *Journal of clinical gastroenterology*. Volume 28, 1999:380-382.
35. Zullo A, Romiti A, Rinaldi V, et al. Gastric pathology in patients with common variable immunodeficiency. *Gut*. Volume 45, 1999:77-81.
36. Bachmeyer C, Monge M, Cazier A, et al. Gastric adenocarcinoma in a patient with X-linked agammaglobulinaemia. *European journal of gastroenterology & hepatology*. Volume 12, 2000:1033-1035.
37. Mellekjaer L, Hammarstrom L, Andersen V, et al. Cancer risk among patients with IgA deficiency or common variable immunodeficiency and their relatives: a combined Danish and Swedish study. *Clinical and experimental immunology*. Volume 130, 2002:495-500.
38. Meyer S, Kingston H, Taylor AMR, et al. Rhabdomyosarcoma in Nijmegen breakage syndrome: strong association with perianal primary site. *Cancer genetics and cytogenetics*. Volume 154, 2004:169-74.
39. Lackmann GM, Wahn V, Poremba C, et al. A teenager with X-linked agammaglobulinemia and vitamin B12 deficiency anemia. *Journal of pediatric gastroenterology and nutrition*. Volume 41, 2005:360-362.
40. Delia M, Liso V, Capalbo S, et al. Common variable immunodeficiency patient with large granular lymphocytosis developing extranodal diffuse large B-cell lymphoma: a case report. *Haematologica*. Volume 91, 2006:ECR61.
41. Desar IME, Keuter M, Raemaekers JMM, et al. Extranodal marginal zone (MALT) lymphoma in common variable immunodeficiency. *The Netherlands*

- journal of medicine. Volume 64, 2006:136-140.
42. Gabizon I, Giladi H, Ovnat A, et al. Small bowel obstruction by a carcinoid tumor in a patient with familial Mediterranean fever. The Israel Medical Association journal : IMAJ. Volume 8, 2006:359-360.
43. Daniels JA, Lederman HM, Maitra A, et al. Gastrointestinal tract pathology in patients with common variable immunodeficiency (CVID): a clinicopathologic study and review. The American journal of surgical pathology. Volume 31, 2007:1800-12.
44. Dunnigan M, Yfantis H, Rapoport AP, et al. Large cell lymphoma presenting as a flare of colitis in a patient with common variable immune deficiency. Digestive diseases and sciences. Volume 52, 2007:830-834.
45. Purim O, Sulkes A, Brenner B. Adjuvant chemotherapy with 5-fluorouracil in a patient with colorectal cancer and Familial Mediterranean Fever. Anti-cancer drugs. Volume 18, 2007:733-735.
46. Brosens LAA, Tytgat KMAJ, Morsink FHM, et al. Multiple colorectal neoplasms in X-linked agammaglobulinemia. Clinical gastroenterology and hepatology : the official clinical practice journal of the American Gastroenterological Association. Volume 6, 2008:115-9.
47. Malhotra RK, Li W. Poorly differentiated gastroenteropancreatic neuroendocrine carcinoma associated with X-linked hyperimmunoglobulin M syndrome. Archives of pathology & laboratory medicine. Volume 132, 2008:847-850.
48. Rosa DD, Pasqualotto AC, Denning DW. Chronic mucocutaneous candidiasis and oesophageal cancer. Medical mycology. Volume 46, 2008:85-91.
49. Thomas ERA, Shanley S, Walker L, et al. Surveillance and treatment of malignancy in Bloom syndrome. Clinical oncology (Royal College of Radiologists (Great Britain)). Volume 20, 2008:375-9.
50. Otabor IA, Abdessalam SF, Erdman SH, et al. Gastric outlet obstruction due to adenocarcinoma in a patient with Ataxia-Telangiectasia syndrome: a case report and review of the literature. World journal of surgical oncology. Volume

7, 2009:29.

51. Yap YL, So JBY. Gastric adenocarcinoma occurring in a young patient with common variable immunodeficiency syndrome. Singapore medical journal. Volume 50, 2009:e201-3.
52. Benjazia E, Turki H, Atig A, et al. Bloom syndrome complicated by colonic cancer in a young Tunisian woman. Clinics and research in hepatology and gastroenterology. Volume 35, 2011:682-684.
53. Dhalla F, da Silva SP, Lucas M, et al. Review of gastric cancer risk factors in patients with common variable immunodeficiency disorders, resulting in a proposal for a surveillance programme. Clinical and experimental immunology. Volume 165, 2011:1-7.
54. Slotta JE, Heine S, Kauffels A, et al. Gastrectomy with isoperistaltic jejunal parallel pouch in a 15-year-old adolescent boy with gastric adenocarcinoma and autosomal recessive agammaglobulinemia. Journal of pediatric surgery. Volume 46, 2011:e21-4.
55. Turkbeyler IH, Pehlivan Y, Comez G, et al. Esophagus cancer and IgA deficiency in a patient with Dubowitz syndrome: a case report. The Tokai journal of experimental and clinical medicine. Volume 36, 2011:29-30.
56. Watkins C, Sahni R, Holla N, et al. Malignancy in Common Variable Immune Deficiency: Report of Two Rare Cases of Gastrointestinal Malignancy and a Review of the Literature. Cardiovascular & Hematological Disorders-Drug Targets. Volume 12: Bentham Science Publishers Ltd., 2012:21-27.
57. Patiroglu T, Eke Gungor H, Arslan D, et al. Gastric signet ring carcinoma in a patient with ataxia-telangiectasia: a case report and review of the literature. Journal of pediatric hematology/oncology. Volume 35, 2013:e341-3.
58. De Petris G, Dhungel BM, Chen L, et al. Gastric adenocarcinoma in common variable immunodeficiency: features of cancer and associated gastritis may be characteristic of the condition. International journal of surgical pathology. Volume 22, 2014:600-606.
59. Loh KP, Hunt JP, Mehendiratta V. Education and imaging. Gastrointestinal:

- Incidentally detected gastric carcinoma in patient with common variable immunoglobulin deficiency. *Journal of gastroenterology and hepatology*. Volume 29, 2014:1127.
60. Staines Boone AT, Torres Martínez MG, López Herrera G, et al. Gastric adenocarcinoma in the context of X-linked agammaglobulinemia: case report and review of the literature. *Journal of clinical immunology*. Volume 34, 2014:134-137.
 61. Vlkova M, Ticha O, Nechvatalova J, et al. Regulatory B cells in CVID patients fail to suppress multifunctional IFN- γ + TNF- α + CD4+ T cells differentiation. *Clinical immunology (Orlando, Fla.)*. Volume 160, 2015:292-300.
 62. Goldberg A, Rose L, McCue P. Common Variable Immunodeficiency Disease and Rectal Squamous Cell Carcinoma: a Case Report of a Rare Syndromic Tumor Type. *Journal of gastrointestinal cancer*. Volume 47, 2016:470-473.
 63. Hayakawa S, Okada S, Tsumura M, et al. A Patient with CTLA-4 Haploinsufficiency Presenting Gastric Cancer. *Journal of Clinical Immunology*. Volume 36: Springer New York LLC, 2016:28-32.
 64. Bratanič N, Kovač J, Pohar K, et al. Multifocal gastric adenocarcinoma in a patient with LRBA deficiency. *Orphanet journal of rare diseases*. Volume 12, 2017:131.
 65. Schober T, Magg T, Laschinger M, et al. A human immunodeficiency syndrome caused by mutations in CARMIL2. *Nature Communications*. Volume 8: Nature Publishing Group, 2017:1-13.
 66. Egg D, Schwab C, Gabrysch A, et al. Increased Risk for Malignancies in 131 Affected CTLA4 Mutation Carriers. *Frontiers in immunology*. Volume 9, 2018:2012.
 67. Pulvirenti F, Pecoraro A, Cinetto F, et al. Gastric Cancer Is the Leading Cause of Death in Italian Adult Patients With Common Variable Immunodeficiency. *Frontiers in immunology*. Volume 9, 2018:2546.
 68. Varricchi G, Pecoraro A, Crescenzi L, et al. Gastroduodenal Disorders in

- Patients with CVID Undergoing Immunoglobulin Therapy. Current pharmaceutical biotechnology. Volume 19, 2018:734-741.
69. Li M, Chen W, Sun X, et al. Metastatic colorectal cancer and severe hypocalcemia following irinotecan administration in a patient with X-linked agammaglobulinemia: a case report. BMC medical genetics. Volume 20, 2019:157.
 70. Antunes PdSL, Tersariol HG, Veiga MMB, et al. NEUROENDOCRINE TUMOR IN A CHILD WITH COMMON VARIABLE IMMUNODEFICIENCY. Revista paulista de pediatria : orgao oficial da Sociedade de Pediatria de Sao Paulo. Volume 38, 2020:e2018146.
 71. HUIZENGA KA, WOLLAEGER EE, GREEN PA, et al. Serum globulin deficiencies in non-tropical sprue, with report of two cases of acquired agammaglobulinemia. The American journal of medicine. Volume 31, 1961:572-80.
 72. KILDEBERG P. The Aldrich syndrome. Report of a case and discussion of pathogenesis. Pediatrics. Volume 27, 1961:362-9.
 73. Hermans PE, Huizenga KA, Hoffman HN, et al. Dysgammaglobulinemia associated with nodular lymphoid hyperplasia of the small intestine. The American journal of medicine. Volume 40, 1966:78-89.
 74. Miller DG. The association of immune disease and malignant lymphoma. Annals of internal medicine. Volume 66, 1967:507-21.
 75. Rádl J, Masopust J, Houstěk J, et al. Paraproteinaemia and unusual dys-gamma-globulinaemia in a case of Wiskott-Aldrich syndrome. An immunochemical study. Archives of disease in childhood. Volume 42, 1967:608-14.
 76. Haerer A. Ataxia- Telangiectasia With gastric adenocarcinoma. Journal of American Medical Association, 1969.
 77. Fraser KJ, Rankin JG. Selective deficiency of IgA immunoglobulins associated with carcinoma of the stomach. Australasian annals of medicine. Volume 19, 1970:165-167.

78. Kersey JH, Spector BD, Good RA. Primary immunodeficiency diseases and cancer: The immunodeficiency-cancer registry. *International Journal of Cancer*. Volume 12, 1973:333-347.
79. Hamoudi AB, Ertel I, Newton WA, et al. Multiple neoplasms in an adolescent child associated with IGA deficiency. *Cancer*. Volume 33, 1974:1134-44.
80. Faraci RP, Hoffstrand HJ, Witebsky FG, et al. Malignant lymphoma of the jejunum in a patient with Wiskott-Aldrich syndrome. Surgical treatment. *Archives of surgery (Chicago, Ill. : 1960)*. Volume 110, 1975:218-220.
81. Hermans PE, Diaz-Buxo JA, Stobo JD. Idiopathic late-onset immunoglobulin deficiency. Clinical observations in 50 patients. *The American journal of medicine*. Volume 61, 1976:221-37.
82. Siegel SE, Hays DM, Romansky S, et al. Carcinoma of the stomach in childhood. *Cancer*. Volume 38, 1976:1781-4.
83. Watanabe A, Hanazono H, Sogawa H, et al. Stomach cancer of a 14-year-old boy with ataxia-telangiectasia. *The Tohoku journal of experimental medicine*. Volume 121, 1977:127-131.
84. Battle WM, Brooks FP. Adenocarcinoma of the stomach with common variable immunodeficiency syndrome. *Archives of internal medicine*. Volume 138, 1978:1682-4.
85. Frai MA. Gastric adenocarcinoma due to ataxia-telangiectasia (Louis-Bar syndrome). *Journal of medical genetics*. Volume 16, 1979:160-161.
86. Lamers CB, Wagener T, Assmann KJ, et al. Jejunal lymphoma in a patient with primary adult-onset hypogammaglobulinemia and nodular lymphoid hyperplasia of the small intestine. *Digestive diseases and sciences*. Volume 25, 1980:553-557.
87. Gonzalez-Vitale JC, Gomez LG, Goldblum RM, et al. Immunoblastic lymphoma of small intestine complicating late-onset immunodeficiency. *Cancer*. Volume 49, 1982:445-449.
88. Mir-Madjlessi SH, Vafai M, Khademi J, et al. Coexisting primary malignant lymphoma and adenocarcinoma of the large intestine in an IgA-deficient boy.

Diseases of the colon and rectum. Volume 27, 1984:822-824.

89. Kinlen LJ, Webster AD, Bird AG, et al. Prospective study of cancer in patients with hypogammaglobulinaemia. *Lancet* (London, England). Volume 1, 1985:263-266.
90. Kiaee F, Azizi G, Rafiemanesh H, et al. Malignancy in common variable immunodeficiency: a systematic review and meta-analysis. *Expert review of clinical immunology*. Volume 15, 2019:1105-1113.
91. Stoffel EM, Boland CR. Genetics and Genetic Testing in Hereditary Colorectal Cancer. *Gastroenterology* 2015;149:1191-1203 e2.
92. Daca Alvarez M, Quintana I, Terradas M, et al. The Inherited and Familial Component of Early-Onset Colorectal Cancer. *Cells* 2021;10.
93. Pedini V, Verga JU, Terrenato I, et al. Incidence of malignancy in patients with common variable immunodeficiency according to therapeutic delay: an Italian retrospective, monocentric cohort study. *Allergy, asthma, and clinical immunology : official journal of the Canadian Society of Allergy and Clinical Immunology*. Volume 16, 2020:54.
94. van Os NJH, Jansen AFM, van Deuren M, et al. Ataxia-telangiectasia: Immunodeficiency and survival. *Clinical immunology (Orlando, Fla.)*. Volume 178, 2017:45-55.
95. Gineau L, Cognet C, Kara N, et al. Partial MCM4 deficiency in patients with growth retardation, adrenal insufficiency, and natural killer cell deficiency. *J Clin Invest* 2012;122:821-32.
96. Cottineau J, Kottemann MC, Lach FP, et al. Inherited GINS1 deficiency underlies growth retardation along with neutropenia and NK cell deficiency. *J Clin Invest* 2017;127:1991-2006.

Figure 1




Figure 2. Gastric cancer reported in patients with inborn errors of immunity




Figure 3. Colon cancer in patients with inborn errors of immunity

