

Title

Describing a complex primary health care population in a learning health system to support future decision support and artificial intelligence initiatives

Authors

Jacqueline K. Kueper MSc,^{1,2} Jennifer Rayner PhD,^{3,4} Merrick Zwarenstein MBBCh,PhD,^{1,3} Daniel J. Lizotte PhD,^{1,2}

1. Department of Epidemiology and Biostatistics, Schulich School of Medicine & Dentistry, Western University, London, Ontario, Canada, N6G 2M1
2. Department of Computer Science, Faculty of Science, Western University, London, Ontario, Canada, N6A 5B7
3. Department of Family Medicine, Schulich School of Medicine & Dentistry, Western University, London, Ontario, Canada, N6G 2M1
4. Alliance for Healthier Communities, Toronto, Ontario, Canada, M6A 3B6

Corresponding author:

Jacqueline K. Kueper
jkueper@uwo.ca

Word Count: 3,782

Figure Count: 4

Table Count: 2

ABSTRACT

Introduction: Learning health systems (LHS) use data to improve care. Descriptive epidemiology to reveal health states and needs of the LHS population is essential for informing LHS initiatives, including development of decision support tools. To properly characterize complex populations, both simple statistical and artificial intelligence techniques can be useful. We present the first large-scale description of the population served by one of the first primary care LHS in North America.

Objectives: Our objective is to describe sociodemographic, clinical, and health care use characteristics of adult primary care clients served by the Alliance for Healthier Communities, which provides team-based primary health care through Community Health Centres (CHCs) across Ontario, Canada.

Methods: Using electronic health record data from 2009-2019 for all CHCs, we perform table-based summaries for each characteristic; and apply unsupervised learning techniques to explore patterns of common condition co-occurrence, care provider teams, and care frequency.

Results: Of the 221,047 eligible clients, those at CHCs that primarily serve those most at risk (homeless, mental health, addictions) tend to have more chronic conditions and social determinants of health, which are also prominent in clients with multimorbidity. Most care is provided by physician and nursing providers, with heterogeneous combinations of other provider types. A subset of clients have many issues addressed within single-visits and there is within- and between-client variability in care frequency. Example methodological considerations learned for future LHS initiatives include the need to carefully consider the level of analysis and associated implications for data quality and target population, heterogeneity in conditions and care characteristics, and non-uniform risk profiles across the care history.

Conclusions: We demonstrate the use of methods from statistics and artificial intelligence, applied with an epidemiological lens, to provide an overview of a complex primary care population. In addition to substantive findings, we discuss implications for future LHS initiatives.

Keywords

Learning Health System, Primary Health Care, Epidemiology, Artificial Intelligence, Unsupervised Machine Learning

Introduction

The recognized potential for analysis of electronic health record (EHR) data to inform health care delivery led to the formalization of the concept of a Learning Health System (LHS) in 2007: a socio-technical system characterized by iterative cycles of data-to-knowledge-to-practice feedback [1, 2]. LHS initiatives target quality improvement, research, or decision support; and usually rely on EHR data from the same population that the findings or end-product are intended to benefit [2–5]. These initiatives can support populations who have historically been excluded from medical research and clinical guideline development, such as those with complex health needs or barriers to participation [6–9].

Primary care, first contact care provided in a community setting over the life course, is inherently complex [10, 11]. The Alliance for Healthier Communities provides team-based primary health care through 72 Community Health Centres (CHCs) across Ontario to clients who face barriers to care and challenges, such as poverty and mental illness, that increase their risk for poor health [12–14]. Population health is a central element of their care model, and the Alliance officially adopted a LHS model in October 2020 [15, 16], making them one of few documented primary care LHSs in North America [5].

A LHS may pursue multiple initiatives to inform and improve care delivery. A first step towards any initiative is identifying needs of clients and providers, which is often driven by internal stakeholders [4]. Descriptive epidemiology is instrumental in outlining health states and needs of populations [17], and may be beneficial to add into these early stages of LHS development both to identify new areas to explore and to support existing ideas. For example, describing how clients are represented in EHR data at a population level may complement clinical experience to identify potential bias or misrepresentation that analyses need to account for to obtain meaningful results [18–20]. In addition to proposed LHS benefits, descriptive studies can contribute towards closing the gap in understanding about the basic functions of primary care in general [21].

To properly understand complex EHR data, we propose using both simple statistical techniques traditionally used in descriptive epidemiology and more complex techniques from artificial intelligence, applied with an epidemiological lens. Simple techniques alone may provide an oversimplified or incorrect view of certain characteristics, which could lead to ineffective or harmful decisions later-on. So, in pursuing our primary purpose of better understanding care provided by the Alliance, we explore the suitability of a variety of techniques for epidemiology of a separate primary care system with its own EHR.

We present the first large-scale descriptive and exploratory study of ongoing primary care clients served by the Alliance using statistical and machine learning methodology. Our *objective* is to summarize sociodemographic, clinical, and health care use characteristics of this population. We use unsupervised learning techniques to identify patterns of multimorbidity, care provider teams, and care access frequency. Findings will provide a foundation for future Alliance LHS initiatives, including those related to their existing interest in using EHR data to segment populations and tailor care. In addition to substantive findings, this work more generally demonstrates the application of an epidemiological lens and use of a variety of methods from statistics and artificial intelligence to effectively describe a complex population and contribute to early stages of a LHS.

Methods

Study population and data source

We use a de-identified extract of the centralized, structured EHR database from all CHCs; clients have unique identifiers to allow tracking of care over time. Issues addressed during care are recorded using Electronic Nomenclature and Classification Of Disorders and Encounters for Family Medicine (ENCODE-FM) [22] and International Classification of Disease (ICD)-10 vocabularies [23]. Primary care EHRs represent an open cohort; Supplementary Appendix 1 (Figure S1) shows the cohort size along calendar- and observation-based time definitions. Clients eligible for inclusion were over 18 years old in 2009, indicated a CHC as their primary care provider, and had at least one encounter at a CHC in 2009 to 2019. Any additional eligibility for specific analyses is described as needed below. We follow RECORD reporting guidelines (Supplementary Appendix 2)[24].

General analysis plan

Sociodemographic, clinical, and health care use characteristics are defined in Supplementary Appendix 3 (Table S1). Methods specific to each category are described below; we perform “table-based summaries” for all, whereby categorical variables are summarized by counts and percentages, and continuous variables by the range, median, mean, and standard deviation. Where specified, findings are stratified by client multimorbidity status (defined below) or CHC “urban at-risk” (UAR) status, which are CHCs located in major urban geographical areas and serve priority populations defined by homelessness and/or mental health and substance use challenges [25]. CHCs without UAR designation still focus on clients with barriers to care but may be in rural or urban settings and do not solely serve clients with the aforementioned complexities [25].

Sociodemographic characteristics

We provide table-based summaries for select fields from the structured EHR client characteristic table and certain ENCODE-FM-derived variables. Missingness of the former occurs at the 1) CHC or provider level, whereby a client is not asked about the characteristic and 2) client level, whereby a client is asked and preferred to not respond. Results are presented overall and stratified by UAR and multimorbidity status.

Clinical characteristics

We describe 20 chronic conditions that define multimorbidity in PC research [26–28] and an additional four conditions of interest identified by Alliance stakeholders. For each condition, clients are assumed to receive related care upon the first record of a relevant code. Conditions are explored in single, composite, and pairwise manners.

Prevalence and incidence

To provide different perspectives on clinical complexity, we calculate two measures of prevalence and one measure of incidence for each of the 24 conditions. We also calculate prevalence of multimorbidity. Our primary multimorbidity definition, including for stratification,

is presence of at least three of the 20 chronic conditions [26–28]. Multimorbidity of at least two conditions is also common and is presented for comparison [27].

- 1) *Eleven-year period prevalence*, based on calendar time, to assess the burden of conditions over the entire observation period (2009–2019). For each condition, the number of clients who ever receive a condition indication is divided by an estimate of the average population size (technical details in Supplementary Appendix 3). Sensitivity analyses include the largest possible denominator: total number of eligible clients, and the smallest reasonable denominator: starting with the middle calendar year (2014), additional clients with at least one visit in adjacent years are added until no prevalence estimate is over 100%. Results are shown overall and UAR-stratified.
- 2) *Observation-based period prevalence*, based on length of client observation, to assess the burden of conditions dependent on the number of years clients have received care at a CHC. To calculate this, clients are separated into 11 sub-cohorts based on the number of years (consecutive 365.25 day intervals, rounded up) between their first and last recorded events. For each sub-cohort and condition, the number of clients who ever receive a condition indication is divided by the number of clients in the sub-cohort. Results are presented as bar graphs.
- 3) *Cumulative incidence*, to assess the rate of condition indications by days of observation. Cumulative incidence curves are plotted using the R package *survival* [29]. To prioritize capture of incident condition-related care, clients with conditions recorded in 2009 are excluded from this analysis.

Condition co-occurrence patterns

To assess co-occurrence for each pair of conditions while adjusting for all of the other conditions, we estimate an *Ising model* using R package *MRFcov* [30, 31] for all conditions except Hepatitis C (Alliance-suggested condition that overlaps with one of the 20 chronic conditions). We convert coefficients, representing the strength of association between each condition pair adjusted for all other conditions, to odds ratios and interpret size using Chen et al. (2010) guidelines [32]. We also view the top frequency-based co-occurrences.

Health care use characteristics

We perform table-based summaries of provider and care access characteristics overall and stratified by UAR CHC, Rural Geography CHC, and client multimorbidity status.

Providers involved

To identify common care provider teams that clients are exposed to across their care histories, we use *non-negative matrix factorization (NMF)* [33] to identify frequently-occurring: 1) “*Ever-seen*” teams whereby dummy variables are used to indicate whether each provider type has ever been involved in care, and 2) *Relative “amount-seen”* teams based on volume of care whereby the number of events associated with each provider type are normalized within clients. For each version, analyses allowing 2,3,5,10, and 15 topics (provider teams) are run with the Python package *sklearn.decomposition.NMF* and the kullback-Leibler divergence distance metric [34]. Resulting topics are interpreted manually. Provider types are maintained as recorded in the EHR except “Other,” “Unknown,” and “Undefined” are combined. We also summarize the top frequency-based provider types involved in care and referrals. Eligible clients require at least one provider type indication in their EHR.

Care access patterns

Complexity of care is measured as the number of events (distinct issues addressed or types of care received) per visit (calendar day of access) to a CHC. *Care frequency* is measured as the number of calendar days at least one event is recorded per year (365.25 day intervals) and per quarter-year (90.30 day intervals). To investigate frequency of care in terms of magnitude and shape (changes in magnitude across care histories), we perform *time series clustering* with the K Medoids algorithm and dynamic time warping distance metric [35] for 1) *short-term clients* with 2-3 observation years and 2) *long-term clients* with 8-10 observation years. For each time interval and cohort, R package *dtwclust* [36] is used to identify 2,3,4, and 5 clusters. Performance is assessed using the silhouette score and visual inspection.

Results

There are 221 047 eligible clients (Supplementary Appendix 3), of whom 64 504 (29.18%) received care at least once in 2009, 141 627 (64.07%) in 2019, and 40 704 (18.4%) received care in both years.

Sociodemographic characteristics

Sociodemographic characteristics are described in **Table 1**, with remaining sub-strata in Supplementary Appendix 3 Table S2. The UAR CHCs tend to provide care to clients who are more commonly male, English-speaking, and have lower levels of education, household income, immigration, stable housing, and/or food security. Clients with multimorbidity tend to be older and more commonly female, reside in rural locations, and have lower levels of education, immigration, stable residence, and/or food security.

Table 1: Sociodemographic characteristics

Characteristic	Values	All Clients n (%)	Urban at Risk CHC ^a n (%)	Multimorbidity n (%)
Number of clients		221 047	35 998	103 172
Age in 2015	25-34	55 505 (25.11)	7976 (22.16)	9346 (9.06)
	35-44	45 646 (20.65)	7540 (20.95)	15 542 (15.06)
	45-54	44 653 (20.2)	8186 (22.74)	23 982 (23.24)
	55-64	37 848 (17.12)	6790 (18.86)	25 578 (24.79)
	65-74	23 162 (10.48)	3644 (10.12)	17 780 (17.23)
	75+	14 233 (6.44)	1862 (5.17)	10 944 (10.61)
Geography	Rural	49 275 (22.29)	6131 (17.03)	26 818 (25.99)
	Urban	167 728 (75.88)	28 538 (79.28)	75 011 (72.70)
	Missing	4044 (1.83)	1329 (3.69)	1343 (1.30)
Sex	Female	127 070 (57.49)	18 699 (51.94)	59 946 (58.10)

Gender	Male	93 294 (42.21)	17 151 (47.64)	43 124 (41.80)
	Other	331 (0.15)	43 (0.12)	19 (0.02)
	Missing	352 (0.16)	105 (0.29)	83 (0.08)
	Female	41 352 (18.71)	5509 (15.30)	21 831 (21.16)
	Gender diverse	340 (0.15)	112 (0.31)	144 (0.14)
	Male	29 366 (13.28)	4585 (12.74)	14 733 (14.28)
Sexual Orientation	Prefer not to answer	1001 (0.45)	51 (0.14)	376 (0.36)
	Missing	148 988 (67.4)	25 741 (71.51)	66 088 (64.06)
	Bisexual	1578 (0.71)	285 (0.79)	690 (0.67)
	Gay	708 (0.32)	192 (0.53)	306 (0.30)
	Heterosexual	57 065 (25.82)	8447 (23.47)	29 105 (28.21)
	Lesbian	485 (0.22)	70 (0.19)	244 (0.24)
	Queer	323 (0.15)	34 (0.09)	91 (0.09)
	Two-Spirit	128 (0.06)	80 (0.22)	61 (0.06)
	Other	246 (0.11)	34 (0.09)	143 (0.14)
	Do not know	924 (0.42)	201 (0.56)	485 (0.47)
Highest Level of Education	Prefer not to answer	7561 (3.42)	877 (2.44)	4078 (3.95)
	Missing	152 029 (68.78)	25 778 (71.61)	67 969 (65.88)
	Post-secondary or equivalent	84 888 (38.4)	12 056 (33.49)	35 763 (34.66)
	Secondary or equivalent	61 831 (27.97)	11 783 (32.73)	32 617 (31.61)
	Less than high school	18 941 (8.57)	3266 (9.07)	10 618 (10.29)
	Other	8507 (3.85)	719 (2.00)	4078 (3.95)
	Do not know	4860 (2.20)	1318 (3.66)	2350 (2.28)
	Prefer not to answer	2950 (1.33)	422 (1.17)	1585 (1.54)
Primary Language	Missing	39 070 (17.67)	6434 (17.87)	16 161 (15.66)
	English	167 163 (75.62)	31 658 (87.94)	79 599 (77.15)
	French	22 547 (10.20)	944 (2.62)	11 091 (10.75)
	Other	26 847 (12.15)	2948 (8.19)	10 710 (10.38)
	Missing	4490 (2.03)	448 (1.24)	1772 (1.72)
Race and Ethnicity	Black	8861 (4.01)	725 (2.01)	3757 (3.64)
	East/Southeast Asian	3739 (1.69)	484 (1.34)	1545 (1.50)

Years Since Arrival in Canada	Indigenous	2944 (1.33)	1577 (4.38)	1641 (1.59)
	Latino	4350 (1.97)	206 (0.57)	1708 (1.66)
	Middle Eastern	2046 (0.93)	344 (0.96)	838 (0.81)
	Other	567 (0.26)	148 (0.41)	306 (0.30)
	South Asian	3597 (1.63)	323 (0.90)	1852 (1.80)
	White	38 464 (17.4)	4531 (12.59)	21 504 (20.84)
	Do not know	838 (0.38)	151 (0.42)	487 (0.47)
	Prefer not to answer	2649 (1.20)	261 (0.73)	1513 (1.47)
	Missing	152 992 (69.21)	27 248 (75.69)	68 021 (65.93)
	0to5yr	13 654 (6.18)	1191 (3.31)	3047 (2.95)
Household Income	6+	51 815 (23.44)	4940 (13.72)	22 722 (22.02)
	None recorded	155 578 (70.38)	29 867 (82.97)	77 403 (75.02)
	\$0 to \$14,999	40 519 (18.33)	8729 (24.25)	17 757 (17.21)
	\$15,000 to \$24,999	21 102 (9.55)	3555 (9.88)	11 081 (10.74)
	\$25,000 to \$39,999	20 877 (9.44)	2988 (8.30)	10 736 (10.41)
	\$40,000 to \$59,999	17 245 (7.80)	2421 (6.73)	8671 (8.40)
	\$60,000 or more	28 494 (12.89)	3862 (10.73)	12 868 (12.47)
	Do not know	15 408 (6.97)	2658 (7.38)	6264 (6.07)
	Prefer not to answer	27 621 (12.50)	4130 (11.47)	14 890 (14.43)
	Missing	49 781 (22.52)	7655 (21.27)	20 905 (20.26)
Household Composition	Couple with children	53 398 (24.16)	6759 (18.78)	20 713 (20.08)
	Couple without child	39 664 (17.94)	5945 (16.51)	22 950 (22.24)
	Extended family	7632 (3.45)	1123 (3.12)	3581 (3.47)
	Grandparents with grandchild(ren)	1746 (0.79)	247 (0.69)	1183 (1.15)
	Siblings	1622 (0.73)	250 (0.69)	669 (0.65)
	Single parent	14 445 (6.53)	2527 (7.02)	6348 (6.15)
	Sole member	32 782 (14.83)	7445 (20.68)	18 597 (18.03)
	Unrelated	8622 (3.90)	1567 (4.35)	2849 (2.76)

	housemates			
	Other	8913 (4.03)	1476 (4.10)	4202 (4.07)
	Do not know	2475 (1.12)	643 (1.79)	1279 (1.24)
	Prefer not to answer	3727 (1.69)	491 (1.36)	1927 (1.87)
	Missing	46 021 (20.82)	7525 (20.90)	18 874 (18.29)
Stable Residence	True	199 349 (90.18)	28 227 (78.41)	90 479 (87.70)
Food Insecurity	True	10 985 (4.97)	2947 (8.19)	7323 (7.10)

^aCHC = Community Health Centre.

Clinical characteristics

Prevalence and incidence

Eleven-year period prevalence estimates range from 1.48% (Hepatitis C) to 80.97% (multimorbidity of two conditions) overall, with generally higher estimates in UAR strata (Table 2). The low sensitivity estimate for the denominator is based on 2012-2015 (n=148 595).

Table 2: Eleven-year period prevalence

Condition	All Clients n (%)	Urban at Risk CHC ^a n (%)
Denominator ^b	165 125	27 256
Hypertension	68 177 (41.29)	12 304 (45.14)
Depression or anxiety	23 828 (14.43)	5533 (20.30)
Chronic musculoskeletal	104 304 (63.17)	18 842 (69.13)
Arthritis	37 201 (22.53)	6906 (25.34)
Osteoporosis	11 462 (6.94)	1950 (7.15)
Asthma or COPD ^c or chronic bronchitis	43 837 (26.55)	9190 (33.72)
Cardiovascular disease	23 311 (14.12)	4673 (17.14)
Heart failure	7994 (4.84)	1564 (5.74)
Stroke or TIA ^d	2967 (1.80)	585 (2.15)
Stomach problem	36 175 (21.91)	7620 (27.96)
Colon problem	24 949 (15.11)	4974 (18.25)
Chronic hepatitis	13 288 (8.05)	2954 (10.84)
Diabetes	35 704 (21.62)	6912 (25.36)
Thyroid disorder	24 793 (15.01)	4217 (15.47)
Any cancer	14 024 (8.49)	2636 (9.67)
Kidney disease or failure	8290 (5.02)	1555 (5.71)
Chronic urinary problem	59 677 (36.14)	11 131 (40.84)

Dementia or Alzheimer's disease	4776 (2.89)	898 (3.29)
Hyperlipidemia	67 175 (40.68)	11 659 (42.78)
Obesity	38 408 (23.26)	6455 (23.68)
Hepatitis C	2436 (1.48)	1173 (4.30)
Smoking or tobacco use	37 355 (22.62)	9597 (35.21)
Substance use	20 853 (12.63)	7508 (27.55)
Lonely or isolated	17 947 (10.87)	5149 (18.89)
Multimorbidity 2+	133 704 (80.97)	24 129 (88.53)
Multimorbidity 3+	103 172 (62.48)	19 237 (70.58)

^aCHC = Community Health Centre

^bDenominator is the approximated average population size across all years (2009-2019)

^cCOPD = Chronic Obstructive Pulmonary Disease

^dTIA = Transient Ischemic Attack

Observation-based period prevalence estimates tend to increase with length of observation; however, cumulative incidence plots for the 156 543 (70.82%) clients without care recorded in 2009 show the rate of condition indications notably decreases after the first year of observation. Sample plots are in **Figure 1**; all are in Supplementary Appendix 1 (Figure S2 and S3).




Figure 1: Example observation-based period prevalence and cumulative incidence plots.

Left column: Observation-based period prevalence. Right column: Cumulative incidence by days of observation.

Condition co-occurrence patterns

Among the 103 172 (46.7%) clients with multimorbidity of at least three chronic conditions, there are 25 162 unique combinations ranging in frequency from 1 (<0.1%) to 845 (0.4%) clients.

Figure 2 presents the *Ising model* results. Pairwise associations between conditions on the log-odds scale range from -0.82 (Osteoporosis—Obesity) to 2.93 (Kidney disease or failure—Chronic urinary problem). There are 1 large, 5 medium, 40 small, and 207 very small associations based on odds ratio magnitude. The five largest positive associations are 1) Kidney Disease or Failure—Chronic Urinary Problem, 2) Smoking or Tobacco Use—Substance Use, 3) Cardiovascular Disease—Heart Failure, 4) Hypertension—Hyperlipidemia, and 5)

Hypertension—Kidney Disease or Failure. In contrast, the top 5 co-occurring conditions based on raw frequency are 1) Hyperlipidemia—Chronic Musculoskeletal, 2) Hypertension—Chronic Musculoskeletal, 3) Hyperlipidemia—Hypertension, 4) Chronic Urinary Problem—Chronic Musculoskeletal, 5) Asthma or COPD or Chronic Bronchitis—Chronic Musculoskeletal. These directly correspond to the conditions with the highest marginal frequencies.




Figure 2: Condition co-occurrence patterns. Heatmap representing the results of the Ising model. Shading is relative to the edge weights or strength of condition co-occurrence. The numbers indicate raw counts in the data; diagonal counts represent clients who only have that single condition. *Legend:* TIA = Transient Ischemic Attack; COPD = Chronic Obstructive Pulmonary Disease.

Health care use characteristics

Table-based summaries of health care use characteristics are in Supplementary Appendix 3 (Table S3). In general, UAR and multimorbidity strata had higher health care use while rural geography CHCs were closer to the overall population.

Providers involved

There are 19 394 unique combinations of the 68 distinct provider types seen across the 220 806 (99.9%) clients with at least one provider type recorded. In terms of referrals, 102 088 (46.2%) clients had at least one internal and 143 922 (65.1%) had at least one external referral recorded. Note internal referrals may not capture “hallway referrals,” whereby a nearby provider provides a quick consult that is not formally recorded.

Figure 3 shows results of the *NMF analysis*, listing the highest-weighted provider types in each topic down to a weight of 3. For the *ever-seen* provider team analysis, physician and nursing provider types emerged most prominently overall. In general, as the number of topics increases, additional provider types emerge and then split apart to dominate separate topics. Exceptions are the high-weighted pairings of nurse and physician and of registered practical nurse and nurse practitioner. Overall, 18 of the 68 possible provider types emerge prominently in at least one topic; only one (respirologist) does not also appear in the amount-seen analysis.

The *amount-seen* provider team analysis has greater weight distributions between provider types within topics. For example, the first of the three-topic analysis has an approximate 1:1:1:6 ratio of care provided by nurse practitioner:nurse:registered practical nurse:physician. In both versions, about half of clients have a non-zero weight for only one of the first two topics; in the amount-seen analysis more clients remain non-zero weight on only one topic as the number of topics increase, e.g. 16.6% versus 2.5% at five topics. In general, results suggests most clients receive the majority of care from physician, nurse practitioner, or nurse provider types, usually in combination with other provider types at a lower volume of care and with heterogeneous co-occurrence. An example of patterns that emerged for other provider types include differences in timing and weight of dietitian/nutritionist and social worker providers between the two analyses. Interpreted alongside the most common provider and referrals types (Supplementary Appendix 3 (Table S4)), findings suggest referrals to dietitian/nutritionist are more common than to social worker, but frequent or longer-term care is more commonly provided by social workers.




Figure 3: Common care provider teams. Boxes represent the topics resulting from the non-negative matrix factorization analysis for A) Ever-seen provider team analysis. B) Relative amount seen provider team analysis. Provider types are listed in order starting with the highest weighted provider; for any given topic, provider types with a weight less than three are not show. *Legend:* NP = Nurse Practitioner, RPN = Registered Practical Nurse.

Care access patterns

Complexity of care from a CHC-perspective is primarily low with 80.4% of client-visits associated with a single-issue and under 1.0% having over five issues addressed (higher intensity); however, from a client-perspective, 24 204 (11.0%) experience at least one visit with over five issues while 38 533 (17.4%) experience a maximum of one issue per visit across their care history. The mean *care access frequency* is 6 days per year (standard deviation=7.4). While 29 191 (13.2%) clients experience at least one year with over 25 days, 7455 (3.4%) average over 25 days per year across their entire care history. There are 8700 (3.94%) clients with at least one frequent care period (year with over 25 days care accessed) and complex care episode (visit with over 5 issues addressed).

For the *time series clustering* analyses, the short-term cohort includes 37 920 clients and 93 625 client-years of observation; the long-term cohort includes 42 855 clients and 387 035 client-years of observation. The silhouette score was always highest for two clusters (Supplementary Appendix 3 (Table S5)). Visual inspection of plots (**Figure 4**) shows high variability within and between clients.

substantive results may be immediately useful to the population of interest, e.g., to inform clinic-level case management and onboarding of new clients.

Sociodemographic characteristics

The CHC EHRs contain rich sociodemographic information, both the presence and absence of which is informative. Social determinants of health were more prevalent in UAR CHC and multimorbidity strata, and there appears to be evidence for the healthy immigrant effect [37]. Completeness rates vary by characteristic and may be due to client, provider, or CHC level decisions. For example, of the 72 059 (32.60%) clients asked about gender only 1001 (1.39%) preferred to not answer. In contrast, more clients, 171 266 (77.48%), were asked about household income but there was a higher tendency to not answer, 27 621 (16.13%). These findings align with a framework to assess selection bias in EHR data that suggests multiple mechanisms are usually responsible for missingness so the focus should be on “what data are observed [instead of missing] and why?” [38] While provider-level decisions may be due to inferring certain characteristics or prioritizing information needed for them to direct care, completeness rates are important for decision support tool performance, which can improve with social determinants of health information [39, 40].

When assessing data quality and completeness, which is emphasized by machine learning for EHR guidelines [2, 4, 19, 41, 42], the implications of pursuing LHS initiatives at different levels should also be considered. For example, a subset of CHCs capture self-reported measures of health, which are valuable research outcomes [43]. While these measures are not suitable for population level analyses, they should be considered for initiatives specific to the collecting CHCs.

Clinical characteristics

Prevalence and incidence

In operationalizing morbidity measures, the denominator must be defined with the intended end-goal in mind. The *eleven-year period prevalence* estimates relate to a CHC-based perspective and are useful for long-term system-level planning, while the *observation-based period prevalence* estimates are more aligned with a client-based perspective and absolute measure of risk. Another consideration is that just as ICD-10 or ENCODE-FM codes do not guarantee true condition presence, the absence of care does not verify absence of conditions [44]. For example, clients may not seek PC when they are healthy, hospitalized, or experiencing barriers to care.

The *cumulative incidence* plots demonstrate that “risk” of condition codes is highest in the first year of observation. Clinically this makes sense, as new clients may have a build-up of unmet care needs. Nonetheless, there are important takeaways for LHS initiatives that require cohort construction. For example, predictive models developed for decision support need to account for the almost qualitative change in risk related to being a new client. Although this care pattern is somewhat unique to PC settings, methods developed for related problems may be useful. For example, accounting for variable lengths of stay in intensive care unit EHRs [45], or handling cold-starts and sparse data for recommender systems [46].

Condition co-occurrence patterns

There is a high prevalence of multimorbidity, but with so many different multimorbidity “compositions” it is hard to see how to make use of the category of multimorbidity. The *Ising model* demonstrates how to go beyond frequency-based comparisons and identify relationships between conditions irrespective of others, but again, this presents as a long tail problem, with very few combinations that are very prominent. PC decision support tools will face the challenge of making recommendations on many different and possibly co-occurring conditions. The majority of existing decision support tools and clinical guidelines focus on a single condition at a time; new techniques for providing evidence-based guidelines or recommendations for these vast numbers of combinations are needed [47–50].

Health care use characteristics

Providers involved

While care for ongoing PC clients is typically led by physicians or nurse practitioners, CHCs include many provider types and LHS initiatives may choose to focus on particular provider type(s). The *NMF analyses* more easily identify reliable patterns of commonly seen provider types and teams than manually sifting through extensive count-based tables. Another use for NMF is dimensionality reduction or data pre-processing, whereby data are summarized to reduce the number of variables that need to be included in an analysis [33]. For example, NMF-derived topics could be used as inputs to a predictive model instead of separate variables to represent each provider type or specific, manually selected combinations.

Care access patterns

Complexity of care from a CHC system-level perspective is primarily low intensity (few problems addressed per visit). The subset of clients who experience higher care complexity do not tend to also have high frequency of care. Sporadic visit patterns may be due to unstable living arrangements or demanding life responsibilities; when there is uncertainty about when a client will return, providers may pack together multiple types of care. The marginal distribution of *care frequency* is right-skewed without a distinct break; most clients experience lower care frequency, but higher frequencies are also observed. In contrast to expectations, we did not identify consistent, distinct client groupings through the time-series clustering, e.g., to indicate a subpopulation of “frequent visitors.” This may be due to restrictions in the types of similarity that dynamic time warping captures. Future analyses could try a different similarity metric or including covariates to account for baseline variability.

Strengths and Limitations

Strengths include the deep interdisciplinary approach used to assess complex, longitudinal EHR data. We used chronic condition definitions recommended for PC research; [26–28] although the algorithms have not been validated for CHCs specifically. Our broad cohort definition supports a high-level overview of the population, but may not be appropriate for specific research questions.

Conclusions

This study demonstrates the use of simple statistics and artificial intelligence techniques, applied with an epidemiological lens, to describe EHR data from a budding LHS. Substantive findings

lay a foundation for future Alliance initiatives and may be informative for other organizations serving complex PC populations.

Key suggestions for future LHS initiatives include the need to carefully deliberate the level of analysis, or who a given initiative should be targeted at (e.g., population or specific CHCs, one or many clinical presentations, all or subset of providers), and the associated implications for how clients will be represented in the data. Representation will depend on analytical-, system-, provider-, and client-level factors. Decision support initiatives need to consider heterogeneity in conditions and care access patterns, including non-uniform risk of condition indications across observation history.

Acknowledgements

This work was supported by the Canadian Institutes of Health Research Canadian Graduate Scholarship-Doctoral to JKK with supervisor DJL.

Statement on conflicts of interest

None declared.

Ethics statement

This study was approved by Western University Review Ethics Board project ID 111353.

Supplementary appendices

Appendix 1 includes extended results presented through figures.

Appendix 2 includes the RECORD reporting guideline checklist.

Appendix 3 includes extended results presented through tables and technical details.

References

- [1] Friedman CP, Allee NJ, Delaney BC, et al. The science of Learning Health Systems: Foundations for a new journal. *Learn Health Syst* 2017; 1: e10020. Doi: 10.1002/lrh2.10020
- [2] Foley T, Horwitz L, Zahran R. *Realising the potential of learning health systems*. Newcastle University: The Learning Healthcare Project. <https://learninghealthcareproject.org/wp-content/uploads/2021/05/LHS2021report.pdf>
- [3] Delaney BC, Peterson KA, Speedie S, et al. Envisioning a Learning Health Care System: The Electronic Primary Care Research Network, a case study. *Ann Fam Med* 2012; 10: 54–59. Doi: 10.1370/afm.1313
- [4] Lindsell CJ, Gatto CL, Dear ML, et al. Learning From What We Do, and Doing What We Learn: A Learning Health Care System in Action. *Academic Medicine* 2021; 96: 1291–1299. Doi: 10.1097/ACM.0000000000004021
- [5] Nash DM, Bhimani Z, Rayner J, et al. Learning health systems in primary care: A systematic scoping review. *BMC Fam Pract* 2021; 22: 126. Doi: 10.1186/s12875-021-01483-z
- [6] Robinson JM, Trochim WMK. An examination of community members', researchers' and health professionals' perceptions of barriers to minority participation in medical research: An

application of concept mapping. *Ethn & Health* 2007; 12: 521–539. Doi: 10.1080/13557850701616987

[7] George S, Duran N, Norris K. A systematic review of barriers and facilitators to minority research participation among African Americans, Latinos, Asian Americans, and Pacific Islanders. *Am J Public Health* 2014; 104: e16–e31. Doi: 10.2105/AJPH.2013.301706

[8] Odierna DH, Schmidt LA. The effects of failing to include hard-to-reach respondents in longitudinal surveys. *Am J Public Health* 2009; 99: 1515–1521. Doi: 10.2105/AJPH.2007.111138

[9] Bonevski B, Randell M, Paul C, et al. Reaching the hard-to-reach: A systematic review of strategies for improving health and medical research with socially disadvantaged groups. *BMC Med Res Methodol* 2014; 14: 42. Doi: 10.1186/1471-2288-14-42

[10] Starfield B. *Primary care. Balancing health needs, services, and technology*. New York, NY: Oxford University Press, Inc., 1998.

[11] CIHR. *CIHR Primary Healthcare Summit 2010 Final Report Summary*. Toronto, Ontario: Canadian Institutes of Health Research, 2010.

[12] Glazier RH, Zagorski B, Rayner J. *Comparison of primary care models in Ontario by demographics, case mix and emergency department use, 2008/09 to 2009/10*. {ICES} {Investigative} {Report}, Toronto, Ont.: Institute for Clinical Evaluative Sciences, <https://www.deslibris.ca/ID/232144> (2012, accessed 6 May 2020).

[13] Booth RG, Richard L, Li L, et al. Characteristics of health care related to mental health and substance use disorders among Community Health Centre clients in Ontario: A population-based cohort study. *cmajo* 2020; 8: E391–E399. Doi: 10.9778/cmajo.20190089

[14] Albrecht D. Community health centres in Canada. *Leadersh Health Serv* 1998; 11: 5–10. Doi: 10.1108/13660759810202596

[15] Alliance for Healthier Communities. Moving Forward as a Learning Health System. *Alliance for Healthier Communities*, <https://myemail.constantcontact.com/EPIC-News--Issue-1.html?soid=1108953382524&aid=uzzy8bphr91U> (2020, accessed 23 November 2020).

[16] Alliance for Healthier Communities. *Towards a Learning Health System: Better Care Tomorrow When We Learn from Today*. Alliance for Healthier Communities. https://www.allianceon.org/sites/default/files/documents/Learning%20Health%20System%20report%202020-10-20%20-%20FINAL_JR.pdf

[17] Cameron D, Jones IG. John Snow, the Broad Street Pump and Modern Epidemiology. *Int J Epidemiol* 1983; 12: 393–396. Doi: 10.1093/ije/12.4.393

[18] Thuraisingam S, Chondros P, Dowsey MM, et al. Assessing the suitability of general practice electronic health records for clinical prediction model development: A data quality assessment. *BMC Med Inform Decis Mak* 2021; 21: 297. Doi: 10.1186/s12911-021-01669-6

[19] Verma AA, Murray J, Greiner R, et al. Implementing machine learning in medicine. *CMAJ* 2021; 193: E1351–E1357. Doi: 10.1503/cmaj.202434

[20] Lee S, Xu Y, D’Souza AG, et al. Unlocking the Potential of Electronic Health Records for Health Research. *Int J Popul Data Sci* 2020; 5: 02. Doi: 10.23889/ijpds.v5i1.1123

[21] Westfall JM, Wittenberg HR, Liaw W. Time to invest in primary care research—commentary on findings from an independent congressionally mandated study. *J Gen Intern Med* 2021; 36: 2117–2120. Doi: 10.1007/s11606-020-06560-0

[22] ENCODE-FM. Electronic Nomenclature and Classification Of Disorders and Encounters for Family Medicine. *ENCODE-FM*, <http://aix1.uottawa.ca/~fammed/fmcencod.htm> (2020, accessed 6 April 2020).

- [23] Organization WH. ICD-10 Version:2019. *World Health Organization*, <https://icd.who.int/browse10/2019/en> (2020, accessed 6 April 2020).
- [24] Benchimol EI, Smeeth L, Guttmann A, et al. The REporting of studies Conducted using Observational Routinely-collected health Data (RECORD) Statement. *PLoS Med* 2015; 12: e1001885. Doi: 10.1371/journal.pmed.1001885
- [25] Glazier RH, Rayner J, Kopp A. *Examining community health centres according to geography and priority populations served, 2011/12 to 2012/13: An ICES chartbook*. Toronto, Ontario: Institute for Clinical Evaluative Sciences in Ontario, <http://www.deslibris.ca/ID/248807> (2015, accessed 20 January 2020).
- [26] Fortin M, Almirall J, Nicholson K. Development of a research tool to document self-reported chronic conditions in primary care. *J Comorb* 2017; 7: 117–123. Doi: 10.15256/joc.2017.7.122
- [27] Lee ES, Lee PSS, Xie Y, et al. The prevalence of multimorbidity in primary care: A comparison of two definitions of multimorbidity with two different lists of chronic conditions in Singapore. *BMC Public Health* 2021; 21: 1409. Doi: 10.1186/s12889-021-11464-7
- [28] Lee YAJ, Xie Y, Lee PSS, et al. Comparing the prevalence of multimorbidity using different operational definitions in primary care in Singapore based on a cross-sectional study using retrospective, large administrative data. *BMJ Open* 2020; 10: e039440. Doi: 10.1136/bmjopen-2020-039440
- [29] Therneau T. A Package for Survival Analysis in R, <https://CRAN.R-project.org/package=survival> (2021).
- [30] van Borkulo CD, Borsboom D, Epskamp S, et al. A new method for constructing networks from binary data. *Sci Rep* 2014; 4: 5918. Doi: 10.1038/srep05918
- [31] Clark NJ, Wells K, Lindberg O. Unravelling changing interspecific interactions across environmental gradients using Markov random fields. *Ecology* 2018; 99: 1277–1283. Doi: 10.1002/ecy.2221
- [32] Chen H, Cohen P, Chen S. How big is a big odds ratio? Interpreting the magnitudes of odds ratios in epidemiological studies. *Commun Stat Simul Comput* 2010; 39: 860–864.
- [33] Wang Y-X, Zhang Y-J. Nonnegative Matrix Factorization: A Comprehensive Review. *IEEE Trans Knowl Data Eng* 2013; 25: 1336–1353. Doi: 10.1080/03610911003650383
- [34] Pedregosa F, Varoquaux G, Gramfort A, et al. Scikit-learn: Machine learning in Python. *J Mach Learn Res* 2011; 12: 2825–2830. <http://jmlr.org/papers/v12/pedregosa11a.html>
- [35] Aghabozorgi S, Seyed Shirkhorshidi A, Ying Wah T. Time-series clustering – A decade review. *Information Systems* 2015; 53: 16–38. Doi: 10.1016/j.is.2015.04.007
- [36] Montero P, Vilar JA. TSclust: An R Package for Time Series Clustering. *J Stat Softw* 2015; 62: 1–43. Doi: 10.18637/jss.v062.i01
- [37] McDonald JT, Kennedy S. Insights into the “healthy immigrant effect”: Health status and health service use of immigrants to Canada. *Soc Sci Med* 2004; 59: 1613–1627. Doi: 10.1016/j.socscimed.2004.02.004
- [38] Haneuse S, Daniels M. A general framework for considering selection bias in EHR-based studies: What data are observed and why? *eGEMs*; 4. Epub ahead of print August 2016. DOI: 10.13063/2327-9214.1203. Doi: 10.13063/2327-9214.1203
- [39] Chen M, Tan X, Padman R. Social determinants of health in electronic health records and their impact on analysis and risk prediction: A systematic review. *J Am Med Inform Assoc* 2020; 27: 1764–1773. Doi: 10.1093/jamia/ocaa143

- [40] Zhao Y, Wood EP, Mirin N, et al. Social determinants in machine learning cardiovascular disease prediction models: A systematic review. *Am J Prev Med*; 0. Epub ahead of print July 2021. DOI: 10.1016/j.amepre.2021.04.016. Doi: 10.1016/j.amepre.2021.04.016
- [41] Wiens J, Saria S, Sendak M, et al. Do no harm: A roadmap for responsible machine learning for health care. *Nat Med* 2019; 25: 1337–1340. Doi: 10.1038/s41591-019-0548-6
- [42] Arbet J, Brokamp C, Meinzen-Derr J, et al. Lessons and tips for designing a machine learning study using EHR data. *J Clin Transl Sci* 2020; 5: 1–10. Doi: 10.1017/cts.2020.513
- [43] CIHI. Patient-reported outcome measures (PROMs). *Canadian Institute for Health Information*, <https://www.cihi.ca/en/patient-reported-outcome-measures-proms> (2022, accessed 7 February 2022).
- [44] Bagley SC, Altman RB. Computing disease incidence, prevalence and comorbidity from electronic medical records. *J Biomed Inform* 2016; 63: 108–111. Doi: 10.1016/j.jbi.2016.08.005
- [45] Zhang L, Chen X, Chen T, et al. DynEHR: Dynamic adaptation of models with data heterogeneity in electronic health records. In: *2021 IEEE EMBS International Conference on Biomedical and Health Informatics (BHI)*. 2021, pp. 1–4. Doi: 10.1109/BHI50953.2021.9508558
- [46] Alyari F, Jafari Navimipour N. Recommender systems: A systematic review of the state of the art literature and suggestions for future research. *Kybernetes* 2018; 47: 985–1017. Doi: 10.1108/K-06-2017-0196
- [47] Moons KGM, Altman DG, Vergouwe Y, et al. Prognosis and prognostic research: Application and impact of prognostic models in clinical practice. *BMJ* 2009; 338: b606. Doi: 10.1136/bmj.b606
- [48] O’Caoimh R, Cornally N, Weathers E, et al. Risk prediction in the community: A systematic review of case-finding instruments that predict adverse healthcare outcomes in community-dwelling older adults. *Maturitas* 2015; 82: 3–21. Doi: 10.1016/j.maturitas.2015.03.009
- [49] Goldstein BA, Navar AM, Pencina MJ, et al. Opportunities and challenges in developing risk prediction models with electronic health records data: A systematic review. *J Am Med Inform Assoc* 2017; 24: 198–208. Doi: 10.1093/jamia/ocw042
- [50] Guthrie B, Boyd CM. Clinical guidelines in the context of aging and multimorbidity. *Public Policy Aging Rep* 2018; 28: 143–149. Doi: 10.1093/ppar/pry038


Abbreviations

CHC: Community Health Centre
 EHR: Electronic Health Record
 ENCODE-FM: Electronic Nomenclature and Classification Of Disorders and Encounters for Family Medicine
 ICD-10: International Classification of Disease - Version 10
 LHS: Learning Health System
 NMF: non-negative matrix factorization
 PC: Primary Care
 UAR: Urban At-Risk

609

Supplementary Material

Appendix 1



Point labels are relative to all eligible clients: % of 221,047.
Horizontal lines are candidate denominators for period prevalence.


611

612


Figure S1: Cohort size by calendar- and observation-based time. Active clients have at least one event during or after the year (calendar- or observation-based) of interest (gap years counted). The number of active observation years refers to the number of 365.25 day periods, counted from the first calendar date that an event was recorded for that client, that clients have at least one event recorded (gap years not counted). Length of observation refers to the number of years from the first to the last year that at least one event is recorded during (gap years counted). Cumulative clients refers to the number of clients who have had at least one event during or before the year of interest. *Legend:* COPD = Chronic Obstructive Pulmonary Disease; TIA = Transient Ischemic Attack; AD = Alzheimer's Disease.

620


621




622



623
624



625
626



627

628






Figure S2: Observation-based period prevalence. Each bar represents the proportion of clients within that observation-based cohort (years are arbitrary 365.25 day consecutive periods between the first and last recorded events) that have at least one indication of the condition of interest across their entire observation history. Conditions are grouped to represent 1) Extra conditions of interest to Alliance stakeholders, 2) 20 chronic conditions, which make up multimorbidity (MM) status, and 3) Overview indicators for the cohorts. *Legend:* COPD = Chronic Obstructive Pulmonary Disease; TIA = Transient Ischemic Attack; AD = Alzheimer's Disease.




640



641

642



643

644

645




Figure S3: Cumulative incidence Cumulative incidence plots by days of observation since the first recorded event. Clients eligible for this analysis must not have any care recorded in the first calendar-year of available data (2009).

651 Appendix 2

652 The RECORD statement – checklist of items, extended from the STROBE statement, that should be reported in observational
653 studies using routinely collected health data.

	Item No.	STROBE items	Location in manuscript where items are reported	RECORD items	Location in manuscript where items are reported
Title and abstract					
	1	(a) Indicate the study's design with a commonly used term in the title or the abstract (b) Provide in the abstract an informative and balanced summary of what was done and what was found	Abstract	<p>RECORD 1.1: The type of data used should be specified in the title or abstract. When possible, the name of the databases used should be included.</p> <p>RECORD 1.2: If applicable, the geographic region and timeframe within which the study took place should be reported in the title or abstract.</p> <p>RECORD 1.3: If linkage between databases was conducted for the study, this should be clearly stated in the title or abstract.</p>	Abstract
Introduction					
Background rationale	2	Explain the scientific background and rationale for the investigation being reported	Introduction		Introduction
Objectives	3	State specific objectives, including any prespecified hypotheses	Introduction		Introduction
Methods					
Study Design	4	Present key elements of study	Introduction,		Introduction,

		design early in the paper	Methods – Study population and data source		Methods – Study population and data source
Setting	5	Describe the setting, locations, and relevant dates, including periods of recruitment, exposure, follow-up, and data collection	Methods – Study population and data source; additional details specific to analyses are presented under the appropriate sub-heading		Methods – Study population and data source; additional details specific to analyses are presented under the appropriate sub-heading
Participants	6	<p>(a) <i>Cohort study</i> - Give the eligibility criteria, and the sources and methods of selection of participants. Describe methods of follow-up</p> <p><i>Case-control study</i> - Give the eligibility criteria, and the sources and methods of case ascertainment and control selection. Give the rationale for the choice of cases and controls</p> <p><i>Cross-sectional study</i> - Give the eligibility criteria, and the sources and methods of selection of participants</p> <p>(b) <i>Cohort study</i> - For matched studies, give matching criteria and number of exposed and unexposed</p>	Methods	<p>RECORD 6.1: The methods of study population selection (such as codes or algorithms used to identify subjects) should be listed in detail. If this is not possible, an explanation should be provided.</p> <p>RECORD 6.2: Any validation studies of the codes or algorithms used to select the population should be referenced. If validation was conducted for this study and not published elsewhere, detailed methods and results should be provided.</p> <p>RECORD 6.3: If the study involved linkage of databases, consider use of a flow diagram or other graphical display to demonstrate the data linkage process, including the number of</p>	Methods

		<i>Case-control study</i> - For matched studies, give matching criteria and the number of controls per case		individuals with linked data at each stage.	
Variables	7	Clearly define all outcomes, exposures, predictors, potential confounders, and effect modifiers. Give diagnostic criteria, if applicable.	Methods, Supplementary Table S1	RECORD 7.1: A complete list of codes and algorithms used to classify exposures, outcomes, confounders, and effect modifiers should be provided. If these cannot be reported, an explanation should be provided.	Supplementary Table S1
Data sources/ measurement	8	For each variable of interest, give sources of data and details of methods of assessment (measurement). Describe comparability of assessment methods if there is more than one group	Methods, Supplementary Table S1		Methods, Supplementary Table S1
Bias	9	Describe any efforts to address potential sources of bias	N/A		N/A
Study size	10	Explain how the study size was arrived at	Methods		Methods
Quantitative variables	11	Explain how quantitative variables were handled in the analyses. If applicable, describe which groupings were chosen, and why	Methods, Supplementary Table S1		Methods, Supplementary Table S1
Statistical methods	12	(a) Describe all statistical methods, including those used to control for confounding (b) Describe any methods used to examine subgroups and interactions (c) Explain how missing data	Methods		Methods

		<p>were addressed</p> <p>(d) <i>Cohort study</i> - If applicable, explain how loss to follow-up was addressed</p> <p><i>Case-control study</i> - If applicable, explain how matching of cases and controls was addressed</p> <p><i>Cross-sectional study</i> - If applicable, describe analytical methods taking account of sampling strategy</p> <p>(e) Describe any sensitivity analyses</p>			
Data access and cleaning methods		..		<p>RECORD 12.1: Authors should describe the extent to which the investigators had access to the database population used to create the study population.</p> <p>RECORD 12.2: Authors should provide information on the data cleaning methods used in the study.</p>	Methods, Supplementary Table S1
Linkage		..		RECORD 12.3: State whether the study included person-level, institutional-level, or other data linkage across two or more databases. The methods of linkage and methods of linkage quality evaluation should be provided.	No linkage
Results					
Participants	13	(a) Report the numbers of individuals at each stage of the	Supplementary Appendix 3	RECORD 13.1: Describe in detail the selection of the persons included in the	Methods

		study (<i>e.g.</i> , numbers potentially eligible, examined for eligibility, confirmed eligible, included in the study, completing follow-up, and analysed) (b) Give reasons for non-participation at each stage. (c) Consider use of a flow diagram		study (<i>i.e.</i> , study population selection) including filtering based on data quality, data availability and linkage. The selection of included persons can be described in the text and/or by means of the study flow diagram.	
Descriptive data	14	(a) Give characteristics of study participants (<i>e.g.</i> , demographic, clinical, social) and information on exposures and potential confounders (b) Indicate the number of participants with missing data for each variable of interest (c) <i>Cohort study</i> - summarise follow-up time (<i>e.g.</i> , average and total amount)	Results, Supplementary Appendix 3		
Outcome data	15	<i>Cohort study</i> - Report numbers of outcome events or summary measures over time <i>Case-control study</i> - Report numbers in each exposure category, or summary measures of exposure <i>Cross-sectional study</i> - Report numbers of outcome events or summary measures	Results, Supplementary Appendix 3		
Main results	16	(a) Give unadjusted estimates and, if applicable, confounder-adjusted estimates and their	Results, Supplementary Appendix 3		

		precision (e.g., 95% confidence interval). Make clear which confounders were adjusted for and why they were included (b) Report category boundaries when continuous variables were categorized (c) If relevant, consider translating estimates of relative risk into absolute risk for a meaningful time period			
Other analyses	17	Report other analyses done—e.g., analyses of subgroups and interactions, and sensitivity analyses	Results, Supplementary Appendix 1,3		
Discussion					
Key results	18	Summarise key results with reference to study objectives	Discussion		Discussion
Limitations	19	Discuss limitations of the study, taking into account sources of potential bias or imprecision. Discuss both direction and magnitude of any potential bias	Discussion	RECORD 19.1: Discuss the implications of using data that were not created or collected to answer the specific research question(s). Include discussion of misclassification bias, unmeasured confounding, missing data, and changing eligibility over time, as they pertain to the study being reported.	Discussion
Interpretation	20	Give a cautious overall interpretation of results considering objectives, limitations, multiplicity of analyses, results from similar studies, and other relevant	Discussion		Discussion

		evidence			
Generalisability	21	Discuss the generalisability (external validity) of the study results	N/A		N/A
Other Information					
Funding	22	Give the source of funding and the role of the funders for the present study and, if applicable, for the original study on which the present article is based	Declarations		Declarations
Accessibility of protocol, raw data, and programming code		..		RECORD 22.1: Authors should provide information on how to access any supplemental information such as the study protocol, raw data, or programming code.	Given the sensitive nature of the data, this information is not shared.

654

655

Appendix 3

Eligible Clients: Of the 881 129 adult clients in the Alliance EHR database in 2009-2019, 232,529 (26.4%) have ongoing primary care client indications, and 221,047 (25.1%) have at least one encounter in 2009-2019 (fully eligible).

Table S1: Characteristic variable definitions

Variable Name	Definition	Source used to guide variable operationalization, if any
Sociodemographic Characteristics		
Age in 2015	2015 minus Year of Birth	
Geography	Geography of place of residence based on Forward Sortation Area: Rural if second digit is 0; Urban if any other valid digit; NA otherwise.	(1)
Sex	Categories as recorded in client characteristic table	
Gender	Collapsed client characteristic table categories	(2)
Sexual Orientation	Categories as recorded in client characteristic table	
Highest Level of Education Completed	Collapsed client characteristic table categories	(2) followed to the extent possible
Primary Spoken Language	Collapsed client characteristic table categories into the two official languages of Canada with remaining languages categorized as Other.	
Race and Ethnicity	Collapsed client characteristic table categories	(3,4)
Year Since Arrival in Canada	Cleaned free text entries from client characteristic table and collapsed into 5 years, 6 or more year, and None Recorded categories. None Recorded cannot differentiate between never-immigrated and never-asked.	
Household Income	Collapsed client characteristic table categories	Note: could not reliably follow guidelines in (2)
Household Composition	Categories as recorded in client characteristic table	
Stable Residence	Collapsed client characteristic table categories into Stable or Unstable (homeless, shelter, other temporary). Additional unstable residence situations were identified as the presence of at least one ENCODE-FM code: 8990, 9433, 9434, 9435, 9436, 9437, 9438, 9439, 9440, 9441, 9442, 9443, 9432, 8982, 8986, 9419, 9424, 8985, 9431, 9415, 9425, 9412, 9414, which were given priority.	List of codes are from an Alliance for Healthier Communities stakeholder
Food Insecurity	At least one ENCODE-FM code: 8972, 9782, 9802, 8971, 9568, 9805	List of codes are from an Alliance for Healthier Communities stakeholder
Clinical Characteristics		
Hypertension	At least one ICD-10 code: i10,i11,i12,i13,i14,i15	(5)

Depression or Anxiety	At least one ICD-10 code: f33,f40,f41	(5)
	At least one ICD-10 code:	(5)
Chronic Musculoskeletal Conditions causing pain or limitation	m40,m41,m42,m43,m44,m45,m46,m47,m48,m49,m50,m51,m52,m53,m54,m60,m61,m62,m63,m65,m66,m67,m68,m70,m71,m72,m73,m74,m75,m76,m77,m78,m79	(5)
Arthritis and/or Rheumatoid Arthritis	At least one ICD-10 code: m05.9,m13.0,m13.9,m15,m16,m17,m18,m19	(5)
Osteoporosis	At least one ICD-10 code: m81	(5)
Asthma, Chronic Obstructive Pulmonary Disease, or Chronic Bronchitis	At least one ICD-10 code: j40,j41,j42,j43,j44,j45,j46	(5)
Cardiovascular Disease (angina, myocardial infarction, atrial fibrillation, poor circulation in the lower limbs)	At least one ICD-10 code: i20,i25,i48,i70,i71,i72,i73,i74,i75,i76,i77,i78,i79	(5)
Heart Failure (including valve problems or replacement)	At least one ICD-10 code: i05,i06,i07,i08,i09,i34,i35,i36,i37,i38,i39,i42,i43,i50	(5)
Stroke and Transient Ischemic Attack	At least one ICD-10 code: g45,i62	(5)
Stomach Problem (irritable bowel, Chron's disease, ulcerative colitis, diverticulosis)	At least one ICD-10 code: k21,k25.7,k29.5	(5)
Colon Problem	At least one ICD-10 code: k50,k51,k52,k57,k58	(5)
Chronic Hepatitis	At least one ICD-10 code: k70,k71,k72,k73,k74,k75,k76,k77	(5)
Diabetes	At least one ICD-10 code: e10,e11,e12,e13,e14	(5)
Thyroid Disorder	At least one ICD-10 code: e00,e01,e02,e03,e04,e05,e06,e07	(5)
Any Cancer (including melanoma, but excluding other skin cancers)	At least one ICD-10 code: c00,c01,c02,c03,c04,c05,c06,c07,c08,c09,c10,c11,c12,c13,c14,c15,c16,c17,c18,c19,c20,c21,c22,c23,c24,c25,c26,c27,c28,c29,c30,c31,c32,c33,c34,c35,c36,c37,c38,c39,c40,c41,c42,c43,c44,c45,c46,c47,c48,c49,c50,c51,c52,c53,c54,c55,c56,c57,c58,c59,c60,c61,c62,c63,c64,c65,c66,c67,c68,c69,c70,c71,c72,c73,c74,c75,c76,c77,c78,c79,c80,c81,c82,c83,c84,c85,c86,c87,c88,c89,c90,c91,c92,c93,c94,c95,c96,c97	(5) modified by removing the 5 year restriction; taking any cancer indication within the 10 year period
Kidney Disease or Failure	At least one ICD-10 code: n18,n19	(5)
Chronic Urinary Problem	At least one ICD-10 code: n03,n11,n18,n20,n21,n22,n23,n25,n26,27,n28,n29,n30,n31,n32,n33,n34,n35,n36,n37,n38,n39,n40,n41,n42,n43,n44,n45,n46,n47,n48,n49,n50,n51+B38	(5)
Dementia or Alzheimer's Disease	At least one ICD-10 code: f00,f01,f02,f03	(5)
Hyperlipidemia (high cholesterol)	At least one ICD-10 code: e78	(5)
Obesity	At least one ICD-10 code: e66	(5) ICD-10 only; no BMI

Hepatitis C	At least one ICD-10 code: b18.2,b19.20,b19.21	(6)
Smoking or Tobacco Use	At least one ENCODE-FM code: 10072,5520,679,9910,5339,5340,5341,5342,5343,5344,5345,5346,5347,5348,5349	JKK selected relevant ENCODE-FM codes based on manual review
Substance Use	At least one ENCODE-FM code: 5304,10004,10005,5305,5306,5307,9754,5308,5309,5310,5311,5312,5313,5314,5315,5316,5317,5318,5319,5320,5321,5322,5323,5324,5325,5326,5327,5328,5329,5330,5331,5332,5333,5334,5335,5336,5337,5338,5350,5351,5352,5353,5354,5355,5356,5357,5358,5359,5360,5361,5362,5363,5364,5365,5366,5367,5368,5369,5370,5371,10007,5372,5373,5374,5375,5376,5377,5378,5379,5380,5381,5382,5383,5384,5385,5386,5387,5388,5389,5390,5391,5392,5393,5394,5395,5396,5397,5398,5399,5400,9845,5401,9844,5401,5402,5403,5404,5405,5406,5407,5408,5409,5410,5411,5412,5413,5414,5415,5416,5417,5418,5419,5420,5421,5422,5423,5424,5425,5426,5427,5428,5429,5430,5431,5432,5433,5434,5435,5436,5437,5438,5439,5440,5441,5442,5443,5444,5445,5446,5447,5448,5449,9277,9278,5450,5451,5452,5453,5454,5455,5456,5457,5458,5459,5460,5461,5462,5463,5464,5465,5466,5467,5468,5469,5470,5471,5472,5473,5474,5475,5476,5477,5478,5479,5480,5481 or recorded in Disabilities Table	JKK selected relevant ENCODE-FM codes based on manual review
Lonely or Isolated	At least one ENCODE-FM code: 5138, 5139, 9265, 9267, 9268, 9512	List of codes are from an Alliance for Healthier Communities stakeholder
Health Care Use Characteristics		
# Years of Observation	Based on records in the service event table: Ceiling of number of days from first to last recorded event divided by 365.25	
# Provider Types Seen	Number of unique provider types recorded in providers involved table. Provider types were maintained as entered except Other, Unknown, and Undefined were collapsed	
# Internal Referrals	Number of records in the internal referrals table	
# External Referrals	Number of records in the external referrals table	
Avg. # Days per Year	Based on records in the service event table: Sum of unique calendar days with at least one event recorded divided by Number of Years of Observation	
Max # Days per Year	Based on records in the service event table: Maximum of number of unique calendar days care is accessed in a single calendar year	
Avg. # Events per Day	Based on records in the service event table: Sum of events divided by number of calendar days care is accessed at least once	
Max # Events per Day	Based on records in the service event table: Maximum number of events recorded in a single calendar day	

Legend: # = Number; Avg. = Average; CHC = Community Health Centre; ENCODE-FM = Electronic Nomenclature and Classification Of Disorders and Encounters for Family Medicine; ICD = International Classification of Disease; SD = Standard Deviation; UAR = Urban at Risk.

References: (1) Canada Post. Addressing guidelines - Forward Sortation Area (FSA) [Internet]. Canada Post. 2022 [cited 2022 Feb 8]. Available from: <https://www.canadapost-postescanada.ca/cpc/en/support/articles/addressing-guidelines/postal-codes.page> (2) CIHI. In Pursuit of Health Equity: Defining Stratifiers for Measuring Health Inequality - A Focus on Age, Sex, Gender, Income, Education and Geographic Location. Ottawa, ON: Canadian Institute for Health Information; 2018 Apr. Available from: <https://www.cihi.ca/sites/default/files/document/defining-stratifiers-measuring-health-inequalities-2018-en-web.pdf> (3) CIHI. Proposed Standards for Race-Based and Indigenous Identity Data Collection and Health Reporting in Canada. Ottawa, ON: Canadian Institute for Health Information; 2020. Available from: <https://www.cihi.ca/en/proposed-standards-for-race-based-and-indigenous-identity-data> (4) Flanagan A, Frey T, Christiansen SL, AMA Manual of Style Committee. Updated guidance on the reporting of race and ethnicity in medical and science journals. JAMA. 2021 Aug 17;326(7):621–7. (5) Fortin M, Almirall J, Nicholson K. Development of a research tool to document self-reported chronic conditions in primary care. J Comorb. 2017 Jan 1;7(1):117–23. (6) Support Path. Hepatitis C ICD-10 Codes. Gilead Sciences; 2015 [cited 2020 Sep 25]. Available from: <https://www.cvph.org/data/files/mysupportpath.pdf>

Table S2: Sociodemographic characteristics with sub-strata

Characteristic	Values	All Clients n (%)	UAR & MM n (%)	UAR & Non-MM n (%)	Non-UAR & MM n (%)	Non-UAR & Non-MM n (%)
Number of clients		221 047	19 237	16761	83 935	101 114
Age in 2015	25-34	55 505 (25.11)	1864 (9.69)	6112 (36.47)	7482 (8.91)	40 047 (39.61)
	35-44	45 646 (20.65)	3154 (16.40)	4386 (26.17)	12 388 (14.76)	25 718 (25.43)
	45-54	44 653 (20.20)	4784 (24.87)	3402 (20.30)	19 198 (22.87)	17 269 (17.08)
	55-64	37 848 (17.12)	4935 (25.65)	1855 (11.07)	20 643 (24.59)	10 415 (10.30)
	65-74	23 162 (10.48)	2952 (15.35)	692 (4.13)	14 828 (17.67)	4690 (4.64)
	75+	14 233 (6.44)	1548 (8.05)	314 (1.87)	9396 (11.19)	2975 (2.94)
Geography	Rural	49 275 (22.29)	3479 (18.08)	2652 (15.82)	23 339 (27.81)	19 805 (19.59)
	Urban	167 728 (75.88)	15 291 (79.49)	13 247 (79.03)	59 720 (71.15)	79 470 (78.59)
	Missing	4044 (1.83)	467 (2.43)	862 (5.14)	876 (1.04)	1839 (1.82)
Sex	Female	127 070 (57.49)	10 647 (55.35)	8052 (48.04)	49 299 (58.73)	59 072 (58.42)
	Male	93 294 (42.21)	8561 (44.50)	8590 (51.25)	34 563 (41.18)	41 580 (41.12)
	Other	331 (0.15)	3 (0.02)	40 (0.24)	16 (0.02)	272 (0.27)
	Missing	352 (0.16)	26 (0.14)	79 (0.47)	57 (0.07)	190 (0.19)
Gender	Female	41 352 (18.71)	3352 (17.42)	2157 (12.87)	18 479 (22.02)	17 364 (17.17)
	Gender diverse	340 (0.15)	52 (0.27)	60 (0.36)	92 (0.11)	136 (0.13)
	Male	29 366 (13.28)	2425 (12.61)	2160 (12.89)	12 308 (14.66)	12 473 (12.34)

Sexual Orientation	Prefer not to answer	1001 (0.45)	37 (0.19)	14 (0.08)	339 (0.40)	611 (0.60)
	Missing	148 988 (67.40)	13371 (69.51)	12 370 (73.80)	52 717 (62.81)	70 530 (69.75)
	Bisexual	1578 (0.71)	141 (0.73)	144 (0.86)	549 (0.65)	744 (0.74)
	Gay	708 (0.32)	94 (0.49)	98 (0.58)	212 (0.25)	304 (0.30)
	Heterosexual	57 065 (25.82)	4703 (24.45)	3744 (22.34)	24 402 (29.07)	24 216 (23.95)
	Lesbian	485 (0.22)	45 (0.23)	25 (0.15)	199 (0.24)	216 (0.21)
	Queer	323 (0.15)	14 (0.07)	20 (0.12)	77 (0.09)	212 (0.21)
	Two-Spirit	128 (0.06)	40 (0.21)	40 (0.24)	21 (0.03)	27 (0.03)
	Other	246 (0.11)	21 (0.11)	13 (0.08)	122 (0.15)	90 (0.09)
	Do not know	924 (0.42)	113 (0.59)	88 (0.53)	372 (0.44)	351 (0.35)
Highest Level of Education	Prefer not to answer	7561 (3.42)	565 (2.94)	312 (1.86)	3513 (4.19)	3171 (3.14)
	Missing	152 029 (68.78)	13501 (70.18)	12 277 (73.25)	54 468 (64.89)	71 783 (70.99)
	Post-secondary or equivalent	84 888 (38.40)	6463 (33.60)	5593 (33.37)	29 300 (34.91)	43 532 (43.05)
	Secondary or equivalent	61 831 (27.97)	6656 (34.60)	5127 (30.59)	25 961 (30.93)	24 087 (23.82)
	Less than high school	18 941 (8.57)	1886 (9.80)	1380 (8.23)	8732 (10.40)	6943 (6.87)
	Other	8507 (3.85)	384 (2.00)	335 (2.00)	3694 (4.40)	4094 (4.05)
	Do not know	4860 (2.20)	734 (3.82)	584 (3.48)	1616 (1.93)	1926 (1.90)
	Prefer not to answer	2950 (1.33)	273 (1.42)	149 (0.89)	1312 (1.56)	1216 (1.20)
	Missing	39 070 (17.67)	2841 (14.77)	3593 (21.44)	13 320 (15.87)	19 316 (19.10)
	English	167 163 (75.62)	17 036 (88.56)	14 622 (87.24)	62 563 (74.54)	72 942 (72.14)
Primary Language	French	22 547 (10.20)	554 (2.88)	390 (2.33)	10 537 (12.55)	11 066 (10.94)
	Other	26 847 (12.15)	1473 (7.66)	1475 (8.80)	9237 (11.00)	14 662 (14.50)
	Missing	4490 (2.03)	174 (0.90)	274 (1.63)	1598 (1.90)	2444 (2.42)
	Black	8861 (4.01)	337 (1.75)	388 (2.31)	3420 (4.07)	4716 (4.66)
	East/SouthEast Asian	3739 (1.69)	248 (1.29)	236 (1.41)	1297 (1.55)	1958 (1.94)
	Indigenous	2944 (1.33)	838 (4.36)	739 (4.41)	803 (0.96)	564 (0.56)
	Latino	4350 (1.97)	102 (0.53)	104 (0.62)	1606 (1.91)	2538 (2.51)
	Middle Eastern	2046 (0.93)	149 (0.77)	195 (1.16)	689 (0.82)	1013 (1.00)
Race and Ethnicity						

	Other	567 (0.26)	79 (0.41)	69 (0.41)	227 (0.27)	192 (0.19)
	South Asian	3597 (1.63)	232 (1.21)	91 (0.54)	1620 (1.93)	1654 (1.64)
	White	38 464 (17.40)	2661 (13.83)	1870 (11.16)	18 843 (22.45)	15 090 (14.92)
	Do not know	838 (0.38)	91 (0.47)	60 (0.36)	396 (0.47)	291 (0.29)
	Prefer not to answer	2649 (1.20)	165 (0.86)	96 (0.57)	1348 (1.61)	1040 (1.03)
	Missing	152 992 (69.21)	14 335 (74.52)	12 913 (77.04)	53 686 (63.96)	72 058 (71.26)
Years Since Arrival in Canada	0 to 5 years	13 654 (6.18)	315 (1.64)	876 (5.23)	2732 (3.25)	9731 (9.62)
	6+ years	51 815 (23.44)	2863 (14.88)	2077 (12.39)	19 859 (23.66)	27 016 (26.72)
	None recorded	155 578 (70.38)	16 059 (83.48)	13 808 (82.38)	61 344 (73.09)	64 367 (63.66)
Household Income	\$0 to \$14,999	40 519 (18.33)	4476 (23.27)	4253 (25.37)	13 281 (15.82)	18 509 (18.31)
	\$15,000 to \$24,999	21 102 (9.55)	2095 (10.89)	1460 (8.71)	8986 (10.71)	8561 (8.47)
	\$25,000 to \$39,999	20 877 (9.44)	1772 (9.21)	1216 (7.25)	8964 (10.68)	8925 (8.83)
	\$40,000 to \$59,999	17 245 (7.80)	1455 (7.56)	966 (5.76)	7216 (8.60)	7608 (7.52)
	\$60,000 or more	28 494 (12.89)	2092 (10.87)	1770 (10.56)	10 776 (12.84)	13 856 (13.7)
	Do not know	15 408 (6.97)	1301 (6.76)	1357 (8.10)	4963 (5.91)	7787 (7.70)
	Prefer not to answer	27 621 (12.50)	2437 (12.67)	1693 (10.10)	12 453 (14.84)	11 038 (10.92)
	Missing	49 781 (22.52)	3609 (18.76)	4046 (24.14)	17 296 (20.61)	24 830 (24.56)
Household Composition	Couple with children	53 398 (24.16)	3280 (17.05)	3479 (20.76)	17 433 (20.77)	29 206 (28.88)
	Couple without child	39 664 (17.94)	3907 (20.31)	2038 (12.16)	19 043 (22.69)	14 676 (14.51)
	Extended family	7632 (3.45)	578 (3.00)	545 (3.25)	3003 (3.58)	3506 (3.47)
	Grandparents with grandchild(ren)	1746 (0.79)	187 (0.97)	60 (0.36)	996 (1.19)	503 (0.50)
	Siblings	1622 (0.73)	140 (0.73)	110 (0.66)	529 (0.63)	843 (0.83)
	Single parent	14 445 (6.53)	1344 (6.99)	1183 (7.06)	5004 (5.96)	6914 (6.84)
	Sole member	32 782 (14.83)	4503 (23.41)	2942 (17.55)	14 094 (16.79)	11 243 (11.12)
	Unrelated housemates	8622 (3.90)	669 (3.48)	898 (5.36)	2180 (2.60)	4875 (4.82)
	Other	8913 (4.03)	788 (4.10)	688 (4.10)	3414 (4.07)	4023 (3.98)
	Do not know	2475 (1.12)	301 (1.56)	342 (2.04)	978 (1.17)	854 (0.84)
	Prefer not to answer	3727 (1.69)	262 (1.36)	229 (1.37)	1665 (1.98)	1571 (1.55)

	Missing	46 021 (20.82)	3278 (17.04)	4247 (25.34)	15 596 (18.58)	22 900 (22.65)
Stable Residence	True	199 349 (90.18)	14813 (77.00)	13 414 (80.03)	75 666 (90.15)	95 456 (94.40)
Food Insecurity	True	10 985 (4.97)	2066 (10.74)	881 (5.26)	5257 (6.26)	2781 (2.75)

Legend: CHC = Community Health Centre; MM = Multimorbidity; UAR = Urban at Risk.

Table S3: Health care use characteristics

Measure	Value	All Clients	UAR CHC	Rural CHC	Multimorbidity
# Years of Observation	Min, Median, Max	(1, 5, 11)	(1, 6, 11)	(1, 7, 11)	(1, 8, 11)
	Mean (SD)	5.6 (3.7)	6.1 (3.8)	6.7 (3.6)	7.4 (3.3)
11 Years of Observation	n (%)	36 724 (16.6)	7976 (22.2)	5374 (25)	29 062 (28.2)
# Provider Types Seen	Min, Median, Max	(0, 4, 19)	(0, 5, 19)	(0, 5, 14)	(0, 6, 19)
	Mean (SD)	4.5 (2.3)	5.1 (2.6)	4.8 (2.1)	5.8 (2.2)
# Internal Referrals	Min, Median, Max	(0, 0, 300)	(0, 1, 300)	(0, 0, 51)	(0, 1, 300)
	Mean (SD)	1.7 (4.3)	2.8 (7.0)	1.4 (2.8)	2.8 (5.7)
# External Referrals	Min, Median, Max	(0, 1, 309)	(0, 2, 309)	(0, 1, 46)	(0, 3, 309)
	Mean (SD)	2.9 (4.5)	3.8 (5.9)	2.5 (3.2)	4.8 (5.5)
Avg. # Days/Year	Min, Median, Max	(0.2, 6, 176.9)	(0.2, 6.9, 129.7)	(0.2, 6.2, 120.3)	(0.3, 9.2, 176.9)
	Mean (SD)	8 (7.4)	9.4 (8.9)	8 (6.7)	11.4 (8.4)
Max # Days/Year	Min, Median, Max	(1, 10, 349)	(1, 12, 245)	(1, 11, 349)	(1, 17, 349)
	Mean (SD)	13.7 (13)	16.8 (16.6)	14.2 (12.1)	20.3 (14.4)
Avg. # Events/Day	Min, Median, Max	(1, 1.2, 66)	(1, 1.2, 29)	(1, 1.2, 31)	(1, 1.2, 31)
	Mean (SD)	1.3 (0.5)	1.3 (0.3)	1.3 (0.3)	1.3 (0.3)
Max # Events/Day	Min, Median, Max	(1, 3, 635)	(1, 3, 635)	(1, 3, 224)	(1, 3, 635)
	Mean (SD)	3.9 (7.8)	4.1 (8.2)	3.8 (6.1)	5.5 (10.7)

Legend: # = Number; Avg. = Average; CHC = Community Health Centre; SD = Standard Deviation; UAR = Urban at Risk.

684 **Table S4: Provider type counts**

Type	Provider Type	Number of Events	% of Events
Provider Involved in Care	Physician	3 693 760	30.1
	Nurse Practitioner (RN-EC)	2 608 238	21.3
	Nurse	2 475 621	20.2
	Registered Practical Nurse (RPN)	990 144	8.1
	Social worker	452 641	3.7
	Other/Unknown/Undefined	448 761	3.7
	Dietitian/Nutritionist	268 395	2.2
	Chiropodist	259 101	2.1
	Counselor	212 799	1.7
	Physiotherapist	171 291	1.4
Internal Referral	Other/Unknown/Undefined	100 649	26.7
	Physician	73 070	19.4
	Nurse Practitioner (RN-EC)	37 333	9.9
	Dietitian/Nutritionist	30 670	8.1
	Nurse	29 326	7.8
	Social worker	28 357	7.5
	Physiotherapist	11 210	3.0
	Chiropractor	9881	2.6
	Chiropodist	9741	2.6
	Counselor	6068	1.6
External Referral	Other/Unknown/Undefined	183 804	28.5
	Dermatologist	41 388	6.4
	Surgeon - general	40 736	6.3
	Gastroenterologist	33 737	5.2
	Surgeon - specialty (eye, heart, brain, etc.)	29 370	4.6
	Physiotherapist	27 639	4.3
	Ear Nose Throat (E.N.T.) specialist	25 791	4.0
	Urologist	22 546	3.5

Gynecologist

21 701 3.4

Cardiologist

20 592 3.2

Table S5: Time series clustering of care access frequency

Analysis	# Clusters (Silhouette Score)	Cluster ID	# Clients	% Clients	Medoid
Short Term by Year	K = 2 (SS = 0.502)	1	11 552	30.5	20, 8
		2	26 368	69.5	6, 2
	K = 3 (SS = 0.301)	1	15 067	39.7	12, 3
		2	16 791	44.3	4, 1
		3	6062	16.0	24, 9
	K = 4 (SS = 0.142)	1	12 931	34.1	5, 1
		2	13 063	34.4	12, 3
		3	5602	14.8	1, 2
		4	6324	16.7	25, 8
	K = 5 (SS = 0.211)	1	3639	9.6	31, 8
		2	11 533	30.4	8, 1
		3	11 155	29.4	3, 2
		4	7722	20.4	12, 5
		5	3871	10.2	17, 11
Short Term by Quarter	K = 2 (SS = 0.541)	1	6068	16.0	10, 5, 5, 7, 3
		2	31 852	84.0	3, 1, 0, 1, 2
	K = 3 (SS = 0.249)	1	10 780	28.4	6, 3, 1, 2, 1
		2	20 431	53.9	2, 0, 0, 0, 1
		3	6709	17.7	6, 1, 3, 4, 3
	K = 4 (SS = 0.044)	1	14 389	37.9	3, 0, 1, 1, 1
		2	8939	23.6	6, 1, 0, 1, 2
		3	9072	23.9	2, 1, 0, 1, 2
		4	5520	14.6	9, 4, 2, 5, 3
	K = 5 (SS = 0.121)	1	4163	11.0	11, 8, 4, 5, 2

Long Term by Year	K = 2 (SS = 0.553)	2	7084	18.7	3, 1, 0, 4, 1
		3	17 282	45.6	3, 1, 0, 1, 2
		4	6111	16.1	5, 2, 1, 0, 1
		5	3280	8.6	6, 0, 1, 6, 1
		1	34 265	80.0	8, 3, 3, 2, 0, 1, 2, 6
	K = 3 (SS = 0.149)	2	8590	20.0	15, 24, 20, 19, 20, 27, 23, 11
		1	15 831	36.9	9, 4, 8, 3, 3, 2, 5, 2
		2	10 557	24.6	24, 9, 13, 19, 12, 12, 16, 6
		3	16 467	38.4	4, 0, 0, 1, 0, 0, 1, 4
		1	2402	5.6	18, 35, 34, 46, 34, 39, 27, 9
	K = 4 (SS = 0.155)	2	23 637	55.2	8, 3, 2, 2, 4, 1, 2, 3
		3	8440	19.7	5, 0, 0, 0, 0, 1, 13, 3
		4	8376	19.5	20, 8, 10, 12, 16, 11, 19, 9
		1	6206	14.5	17, 7, 1, 1, 2, 2, 4, 2
		2	5166	12.1	9, 13, 11, 11, 15, 21, 22, 9
Long Term by Quarter	K = 5 (SS = 0.136)	3	4716	11.0	27, 16, 11, 7, 10, 10, 13, 5
		4	11 254	26.3	6, 2, 6, 6, 7, 7, 12, 3
		5	15 513	36.2	3, 0, 0, 0, 0, 1, 5, 2
		1	36 775	85.8	4, 1, 0, 2, 0, 0, 0, 0, 0, 1, 4, 0, 0, 0, 0, 1, 2, 0, 0, 0, 0, 0, 0, 0, 0, 0, 1, 0, 2
		2	6080	14.2	7, 1, 6, 2, 3, 10, 6, 4, 5, 6, 5, 5, 2, 4, 4, 7, 6, 4, 5, 3, 5, 4, 7, 6, 9, 11, 6, 8, 3, 4
	K = 3 (SS = 0.007)	1	16 528	38.6	1, 0, 5, 0, 1, 2, 0, 0, 0, 3, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 1, 0, 1
		2	9729	22.7	5, 3, 1, 1, 1, 3, 2, 1, 3, 3, 8, 3, 1, 1, 2, 1, 1, 1, 0, 0, 1, 1, 1, 0, 1, 3, 4, 7, 2
		3	16 598	38.7	3, 0, 1, 0, 0, 0, 0, 0, 0, 2, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 1, 3, 0, 1, 1, 4
		1	998	2.3	7, 0, 3, 4, 1, 2, 2, 1, 1, 1, 1, 1, 1, 0, 1, 2, 1, 1, 0, 2, 2, 1, 6, 2, 3, 9, 20, 10, 13, 9, 12, 13, 2
		2	26 775	62.5	3, 0, 1, 0, 2, 0, 0, 0, 1, 2, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 2, 1, 1, 0, 1
	K = 4 (SS = 0.236)				

K = 5 (SS = -0.031)

Eleven-year period prevalence technical details:

Numerator: number of clients with at least one relevant code at any point from 2009 through 2019.

The same process was followed to get estimates for the entire eligible population and for the subset of clients who receive care from urban at risk community health centres.