

## **Symptoms of depression and anxiety among Vietnamese immigrants in Japan during the COVID-19 pandemic: a cross-sectional web-based study**

**Short title: Vietnamese mental health in Japan during COVID-19**

Tadashi Yamashita<sup>a</sup>, Pham Nguyen Quy<sup>b</sup>, Emi Nogami<sup>c</sup>, Erina Seto-Suh<sup>d</sup>, Chika Yamada<sup>e</sup>, Saori Iwamoto<sup>a</sup>, Kyoko Shimazawa<sup>f</sup>, Kenji Kato<sup>g</sup>

<sup>a</sup> Faculty of Nursing, Kobe City College of Nursing, Kobe City, Japan

<sup>b</sup> Department of Medical Oncology, Kyoto Miniren Central Hospital, Kyoto City, Japan

<sup>c</sup> Department of Literature, Mukogawa Women's University, Nishinomiya City, Japan

<sup>d</sup> Human Rights Research Institute, Kindai University, Higashi Osaka City, Japan

<sup>e</sup> Department of Environmental Coexistence, Center for Southeast Asian Studies, Kyoto University, Kyoto City, Japan

<sup>f</sup> Faculty of Global Nursing, Otemae University, Osaka City, Japan

<sup>g</sup> Faculty of Nursing, Kobe Women's University, Kobe City, Japan

Corresponding author: Tadashi Yamashita. E-mail: [yamashita@tr.kobe-ccn.ac.jp](mailto:yamashita@tr.kobe-ccn.ac.jp)

Word Count: 5286

## ABSTRACT

### Background:

Mental health among immigrants in Japan may be related to aspects of COVID-19 infection as well as pandemic-related lifestyle changes, investigating mental health status among Vietnamese residents remains an important public health concern. The mental health status of Vietnamese immigrants in Japan during the COVID-19 epidemic remains unclear. We aimed to examine the mental health status and related factors among Vietnamese immigrants in Japan during the coronavirus disease 2019 (COVID-19) pandemic using data from an online cross-sectional survey administered from September 21 to October 21, 2021.

**Methods:** Patient Health Questionnaire-9 (PHQ-9) and Generalized Anxiety Disorder 7-item (GAD-7) scores were considered the main outcome measures. Multivariable logistic regression was used to identify factors related to symptoms of depression and anxiety, and the chosen variables were entered simultaneously in the survey.

**Results:** Among 621 participants (age: 26.0±4.8 years; male: 347 [55.9%]) who completed the questionnaire, 73.7% reported a decrease in income when compared with the period before the COVID-19 pandemic, and 60.4% reported being recently affected by poor socioeconomic status. Moderate to severe symptoms of depression (PHQ-9 score  $\geq$  10 points) and mild-to-severe symptoms of anxiety (GAD-7 score  $\geq$  5 points) were observed in 203 (32.7%) and 285 (45.9%) individuals, respectively. Factors related to depressive symptoms were age (odds ratio

[OR]=0.94,  $p=0.043$ ), pre-existing health conditions (OR=2.46,  $p<0.001$ ), and subjective socioeconomic status (OR=2.47,  $p<0.001$ ). Factors related to anxiety symptoms were being single (OR=1.72,  $p=0.044$ ), pre-existing health conditions (OR=2.52,  $p<0.001$ ), subjective socioeconomic status (OR=2.72,  $p<0.001$ ), and a partner with whom to discuss one's health (OR=1.66,  $p=0.013$ ).

**Conclusions:** The current findings demonstrate that, when compared with non-Vietnamese Japanese people, people with Vietnamese backgrounds experienced a decrease in income, worsening working conditions, and poor mental health status during the COVID-19 pandemic. These findings highlight the potential contribution of low socioeconomic status and social isolation to poor mental health status.

### **Keywords**

Mental health, Depression, Anxiety, Immigrants, Vietnamese in Japan, COVID-19

## Background

The number of foreign residents in Japan has continued to increase annually, totalling approximately 2,760,000 as of December 2021. Specifically, there has been a rapid increase in the number of Vietnamese immigrants, the population of which has increased from 10,000 in 1990 to 430,000 in 2021. [1] The Immigration Services Agency of Japan recently reported that the country includes the second-largest number of Vietnamese immigrants after China, which can be attributed to the increase of technical interns and skilled labour positions adopted in Japan. [2]

The coronavirus disease 2019 (COVID-19) pandemic has had a severe impact on health and society worldwide. In Japan, approximately 17 million people have been infected with COVID-19, resulting in approximately 36,000 deaths by July 2022. [3] The pandemic has also affected the lives of Vietnamese immigrants living in Japan, who already hold a marginalised and precarious position, by reducing working hours and increasing layoffs. Furthermore, temporary restrictions on returning to Vietnam may have significantly affected mental health among immigrants.

Studies on mental health due to the COVID-19 pandemic showed that people with chronic illnesses and poor job prospects [4] and women suffered more mental distress than men. [5] On the other hand, social capital has been reported to play a protective function in mental health. [6] Studies of immigrants during the COVID-19 pandemic have reported mental health

crises among immigrants in the US, [7] Canada, [8] Europe, [9] India, [10] and South Korea. [11] A web-based survey on mental health of immigrants in 2021 reported most of students from China in Japan have worries about being infected regarding COVID-19 prevention, [12] and approximately 70% of Chinese immigrants in Japan have expressed concerns regarding COVID-19 infection. [13] However, the mental health status of Vietnamese immigrants in Japan during the COVID-19 epidemic remains unclear. Vietnamese immigrants in Japan may have similar concerns regarding COVID-19.

Since mental health among immigrants in Japan may be related to aspects of COVID-19 infection as well as pandemic-related lifestyle changes, investigating mental health status among Vietnamese residents—the second-largest foreign community in Japan—remains an important public health concern. Examining the relationship between socioeconomic factors and mental health among Vietnamese residents in Japan may also aid in the modification and development of health care policies. Therefore, to identify factors that may influence mental health status, the present study aimed to investigate symptoms of depression and anxiety among Vietnamese immigrants in Japan during the COVID-19 pandemic.

## **Methods**

This study was approved by the Kobe City College of Nursing Ethics Research Committee (approval number: 20124-05).

### ***Design and sampling***

This cross-sectional survey was conducted among Vietnamese immigrants in Japan from September 21 to October 21, 2021. During the survey, Japan was in the latter half of the fifth COVID-19 wave. To prevent COVID-19 spread through contact, we conducted an online survey using SurveyMonkey. Since more than 70% of Vietnamese people use the Internet, [14] we conducted the survey using social networks as the main distribution platform. A one-page participants information statement, which also served as a recruitment poster, was posted in Facebook groups and on the personal accounts of the researchers. This statement briefly explained the study background, purpose, and procedure; voluntary nature of participation; anonymity of the questionnaire; strict privacy protection practices; and how to complete the questionnaire. It also included the statement, "By submitting the online questionnaire, the participant agrees that they have read and understand the participants information statement and agree to participate in the survey." The inclusion criteria for participation were as follows: Vietnamese citizenship or Japanese citizenship in those of Vietnamese descent, current residence in Japan, and age  $\geq 18$  years. Participants received an online token for 200 Japanese yen (\$1.8) after participation. To avoid duplicate responses from participants, SurveyMonkey was used to allow only one response from the same terminal.

### ***Data collection tools***

The survey involved a self-administered questionnaire, which was used to collect data regarding demographic variables such as sex, age, duration of residence in Japan, area of residence, marital status, education level, birth country, visa status, Japanese language level, and pre-existing health conditions. Based on the opinions of Japanese infectious disease experts, we also classified residence into three categories based on the number of COVID-19 infections as of December 2021 (>2,000 per 100,000: Tokyo, Osaka, Okinawa; 1,999–1,000 per 100,000: Hokkaido, Saitama, Chiba, Kanagawa, Aichi, Kyoto, Nara, Hyogo, Fukuoka; <999 per 100,000: Others).

Economic and employment status was assessed based on national health insurance, public assistance, subjective socioeconomic status, and monthly income compared with those before the COVID-19 pandemic. International students were included in this survey. Under Japanese law, international students are allowed to work part-time or up to 28 hours per week outside their visa status. This income may account for a large portion of their living expenses. Therefore, in this survey, we also ascertained the income and working conditions of international students. Social support status was assessed based on whether patients had partners with whom they can talk about their health. Levels of depression and anxiety were assessed using the Patient Health Questionnaire-9 (PHQ-9) and Generalized Anxiety Disorder 7-item (GAD-7), respectively, which have been validated for assessing mental health status in

Vietnam. [15-20] In our study, the Cronbach's alpha values for the reliability of the PHQ-9 and GAD-7 were 0.844 and 0.933, respectively.

### ***Statistical analysis***

We performed descriptive analyses using means and standard deviations (SD) for continuous variables as well as counts and percentages for binary or categorical variables. Each PHQ-9 item was rated on a four-point Likert scale ranging from 0 (not at all) to 3 (almost every day), with the total score ranging from 0 to 27. The severity of depression was divided into five categories based on PHQ-9 scores, as follows: none (0–4 points), mild (5–9 points), moderate (10–14 points), severe (15–19 points), and extremely severe (20–27 points). [21] A PHQ-9 score  $\geq 10$  points was considered to indicate clinically relevant depressive symptoms, based on a previous survey of COVID-19 in Vietnam. [15, 17-19] The severity of anxiety was classified into four categories based on GAD-7 scores: no anxiety (0–4 points), mild (5–9 points), moderate (10–14 points), and severe (15–21 points). [22] Each question item ranges from 0 (not at all) to 3 (almost every day) on a 4-point Likert scale, and the total score ranged from 0 to 21. GAD-7 scores  $\geq 5$  were considered to indicate clinically relevant symptoms of anxiety, as previously reported. [16, 20]

We conducted multivariable logistic regression analysis (forced-entry method) to identify factors related to symptoms of depression and anxiety, which were used as the

dependent variables. The potential predictors included sociodemographic variables such as age, sex, marital status, education level, pre-existing health conditions; immigrant-specific variables such as the duration of residence in Japan, visa status, Japanese language proficiency; economic variables; variables related to the COVID-19 pandemic such as the rate of COVID-19 infection in the area; and variables related to social connectedness, such as the availability of a conversation partner. As employment status affects economic status, economic status, not employment status, was used as a predictor in this analysis. Inclusion of variables was based on previous studies of mental health among immigrants, [23, 24] and the chosen variables were entered simultaneously for multivariable regression. We created a correlation matrix before entering the independent variables; further, we confirmed that there were no strong correlations among the independent variables with  $r > 0.80$ . Statistical analyses were performed using the SPSS software (version 19.0; IBM Corp., Armonk, NY, USA). Statistical significance was set at  $P < 0.05$  (2-sided tests).

## Results

A total of 1,046 people participated in the online survey, with 652 participants providing complete responses. After deleting responses with missing values or outliers, we included 621 responses. All participants were Vietnamese. The mean age ( $\pm$ SD) was  $26.0 \pm 4.8$  years; further, the mean duration of residence in Japan ( $\pm$  SD) was  $3.4 \pm 3.1$  years.

Approximately 30% of the respondents lived in areas with high infection rates. Most respondents (79.2%) were single (including divorced or bereaved), and almost all were born in Vietnam (99.0%). The most common visa status was "technical intern training (Ginjojisshu)" (29.5%), followed by "international student" (29.3%) and "status of residence based on employment" (26.9%). The most common educational level was high school (38.3%), followed by college or university (34.5%) and technical school (21.3%). Further, 21.3% of respondents had a pre-existing health conditions, and the most common was gastrological (stomach/oesophagus/duodenum) conditions (5.5%), followed by otorhinolaryngological (5.2%) and musculoskeletal/joint disease (5.2%) conditions (Table 1).

<Please Insert Table 1 here>

Regarding economic status, <5.5% and 6.9% of respondents lacked national health insurance and used public assistance, respectively. Approximately 34.6% of respondents reported a decrease in monthly income when compared with the period before the COVID-19 pandemic, with approximately 39.1% reporting a slight decrease. Approximately 14.2% and 46.2% of participants reported very poor and poor subjective socioeconomic status, respectively. Additionally, 18.7% of respondents reported dismissal from work/unemployment during the COVID-19 pandemic, while 64.1% reported a reduction in the number of working

days (Table 2).

**Table 2:** Economic and employment status\* (n = 621)

Variable	Item	n	%
National health insurance	Yes	587	94.5
	No	34	5.5
Public assistance	Used	43	6.9
	Not used	578	93.1
Monthly income when compared with that before COVID-19	Decrease of $\geq 40\%$	215	34.6
	Decrease of 10–40%	243	39.1
	Decrease $< 10\%$	146	23.5
	Increase $> 10\text{--}40\%$	15	2.4
	Increase $> 40\%$	2	0.3
Subjective socioeconomic status	Really good	5	0.8
	Slightly better	47	7.6
	General	194	31.2
	Slightly difficult	287	46.2
	Very difficult	88	14.2
Employment status (multiple answers)	No change/no impact	93	15.0
	Dismissal/Unemployment	116	18.7
	Change in occupation type	76	12.2
	Change in work content	101	16.3
	Reduction in working days	398	64.1
	Increase in working days	17	2.7
	Other	44	7.1

\* International students were included in the statistical analysis for “Economic and employment status” since they are allowed to have part-time jobs in Japan.

Further, 30.3% of participants reported having a partner with whom to discuss their health, with family (21.9%) being the most common, followed by Vietnamese friends (9.4%), work colleagues (8.7%), and medical professionals (including through online consultation)

(4.3%) (Table 3).

**Table 3:** Social support status (n = 621)

Variable	Item	n	%
Partner with whom to discuss health status	Yes	188	30.3
	No	433	69.7
Who is the partner with whom you can discuss your health? (Multiple answers)	Family	136	21.9
	Japanese friend	21	3.4
	Vietnamese friend	59	9.5
	People at the same office	54	8.7
	People at the same school	15	2.4
	Medical professionals (including through online consultation)	27	4.3
	Religious facilities (temples, churches, etc.)	4	0.6
	Other	23	3.7

The mean PHQ-9 score was  $7.9 \pm 6.0$ . Mild, moderate, severe, and very severe depressive symptoms were observed in 212 (34.1%), 117 (18.8%), 52 (8.4%), and 34 (5.5%) participants, respectively. Further, 203 (32.7%) individuals had a PHQ-9 score  $\geq 10$  points. The mean GAD-7 score was  $5.4 \pm 5.3$ . Mild, moderate, and severe symptoms of anxiety were observed in 169 (27.2%), 69 (11.1%), and 47 (7.6%) participants, respectively. Moreover, 285 (45.9%) individuals had a GAD-7 score  $\geq 5$  points (Table 4).

**Table 4:** Symptoms of depression and anxiety (n = 621)

Variable	Item	n	%
----------	------	---	---

Depressive symptoms (PHQ-9)	Mean $\pm$ SD	7.9 $\pm$ 6.0	
	No depression (0–4 points)	206	33.2
	Mild (5–9 points)	212	34.1
	Moderate (10–14 points)	117	18.8
	Severe (15–19 points)	52	8.4
	Very severe (20–27 points)	34	5.5
	10–27 points	203	32.7
Anxiety symptoms (GAD-7)	Mean $\pm$ SD	5.4 $\pm$ 5.3	
	No anxiety (0–4 points)	336	54.1
	Mild (5–9 points)	169	27.2
	Moderate (10–14 points)	69	11.1
	Severe (15–21 points)	47	7.6
		5–21 points	285

We performed a multivariable logistic regression analysis by the forced-entry method. Twenty-five participants with missing responses for residency status were considered as missing values. The results of the  $\chi^2$  test were significant ( $P < 0.001$ ) for the PHQ-9 and GAD-7 regression models. The Hosmer and Lemeshow's test for PHQ-9 and GAD-7 scores revealed  $P$  values of 0.671 and 0.194, respectively, and the goodness-of-fit for the model was acceptable. Additionally, there were no outliers in which the predicted values exceeded the measured values by  $\pm 3$  SD. Depressive symptoms were associated with age (odds ratio [OR]=0.94, 95% confidence interval [CI]=0.89–0.99), pre-existing health conditions (OR=2.46, 95% CI=1.61–3.76), and low socioeconomic status (OR=2.47, 95% CI=1.64–3.71). Factors related to anxiety included being single (OR=1.72, 95% CI=1.01–2.93), pre-existing health conditions (OR=2.52, 95% CI=1.63–3.88), low socioeconomic status (OR=2.72, 95% CI=1.87–3.97), and not having

a partner with whom to discuss one's health (OR=1.66, 95% CI=1.11–2.47) (Table 5).

<Please Insert Table 5 here>

## Discussion

Our findings indicated that many Vietnamese immigrants in Japan experienced a decrease in income and worsening working conditions during the COVID-19 pandemic. In addition, approximately half of respondents reported symptoms of depression and anxiety, which were significantly associated with pre-existing health conditions, and low socioeconomic status in the multivariable regression analysis. Beyond these associations, symptoms of depression were associated with age, and the symptoms of anxiety were associated with the availability of partners with whom to discuss one's health. These results demonstrate that there is a mental health problem among Vietnamese immigrants in Japan, highlighting the need for financial support and measures for preventing social isolation.

We observed a high prevalence of depression symptoms, with 61.3% and 32.7% of respondents reporting mild-to-severe and moderate-to-very severe symptoms, respectively. This prevalence is higher than previously reported values. The prevalence of depressive symptoms was greater in our study than in other studies conducted after the start of the pandemic. Porter et al. [25] reported that the prevalence of mild-to-severe depression as measured using the PHQ-8 was 9.5% among the young population, while Luong et al. [26] reported a prevalence rate of 18.4% for moderate-to-severe depression (PHQ-9 scores) among

pregnant women in Vietnam. Other studies using the PHQ-9 have reported that the rate of moderate-to-severe depressive symptoms was higher among Vietnamese immigrants than among Japanese citizens: 17.9% among adults aged 18 years and older in 2020; [27] 11.6% in 2020 and 16.6% in 2021 among college students; [28, 29] and 14.8% among hospital workers in 2020. [30] The current evidence indicates that depressive symptoms among Vietnamese immigrants in Japan appear to be more severe than those among Japanese and Vietnamese individuals living in their home countries.

We observed a high prevalence of anxiety symptoms in the current study, with 45.9% and 18.7% of respondents reporting mild-to-severe and moderate-to-severe anxiety symptoms, respectively. A previous study reported that only 33.1% and 5.5% of clinical health care workers in Vietnam experienced mild-to-severe and moderate-to-severe symptoms of anxiety, respectively. [20] Another recent study noted that 25.5% and 4.3% of healthcare workers in Vietnam exhibited mild-to-severe and moderate-to-severe anxiety symptoms, respectively. [16] Ueda et al. [31] reported mild-to-severe symptoms of anxiety in 31.6% of Japanese people 18 years and older, based on GAD-7 scores. As observed for depressive symptoms, the prevalence of anxiety appears to be similar between Vietnamese immigrants and other immigrant populations in Japan, with immigrants experiencing more severe symptoms than Japanese and Vietnamese individuals living in their home countries.

The World Health Organization recently reported that economic problems have affected the

mental health of migrants during the COVID-19 pandemic. [32] In December 2020, approximately 46,000 foreigners living in Japan were laid off, unable to continue working, or unable to return home, [4] many of whom were likely Vietnamese. In this survey, 60.4% of participants reported a slightly difficult or worse economic situation relative to the pre-pandemic period, while 14.2% reported a severe change in economic status, indicating that Vietnamese immigrants in Japan remain anxious about their current and future lives in Japan. The Ministry of Health, Labour, and Welfare (Japan) has been providing financial support including a welfare fund loan system, public assistance, counselling, support for finding employment, and support for early return to the home country; [4, 33, 34] however, there have been no positive outcomes. Nonetheless, there is a need to continue supporting the economic and daily life of Vietnamese immigrants in Japan since worsening economic situations and instability in daily life are strongly associated with mental health status.

Our findings additionally demonstrated that the presence of pre-existing health conditions significantly impacted mental health among Vietnamese immigrants in Japan. Among the most important health issues in Japan is the health disparity among immigrants living in Japan given the lack of access to medical services. [35] Another report on immigrants in Japan who participated in free medical check-ups conducted by a Non-Governmental Organization revealed that many were not connected to medical services. [36] Fear of COVID-19 has also generally reduced the number of people receiving medical services in Japan. [37]

The COVID-19 pandemic further restricted access to medical services among immigrants in Japan, including Vietnamese residents, which might contribute to increases in their anxiety.

Our analysis also revealed that age differences have a significant impact on depressive symptoms. The result was that depressive symptoms were less likely to occur with increasing age. This means that young people are at high risk. Similar reports have been seen in Vietnam and Japan under the COVID-19 pandemic; [25, 38] therefore, this result may not be a characteristic issue for Vietnamese immigrants in Japan. However, young Vietnamese individuals in Japan find it difficult to respond to changes in their living environment due to COVID-19 pandemic, which is expected to lead to an increase in their depressive symptoms. Our analysis also revealed that being single and absence of partners with whom to discuss one's health significantly affected the risk of experiencing symptoms of anxiety. The Immigration Services Agency of Japan reported that foreigners living in Japan had few connections with others outside of work. [39] A South Korean study conducted in October 2020 reported that immigrants who live alone have moderate-to-extreme anxiety symptoms when compared with those living with their families. [11] These findings indicate that connections with others, especially family members, are important for maintaining mental health among immigrants. Thus, Vietnamese immigrants in Japan will require further support to connect with their families, Vietnamese friends, and others during the remainder of the COVID-19 pandemic.

The present study had some limitations, including its self-reported and retrospective

nature, which may have introduced recall bias. In addition, the survey may not be fully representative of the Vietnamese population living in Japan since some older adults of Vietnamese people may not have been able to participate using the online format. Since this survey was collected via social media, responses may have been biased toward Vietnamese residents in Japan who are interested in mental health. Therefore, there is a possibility of participant selection bias. Finally, the study's research design did not consider the association between family networks and mental health at the time of immigration noted by Dykxhoorn et al. [40] Further research is needed to clarify the actual mental health status of Vietnamese people living in Japan by re-examining the selection of participants and confounding factors.

## **Conclusions**

In this cross-sectional survey, we examined the mental health status of 621 Vietnamese immigrants living in Japan during the COVID-19 pandemic. Our analysis yielded the following five main findings: (1) many Vietnamese immigrants experienced economic difficulties; (2) symptoms of depression (moderate-to-very-severe) and anxiety (mild-to-severe) were observed in 32.7% and 45.9% of respondents; (3) low subjective socioeconomic status and pre-existing health conditions were associated with depression and anxiety; (4) younger people are more likely to have depressive symptoms; and (5) being single and the absence of partners with whom to discuss one's health were associated with anxiety. Together, our findings demonstrated that mental health status was worse among Vietnamese immigrants than among

Japanese citizens during the COVID-19 pandemic, and that there is a need for financial support and measures that can prevent poor mental health status among them.

### **List of Abbreviations**

CI: confidence interval

GAD-7: Generalized Anxiety Disorder-7

OR: odds ratio

PHQ-9: Patient Health Questionnaire-9

SD: standard deviation

### **Acknowledgments**

We wish to extend special thanks to all the respondents who participated in the survey. We also thank Professor Hiroya Matsuo for his helpful advice.

### **Sources of Funding**

This work was supported by JSPS KAKENHI (Grant Number JP19K11277 and 22H03420).

### **Author contributions**

T. Y. conceived the original idea for the study. T. Y., P. N. Q., S. I., and K. K. designed the questionnaire. T. Y., C. Y., and K. K. analysed the data and wrote the first draft of the manuscript.

P. N. Q., N. E., E. S., C. Y., S. I., K. S., and K. K. contributed to the interpretation of the results.

All authors made substantial intellectual contributions to the study and approved the final draft of the manuscript.

### **Data availability**

The data that support the findings of this study are available from the corresponding author upon reasonable request.

### **Declaration of interest**

None.

### **Ethics approval and consent to participate**

This study was approved by the Kobe City College of Nursing Ethics Research Committee (approval number: 20124-05).

### **Consent for publication**

Not applicable.

## References

- 1 Immigration Services Agency of Japan. Number of foreign residents as of the End of December 2021; 2021.  
[https://www.moj.go.jp/isa/policies/statistics/toukei\\_ichiran\\_touroku.html](https://www.moj.go.jp/isa/policies/statistics/toukei_ichiran_touroku.html) Accessed 23 Aug 2022
- 2 Immigration Services Agency of Japan. Application status of the specific technology system; 2022. <https://www.moj.go.jp/isa/content/001359454.pdf> Accessed 27 Aug 2022
- 3 Ministry of Health, Labour and Welfare (Japan). Incidence of COVID-19 in Japan; 2022.  
<https://www.mhlw.go.jp/stf/covid-19/kokunainohasseijoukyou.html> Accessed 21 Aug 2022
- 4 Kim HH, Laurence J. COVID-19 restrictions and mental distress among American adults: Evidence from Corona Impact Survey (W1 and W2). *J Public Health (Oxf)*. 2020;42:704–11.  
<https://doi.org/10.1093/pubmed/fdaa148>, PMID: 32880640, PMCID: PMC7499737
- 5 Sønderskov KM, Dinesen PT, Santini ZI, Østergaard SD. The depressive state of Denmark during the COVID-19 pandemic. *Acta Neuropsychiatr*. 2020;32:226–8.  
<https://doi.org/10.1017/neu.2020.15>. Epub 22 Apr 2020. PMID: 32319879, PMCID: PMC7176490
- 6 Chan SM, Chung GK, Chan YH, Woo J, Yeoh EK, Chung RY, et al. The mediating role of individual-level social capital among worries, mental health and subjective well-being among adults in Hong Kong during the COVID-19 pandemic. *Curr Psychol*. 2021:1–11.

<https://doi.org/10.1007/s12144-021-02316-z> [Epub ahead of print]. PMID: 34580570,  
PMCID: PMC8459135

7 Galletly CL, Lechuga J, Dickson-Gomez JB, Glasman LR, McAuliffe TL, Espinoza-Madrigal I. Assessment of COVID-19–Related immigration concerns among Latinx immigrants in the US. *JAMA Netw Open*. 2021;4:e2117049.  
<https://doi.org/10.1001/jamanetworkopen.2021.17049>

8 Evra R, Mongrain E. Mental health status of Canadian immigrants during the COVID-19 pandemic; 2020. <https://www150.statcan.gc.ca/n1/pub/45-28-0001/2020001/article/00050-eng.htm> Accessed 28 Aug 2022

9 Spiritus-Beerden E, Verelst A, Devlieger I, Langer Primdahl N, Botelho Guedes F, Chiarenza A, et al. Mental health of refugees and migrants during the COVID-19 pandemic: The role of experienced discrimination and daily stressors. *Int J Environ Res Public Health*. 2021;18:6354.  
<https://doi.org/10.3390/ijerph18126354>

10 Choudhari R. COVID 19 Pandemic: Mental health challenges of internal migrant workers of India. *Asian J Psychiatr*. 2020;54:102254. <https://doi.org/10.1016/j.ajp.2020.102254>

11 Acharya SR, Moon DH, Chun JH, Shin YC. COVID-19 and mental health: Anxiety disorders among immigrants due to COVID-19 outbreak in South Korea. *Int J Psychiatry Med*. 2022;912174211042695;57:323–37. <https://doi.org/10.1177/00912174211042695>

12 Teng Y, Takemoto K. Associations between psychological conditions and social capital

- among Chinese international students in Japan during the COVID-19 pandemic. *J Disaster Res.* 2022;17:136–43. <https://doi.org/10.20965/jdr.2022.p0136>
- 13 Hu Q, Umeda M. Stress, anxiety, and depression for Chinese residents in Japan during the COVID-19 pandemic. *Int J Environ Res Public Health.* 2021;18:4958. <https://doi.org/10.3390/ijerph18094958>
- 14 The World Bank. Individuals using the Internet (% of population). Vietnam. 2022. <https://data.worldbank.org/indicator/IT.NET.USER.ZS?locations=VN> Accessed 28 Aug 2022
- 15 Do BN, Nguyen PA, Pham KM, Nguyen HC, Nguyen MH, Tran CQ, et al. Determinants of health literacy and its associations with health-related behaviors, depression among the older people with and without suspected COVID-19 symptoms: A multi-institutional study. *Front Public Health.* 2020;8:581746. <https://doi.org/10.3389/fpubh.2020.581746>
- 16 Doan QH, Tran NN, Than MH, Nguyen HT, Bui VS, Nguyen DH, et al. Depression, anxiety and associated factors among frontline hospital healthcare workers in the fourth wave of COVID-19: Empirical findings from Vietnam. *Trop Med Infect Dis.* 2021;7:3. <https://doi.org/10.3390/tropicalmed7010003>
- 17 Nguyen HC, Nguyen MH, Do BN, Tran CQ, Nguyen TTP, Pham KM, et al. People with suspected COVID-19 symptoms were more likely depressed and had lower health-related quality of life: The potential benefit of health literacy. *J Clin Med.* 2020;9:965. <https://doi.org/10.3390/jcm9040965>

- 18 Pham T, Bui L, Nguyen A, Nguyen B, Tran P, Vu P, et al. The prevalence of depression and associated risk factors among medical students: An untold story in Vietnam. PLOS ONE. 2019;14:e0221432. <https://doi.org/10.1371/journal.pone.0221432>
- 19 Pham KM, Pham LV, Phan DT, Tran TV, Nguyen HC, Nguyen MH, et al. Healthy dietary intake behavior potentially modifies the negative effect of COVID-19 lockdown on depression: A hospital and Health Center Survey. Front Nutr. 2020;7:581043. <https://doi.org/10.3389/fnut.2020.581043>
- 20 Quang N, Kien NT, Anh PN, Anh DTV, Nghi TDB, Lan PP, et al. The level of expression of anxiety and depression in clinical health care workers during the COVID-19 outbreak in 2 hospitals in Hanoi, Vietnam. Health Serv Insights. 2021;14:11786329211033245. <https://doi.org/10.1177/11786329211033245>
- 21 Spitzer RL, Williams JB, Kroenke KJ. Test review: Patient health Questionnaire–9 (PHQ-9). Rehabil Couns Bull. 2014;57:246–8. <https://doi.org/10.1177/0034355213515305>
- 22 Spitzer RL, Kroenke K, Williams JB, Löwe B. A brief measure for assessing generalized anxiety disorder: The GAD-7. Arch Intern Med. 2006;166:1092–7. <https://doi.org/10.1001/archinte.166.10.1092>
- 23 Breslau J, Borges G, Tancredi D, Saito N, Kravitz R, Hinton L, et al. Migration from Mexico to the United States and subsequent risk for depressive and anxiety disorders: A cross-national study. Arch Gen Psychiatry. 2011;68:428–33.

<https://doi.org/10.1001/archgenpsychiatry.2011.21>

24 Zhang J, Fang L, Wu YW, Wiecek WF. Depression, anxiety, and suicidal ideation among Chinese Americans: A study of immigration-related factors. *J Nerv Ment Dis.* 2013;201:17–22. <https://doi.org/10.1097/NMD.0b013e31827ab2e2>

25 Porter C, Favara M, Hittmeyer A, Scott D, Sánchez Jiménez A, Ellanki R, et al. Impact of the COVID-19 pandemic on anxiety and depression symptoms of young people in the global south: Evidence from a four-country cohort study. *BMJ, (Open).* 2021;11:e049653. <https://doi.org/10.1136/bmjopen-2021-049653>

26 Luong TC, Pham TTM, Nguyen MH, Do AQ, Pham LV, Nguyen HC, et al. Fear, anxiety and depression among pregnant women during COVID-19 pandemic: Impacts of healthy eating behaviour and health literacy. *Ann Med.* 2021;53:2120–31. <https://doi.org/10.1080/07853890.2021.2001044>

27 Yamamoto T, Uchiumi C, Suzuki N, Yoshimoto J, Murillo-Rodriguez E. The psychological impact of “mild lockdown” in Japan during the COVID-19 pandemic: A nationwide survey under a declared state of emergency. *Int J Environ Res Public Health.* 2020;17:9382. <https://doi.org/10.3390/ijerph17249382>

28 Nomura K, Minamizono S, Maeda E, Kim R, Iwata T, Hirayama J, et al. Cross-sectional survey of depressive symptoms and suicide-related ideation at a Japanese national university during the COVID-19 stay-home order. *Environ Health Prev Med.* 2021;26:30.

<https://doi.org/10.1186/s12199-021-00953-1>

- 29 Nomura K, Yamazaki T, Maeda E, Hirayama J, Ono K, Fushimi M, et al. Longitudinal survey of depressive symptoms among university students during the COVID-19 pandemic in Japan. *Front Psychol.* 2022;13:863300. <https://doi.org/10.3389/fpsyg.2022.863300>
- 30 Miki T, Yamamoto S, Fukunaga A, Inoue Y, Ishiwari H, Ishii M, et al. Association between eating balanced meals and depressive symptoms in Japanese hospital workers during the COVID-19 pandemic. *Neuropsychopharmacol Rep.* 2022;42:109–13. <https://doi.org/10.1002/npr2.12230>
- 31 Ueda M, Stickley A, Sueki H, Matsubayashi T. Mental health status of the general population in Japan during the COVID-19 pandemic. *Psychiatr Clin Neurosci.* 2020;74:505–6. <https://doi.org/10.1111/pcn.13105>
- 32 World Health Organization. Mental health and forced displacement; 2021. <https://www.who.int/news-room/fact-sheets/detail/mental-health-and-forced-displacement>  
Accessed 28 Aug 2022
- 33 Ministry of Health, Labour and Welfare (Japan). Guidance on temporary loan emergency funds; 2021. <https://corona-support.mhlw.go.jp/seikatsufukushi/en/index.html> Accessed 24 Feb 2022
- 34 Ministry of Health, Labour and Welfare (Japan). Monetary assistance for everyday lives; 2021. <https://www.mhlw.go.jp/content/000743778.pdf> Accessed 24 Feb 2022

- 35 Yasukawa K, Sawada T, Hashimoto H, Jimba M. Health-care disparities for foreign residents in Japan. *Lancet*. 2019;393:873–4. [https://doi.org/10.1016/S0140-6736\(19\)30215-6](https://doi.org/10.1016/S0140-6736(19)30215-6)
- 36 Higuchi M, Endo M, Yoshino A. Factors associated with access to health care among foreign residents living in Aichi Prefecture, Japan: Secondary data analysis. *Int J Equity Health*. 2021;20:135. <https://doi.org/10.1186/s12939-021-01465-8>
- 37 Makiyama K, Kawashima T, Nomura S, Eguchi A, Yoneoka D, Tanoue Y, et al. Trends in healthcare access in Japan during the first wave of the COVID-19 pandemic, up to June 2020. *Int J Environ Res Public Health*. 2021;18:3271. <https://doi.org/10.3390/ijerph18063271>
- 38 Fukase Y, Ichikura K, Murase H, Tagaya H. Age-related differences in depressive symptoms and coping strategies during the COVID-19 pandemic in Japan: A longitudinal study. *J Psychosom Res*. 2022;155:110737. <https://doi.org/10.1016/j.jpsychores.2022.110737>
- 39 Immigration Services Agency of Japan. Basic survey on foreign residents in Japan Report in 2020. <https://www.moj.go.jp/isa/content/001341984.pdf> Accessed 24 Feb 2022; 2021
- 40 Dykxhoorn J, Hollander A-C, Lewis G, Dalman C, Kirkbride JB. Family networks during migration and risk of non-affective psychosis: A population-based cohort study. *Schizophr Res*. 2019;208:268–75. <https://doi.org/10.1016/j.schres.2019.01.044>

## Tables

**Table 1:** Participant characteristics (n = 621)

Variable	Item	n	%
Sex	Male	347	55.9
	Female	274	44.1
Age	(Mean ± SD)	26.0 ± 4.8	
Duration of residence in Japan	(Mean ± SD)	3.4 ± 3.1	
COVID-19 infection rate in the area of residence	Low	181	29.1
	Middle	251	40.4
	High	189	30.4
Marital status	Married	129	20.8
	Single, divorced, or bereaved	492	79.2
Country of birth	Vietnam	615	99.0
	Japan	4	0.6
	Other	2	0.3
Visa status	Residence based on the nature of activities	26	4.2
	Residence based on employment except Technical Intern Training and Skilled Labour	167	26.9
	International student	182	29.3
	Technical Intern Training (Ginojishshu)	183	29.5
	Skilled Labour (Tokuteiginou)	38	6.1
	No response	25	4.0
Japanese language proficiency	Ability to speak fluently and confidently	71	11.4
	Ability to speak at a level that does not affect work or study	273	44.0
	Adequate ability to speak without impacting daily life	217	34.9
	Barely able to speak Japanese	60	9.7
Education level	Junior high school and under	5	0.8
	High school	238	38.3
	Technical school	132	21.3

	College (two years) or University	214	34.5
	Graduate school	32	5.2
Pre-existing health conditions	Yes	132	21.3
	No	489	78.7
Pre-existing health conditions (Multiple answers)	Stomach / oesophagus / duodenum	34	5.5
	Otorhinolaryngology	32	5.2
	Musculoskeletal / joint	32	5.2
	Dental	31	5.0
	Psychiatric disorders	25	4.0
	Blood	12	1.9
	Hyperlipidaemia	10	1.6
	High blood pressure	9	1.4
	Enteropathy	8	1.3
	Gynaecology	8	1.3
	Hepatobiliary tract	7	1.1
	Male reproductive organs	7	1.1
	Kidney disease	5	0.8
	Urinary system	5	0.8
	Heart disease	4	0.6
	Lung disease	4	0.6
Diabetes mellitus	3	0.5	
Neuropathy	3	0.5	
Cancer	2	0.3	

**Table 5:** Factors associated with symptoms of depression and anxiety (n = 596)

		Depressive symptoms (PHQ-9)			Anxiety symptoms (GAD-7)		
		Odds [95% CI]	Beta	P	Odds [95% CI]	Beta	P
Age		0.94 [0.89, 0.99]	-0.06	0.043 *	0.97 [0.92, 1.03]	-0.03	0.348
Duration of residence in Japan		1.08 [0.99, 1.17]	0.08	0.084	1.03 [0.95, 1.12]	0.03	0.441
Sex	Female	1.00 (reference)			1.00 (reference)		
	Male	0.92 [0.63, 1.34]	-0.09	0.656	1.09 [0.76, 1.57]	0.09	0.640
Marital status	Married	1.00 (reference)			1.00 (reference)		
	Single, divorced, or bereaved	1.61 [0.90, 2.88]	0.48	0.109	1.72 [1.01, 2.93]	0.55	0.044 *
Visa status	Residence based on status and position	1.00 (reference)			1.00 (reference)		
	Residence based on employment	2.46 [0.56, 10.82]	0.90	0.233	2.51 [0.67, 9.37]	0.92	0.172
Japanese language proficiency	Ability to speak well/to some extent	1.00 (reference)			1.00 (reference)		
	Barely able to speak	1.01 [0.50, 2.00]	0.01	0.990	0.78 [0.40, 1.53]	-0.24	0.474
Education level	College, University, or higher	1.00 (reference)			1.00 (reference)		
	Technical school	1.16 [0.71, 1.91]	0.15	0.549	1.05 [0.66, 1.69]	0.05	0.831
	High school	0.94 [0.59, -]	-	0.784	1.20 [0.77, -]	0.18	0.423

	or lower	1.49]	0.07		1.86]		
Pre-existing health conditions	No	1.00 (reference)			1.00 (reference)		
	Yes	2.46 [1.61, 3.76]	0.90 **	<0.001	2.52 [1.63, 3.88]	0.92 **	<0.001
Subjective socioeconomic status	General or better	1.00 (reference)			1.00 (reference)		
	Difficult or very difficult	2.47 [1.64, 3.71]	0.90 **	<0.001	2.72 [1.87, 3.97]	1.00 **	<0.001
COVID-19 infection rate in the area of residence	Low	1.00 (reference)			1.00 (reference)		0.654
	Middle	0.97 [0.62, 1.52]	- 0.03	0.895	0.83 [0.54, 1.27]	- 0.19	0.391
	High	1.11 [0.70, 1.77]	0.10	0.663	0.96 [0.61, 1.51]	- 0.04	0.859
Partner with whom to discuss one's health	Yes	1.00 (reference)			1.00 (reference)		
	No	1.23 [0.81, 1.88]	0.21	0.339	1.66 [1.11, 2.47]	0.51	0.013 *

CI, confidence interval

\* p-value < 0.05

\*\* p-value < 0.01