1 Title

- 2 Modifiable lifestyle activities affect cognition in cognitively healthy middle-aged individuals
- 3 at risk for late-life Alzheimer's Disease
- 4 **Running title:** Midlife lifestyle, AD risk and cognition

5 Authors

- 6 Amy Heneghan ^{ab1}, Feng Deng ^{ab1}, Katie Wells ^c, Karen Ritchie ^{cd}, Graciela Muniz-Terrera
- ⁷, Craig W **Ritchie**^c, Brian **Lawlor**^{ab}, Lorina **Naci**^{ab*}
- ^a Trinity College Institute of Neuroscience, School of Psychology, Trinity College Dublin,
- 9 Dublin, Ireland.
- ^b Global Brain Health Institute, Trinity College Dublin, Dublin, Ireland
- ^c Edinburgh Dementia Prevention, University of Edinburgh, Edinburgh, UK
- ^d INSERM and University of Montpellier, Montpellier, France.
- ¹ These authors contributed equally to this work

^{*}Corresponding author:

- 15 Lorina Naci
- 16 School of Psychology
- 17 Trinity College Institute of Neuroscience
- 18 Global Brain Health Institute
- 19 Trinity College Dublin
- 20 Dublin, Ireland

21 Telephone: +353 (0)87 688 5642

22 Email: <u>nacil@tcd.ie</u>

23 Abstract

24	It is now acknowledged that Alzheimer's Disease (AD) processes are present decades before
25	the onset of clinical symptoms, but it remains unknown whether lifestyle factors can protect
26	against these early AD processes in mid-life. We asked whether modifiable lifestyle activities
27	impact cognition in middle-aged individuals who are cognitively healthy, but at risk for late
28	life AD. Participants (40-59 years) completed cognitive and clinical assessments at baseline
29	(N = 206) and two years follow-up (N = 174). Mid-life activities were measured with the
30	Lifetime of Experiences Questionnaire. We assessed the impact of lifestyle activities, known
31	risk factors for sporadic late-onset AD (Apolipoprotein E $_{\Box}4$ allele status, family history of
32	dementia, and the Cardiovascular Risk Factors Aging and Dementia score), and their
33	interactions on cognition. More frequent engagement in physically, socially and intellectually
34	stimulating activities was associated with better cognition (verbal, spatial and relational
35	memory), at baseline and follow-up. Critically, more frequent engagement in these activities
36	was associated with stronger cognition (verbal and visuospatial functions, and conjunctive
37	short-term memory binding) in individuals with family history of dementia. Impaired
38	visuospatial function is one of the earliest cognitive deficits in AD and has previously
39	associated with increased AD risk in this cohort. Additionally, conjunctive memory functions
40	have been found impaired in the pre-symptomatic stages of AD. These findings suggest that
41	modifiable lifestyle activities offset cognitive decrements due to AD risk in mid-life and
42	support the targeting of modifiable lifestyle activities for the prevention of Alzheimer's
43	Disease.

44

45 Keywords: Cognitively Healthy; Mid-life; Cognition; Lifestyle Factors; Risk Factors;
46 Alzheimer's Disease

47 Introduction

48	Dementia is a growing pandemic that presents profound challenges to health care systems,
49	families, and societies throughout the world. Alzheimer's Disease (AD), the most common
50	etiology of dementia, is characterized by relentless neurodegeneration and accelerated
51	cognitive decline in the years following presentation of clinical symptoms. It is now accepted
52	that AD pathological processes are present decades before clinically relevant symptoms
53	appear [1-2]. In the absence of effective pharmacological treatments, there is an immediate
54	need for early identification, intervention and risk burden modification [3-4] for AD.
55	Growing evidence suggests that up to 40% of all dementia cases are associated with known
56	lifestyle-related modifiable risk factors, such as alcohol consumption, obesity and
57	hypertension among others [4]. As exposure to most of these risk factors begins decades
58	before dementia onset, interventions must be implemented in mid-life [5-7]. Mid-life, thus,
59	presents a critical and unique window for disease-altering interventions, before the
60	manifestation of substantial brain damage. However, the indicators of AD in mid-life, and the
61	impact of modifiable lifestyle factors on the incipient disease process remain poorly
62	understood.
63	AD is neuropathologically characterized by the accumulation of beta-amyloid (A β) and
64	hyperphosphorylated TAU (pTau) [8-9]. The etiology of the sporadic form of AD remains
65	poorly understood. However, recent studies suggest that tau deposition, is a key etiological
66	factor that presages sporadic AD [10-12]. Analyses of thousands human brains across the
67	lifespan show that tau pathology begins about a decade before formation of A β plaques [13].
68	Additionally, tau pathology, but not $A\beta$, correlates with progressive grey matter loss [14] and
69	cognitive impairment [15].Neuropathological findings show that tau pathology starts in late

young adulthood and early midlife [30–40years], in subcortical nucleus locus coeruleus (LC)

71	[9] —the key brain site for the production of noradrenaline— then spreads initially to the
72	transentorhinal and entorhinal cortex [8], to the hippocampal and neocortical association
73	cortex, and finally throughout neocortex. By contrast, A β plaques are usually first deposited
74	in the association neocortex of the temporal lobe, and then extend throughout the cortex,
75	including subcortical structures, with disease progression [13]. Therefore, brain changes
76	related to sporadic AD in early midlife (i.e. 40-50 years) are likely associated with
77	neurodegeneration from to pTau deposition and, in later age groups, A β plaques additionally
78	contribute strongly to underlying neurodegeneration.
79	Studies on preclinical AD have used risk stratification approaches to investigate early,
80	preclinical changes. Key risk factors include the Apolipoprotein E (APOE) 14 genotype [16-
81	17], the main genetic risk factor for sporadic late onset AD in the Indo-European population
82	[18], and family history (FH) of dementia [16, 19-20]. Several dementia risk scores
83	incorporating lifestyle risk factors have been devised [21-23]. Amongst them, the
84	Cardiovascular Risk Factors Aging and Dementia (CAIDE) score has been optimized for
85	middle-aged populations [23] and has been validated in a large US population followed
86	longitudinally over 40 years [24].
87	The PREVENT Dementia Program [25] initiated in 2013, is a prospective study of the
88	cognitively healthy middle-aged children of persons with dementia, designed to seek out
89	clinical and biological changes, which may subsequently be used as short-term outcome
90	measures for midlife secondary preventions. Studies of this and other similar cohorts of
91	cognitively healthy midlife individuals have related these three risk factors to a range of
92	structural and functional brain changes, including APOE $\Box 4$ genotype to loss of volume in the
93	hippocampal molecular layer [26], cerebral hyperperfusion [27-28], reduced grey matter
94	volume in the right hippocampus, precentral gyrus, and cerebellar cortex [29], and to

95	decreased cortical thickness in the frontal cortex [30]; FH to volumetric alterations in
96	hippocampal subfields and to disrupted white matter integrity [27, 31]; and, CAIDE to whole
97	brain atrophy [32-34] and to hippocampal volume loss [31, 34]. All three risk groups have
98	also been found to impact cognition in mid-life. APOE $_{\Box}4$ genotype has been significantly
99	associated with improvements in verbal, spatial and relational memory [35], immediate recall
100	[36] and form perception [34], FH with poorer verbal processing and memory performance in
101	participants under 65 years [37], and poorer executive function [38], and higher CAIDE
102	scores with impaired verbal and visuospatial functions, and short-term (conjunctive) memory
103	[34-35], and impaired executive function [39]. Taken together, these studies have established
104	that risk for late-life AD, including risk that incorporates lifestyle factors, has a significant
105	impact on the brain health and cognition of middle-aged individuals who are presently
106	cognitively healthy.
107	As a multidimensional construct, lifestyle has a multipronged impact on cognition and the

108 brain. By contrast to aforementioned cardiovascular risk factors captured by the CAIDE 109 score, several lifestyle activities have been found to protect brain health and cognition in later 110 life. Factors, such as education and occupational complexity have been associated with 111 preservation of cognitive function in order adults [40] and reduced symptom severity in 112 Alzheimer's disease [4, 41-43], a phenomenon known as "cognitive resilience". These factors 113 are thought to explain why, in late-life Alzheimer's disease, the level of cognitive impairment 114 shows substantial variability even when accounting for key pathologies including A β and 115 pathological tau [1, 44]. One prominent account of the biological mechanisms mediating the 116 relationship between stimulating lifestyle activities and cognitive resilience in AD [45-46] 117 suggests that environmental enrichment upregulates the noradrenergic system which 118 originated in the locus coeruleus, that otherwise depletes with age [47] and AD pathology 119 [10], leading to compensatory brain mechanisms for cognitive function.

120	While epidemiological evidence strongly suggests that education and occupation contribute
121	to cognitive resilience [48], there is a renewed interest in the additional contribution of other
122	activities undertaken in mid-life, given their potential modifiability. For example, Gow et al.
123	(2017) [49] and Chan et al. (2018) [40] used the Lifetime of Experiences Questionnaire
124	(LEQ) and found that physically, socially and intellectually stimulating lifestyle activities
125	undertaken in mid-life, independently of education, help to maintain late-life cognitive
126	performance in older adults, after adjusting for childhood cognitive ability [49]. However, it
127	is not known whether the protection conferred by lifestyle activities, other than education,
128	offsets the impact of AD risk in mid-life, or whether it builds up gradually over time with its
129	benefits viable only in late life or relative to established AD pathology. The answer to this
130	question is critical for identifying interventions that target modifiable factors for the
131	prevention of Alzheimer's disease from the earliest life stages.
131 132	prevention of Alzheimer's disease from the earliest life stages. To address this gap, we investigated the impact of interactions between three risk factors for
131 132 133	prevention of Alzheimer's disease from the earliest life stages. To address this gap, we investigated the impact of interactions between three risk factors for late-life AD (APOE _4 genotype, FH and CAIDE) with lifestyle activities on cognition,
131 132 133 134	prevention of Alzheimer's disease from the earliest life stages. To address this gap, we investigated the impact of interactions between three risk factors for late-life AD (APOE 4 genotype, FH and CAIDE) with lifestyle activities on cognition, independently of sex, age and years of education. These relationships were investigated in a
131 132 133 134 135	prevention of Alzheimer's disease from the earliest life stages. To address this gap, we investigated the impact of interactions between three risk factors for late-life AD (APOE 4 genotype, FH and CAIDE) with lifestyle activities on cognition, independently of sex, age and years of education. These relationships were investigated in a large cohort of cognitively healthy middle-aged individuals, assessed at baseline (N=206),
131 132 133 134 135 136	prevention of Alzheimer's disease from the earliest life stages. To address this gap, we investigated the impact of interactions between three risk factors for late-life AD (APOE 4 genotype, FH and CAIDE) with lifestyle activities on cognition, independently of sex, age and years of education. These relationships were investigated in a large cohort of cognitively healthy middle-aged individuals, assessed at baseline (N=206), and at two years follow-up (N=174). To relate our findings to previous studies, we used the
131 132 133 134 135 136 137	prevention of Alzheimer's disease from the earliest life stages. To address this gap, we investigated the impact of interactions between three risk factors for late-life AD (APOE dependently, FH and CAIDE) with lifestyle activities on cognition, independently of sex, age and years of education. These relationships were investigated in a large cohort of cognitively healthy middle-aged individuals, assessed at baseline (N=206), and at two years follow-up (N=174). To relate our findings to previous studies, we used the LEQ [50], the same instrument as in aforementioned studies, to evaluate lifestyle activities
131 132 133 134 135 136 137 138	prevention of Alzheimer's disease from the earliest life stages. To address this gap, we investigated the impact of interactions between three risk factors for late-life AD (APOE 4 genotype, FH and CAIDE) with lifestyle activities on cognition, independently of sex, age and years of education. These relationships were investigated in a large cohort of cognitively healthy middle-aged individuals, assessed at baseline (N=206), and at two years follow-up (N=174). To relate our findings to previous studies, we used the LEQ [50], the same instrument as in aforementioned studies, to evaluate lifestyle activities specific to mid-life, yielding two composite factors, (a) occupation and managerial
131 132 133 134 135 136 137 138 139	prevention of Alzheimer's disease from the earliest life stages. To address this gap, we investigated the impact of interactions between three risk factors for late-life AD (APOE []4 genotype, FH and CAIDE) with lifestyle activities on cognition, independently of sex, age and years of education. These relationships were investigated in a large cohort of cognitively healthy middle-aged individuals, assessed at baseline (N=206), and at two years follow-up (N=174). To relate our findings to previous studies, we used the LEQ [50], the same instrument as in aforementioned studies, to evaluate lifestyle activities specific to mid-life, yielding two composite factors, (a) occupation and managerial responsibility, and (b) physical, social and intellectual activities. Our hypothesis was that
131 132 133 134 135 136 137 138 139 140	prevention of Alzheimer's disease from the earliest life stages. To address this gap, we investigated the impact of interactions between three risk factors for late-life AD (APOE 4 genotype, FH and CAIDE) with lifestyle activities on cognition, independently of sex, age and years of education. These relationships were investigated in a large cohort of cognitively healthy middle-aged individuals, assessed at baseline (N=206), and at two years follow-up (N=174). To relate our findings to previous studies, we used the LEQ [50], the same instrument as in aforementioned studies, to evaluate lifestyle activities specific to mid-life, yielding two composite factors, (a) occupation and managerial responsibility, and (b) physical, social and intellectual activities. Our hypothesis was that there would be a significant association between mid-life lifestyle factors and cognitive.

142 Methods

143 Participants

144	PREVENT is an ongoing longitudinal multi-site research programme based across the UK
145	and Ireland, seeking to identify early biomarkers of AD and elaborate on risk-mechanism
146	interactions for neurodegenerative diseases decades before the cardinal symptoms of
147	dementia emerge. Its protocol has been described in detail elsewhere [25]. In the first
148	PREVENT programme phase, participants were recruited at a single site, via the dementia
149	register database held at the West London National Health Service (NHS) Trust, of the UK
150	National Health Service, the Join Dementia Research website
151	(https://www.joindementiaresearch.nihr.ac.uk/), through public presentations, social media
152	and word of mouth. Procedures involving experiments on human subjects were done in
153	accord with the ethical standards of the Institutional Review Board of Imperial College
154	London and in accord with the Helsinki Declaration of 1975. Approval for the study was
155	granted by the NHS Research Ethics Committee London Camberwell St Giles. Consented
156	participants were seen at the West London NHS Trust, where they underwent a range of
157	clinical and cognitive assessments [25]. The cohort comprised cognitively healthy volunteers
158	aged 40-59 years. The West London dataset was used here to avail of both baseline and
159	follow-up testing data acquired for this cohort. 210 individuals (62 male; 148 female) were
160	tested at baseline, with 188 (89.5%) (55 male; 133 female) retained at 2 years follow-up
161	(Table 1).

163 Assessments

164 Risk factors

165 $APOE_{\Box}4$ Genotyping

- 166 The process of APOE $_{\Box}4$ allele identification is outlined in detail in Ritchie et al. (2017) [34].
- 167 In brief, genomic DNA was isolated from blood samples and APOE genotyping was
- 168 performed. All members of the research and clinical teams were blind to the result of APOE
- 169 genotyping. In this study, APOE $\Box 4$ risk is determined by ≥ 1 APOE $\Box 4$ allele. 75/210 carried
- 170 ≥ 1 APOE $\Box 4$ allele (See Table 1).

171 *Family History*

- 172 FH was determined by a 'yes'/'no' question during clinical visits, which asked participants
- 173 whether a parent had a diagnosis of dementia. Participants were asked to include the
- dementia subtype if known, but answering 'yes' alone categorized a participant as FH+. The
- answer 'no' likely captured both participants with no family history of dementia, and
- 176 participants for whom FH was unknown. In summary, participants were defined as FH+ if at
- 177 least one parent was diagnosed with dementia. Cases where the FH was unknown or partially
- known were not recorded outside of the binary yes/no scoring. 103/210 were FH+.

179 Cardiovascular Risk Factors, Aging, and Incidence of Dementia (CAIDE) score

- 180 CAIDE is a composite scale of estimated future dementia risk based on mid-life
- 181 cardiovascular measures [51-52]. It takes into consideration the individual's age, sex,
- educational attainment, APOE $_{\Box}4$ genotype, activity level, BMI, cholesterol and systolic
- blood pressure [23] and is scored on a range of 0 18. A higher score indicates greater risk.
- 184 The CAIDE dementia risk score was calculated for each participant at baseline and follow-
- 185 up.

186

187 Cognitive testing

- 188 The Addenbrooke's Cognitive Assessment (ACE) III [53] was recorded at the follow-up
- session, but not at the baseline. Subsequent analyses that focused on the subset of
- assessments performed at both testing session did not include the ACE) III.

191 Cognitive function was assessed at baseline and follow up with the COGNITO

neuropsychological battery [54], designed to examine information processing across a widerange of cognitive functions in adults of all ages and not restricted to those functions usually

194 implicated in dementia detection in the elderly. Tests are administered using a tactile screen

to capture information processing time as well as response accuracy and require about 40

196 minutes to complete. The tests, by order of presentation, are: reaction time; reading;

197 comprehension of phonemes, phrases, and syntax; focused and divided attention in both

198 visual and auditory modalities; visual working memory (visual tracking with auditory

interference); the Stroop test; immediate, delayed, and recognition trials for verbal recall

200 (name list); delayed recognition of spatial stimuli (faces); visuospatial associative learning;

visuospatial span; form perception; denomination of common objects; spatial reasoning;

202 copying of meaningful and meaningless figures; verbal fluency with se-mantic and phonetic

203 prompts; immediate recall of a narrative; immediate recall of a description of the relative

204 position of objects; vocabulary; implicit memory (recognition of new and previously learned205 material).

206 The COGNITO tests are designed to test several aspects of cognition, including attention

207 (task: visual attention), memory (tasks: narrative recall, description recall, implicit memory,

208 name-face association, working memory), language (tasks: phoneme comprehension, verbal

fluency) and visuospatial abilities (task: geometric figure recognition) [54-55]. Based on

210	previous studies [54-55], 11 summary variables from the COGNITO battery capturing the
211	above functions were used here [for a list see Supporting Information (SI)].
212	Additionally, we used the Visual Short-Term Memory Binding task (VSTMBT) [56], a
213	computer-based task that assesses visual short-term memory binding of single features, e.g.,
214	complex shape or color combinations, or feature conjunctions, e.g., shape and color
215	combinations. In the single feature condition, participants must identify whether the test
216	stimuli (three random 6-sided polygons) are the "same" as or "different" to the studied
217	stimuli in terms of shape (shape only) or color (color only). In the binding condition,
218	participants are required to correctly identify if both the shape and color of the test stimuli
219	match studied stimuli. Two summary variables from the VSTMBT were the percentage of
220	correctly recognized items from the two conditions.

221

222 Measurement of lifestyle activities

223 The Lifetime of Experiences Questionnaire (LEQ) [50], designed to take a lifespan approach 224 to the measurement of cognitive resilience [57-59] and mental activity, measures engagement 225 in a broad range of lifestyle activities across three stages of life: young adulthood (13-29 226 years), mid-life (30-64 years) and late life (65 years onwards). Therefore, the LEQ is 227 preferable for looking at a mid-life cohort compared to other scales that capture dementia-228 specific risk related to modifiable lifestyle factors (e.g., LIBRA [60]). The LEQ comprises 229 sub-scores capturing "specific" activities, reflecting the primary activity undertaken in each 230 life stage and "non-specific" activities, reflecting engagement in physical, social and 231 intellectual activities in any stage. For the purpose of this paper, we define mid-life 'lifestyle' 232 as all the activities captured by the LEQ (below).

233

234 Mid-life specific score

235	The mid-life specific component score centers on occupation and comprises two sub-scores.
236	For the first occupational sub-score, participants were asked to record their primary
237	occupation in each 5-year interval from age 30 to age at assessment. Each reported
238	occupation was scored according to the International Standard Classification of Occupations
239	(ISCO 08) guidelines (https://www.ilo.org/public/english/bureau/stat/isco/isco08/), and
240	scores were inverted and summed. The second sub-score is a measure of the managerial
241	responsibility associated with reported occupations. If participants indicated that they were
242	employed in a managerial capacity, the number of people that they oversaw in four of their
243	reported occupations was documented. Managerial responsibility was scored as follows; 0
244	people = 8, 1-5 people = 16, 5-10 people = 24 and $10+$ people = 32. The highest score is
245	recorded as the managerial responsibility sub-score. The occupational history and managerial
246	sub-scores were summed and multiplied by a normalization factor of 0.25. Normalization
247	ensures that the mid-life specific and non-specific scores have comparable mean values [50].

248

249 Mid-life non-specific score

250 The non-specific score assesses frequency of engagement in 7 activities, capturing those of a 251 physically, socially and intellectually stimulating nature, scored on a 5-point Likert scale of 252 frequency (never, less than monthly, monthly, fortnightly, weekly, daily). Scores range from 0-35, with higher scores reflecting more frequent engagement in such activities. The items 253 254 included in the scale are socializing with family or friends, practicing a musical instrument, 255 practicing an artistic pastime, engagement in physical activity that is mildly, moderately, or 256 vigorously energetic, reading, practicing a second language and travel. The travel item asks 257 participants if they have visited any of a list of continents between the ages of 30-54.

258 Responses were scored on a 5-point scale as follows; none, 1-2 regions, 3-4 regions, 5

259 regions, 6-7 regions, all regions.

261 Statistical analyses

262	To reduce the number of multiple comparisons between the cognitive tests, given the
263	multitude of cognitive assessments obtained for this cohort (13 summary variables, see SI and
264	Table S1), in an independent study of this dataset [35], three composite cognitive components
265	were extracted from the above cognitive assessments, by using rotated principal component
266	analysis (rPCA) (Figure S1). In subsequent analyses, we used these cognitive components,
267	rather than the individual tests to measure the impact of risk and protective lifestyle factors
268	on cognition.
269	At baseline, the three components were: 1) C1: verbal, spatial and relational memory; 2) C2:
270	working and short-term (single-feature) memory; 3) C3: verbal and visuospatial functions,
271	and short-term (conjunctive) memory (Figure S1 A). Higher scores in C1 and C2 reflected
272	better performance, whereas higher scores in C3 reflected poorer performance. For ease of
273	visualization and comparability to C1 and C2, values of C3 were recoded with reversed sign.
274	The highest loading tasks and their weights for each component differed slightly for the
275	cross-sectional and longitudinal data. There was high similarity between baseline and follow-
276	up for C1 ($\phi = 0.90$), and C3 ($\phi = 0.76$), with relatively low similarly for C2 ($\phi = 0.40$)
277	(Figure S1 B). Although C1 and C3 had relatively high similarity across time, the
278	longitudinal changes in these components could not be quantified by direct comparison,
279	because they captured different cognitive functions at each time-point and any longitudinal
280	change would be hard to interpret.

281	We used the Statistical Package for Social Sciences (SPSS V.26) and R software for all
282	statistical analyses. The normality of the data was assessed by combining the visualization of
283	a quantile-quantile plot and the Shapiro-Wilk test. Demographic and clinical information of
284	the study cohort was analyzed across risk groups using chi-square (χ^2 tests) for categorical
285	variables and Mann Whitney U tests for continuous (discrete) variables, given that they were
286	not normally distributed in this cohort. Subsequently, we used hierarchical regression models
287	to look at the contribution of mid-life lifestyle factors (LEQ specific and non-specific scores)
288	risk factors (APOE $_{\Box}4$ genotype and FH) and of their interactions on cognitive performance,
289	at baseline and follow-up. The effect of APOE $_{\square}4$ and CAIDE risk factors was modelled
290	independently, in order to avoid modelling the variance associated with APOE
291	genotype in the same model twice. In each case, dependent variables were the
292	aforementioned composite cognitive components, each assessed in a separate model.
293	The variance of some parts of physical activity is accounted for twice in the model, in the
294	CAIDE score and in the non-specific LEQ factor, resulting in weakening of its statistical
295	contribution that can by captured by the lifestyle factor. Therefore, any effect of physical
296	activity that can be captured by the lifestyle factor (independent variable) in the same model
297	as CAIDE (independent variable), can only provide a conservative estimate of the
298	contribution of physical activity to the dependent variable of cognitive performance.
299	

As aforementioned, the baseline and follow-up datasets were analyzed independently, due to the slightly differing cognitive domains for each visit. Although the two datasets were not independent, we considered each of them in its own right, because during 2 years, a proportion of our participants may have had substantial brain health changes that are yet subthreshold to clinical manifestations. Therefore, the follow-up dataset has the potential to

reveal the impact of age on variables of interest. Corrections for multiple comparisons wereperformed across all analyses.

307	To avoid multicollinearity, we mean centered continuous variables (specific and non-specific
308	scores). Age, sex and years of education were included as covariates in hierarchical models.
309	A significant interaction effect between a risk and a lifestyle factor would indicate that the
310	effect of the lifestyle factor on cognitive performance differs across levels/values of the risk
311	factor. For any observed interactions, we plotted the regression of the lifestyle factor on
312	cognitive performance for each level/value of the risk factor [61], to interpret the effect. We
313	then tested the significance of the slopes of the simple regression lines, to investigate in
314	which level/value of the risk factor we found an effect of lifestyle on cognitive performance.

315 **Results**

316 Demographic characteristics

317	In the first PREVENT study phase, 210 participants were recruited from a single site. All 210
318	completed several cognitive tests and the LEQ questionnaire. 188 were also assessed at the
319	follow up, two years later. At baseline, 2 participants were missing information relating to
320	their APOE $_{\Box}4$ genotype, with a further 2 missing cognitive data. The final baseline cohort
321	that could be used for the analyses was $N = 206$. At follow-up, 12 participants were missing
322	cognitive data, alongside the 2 participants who were missing APOE $_{\Box}4$ genotype data.
323	Therefore, the final follow-up cohort was $N = 174$. Mild cognitive impairment and dementia
324	were ruled out based on detailed clinical assessment at baseline and follow-up. Please see the
325	Supplementary Information and Table S1 for a list of all the outcome variables collected at
326	baseline and follow-up.
327	Demographic specifications of the cohort at baseline and follow-up, stratified by APOE $_{\Box}4$

genotype and family history of dementia, are shown in Table 1. There were no significant differences in age, sex or years of education between the groups. As expected, APOE $\Box 4$ allele genotype was more frequently found in the FH+ than FH- group at baseline (p = 0.01) and follow up (p = 0.02). CAIDE scores (including APOE status) were significantly higher for the FH+ than FH- group at baseline (p = 0.03) and follow up (p = 0.003). Naturally, CAIDE scores including APOE status were significantly higher for the APOE $\Box 4$ + than

- APOE \square 4- group at baseline (p = 0.0003) and follow up (p = 0.0002), but when APOE status
- 335 was excluded the CAIDE scores did not differ between the two groups (Table 1).

336

337

Table 1 about here please

339 Verbal, Spatial and Relational memory

358

340	At baseline, a hierarchical regression model with lifestyle factors, i.e., the specific and non-
341	specific LEQ scores, risk factors (APOE $_{\square}4$ and FH), and age, sex and years of education as
342	covariates, and verbal, spatial and relational memory as the dependent variable was
343	significant overall [F (7, 198) = 5.84, $R^2 = 0.17$, p < 0.0001] (Supplementary Table 2a), and
344	showed a significant positive association between verbal, spatial and relational memory and
345	education [β (SE) = 0.09 (0.02), p < 0.0001] (Figure 1a). Similarly, at follow up the model's
346	performance was significant [F (7, 166) = 5.52, $R^2 = 0.19$, p < 0.0001], and there was a
347	significant positive association between verbal, spatial and relational memory and education
348	$[\beta$ (SE) = 0.10 (0.02), p < 0.0001] (Figure 1b). Higher education values were significantly
349	associated with higher performance at both timepoints. The inclusion of risk by lifestyle
350	interaction terms (FH \times specific score, FH \times non-specific score, APOE $_{\Box}4$ \times specific score,
351	APOE $_{\Box}4 \times \text{non-specific score}$) into the hierarchical regression model (Supplementary Table
352	2b) did not show any significant associations between lifestyle x risk interaction terms and
353	cognitive performance, at baseline or follow-up.
354	
355	Figure 1 about here please
555	rigare r ubbai nere picase
356	
357	The hierarchical regression models with lifestyle factors and CAIDE as independent

baseline [F (3, 202) = 3.11, $R^2 = 0.04$, p = 0.03] and at follow-up [F (3, 166) = 3.52, $R^2 =$

0.04, p = 0.02] (Supplementary Table 2c). At both timepoints, we found a significant positive

variables, and verbal, spatial and relational memory as the dependent was significant at

association between the non-specific LEQ factor and verbal, spatial and relational memory

362 (baseline: [β (SE) = 0.04 (0.02), p = 0.02], figure 2a) and (follow up: [β (SE) = 0.07 (0.02), p

363	= 0.002], figure 2b). More frequent engagement in physically, socially and intellectually
364	stimulating activities was associated with better verbal, spatial and relational memory. This
365	association was independent of CAIDE and therefore of age, sex and years of education, as
366	included in this score. The inclusion of CAIDE by lifestyle interaction terms (CAIDE \times
367	specific score, CAIDE \times non-specific score) into the hierarchical regression model
368	(Supplementary Table 2d) did not show any significant associations between lifestyle x risk
369	interaction terms and cognitive performance, at baseline or follow-up.
370	
271	Eisense 2. shout have showed
3/1	Figure 2 about nere please
372	
373	Working and Short-Term (Single-Feature) Memory
374	Hierarchical regression analyses showed no significant associations of any of the independent
375	variables and performance on the working and short-term (single-feature) memory, at
376	baseline or follow up (Supplementary Table 3a, 3c) Furthermore, no significant associations
377	with interaction terms were observed at either timepoint (Supplementary Table 3b, 3d).
378	
279	Verbal and Visuospatial Functions, and Short-Term (Conjunctive) Memory
	(1 + 1) = (1 +
380	At baseline, the hierarchical regression model was significant [F (7, 198) = 2.39, $R = 0.08$, p
381	= 0.02], and showed a significant negative association between verbal and visuospatial
382	function and short-term (conjunctive) memory with age [β (SE) = -0.04 (0.01), p = 0.006]
383	(Supplementary Table 4a). Higher age was associated with poorer performance. At follow-
384	up, there were no significant associations between the main independent variables with verbal
385	and visuospatial function and short-term (conjunctive) memory. The inclusion of risk by
386	lifestyle interaction terms in the hierarchical model revealed a significant association between

387	the non-specific LEQ score x FH interaction term and cognitive performance [β (SE) = 0.11
388	(0.05), p = 0.01] (Supplementary Table 4b). To interpret this interaction, we investigated the
389	relationship between cognitive performance and the non-specific LEQ score for the FH+ and
390	FH- groups independently (Figure 3). We found a significant negative relationship between
391	non-specific lifestyle activities and cognition in the FH+ group [β (SE)= 0.08 (0.03), p =
392	0.02), that was independent of age, sex, or years of education. We did not observe a
393	relationship between non-specific lifestyle activities and cognition in the FH- group [β (SE) =
394	-0.03 (0.03), $p = 0.34$]. These results suggested that for individuals with positive family
395	history, more frequent engagement in the non-specific LEQ factor — namely physically,
396	socially and intellectually engaging activities — was associated with better performance in
397	verbal and visuospatial function and short-term (conjunctive) memory. Finally, we also
398	observed trend associations between cognitive performance and the specific LEQ factor score
399	x FH interaction term [β (SE) = -0.07 (0.04), p = 0.06]. This association, while suggestive of
400	a role for occupational complexity on cognition, stratified by family history risk group, is
401	weak and needs to be investigated further in future studies. Additionally, as previously
402	reported [35] we observed a significant negative association between CAIDE and verbal and
403	visuospatial functions and short-term (conjunctive) memory at baseline (Supplementary
404	Table 4c). There were no associations between interactions of CAIDE x lifestyle and
405	cognition at either baseline or follow up (Supplementary Table 4d).
406	

407

Figure 3 about here please

409 Conclusion

410	It is now acknowledged that Alzheimer's Disease processes are present decades before the					
411	onset of clinical symptoms [2, 62, but, to date, it has remained unknown whether lifestyle					
412	factors can already protect against these early AD processes in mid-life. We asked whether					
413	these modifiable lifestyle activities impact cognition in middle-aged individuals who are					
414	cognitively healthy but at risk for late life AD. Lifestyle activities significantly impacted					
415	cognition in mid-life. Individuals with greater educational attainment showed stronger					
416	cognition in a composite dimension capturing verbal, spatial and relational memory. This					
417	result is consistent with a previous study [34] of this cohort showing that visuospatial abilities					
418	were positively associated with education. It is also consistent with epidemiological studies					
419	on older adults showing that education contributes to cognitive resilience in older life [48].					
420	Our cohort is highly educated, with approximately 30% reaching a post-graduate					
421	qualification [34]. As the LEQ education measure captures total years of education, the effect					
422	of education we observed, reflects long-term effects set in motion from early life and young					
423	adulthood.					
424	The key question in this study, however, was to investigate any additional contribution of					
425	activities undertaken in mid-life, independently of education. The first novel finding of this					
426	study was that more frequent engagement in physically, socially and intellectually stimulating					
427	activities in mid-life was associated with stronger cognition in a composite dimension					
428	capturing verbal, spatial and relational memory, both at baseline and at follow-up. This effect					
429	was independent of sex, age, years of education and cardiovascular factors captured by the					
430	CAIDE score. The second novel finding was that physically, socially and intellectually					
431	stimulating activities undertaken in mid-life had a significant effect in the cognition of					

432 middle-aged individuals at risk for late-life AD. Specifically, higher engagement in these

433 activities was associated with significantly stronger cognition in another composite 434 dimension, capturing verbal and visuospatial functions and short-term (conjunctive) memory 435 in cognitively healthy individuals, who were at risk for late life AD though family history of 436 dementia, at follow up. Importantly, this effect was independent of age, sex, and years of 437 education. The presence of this effect in the follow-up but not the baseline dataset is likely 438 due to the older age of the cohort at follow-up. Short-term (conjunctive) memory functions 439 have been found impaired in the pre-symptomatic stages of AD [56]. Additionally, impaired 440 visuospatial function is one of the earliest cognitive deficits observed in AD [63-64] and has 441 previously been linked to increased AD risk in this cohort [34-35]. In similarly cognitively 442 healthy, but older (> 65 years) North American cohorts [65-66], changes in cerebrospinal 443 fluid biomarkers related to inflammation have been associated with changes in visuospatial 444 cognitive performance, thereby suggesting a biochemically-mediated effect of early 445 pathology in presymptomatic individuals. 446 Taken together, our results suggest that stimulating lifestyle activities may boost cognitive 447 functions that are very vulnerable to AD risk and early AD neuropathology [56, 63-64]. 448

449 individual function is limited by their composite assessment in this study and requires

However, we caution that the interpretability of the effect of lifestyle activities on each

450 individuation in future studies with a longer longitudinal follow-up window. Physically,

451 socially and intellectually stimulating mid-life activities have been shown to influence late-

452 life cognition, suggesting that they contribute to the cognitive resilience that mitigates the

453 effect of age-related cognitive decline and AD neuropathology [40, 49]. Our finding advances

454 understanding by showing that engagement with these activities contributes to cognitive

455 resilience to risk of AD, or even incipient AD neuropathology, from mid-life, in individuals

456 who are presently cognitively healthy.

457	Family history of dementia is a well-established risk factor [16, 19-20], independent of the
458	genetic risk bestowed by APOE $_{\Box}4$ genotype, that captures both genetic and environmental
459	risk influences. For example, an individual who has one or more parent with dementia, may,
460	independently to the presence of the APOE $\Box 4$ genotype, be exposed to negative
461	environmental influences that contribute to neuropathology, including increased stress [67],
462	caregiver burden [68-69] and reduced participation in enriching environments due to
463	caregiving duties [70]. Therefore, alongside contributing to cognitive resilience, enhanced
464	engagement with physically, socially and intellectually stimulating activities may positively
465	impact individuals with a family history of dementia by counteracting these negative
466	environmental influences.
467	What might had the machanism by which stimulating activities impact acquition in midlife?
467	what might be the mechanism by which stimulating activities impact cognition in midnle?
468	Models of AD disease delineated by neuropathological staging [9] place the locus coeruleus,
469	a small nucleus in the pontine tegmentum region of the brainstem, at the pathogenesis of AD
470	[10-11, 71]. The LC is responsible for the production of neurotransmitter noradrenaline (NA)
471	[72], a major driver of the brain's arousal system, which strongly modulates high-order
472	cognition. Novelty provides a key trigger for arousal, and, thus, the LC-NA activity occurs
473	strongly in response to novel stimuli. Therefore, stimulating activities upregulate the LC and
474	may protect it from early neurodegeneration due to pTau deposition. Furthermore, as the
475	brain site of noradrenaline, the LC is key to the protective effect of lifestyle factors on the
476	whole brain. Studies suggests that environmental enrichment, through activities such as those
477	measured in this study, upregulates the noradrenergic system, which otherwise depletes with
478	age [47] and AD pathology [10], leading to compensatory brain mechanisms for cognitive
479	function, such as the strengthening of the fronto-parietal brain and other large-scale brain
480	networks [45-46].

481	In summary, our findings suggest that modifiable lifestyle activities may offset AD risk-
482	related cognitive decrements in mid-life and support the targeting of certain modifiable
483	lifestyle activities for the prevention of Alzheimer's Disease, especially in those with a family
484	history. These activities include socializing with family or friends, practicing a musical
485	instrument, practicing an artistic pastime, engagement in energetic physical activities,
486	reading, practicing a second language and travel. Given the apparent lifestyle contributions
487	both as risk and protective factors of dementia [4], a healthy lifestyle may be the individual's
488	current best defense against sporadic late onset AD. The modifiability of the lifestyle
489	activities identified here renders them promising cost-effective candidates for intervention
490	and prevention strategies from early life. These activities may be particularly significant for
491	non-pharmacological interventions for AD in low and middle-income countries (LMIC),
492	where barriers to education are more prevalent than in high income countries [4].

493

494 Methodological considerations

495 As education is strongly positively linked to IQ [73] and our cohort was highly educated, the 496 question of reverse causation arises [40, 49, 74]. This points to the possibility that cognitive 497 abilities may determine engagement in stimulating activities, rather than the inverse. 498 However, the effect on mid-life verbal and visuospatial functions, and short-term 499 (conjunctive) memory is independent of the total years of education, which shows that 500 education does not directly drive this effect. Furthermore, we found that mid-life lifestyle 501 activities were associated with improved cognitive performance only in the family history 502 positive group, which is not, at least not prima facie, more educated than the family history 503 negative group. Thus, this effect is likely independent of any indirect effects of education.

504	The use of composite cognitive domains that capture slightly different functions in the two
505	assessment points [35] prevents investigation of the impact of lifestyle on cognitive changes
506	over time in this study. Nevertheless, previous studies from this cohort have shown only
507	subtle changes over the two-year period [75-76], possibly due to the relatively young age
508	range of the sample and the short follow up window [34, 55]. Therefore, future studies that
509	follow this cohort over a longer period and test hypotheses informed by the previous study
510	waves are needed to determine the longitudinal impact of lifestyle activities in cognitively
511	healthy middle-aged individuals at risk for late life AD.
512	It is worth noting that we found no evidence that lifestyle factors can mitigate the impact of
513	the CAIDE risk or for interactions between CAIDE lifestyle and cognition as reported in
515	the CALDE Hisk, of for interactions between CALDE, mestyle and cognition, as reported in
514	some previous studies (see Kivipelto et al. 2018 [77] for a review). Unlike the CAIDE score
515	(including systolic blood pressure, BMI, total cholesterol, physical activity), the lifestyle
516	factors captured by our instrument (the LEQ) are not primarily related to cardiovascular
517	wellbeing (with the exception of exercise, 1/9 items), and therefore not likely to impact
518	greatly on cardiovascular health in this relatively young cohort (i.e., 40–59 years). By
519	contrast, the studies reviewed by Kivipelto et al. (2018) include older populations (> 60–70
520	years), where cardiovascular factors are more prevalent drivers of health vulnerabilities, and
521	furthermore those studies, unlike the present one, had a strong focus on cardiovascular health,
522	including nutritional guidance, exercise and monitoring and management of metabolic and
523	vascular risk factors.

524

525

527 Acknowledgement

- 528 F.D. was funded by the Provost PhD Award Scheme from Trinity College Dublin, to L.N.
- 529 L.N. was also funded by a L'Oréal-UNESCO for Women In Science International Rising
- 530 Talent Award, the Welcome Trust Institutional Strategic Support grant, and the Global Brain
- 531 Health Institute Project Grant.
- 532 The PREVENT-Dementia study is supported by the UK Alzheimer's Society (grant numbers
- 533 178 and 264), the US Alzheimer's Association (grant number TriBEKa-17-519007) and
- 534 philanthropic donations.
- 535 We thank all PREVENT-Dementia participants for their enthusiastic participation in this
- study. We also thank the Research Delivery service at West London NHS Trust and the
- 537 Wolfson Clinical Imaging Facility at Imperial College London for their support in running

538 the study.

- 540 Conflict of interest
- 541 The authors declare no conflict of interest.

542 References

- 543 [1] Jack, C. R., Jr, Knopman, D. S., Jagust, W. J., Petersen, R. C., Weiner, M. W., Aisen, P.
- 544 S., Shaw, L. M., Vemuri, P., Wiste, H. J., Weigand, S. D., Lesnick, T. G., Pankratz, V. S.,
- 545 Donohue, M. C., & Trojanowski, J. Q. (2013). Tracking pathophysiological processes in
- 546 Alzheimer's disease: an updated hypothetical model of dynamic biomarkers. *The Lancet*.
- 547 Neurology, 12(2), 207–216. <u>https://doi.org/10.1016/S1474-4422(12)70291-0</u>
- 548 [2] Ritchie, K., Ritchie, C. W., Yaffe, K., Skoog, I., & Scarmeas, N. (2015). Is late-onset
- 549 Alzheimer's disease really a disease of midlife?. *Alzheimer's & dementia (New York, N.*
- 550 *Y.*), *1*(2), 122–130. <u>https://doi.org/10.1016/j.trci.2015.06.004</u>
- 551 [3] Ritchie, C.W., Waymont, J.M.J., Pennington, C. et al. The Scottish Brain Health Service
- 552 Model: Rationale and Scientific Basis for a National Care Pathway of Brain Health Services
- 553 in Scotland. J Prev Alzheimers Dis (2021). https://doi.org/10.14283/jpad.2021.63
- [4] Livingston, G., Huntley, J., Sommerlad, A., Ames, D., Ballard, C., Banerjee, S., Brayne,
- 555 C., Burns, A., Cohen-Mansfield, J., Cooper, C., Costafreda, S. G., Dias, A., Fox, N., Gitlin,
- L. N., Howard, R., Kales, H. C., Kivimäki, M., Larson, E. B., Ogunniyi, A., Orgeta, V., ...
- 557 Mukadam, N. (2020). Dementia prevention, intervention, and care: 2020 report of the Lancet
- 558 . Lancet (London, England), 396(10248), 413–446. <u>https://doi.org/10.1016/S0140-</u>
- 559 <u>6736(20)30367-6</u>
- 560 [5] Irwin, K., Sexton, C., Daniel, T., Lawlor, B., & Naci, L. (2018). Healthy Aging and
- 561 Dementia: Two Roads Diverging in Midlife?. *Frontiers in aging neuroscience*, 10, 275.
- 562 https://doi.org/10.3389/fnagi.2018.00275
- [6] Lachman, M. E., Teshale, S., & Agrigoroaei, S. (2015). Midlife as a Pivotal Period in the
- Life Course: Balancing Growth and Decline at the Crossroads of Youth and Old

- 565 Age. International journal of behavioral development, 39(1), 20–31.
- 566 https://doi.org/10.1177/0165025414533223
- 567 [7] Gottesman, R. F., Schneider, A. L., Zhou, Y., Coresh, J., Green, E., Gupta, N., Knopman,
- 568 D. S., Mintz, A., Rahmim, A., Sharrett, A. R., Wagenknecht, L. E., Wong, D. F., & Mosley,
- 569 T. H. (2017). Association Between Midlife Vascular Risk Factors and Estimated Brain
- 570 Amyloid Deposition. JAMA, 317(14), 1443–1450. <u>https://doi.org/10.1001/jama.2017.3090</u>
- 571 [8] Braak, H., Thal, D. R., Ghebremedhin, E., & Del Tredici, K. (2011). Stages of the
- pathologic process in Alzheimer disease: age categories from 1 to 100 years. *Journal of*
- 573 *neuropathology and experimental neurology*, *70*(11), 960–969.
- 574 https://doi.org/10.1097/NEN.0b013e318232a379
- 575 [9] Braak, H., & Braak, E. (1991). Neuropathological stageing of Alzheimer-related changes.
- 576 Acta neuropathologica, 82(4), 239–259. https://doi.org/10.1007/BF00308809
- 577 [10] Jacobs, H., Becker, J. A., Kwong, K., Engels-Domínguez, N., Prokopiou, P. C., Papp, K.
- 578 V., Properzi, M., Hampton, O. L., d'Oleire Uquillas, F., Sanchez, J. S., Rentz, D. M., El
- 579 Fakhri, G., Normandin, M. D., Price, J. C., Bennett, D. A., Sperling, R. A., & Johnson, K. A.
- 580 (2021). In vivo and neuropathology data support locus coeruleus integrity as indicator of
- 581 Alzheimer's disease pathology and cognitive decline. *Science translational medicine*,
- 582 *13*(612), eabj2511. https://doi.org/10.1126/scitranslmed.abj2511
- 583 [11] Arnsten, A., Datta, D., Del Tredici, K., & Braak, H. (2021). Hypothesis: Tau pathology
- is an initiating factor in sporadic Alzheimer's disease. Alzheimer's & dementia : the journal of
- 585 the Alzheimer's Association, 17(1), 115–124. https://doi.org/10.1002/alz.12192

- 586 [12] Kametani, F., & Hasegawa, M. (2018). Reconsideration of Amyloid Hypothesis and Tau
- 587 Hypothesis in Alzheimer's Disease. *Frontiers in neuroscience*, *12*, 25.
- 588 <u>https://doi.org/10.3389/fnins.2018.00025</u>
- 589 [13] Braak, H., & Del Tredici, K. (2015). Neuroanatomy and pathology of sporadic
- 590 Alzheimer's disease. *Advances in anatomy, embryology, and cell biology, 215*, 1–162.
- 591 [14] La Joie, R., Ayakta, N., Seeley, W. W., Borys, E., Boxer, A. L., DeCarli, C., Doré, V.,
- 592 Grinberg, L. T., Huang, E., Hwang, J. H., Ikonomovic, M. D., Jack, C., Jr, Jagust, W. J., Jin,
- 593 L. W., Klunk, W. E., Kofler, J., Lesman-Segev, O. H., Lockhart, S. N., Lowe, V. J., Masters,
- 594 C. L., ... Rabinovici, G. D. (2019). Multisite study of the relationships between antemortem
- 595 [11C]PIB-PET Centiloid values and postmortem measures of Alzheimer's disease
- neuropathology. Alzheimer's & dementia : the journal of the Alzheimer's Association, 15(2),
- 597 205–216. <u>https://doi.org/10.1016/j.jalz.2018.09.001</u>
- 598 [15] Giannakopoulos, P., Herrmann, F. R., Bussière, T., Bouras, C., Kövari, E., Perl, D. P.,
- 599 Morrison, J. H., Gold, G., & Hof, P. R. (2003). Tangle and neuron numbers, but not amyloid
- 600 load, predict cognitive status in Alzheimer's disease. *Neurology*, 60(9), 1495–1500.
- 601 https://doi.org/10.1212/01.wnl.0000063311.58879.01
- 602 [16] Donix, M., Small, G. W., & Bookheimer, S. Y. (2012). Family history and APOE-4
- 603 genetic risk in Alzheimer's disease. *Neuropsychology review*, 22(3), 298–309.
- 604 <u>https://doi.org/10.1007/s11065-012-9193-2</u>
- 605 [17] Liu, C. C., Liu, C. C., Kanekiyo, T., Xu, H., & Bu, G. (2013). Apolipoprotein E and
- Alzheimer disease: risk, mechanisms and therapy. *Nature reviews*. *Neurology*, 9(2), 106–118.
- 607 <u>https://doi.org/10.1038/nrneurol.2012.263</u>

- [18] Lambert, J. C., Ibrahim-Verbaas, C. A., Harold, D., Naj, A. C., Sims, R., Bellenguez, C.,
- 609 DeStafano, A. L., Bis, J. C., Beecham, G. W., Grenier-Boley, B., Russo, G., Thorton-Wells,
- 610 T. A., Jones, N., Smith, A. V., Chouraki, V., Thomas, C., Ikram, M. A., Zelenika, D.,
- 611 Vardarajan, B. N., Kamatani, Y., ... Amouyel, P. (2013). Meta-analysis of 74,046 individuals
- 612 identifies 11 new susceptibility loci for Alzheimer's disease. *Nature genetics*, 45(12), 1452–
- 613 1458. <u>https://doi.org/10.1038/ng.2802</u>
- [19] Berti, V., Mosconi, L., Glodzik, L., Li, Y., Murray, J., De Santi, S., Pupi, A., Tsui, W.,
- & De Leon, M. J. (2011). Structural brain changes in normal individuals with a maternal
- 616 history of Alzheimer's. *Neurobiology of aging*, *32*(12), 2325.e17–2325.e2.325E26.
- 617 <u>https://doi.org/10.1016/j.neurobiolaging.2011.01.001</u>
- [20] Scarabino, D., Gambina, G., Broggio, E., Pelliccia, F., & Corbo, R. M. (2016). Influence
- of family history of dementia in the development and progression of late-onset Alzheimer's
- 620 disease. American Journal of Medical Genetics Part B: Neuropsychiatric Genetics, 171(2),
- 621 250-256. <u>https://doi.org/10.1002/ajmg.b.32399</u>
- 622 [21] Barnes, D. E., Beiser, A. S., Lee, A., Langa, K. M., Koyama, A., Preis, S. R., Neuhaus,
- 623 J., McCammon, R. J., Yaffe, K., Seshadri, S., Haan, M. N., & Weir, D. R. (2014).
- 624 Development and validation of a brief dementia screening indicator for primary
- 625 care. *Alzheimer's & dementia : the journal of the Alzheimer's Association*, *10*(6), 656–665.e1.
- 626 <u>https://doi.org/10.1016/j.jalz.2013.11.006</u>
- 627 [22] Deckers, K., van Boxtel, M. P., Schiepers, O. J., de Vugt, M., Muñoz Sánchez, J. L.,
- Anstey, K. J., Brayne, C., Dartigues, J. F., Engedal, K., Kivipelto, M., Ritchie, K., Starr, J.
- 629 M., Yaffe, K., Irving, K., Verhey, F. R., & Köhler, S. (2015). Target risk factors for dementia
- 630 prevention: a systematic review and Delphi consensus study on the evidence from

- 631 observational studies. *International journal of geriatric psychiatry*, *30*(3), 234–246.
- 632 <u>https://doi.org/10.1002/gps.4245</u>
- [23] Kivipelto, M., Ngandu, T., Laatikainen, T., Winblad, B., Soininen, H., & Tuomilehto, J.
- 634 (2006). Risk score for the prediction of dementia risk in 20 years among middle aged people:
- a longitudinal, population-based study. *The Lancet. Neurology*, 5(9), 735–741.
- 636 <u>https://doi.org/10.1016/S1474-4422(06)70537-3</u>
- [24] Exalto, L. G., Quesenberry, C. P., Barnes, D., Kivipelto, M., Biessels, G. J., & Whitmer,
- 638 R. A. (2014). Midlife risk score for the prediction of dementia four decades later. *Alzheimer's*
- 639 & dementia : the journal of the Alzheimer's Association, 10(5), 562–570.
- 640 https://doi.org/10.1016/j.jalz.2013.05.1772
- [25] Ritchie, C. W., & Ritchie, K. (2012). The PREVENT study: a prospective cohort study
- to identify mid-life biomarkers of late-onset Alzheimer's disease. *BMJ open*, 2(6), e001893.
- 643 <u>https://doi.org/10.1136/bmjopen-2012-001893</u>
- [26] Dounavi, M. E., Mak, E., Wells, K., Ritchie, K., Ritchie, C. W., Su, L., & O'Brien, J. T.
- 645 (2020). Volumetric alterations in the hippocampal subfields of subjects at increased risk of
- 646 dementia. *Neurobiology of aging*, 91, 36-44.
- 647 https://doi.org/10.1016/j.neurobiolaging.2020.03.006
- 648 [27] Mak, E., Dounavi, M. E., Low, A., Carter, S. F., McKiernan, E., Williams, G. B., ... & T
- 649 O'Brien, J. (2021). Proximity to dementia onset and multi-modal neuroimaging changes: The
- 650 prevent-dementia study. *NeuroImage*, 229, 117749.
- 651 https://doi.org/10.1016/j.neuroimage.2021.117749
- [28] McKiernan, E. F., Mak, E., Dounavi, M. E., Wells, K., Ritchie, C., Williams, G., ... &
- 653 O'brien, J. (2020). Regional hyperperfusion in cognitively normal APOE ε4 allele carriers in

- 654 mid-life: analysis of ASL pilot data from the PREVENT-Dementia cohort. Journal of
- *Neurology, Neurosurgery & Psychiatry*, *91*(8), 861-866. <u>https://doi.org/10.1136/jnnp-2020-</u>
 322924
- [29] Cacciaglia, R., Molinuevo, J. L., Falcón, C., Brugulat-Serrat, A., Sánchez-Benavides, G.,
- 658 Gramunt, N., Esteller, M., Morán, S., Minguillón, C., Fauria, K., Gispert, J. D., & ALFA
- 659 study (2018). Effects of APOE-ε4 allele load on brain morphology in a cohort of middle-aged
- 660 healthy individuals with enriched genetic risk for Alzheimer's disease. Alzheimer's &
- 661 *dementia : the journal of the Alzheimer's Association*, *14*(7), 902–912.
- 662 <u>https://doi.org/10.1016/j.jalz.2018.01.016</u>
- [30] Fennema-Notestine, C., Panizzon, M. S., Thompson, W. R., Chen, C. H., Eyler, L. T.,
- 664 Fischl, B., Franz, C. E., Grant, M. D., Jak, A. J., Jernigan, T. L., Lyons, M. J., Neale, M. C.,
- 665 Seidman, L. J., Tsuang, M. T., Xian, H., Dale, A. M., & Kremen, W. S. (2011). Presence of
- 666 ApoE £4 allele associated with thinner frontal cortex in middle age. Journal of Alzheimer's
- 667 *disease : JAD, 26 Suppl 3*(Suppl 3), 49–60. <u>https://doi.org/10.3233/JAD-2011-0002</u>
- [31] McKeever, A., Paris, A. F., Cullen, J., Hayes, L., Ritchie, C. W., Ritchie, K., ... & Su, L.
- 669 (2020). Hippocampal Subfield volumes in middle-aged adults at risk of dementia. Journal of
- 670 *Alzheimer's Disease*, 75(4), 1211-1218. <u>https://doi.org/10.3233/JAD-200238</u>
- 671 [32] Liu, X., Dounavi, M. E., Ritchie, K., Wells, K., Ritchie, C. W., Su, L., Muniz-Terrera,
- 672 G., & O'Brien, J. T. (2021). Higher midlife CAIDE score is associated with increased brain
- atrophy in a cohort of cognitively healthy middle-aged individuals. *Journal of*
- 674 *neurology*, 268(5), 1962–1971. <u>https://doi.org/10.1007/s00415-020-10383-8</u>
- [33] O'Brien, J. T., Firbank, M. J., Ritchie, K., Wells, K., Williams, G. B., Ritchie, C. W., &
- 676 Su, L. (2020). Association between midlife dementia risk factors and longitudinal brain

- atrophy: the PREVENT-Dementia study. *Journal of neurology, neurosurgery, and*
- 678 *psychiatry*, 91(2), 158–161. <u>https://doi.org/10.1136/jnnp-2019-321652</u>
- [34] Ritchie, K., Carrière, I., Su, L., O'Brien, J. T., Lovestone, S., Wells, K., & Ritchie, C. W.
- 680 (2017). The midlife cognitive profiles of adults at high risk of late-onset Alzheimer's disease:
- 681 The PREVENT study. Alzheimer's & dementia : the journal of the Alzheimer's
- 682 Association, 13(10), 1089–1097. <u>https://doi.org/10.1016/j.jalz.2017.02.008</u>
- [35] Deng, F., Wells, K., Ritchie, K., Muniz-Terrera, G., Ritchie, C. W., O'Brien, J.T.,
- Lawlor, B., Naci, L. (2022) Functional connectivity of the locus coeruleus with the
- hippocampus and the cognitive profiles of asymptomatic mid-life individuals at risk for
- 686 Alzheimer's disease. *bioRxiv*.
- 687 [36] Jochemsen, H. M., Muller, M., van der Graaf, Y., & Geerlings, M. I. (2012). APOE ε4
- differentially influences change in memory performance depending on age. The SMART-MR
- 689 study. *Neurobiology of Aging*, *33*(4), 832.e15–832.e8.32E22.
- 690 <u>https://doi.org/10.1016/j.neurobiolaging.2011.07.016</u>
- [37] Talboom, J. S., Håberg, A., De Both, M. D., Naymik, M. A., Schrauwen, I., Lewis, C.
- 692 R., Bertinelli, S. F., Hammersland, C., Fritz, M. A., Myers, A. J., Hay, M., Barnes, C. A.,
- 693 Glisky, E., Ryan, L., & Huentelman, M. J. (2019). Family history of Alzheimer's disease
- alters cognition and is modified by medical and genetic factors. *eLife*, 8, e46179.
- 695 <u>https://doi.org/10.7554/eLife.46179</u>
- [38] Debette, S., Wolf, P. A., Beiser, A., Au, R., Himali, J. J., Pikula, A., Auerbach, S.,
- 697 Decarli, C., & Seshadri, S. (2009). Association of parental dementia with cognitive and brain
- 698 MRI measures in middle-aged adults. *Neurology*, *73*(24), 2071–2078.
- 699 https://doi.org/10.1212/WNL.0b013e3181c67833

- [39] Stephen, R., Liu, Y., Ngandu, T., Rinne, J. O., Kemppainen, N., Parkkola, R.,
- Laatikainen, T., Paajanen, T., Hänninen, T., Strandberg, T., Antikainen, R., Tuomilehto, J.,
- 702 Keinänen Kiukaanniemi, S., Vanninen, R., Helisalmi, S., Levälahti, E., Kivipelto, M.,
- 703 Soininen, H., & Solomon, A. (2017). Associations of CAIDE Dementia Risk Score with
- 704 MRI, PIB-PET measures, and cognition. Journal of Alzheimer's disease : JAD, 59(2), 695–
- 705 705. https://doi.org/10.3233/JAD-170092
- [40] Chan, D., Shafto, M., Kievit, R., Matthews, F., Spink, M., Valenzuela, M., Cam-CAN,
- 707 & Henson, R. N. (2018). Lifestyle activities in mid-life contribute to cognitive reserve in late-
- ⁷⁰⁸ life, independent of education, occupation, and late-life activities. *Neurobiology of aging*, 70,
- 709 180–183. <u>https://doi.org/10.1016/j.neurobiolaging.2018.06.012</u>
- 710 [41] Livingston, G., Sommerlad, A., Orgeta, V., Costafreda, S. G., Huntley, J., Ames, D.,
- 711 Ballard, C., Banerjee, S., Burns, A., Cohen-Mansfield, J., Cooper, C., Fox, N., Gitlin, L. N.,
- Howard, R., Kales, H. C., Larson, E. B., Ritchie, K., Rockwood, K., Sampson, E. L., Samus,
- 713 Q., ... Mukadam, N. (2017). Dementia prevention, intervention, and care. Lancet (London,
- 714 England), 390(10113), 2673–2734. https://doi.org/10.1016/S0140-6736(17)31363-6
- [42] Wang, H. X., MacDonald, S. W., Dekhtyar, S., & Fratiglioni, L. (2017). Association of
- 716 lifelong exposure to cognitive reserve-enhancing factors with dementia risk: A community-
- based cohort study. *PLoS medicine*, *14*(3), e1002251.
- 718 <u>https://doi.org/10.1371/journal.pmed.1002251</u>
- [43] Dekhtyar, S., Marseglia, A., Xu, W., Darin-Mattsson, A., Wang, H. X., & Fratiglioni, L.
- 720 (2019). Genetic risk of dementia mitigated by cognitive reserve: A cohort study. Annals of
- 721 *neurology*, 86(1), 68–78. <u>https://doi.org/10.1002/ana.25501</u>

- [44] Franzmeier, N., Koutsouleris, N., Benzinger, T., Goate, A., Karch, C. M., Fagan, A. M.,
- 723 McDade, E., Duering, M., Dichgans, M., Levin, J., Gordon, B. A., Lim, Y. Y., Masters, C. L.,
- 724 Rossor, M., Fox, N. C., O'Connor, A., Chhatwal, J., Salloway, S., Danek, A., Hassenstab, J.,
- 725 ... Ewers, M. (2020). Predicting sporadic Alzheimer's disease progression via inherited
- 726 Alzheimer's disease-informed machine-learning. Alzheimer's & dementia : the journal of the
- 727 Alzheimer's Association, 16(3), 501–511. <u>https://doi.org/10.1002/alz.12032</u>
- [45] Robertson I. H. (2014). Right hemisphere role in cognitive reserve. *Neurobiology of*
- 729 aging, 35(6), 1375–1385. <u>https://doi.org/10.1016/j.neurobiolaging.2013.11.028</u>
- [46] Robertson I. H. (2013). A noradrenergic theory of cognitive reserve: implications for
- Alzheimer's disease. *Neurobiology of aging*, *34*(1), 298–308.
- 732 https://doi.org/10.1016/j.neurobiolaging.2012.05.019
- [47] Liu, K. Y., Kievit, R. A., Tsvetanov, K. A., Betts, M. J., Düzel, E., Rowe, J. B., Cam-
- 734 CAN, Howard, R., & Hämmerer, D. (2020). Noradrenergic-dependent functions are
- associated with age-related locus coeruleus signal intensity differences. Nature
- *communications*, *11*(1), 1712.
- 737 [48] Richards, M., & Deary, I. J. (2005). A life course approach to cognitive reserve: a model
- for cognitive aging and development?. *Annals of neurology*, 58(4), 617–622.
- 739 <u>https://doi.org/10.1002/ana.20637</u>
- [49] Gow, A. J., Pattie, A., & Deary, I. J. (2017). Lifecourse Activity Participation From
- 741 Early, Mid, and Later Adulthood as Determinants of Cognitive Aging: The Lothian Birth
- 742 Cohort 1921. The journals of gerontology. Series B, Psychological sciences and social
- 743 sciences, 72(1), 25–37. <u>https://doi.org/10.1093/geronb/gbw124</u>

- [50] Valenzuela, M. J., & Sachdev, P. (2007). Assessment of complex mental activity across
- the lifespan: development of the Lifetime of Experiences Questionnaire
- 746 (LEQ). Psychological medicine, 37(7), 1015–1025. <u>https://doi.org/10.1017/S003329170600938X</u>
- [51] Fayosse, A., Nguyen, D. P., Dugravot, A., Dumurgier, J., Tabak, A. G., Kivimäki, M.,
- 748 Sabia, S., & Singh-Manoux, A. (2020). Risk prediction models for dementia: role of age and
- 749 cardiometabolic risk factors. *BMC medicine*, 18(1), 107. <u>https://doi.org/10.1186/s12916-020-</u>
- 750 <u>01578-x</u>
- [52] Sindi, S., Calov, E., Fokkens, J., Ngandu, T., Soininen, H., Tuomilehto, J., & Kivipelto,
- 752 M. (2015). The CAIDE Dementia Risk Score App: The development of an evidence-based
- mobile application to predict the risk of dementia. Alzheimer's & dementia (Amsterdam,
- 754 *Netherlands*), *1*(3), 328–333. <u>https://doi.org/10.1016/j.dadm.2015.06.005</u>
- [53] Hsieh, S., Schubert, S., Hoon, C., Mioshi, E., & Hodges, J. R. (2013). Validation of the
- 756 Addenbrooke's Cognitive Examination III in frontotemporal dementia and Alzheimer's
- disease. *Dementia and geriatric cognitive disorders*, *36*(3-4), 242–250.
- 758 https://doi.org/10.1159/000351671
- [54] Ritchie, K., de Roquefeuil, G., Ritchie, C., Besset, A., Poulain, V., Artero, S., &
- Ancelin, M. L. (2014). COGNITO: computerized assessment of information
- 761 processing. Journal of Psychology and Psychotherapy, 4(2). <u>https://doi.org/10.4172/2161-</u>
- 762 <u>0487.1000136</u>
- [55] Ritchie, K., Carrière, I., Howett, D., Su, L., Hornberger, M., O'Brien, J. T., Ritchie, C.
- 764 W., & Chan, D. (2018). Allocentric and Egocentric Spatial Processing in Middle-Aged
- 765 Adults at High Risk of Late-Onset Alzheimer's Disease: The PREVENT Dementia

- 766 Study. Journal of Alzheimer's disease : JAD, 65(3), 885–896. https://doi.org/10.3233/JAD-
- 767 <u>180432</u>
- [56] Parra, M. A., Abrahams, S., Logie, R. H., Méndez, L. G., Lopera, F., & Della Sala, S.
- 769 (2010). Visual short-term memory binding deficits in familial Alzheimer's disease. Brain : a
- 770 *journal of neurology*, 133(9), 2702–2713. <u>https://doi.org/10.1093/brain/awq148</u>
- 771 [57] Stern Y. (2009). Cognitive reserve. *Neuropsychologia*, 47(10), 2015–2028.
- 772 https://doi.org/10.1016/j.neuropsychologia.2009.03.004
- [58] Stern Y. (2012). Cognitive reserve in ageing and Alzheimer's disease. *The Lancet*.
- 774 Neurology, 11(11), 1006–1012. <u>https://doi.org/10.1016/S1474-4422(12)70191-6</u>
- 775 [59] Stern, Y., Barnes, C. A., Grady, C., Jones, R. N., & Raz, N. (2019). Brain reserve,
- cognitive reserve, compensation, and maintenance: operationalization, validity, and
- mechanisms of cognitive resilience. *Neurobiology of aging*, 83, 124–129.
- 778 https://doi.org/10.1016/j.neurobiolaging.2019.03.022
- [60] Schiepers, O., Köhler, S., Deckers, K., Irving, K., O'Donnell, C. A., van den Akker, M.,
- Verhey, F., Vos, S., de Vugt, M. E., & van Boxtel, M. (2018). Lifestyle for Brain Health
- 781 (LIBRA): a new model for dementia prevention. *International journal of geriatric*
- 782 psychiatry, 33(1), 167–175. <u>https://doi.org/10.1002/gps.4700</u>
- [61] Aiken, L. S., & West, S. G. (1991). Multiple regression: Testing and interpreting
- 784 interactions. Newbury Park, CA: Sage.
- 785 [62] Jack, C. R., Jr, Knopman, D. S., Jagust, W. J., Shaw, L. M., Aisen, P. S., Weiner, M. W.,
- 786 Petersen, R. C., & Trojanowski, J. Q. (2010). Hypothetical model of dynamic biomarkers of

- the Alzheimer's pathological cascade. *The Lancet. Neurology*, 9(1), 119–128.
- 788 https://doi.org/10.1016/S1474-4422(09)70299-6
- [63] Williams, O. A., An, Y., Armstrong, N. M., Kitner-Triolo, M., Ferrucci, L., & Resnick,
- 790 S. M. (2020). Profiles of Cognitive Change in Preclinical and Prodromal Alzheimer's Disease
- 791 Using Change-Point Analysis. *Journal of Alzheimer's disease : JAD*, 75(4), 1169–1180.
- 792 <u>https://doi.org/10.3233/JAD-191268</u>
- 793 [64] Laukka, E. J., Macdonald, S. W., Fratiglioni, L., & Bäckman, L. (2012). Preclinical
- rognitive trajectories differ for Alzheimer's disease and vascular dementia. Journal of the
- 795 International Neuropsychological Society : JINS, 18(2), 191–199.
- 796 https://doi.org/10.1017/S1355617711001718
- 797 [65] Miron, J., Picard, C., Lafaille-Magnan, M. É., Savard, M., Labonté, A., Breitner, J.,
- 798 Rosa-Neto, P., Auld, D., Poirier, J., & PREVENT-AD research group (2019). Association of
- 799 TLR4 with Alzheimer's disease risk and presymptomatic biomarkers of inflammation.
- Alzheimer's & dementia : the journal of the Alzheimer's Association, 15(7), 951–960.
- 801 <u>https://doi.org/10.1016/j.jalz.2019.03.012</u>
- [66] Noble, J. M., Manly, J. J., Schupf, N., Tang, M. X., Mayeux, R., & Luchsinger, J. A.
- 803 (2010). Association of C-reactive protein with cognitive impairment. Archives of neurology,
- 804 67(1), 87–92. https://doi.org/10.1001/archneurol.2009.308
- 805 [67] Franks, K. H., Bransby, L., Saling, M. M., & Pase, M. P. (2021). Association of Stress
- 806 with Risk of Dementia and Mild Cognitive Impairment: A Systematic Review and Meta-
- Analysis. Journal of Alzheimer's disease : JAD, 82(4), 1573–1590.
- 808 <u>https://doi.org/10.3233/JAD-210094</u>

- [68] Chiao, C. Y., Wu, H. S., & Hsiao, C. Y. (2015). Caregiver burden for informal
- 810 caregivers of patients with dementia: A systematic review. International nursing
- 811 review, 62(3), 340–350. <u>https://doi.org/10.1111/inr.12194</u>
- 812 [69] Yu, H., Wang, X., He, R., Liang, R., & Zhou, L. (2015). Measuring the Caregiver
- 813 Burden of Caring for Community-Residing People with Alzheimer's Disease. *PloS*
- 814 one, 10(7), e0132168. <u>https://doi.org/10.1371/journal.pone.0132168</u>
- 815 [70] Ross, K. M., & Carroll, J. E. (2017). Alzheimer's Disease Caregivers' Health and
- 816 Important Resources: Next Steps in Intervention Research. *Psychosomatic medicine*, 79(7),
- 817 732–734. <u>https://doi.org/10.1097/PSY.000000000000518</u>
- 818 [71] Márquez, F., & Yassa, M. A. (2019). Neuroimaging Biomarkers for Alzheimer's
- 819 Disease. *Molecular neurodegeneration*, 14(1), 21. <u>https://doi.org/10.1186/s13024-019-0325-</u>
- 820 <u>5</u>
- [72] Amaral, D. G., & Sinnamon, H. M. (1977). The locus coeruleus: neurobiology of a
- 822 central noradrenergic nucleus. *Progress in neurobiology*, 9(3), 147–196.
- 823 https://doi.org/10.1016/0301-0082(77)90016-8
- [73] Ritchie, S. J., & Tucker-Drob, E. M. (2018). How Much Does Education Improve
- 825 Intelligence? A Meta-Analysis. *Psychological science*, 29(8), 1358–1369.
- 826 https://doi.org/10.1177/0956797618774253
- [74] Borgeest, G. S., Henson, R. N., Shafto, M., Samu, D., Cam-CAN, & Kievit, R. A.
- 828 (2020). Greater lifestyle engagement is associated with better age-adjusted cognitive
- abilities. *PloS one*, 15(5), e0230077. <u>https://doi.org/10.1371/journal.pone.0230077</u>

- 830 [75] Dounavi, M. E., Low, A., McKiernan, E. F., Mak, E., Muniz-Terrera, G., Ritchie, K.,
- 831 Ritchie, C. W., Su, L., & O'Brien, J. T. (2021). Evidence of cerebral hemodynamic
- 832 dysregulation in middle-aged APOE ε4 carriers: The PREVENT-Dementia study. *Journal of*
- 833 *cerebral blood flow and metabolism : official journal of the International Society of Cerebral*
- 834 *Blood Flow and Metabolism*, *41*(11), 2844–2855.
- 835 https://doi.org/10.1177/0271678X211020863
- [76] Low, A., Su, L., Stefaniak, J. D., Mak, E., Dounavi, M. E., Muniz-Terrera, G., Ritchie,
- 837 K., Ritchie, C. W., Markus, H. S., & O'Brien, J. T. (2021). Inherited risk of dementia and the
- 838 progression of cerebral small vessel disease and inflammatory markers in cognitively healthy
- midlife adults: the PREVENT-Dementia study. *Neurobiology of aging*, 98, 124–133.
- 840 <u>https://doi.org/10.1016/j.neurobiolaging.2020.10.029</u>
- [77] Kivipelto, M., Mangialasche, F., & Ngandu, T. (2018). Lifestyle interventions to prevent
- 842 cognitive impairment, dementia and Alzheimer disease. *Nature reviews. Neurology*, 14(11),
- 843 653–666. <u>https://doi.org/10.1038/s41582-018-0070-3</u>

844

Figures and Legends

	Baseline (n=210)			Follow-up (n=188)		
	FH- (n=107)	FH+ (n=103)	<i>p</i> (Mann-Whitney U)	FH- (n=89)	FH+ (n=99)	p (Mann-Whitney U)
Age (y)	52.0 ± 11.0	53.0 ± 7.0	0.54	54.0 ± 11.0	55.0 ± 6.0	0.22
Years of Education	16.0 ± 6.0	16.0 ± 5.0	0.34	17.0 ± 5.0	16.0 ± 5.0	0.21
CAIDE	5.0 ± 4.0	6.0 ± 2.0	0.03*	6.0 ± 4.0	7.0 ± 3.0	0.003**
			p (Chi-Square)			p (Chi-Square)
Sex (%f)	69.2%	71.8%	0.67	69.7%	71.7%	0.76
APOE e4 (% Carriers) §	28.0%	44.6%	0.01*	28.1%	44.3%	0.02*
	APOEe4- (n=133)	APOEe4+ (n=75)	p (Mann-Whitney U)	APOEe4- (n=118)	APOEe4+ (n=68)	<i>p</i> (Mann-Whitney U)
Age (y)	53.0 ± 8.0	52.0 ± 8.0	0.06	55.0 ± 8.0	54.5 ± 8.0	0.08
Years of Education	16.0 ± 5.0	17.0 ± 5.0	0.50	16.0 ± 5.0	17.0 ± 5.0	0.47
CAIDE (incl. APOE status)	5.0 ± 3.0	7.0 ± 3.0	0.0003***	6.0 ± 3.0	7.0 ± 4.0	0.0002***
CAIDE (excl. APOE status)	5.0 ± 2.0	4.0 ± 3.0	0.12	5.0 ± 2.0	4.5 ± 3.0	0.25
			p (Chi-Square)			p (Chi-Square)
Sex (%f)	69.9%	70.7%	0.91	71.2%	69.1%	0.77

Table 1. Demographic specifications of the cohort at baseline and follow-up based on the dementia family history and on APOE genotype

Note: median \pm interquartile range (IQR) was reported for continuous variables. Abbreviations: FH-, Negative family history of dementia; FH+, positive family history of dementia; APOE $_{\Box}4$ +, Apolipoprotein $_{\Box}4$ genotype positive: APOE $_{\Box}4$ -, Apolipoprotein $_{\Box}4$ genotype negative. [§] Information for the APOE $_{\Box}4$ genotype was not collected for 2 subjects.



Figure 1. Association of years of education with verbal, spatial and relational memory performance at baseline (a) and follow-up (b). The x axis displays total reported years of education attained. On the y axis, higher scores represent better verbal, spatial and relational memory performance.



Figure 2. Association of physical, social and intellectual activities with verbal, spatial and relational memory performance at baseline (a) and follow-up (b). Physical, social and intellectual activities are mean-centred. On the x axis, higher scores represent more frequent engagement in physical, social and intellectual lifestyle activities, and on the y axis, higher scores represent better verbal, spatial and relational memory performance.



Figure 3. Interaction of physical, social and intellectual activities and family history of dementia on verbal and visuospatial functions, and short-term (conjunctive) memory at follow-up. On the x axis, higher scores represent more frequent engagement in physical, social and intellectual lifestyle activities, and on the y axis, higher scores represent poorer verbal and visuospatial functions, and short-term (conjunctive) memory. Physical, social and intellectual activities are mean-centred. The positive family history of dementia group showed a significant positive association between engagement in these activities and improved verbal and visuospatial functions, and short-term (conjunctive) memory. No significant association was seen for the negative family history group. Abbreviations: FH, family history; FH+, family history positive; FH-, family history negative.