1	The Co-production of the Roots Framework: A Reflective
2	Framework for Mapping the Implementation Journey of Trauma-
3	informed Care
4	
5	Steven Anthony Thirkle ^a [*] , Angela Kennedy ^b , Petia Sice ^c and Paras Patel ^b
6	^a Population Health Sciences Institute, Faculty of Medical Sciences, Baddiley Clark
7	Building, Richardson Road, Newcastle University, Newcastle upon Tyne, NE2 4AX,
8	United Kingdom
9	^b St. Nicholas Hospital, Jubilee Rd, Cumbria, Northumberland Tyne and Wear NHS
10	Foundation Trust, Newcastle-upon Tyne NE3 3XT, United Kingdom
11	^c Department of Computer and Information Sciences, Faculty of Engineering and
12	Environment, Ellison Building, Ellison Place, Northumbria University, Newcastle City
13	Campus, Newcastle-upon-Tyne, NE2 1XE, United Kingdom
14	
15	*Corresponding author: Steven Anthony Thirkle, Population Health Sciences
16	Institute, Faculty of Medical Sciences, Newcastle University, Newcastle upon Tyne,
17	NE2 4AX, United Kingdom. ORCID ID - 0000-0002-6115-6637. E-mail:
18	steven.thirkle@newcastle.ac.uk Telephone: 07955593737
19	Abstract
20	Background: The trauma-informed care programme at the Tees, Esk and Wear
21	Valleys Foundation NHS Trust identified a need to evaluate the ongoing service-wide
22	trauma-informed care implementation effort. An absence of staff, service user and
23	system-related outcomes specific to trauma-informed care presented barriers to
24	monitoring the adoption of trauma-informed approaches and progress over time across

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

25	the Tees, Esk and Wear Valleys Foundation NHS Trust. This paper describes the co-
26	production of a new self-assessment tool, Roots, a discussion-based framework that
27	facilitates learning and improvement by reflecting on positive or negative examples of
28	trauma-informed services.
29	Methods: Using secondary data obtained from an affiliated national trauma summit
30	and instruments found in literature, domains and items were co-produced with the
31	help of trauma-informed care leads, NHS staff and service users. The research design
32	consisted of community-based co-production methods such as surveys, focus groups,
33	and expert consultations.
34	Results: Adopting trauma-informed care requires enthusiasm and commitment from
35	all members of the organisation. Services must adapt to meet the dynamic needs of
36	staff and service users to ensure they remain trauma-informed; this must be done as a
37	community.
38	Conclusions: Following an extensive co-production process, the Roots framework
39	was published open-access and accompanied by a user manual. Roots can provide
40	both qualitative and quantitative insights on trauma-informed care implementation by
41	provoking the sharing of experience across services.
42	Keywords: Trauma-informed Care; Implementation; Evaluation Framework; Self-
43	assessment; Co-production.
44	
45	
46	
47	
48	

49 **1. Introduction**

50	Trauma-informed care has seen various implementation efforts; many have seen local
51	success; others have failed to provide sufficient evidence (1-4). Trauma-informed
52	care is an evolving organic system model (5,6). It requires a paradigm shift in thinking
53	for service providers to deliver care that is rooted in the understanding of the
54	widespread prevalence and effects of trauma on people (7). An informed
55	conceptualisation of trauma is threefold: the trauma event, acknowledging that
56	traumatisation can occur when psychological/social integrity is threatened; how the
57	event is <i>experienced</i> ; and the <i>effects</i> of the event (7). This shift in thinking and
58	conceptualisation of trauma constitutes trauma-informed care. All members of the
59	organisation must embrace the constituents of trauma-informed care for the system
60	model to be of significant benefit; it must not be imposed upon individuals but rather
61	emerge from individuals who experience the organisation (8,9). For systemic
62	adoption, many organisations require concrete evidence of the benefits that trauma-
63	informed care provides (10,11). These benefits are numerous; however, they are often
64	not quantifiable. This does not serve the implementation of trauma-informed care
65	well, as service providers are requesting effective techniques for implementing the
66	necessary changes and specific examples of what it means in practice (10).
67	Establishing the need to develop metrics that can be used in qualitative and
68	quantitative ways to demonstrate the effective implementation of trauma-informed
69	care and realise the benefits that this systemic and individualistic change can provide
70	(11). Calling for a development process bespoke to the organisation, as trauma-
71	informed care is an emergent paradigm (3,8).
72	Organisational culture change demands individuals to follow suit and to sometimes
73	abandon personal principles (12-14). A frictionless change requires the consideration
74	of affect, sub-groups, personal and existing organisational values, and the quantity and
75	quality of support from leadership (8,9). Factoring in complexity concepts such as

76	emergence, self-organisation, and the sensitivity to initial conditions can assist in the
77	understanding of human systems and utilising previous developments can save time
78	and help co-construct a bigger picture of what it means to be trauma-informed (15,16).
79	This article provides basic information on the Roots framework so that future work
80	can benefit; it uses a similar reporting structure provided by Jung et al. (2009); as a
81	result, readers are given a topographical view of the framework (9).

82 2. Background

83 At the Tees, Esk and Wear Valleys (TEWV) Foundation NHS Trust, Dr Angela 84 Kennedy, with the help of other key figures, established the trauma-informed care 85 programme to implement trauma-informed care into services. Large-scale training and 86 service-change efforts were proving successful. However, an efficient implementation 87 and evaluation method was seen to be missing. Research and development efforts 88 were identified as being necessary to investigate potential solutions. The project that 89 emerged had one aim: To co-produce an integrative framework for data collection, 90 analysis, and interpretation. The objectives to reach this aim were threefold. The first 91 was to identify a relevant body of knowledge and investigate similar approaches that 92 have sought to evaluate trauma-informed care. The second was to co-produce an 93 evaluation framework that is bespoke to the United Kingdom. The third was to 94 produce documentation for practical use. Three fields from the literature helped 95 construct the narrative. These were organisational culture change, complexity theory, 96 and trauma-informed care. The organisational culture change literature is highly 97 applicable to trauma-informed care (8,9). Trauma-informed care can be viewed as a 98 culture and the change required is a cultural one. Complexity theory can offer an 99 informative view of culture - recognising that culture is a complex phenomenon that 100 emerges in the interactions between actors is conducive to successful change 101 (15,16,18). Within the trauma-informed care literature, previous frameworks were 102 identified; these were systematically evaluated (1-3,19). The Roots tool emerged from

103	a co-production approach to development with staff and service-users being consulted
104	at each stage.

105 2.1. The Implementation of Trauma-Informed Care

106 Trauma-informed care is socially constructed in the environment (9). It is manifested 107 and brought to life by the inhabitants of the service. Significant system change is 108 required for the implementation of trauma-informed care. This change marks a shift in 109 thinking from what is wrong with you to what happened to you. This is an essential 110 step to begin the narrative of care (7). All facets of the organisation must engage in 111 this transformation for trauma-informed care to recognise success. Adaptations to the 112 physical environment, raising awareness, and training staff are relatively minor steps 113 in comparison to the tidal-wave requirements of change on systems and care processes 114 (20).

- 115
 2.2. Roots A Discussion-Based Framework That Facilitates Learning and

 116
 Improvement by Reflecting on Positive or Negative Examples of Trauma
- 117 informed Services.

118 Roots is a developmental framework that uses insights from organisational culture 119 change, human behaviour, complexity theory, and trauma-informed care evaluation. 120 Roots was developed in the United Kingdom and was released in April 2021. There 121 are currently two versions available. One for staff and another for service users. Both 122 versions are identical with only changes made to terminology for accessibility 123 purposes. The definition or conceptual model of Roots is as follows: for an 124 organisation to be trauma-informed, it needs to apply trauma-informed principles and 125 culture in practice. Adopting a systems-wide value model requires enthusiasm and 126 commitment from all members of the organisation. As organisations and individuals 127 within the organisation change, the service must adapt to meet the dynamic needs of 128 staff and service users to ensure they remain trauma-informed (9). Communication

129	and clarification of these values, across teams, departments, buildings, or trusts will
130	assist in providing individuals with the self-knowledge that is often missing, i.e., how
131	are we adhering to the principles of trauma-informed care that we, as a group, have
132	selected as being relevant right now? and how are we able to improve in the areas
133	that we are not doing so well in? The intended purpose of Roots is to map the
134	implementation journey of trauma-informed care. The tool is for staff, service users,
135	and teams to think about what might make up trauma-informed care in their areas. It
136	takes the form of a word document that is completed by the facilitator on the
137	discussion of each item. Roots is comprised of 54 items answered quantitatively using
138	a RAG (red, amber, and green) rating, and qualitatively by prompting for reasons of
139	applicability and examples in practice. These items are shared among seven domains:
140	Safety (11 items), Language (8 items), Social (7 items), Trauma-specific Interventions
141	(7 items), Empowerment (7 items), Whole System (6 items), and Compassionate
142	Leadership (8 items). The set of practice points for reflection by the group challenges
143	thinking and enables discussion. The applicable to service column is asking for the
144	reason why this item needs to be applied in service to facilitate trauma-informed care.
145	However, each item is indicative and may not be applicable in every setting. The
146	implementation column is the RAG rating which asks the user how trauma-informed
147	they believe their service is with regards to the item in question. The <i>example</i> column
148	asks for examples as to why the service may or may not be delivering trauma-
149	informed care. Reflecting on each practice point can stimulate positive or negative
150	examples and provide meaningful information. The act of assigning a colour can
151	allow the individual or service to reflect on their current standing with trauma-
152	informed service delivery. This can also prompt and motivate individuals and services
153	to improve delivery. Providing examples can be useful for clarity and comparison.
154	The level of measurement used in Roots is ordinal.

155	Items were generated by using a combination of surveys, focus groups, expert
156	meetings, and secondary data. The domains within Roots were obtained from the
157	National Trauma-informed Care Community of Action's implementation report titled:
158	Creating a Narrative for Trauma-Informed Service Transformation which emerged
159	from a summit of clinicians, managers, leaders, people with lived experience,
160	researchers, and other interested parties on Thursday 28th March 2019 (21). An
161	investigation into the literature uncovered previous approaches. These instruments and
162	frameworks were studied, and domains were taken from four prominent frameworks
163	which were then compared at a trauma leads meeting with the domains that emerged
164	from the national trauma-informed care summit. At the same meeting, trauma leads
165	were provided with definitions on the summit domains and asked to provide a set of
166	standardised questions (items). These items were then shared with psychologists to
167	translate for staff accessibility purposes. Further translation was required for the
168	service-user version and so the staff version was shared with an expert by experience
169	group. Both staff and service-user versions of the framework were then issued to
170	wider audiences at the NHS trust using surveys to further articulate the forms. Trauma
171	leads meetings were used to discuss the results and changes were decided on. Focus
172	groups were then arranged to pilot test the framework. These were conducted with
173	staff; trauma leads and their staff members in their respective services, and a small
174	group of service-users. Prior to the focus groups, the service-user framework was
175	modified to reflect a third-person perspective rather than what was seen to be a
176	confusing first-person one; this was the only significant modification made to the
177	framework since the initial translation. Items were not reduced or modified further as
178	they were voted as being essential to remain in the framework. Due to resource
179	restrictions, it was not possible to conduct a follow-up assessment on the
180	psychometric properties of an instrument (reliability and validity). However, focus
181	group data would suggest strong face validity, acceptability, and feasibility. The

182	framework does have the potential to be susceptible to bias if the facilitator is part of
183	the team or has an agenda. This can be mitigated by using external facilitators. There
184	are no norms attached to the Roots tool. However, these can be created locally if
185	group comparisons are desired. Roots has no formal method of calibration. Informal
186	methods can be set by users of the tool to ascertain accuracy. There is currently no
187	data recorded for further tests of reliability such as internal consistency or
188	reproducibility (test-retest and inter-observer). The content, criterion, predictive,
189	convergent, discriminant, cross-cultural, and dimensional validity of Roots remains
190	untested. Roots is designed to be sensitive and open to change, the tool must reflect
191	the needs of the individuals it serves.
192	The Roots tool was co-developed, and pilot tested in mental health care settings.
193	These services included the community, prison services, adult wards, and CAMHS
194	(Child and Adolescent Mental Health Services). However, Roots has the potential to
195	be deployed in any setting, with changes made to the language and the exercise.
195 196	be deployed in any setting, with changes made to the language and the exercise.3. Statement of the Problem
195 196 197	be deployed in any setting, with changes made to the language and the exercise.3. Statement of the ProblemThe Tees, Esk and Wear Valleys NHS Foundation Trust has been implementing
195 196 197 198	 be deployed in any setting, with changes made to the language and the exercise. 3. Statement of the Problem The Tees, Esk and Wear Valleys NHS Foundation Trust has been implementing trauma-informed care for many years. Early implementation efforts took the form of
195 196 197 198 199	 be deployed in any setting, with changes made to the language and the exercise. 3. Statement of the Problem The Tees, Esk and Wear Valleys NHS Foundation Trust has been implementing trauma-informed care for many years. Early implementation efforts took the form of care pathways. In 2009, the trauma-informed pathway was designated as the first
 195 196 197 198 199 200 	be deployed in any setting, with changes made to the language and the exercise. 3. Statement of the Problem The Tees, Esk and Wear Valleys NHS Foundation Trust has been implementing trauma-informed care for many years. Early implementation efforts took the form of care pathways. In 2009, the trauma-informed pathway was designated as the first clinical link pathway. This pathway differed from clinical routes in that it was relevant
 195 196 197 198 199 200 201 	be deployed in any setting, with changes made to the language and the exercise. 3. Statement of the Problem The Tees, Esk and Wear Valleys NHS Foundation Trust has been implementing trauma-informed care for many years. Early implementation efforts took the form of care pathways. In 2009, the trauma-informed pathway was designated as the first clinical link pathway. This pathway differed from clinical routes in that it was relevant regardless of the diagnosis of trauma. Instead of encouraging talents that the
 195 196 197 198 199 200 201 202 	be deployed in any setting, with changes made to the language and the exercise. 3. Statement of the Problem The Tees, Esk and Wear Valleys NHS Foundation Trust has been implementing trauma-informed care for many years. Early implementation efforts took the form of care pathways. In 2009, the trauma-informed pathway was designated as the first clinical link pathway. This pathway differed from clinical routes in that it was relevant regardless of the diagnosis of trauma. Instead of encouraging talents that the workforce cannot supply, the trauma-informed pathway encourages staff to use their
 195 196 197 198 199 200 201 202 203 	be deployed in any setting, with changes made to the language and the exercise. 3. Statement of the Problem The Tees, Esk and Wear Valleys NHS Foundation Trust has been implementing trauma-informed care for many years. Early implementation efforts took the form of care pathways. In 2009, the trauma-informed pathway was designated as the first clinical link pathway. This pathway differed from clinical routes in that it was relevant regardless of the diagnosis of trauma. Instead of encouraging talents that the workforce cannot supply, the trauma-informed pathway encourages staff to use their skills in trauma-informed ways. This empowered staff when they realised that this
 195 196 197 198 199 200 201 202 203 204 	be deployed in any setting, with changes made to the language and the exercise. 3. Statement of the Problem The Tees, Esk and Wear Valleys NHS Foundation Trust has been implementing trauma-informed care for many years. Early implementation efforts took the form of care pathways. In 2009, the trauma-informed pathway was designated as the first clinical link pathway. This pathway differed from clinical routes in that it was relevant regardless of the diagnosis of trauma. Instead of encouraging talents that the workforce cannot supply, the trauma-informed pathway encourages staff to use their skills in trauma-informed ways. This empowered staff when they realised that this meant they could offer something critical to service users. A business case was
 195 196 197 198 199 200 201 202 203 204 205 	be deployed in any setting, with changes made to the language and the exercise. 3. Statement of the Problem The Tees, Esk and Wear Valleys NHS Foundation Trust has been implementing trauma-informed care for many years. Early implementation efforts took the form of care pathways. In 2009, the trauma-informed pathway was designated as the first clinical link pathway. This pathway differed from clinical routes in that it was relevant regardless of the diagnosis of trauma. Instead of encouraging talents that the workforce cannot supply, the trauma-informed pathway encourages staff to use their skills in trauma-informed ways. This empowered staff when they realised that this meant they could offer something critical to service users. A business case was developed by the trauma-informed care lead, Dr Angela Kennedy, for a formally
 195 196 197 198 199 200 201 202 203 204 205 206 	be deployed in any setting, with changes made to the language and the exercise. 3. Statement of the Problem The Tees, Esk and Wear Valleys NHS Foundation Trust has been implementing trauma-informed care for many years. Early implementation efforts took the form of care pathways. In 2009, the trauma-informed pathway was designated as the first clinical link pathway. This pathway differed from clinical routes in that it was relevant regardless of the diagnosis of trauma. Instead of encouraging talents that the workforce cannot supply, the trauma-informed pathway encourages staff to use their skills in trauma-informed ways. This empowered staff when they realised that this meant they could offer something critical to service users. A business case was developed by the trauma-informed care lead, Dr Angela Kennedy, for a formally funded project to embed trauma-informed care into services. A goal of the project was

208	contribute to the evidence-base for trauma-informed care. The trauma-informed care
209	programme at the Tees, Esk, and Wear Valleys NHS Foundation Trust realised the
210	need for an evaluation and implementation framework to make progress with this
211	trauma-informed system change. Checklists have been created and measurements
212	have been conducted previously regarding trauma-informed change, but never in the
213	United Kingdom. The NHS is a unique system in that it represents a large number of
214	individuals spread out over many localities. The TEWV NHS Foundation Trust is one
215	of the largest specialist mental health and learning disabilities trusts in the country,
216	with an annual income of £380 million and a workforce of over 6700 staff operating
217	from around 100 sites in Durham, Teeside, North Yorkshire, and York and Selby.
218	TEWV NHS provide a range of inpatient and community services to 2 million people
219	living in County Durham, the Tees Valley, Scarborough, Whitby, Ryedale, Harrogate,
220	Hambleton and Richmondshire. TEWV NHS services are spread out over a wide
221	geographical area of around 3600 square miles, inclusive of coastal, rural, and
222	industrial areas (22). The geographical nature of the TEWV NHS Foundation Trust is
223	conducive to complexity. All NHS trusts are subject to stringent ethical procedures
224	and research involving staff and service-users undergo strict ethical clearance
225	procedures. An implementation and evaluation framework unique to these particular
226	circumstances that was able to navigate strict ethical boundaries was essential.
227	4. Methods Used During the Development of the Roots Framework
228	The collaborative development of the Roots framework was a five-step process.
229	Firstly, a literature review was undertaken to investigate similar approaches. These
230	other approaches were evaluated using an existing culture instrument review
231	framework (17), and were then compared with the domains that emerged from the
232	national trauma summit at a trauma leads meeting. Items were generated at the same
233	meeting and were taken to surveys with staff and service users to confirm articulation.
234	Focus groups were held with one service user group and five different staff services.

These were used to further articulate and pilot test the framework in practice. Theprocess can be seen in Figure 1.

237

238	Fig.1 – Methods used during the co-production process of Roots
239	The design of the Roots framework stemmed from the ongoing work of the trauma-
240	informed care programme. This meant that the research design evolved alongside
241	programme implementation. Monthly meetings with trauma leads contributed
242	significantly to the research design. The trauma leads are select individuals who
243	represent trauma-informed care in their respective services. They were instrumental in
244	the facilitation of various developmental facets, including access to various staff and
245	service user groups. Approximately twelve trauma leads would usually be present at
246	each meeting. The first phase of this study was an evaluation of other similar
247	instruments and frameworks (9). The literature contributed to expectations and other
248	considerations for trauma-informed service evaluation; including the use of principles
249	or domains for which both implementation and evaluation can revolve. Meanwhile,
250	the trauma-informed programme was a joint-organiser of a national trauma summit
251	titled: Creating a Narrative for Trauma-informed Service Transformation (21). The
252	goal of this summit was to respond to calls from people with lived experience of
253	trauma in services and the challenge of trauma being included in the new NHS 10-
254	year plan. It was arranged as a forum to share ideas for good practice and an
255	opportunity to network with others who are motivated towards similar goals. This was
256	held on Thursday 28th March and brought together clinicians, managers, leaders,
257	people with lived experience, researchers, and many others. Attendees were asked to
258	think of a specific positive example that they have experienced, witnessed, or been
259	involved in. The aim of this was to tap into individual wisdom and creativity and to
260	explore the multiple dimensions of what it means to be trauma-informed and share

261	concrete examples that can be replicated by others. A report was drafted titled:
262	Developing Real-World System Capability in Trauma-informed Care: Learning from
263	Good Practice (21). In this report, seven domains were observed as having emerged
264	from the summit, these were: 1) Safety, 2) Human Experience Language, 3)
265	Empowerment, 4) Healing Interventions, 5) Responsive System Design, 6)
266	Compassionate and Transformational Leadership, and 7) Relational Reparation (21).
267	These domains were significant to the Roots framework as they had been developed
268	with individuals that have experienced UK health systems. The domains were used
269	but their wording was amended to better reflect local services. The revised domains
270	were: 1) Safety, 2) Language, 3) Social, 4) Trauma-specific Interventions, 5)
271	Empowerment, 6) Whole System, and 7) Compassionate Leadership.
272	5. The Co-Production of Domains and Items
273	At a trauma leads meeting, the seven domains were judged against the domains taken
274	from the literature. Four instruments were selected for this process, the Attitudes
275	Related to Trauma-Informed Care (ARTIC) (1), the Creating Cultures of Trauma-
276	informed Care (CCTIC) (4), the TICOMETER (2), and the Trauma-informed Practice
277	(TIP) Scales (19). These were identified as being relevant and empirically tested.
278	During the meeting, the trauma leads worked through a slideshow that presented the
279	domains from the literature alongside the domains from the summit and were asked to
280	determine encapsulation. Encapsulation was confirmed if one or more of the summit
281	domains captured the meaning used in the language of the domains from the literature.
282	Table 1 presents the ARTIC comparison, Table 2 presents the CCTIC comparison,
283	Table 3 presents the TICOMETER comparison, and Table 4 presents the TIP Scales
284	comparison. The left column represents the domains from the instruments found in the
285	literature, and the right column represents which domain from the summit that trauma
286	leads felt captured the meaning of the corresponding domain from the literature. This
287	process was held to determine if the summit domains were appropriate for use. The

288	trauma leads all agreed through consensus that all domains from the four instruments
289	were captured by one or more of the domains from the summit. However, trauma
290	leads felt as though staff wellbeing should represent the eighth domain.
291	Table 1 – ARTIC Comparison

ARTIC	SUMMIT
Underlying Causes of Problem	Social Context, Language
Behaviour and Symptoms	
Responses To Problem Behaviour and	Safety, Social Context, Trauma-Specific
Symptoms	Interventions
On-The-Job Behaviour	All
Self-Efficacy at Work	Safety, Whole System, Compassionate
	Leadership, Empowerment
Reactions to the Work	Trauma-Specific Interventions
Personal Support of TIC	Safety, Whole System
System-Wide Support for TIC	Whole System, Compassionate
	Leadership, Safety, Social Context

Table 2 – CCTIC Comparison

CCTIC	SUMMIT
Safety	All
Trustworthiness	Safety, Language, Empowerment, Whole
	System, Social Context
Choice	Empowerment, Safety, Social Context
Collaboration	Empowerment, Whole System, Social
	Context, Safety
Empowerment	Empowerment, Language, Whole System

Trauma Screening Process	Safety, Language, Empowerment,
	Trauma-Specific Interventions, Whole
	System
Table 3 – TICOM	ETER Comparison
TICOMETER	SUMMIT
Building Trauma-Informed Knowledge	All
and Skills	
Establishing Trusting Relationships	All
Respecting Service Users	Whole System, Safety, Language
Fostering Trauma-Informed Service	All
Delivery	
Promoting Trauma-Informed Policies and	Whole System, Language
Procedures	
Table 4 TID Co	alas Comparison
Table 4 – Tir Sc	cales Comparison
TIP SCALES	SUMMIT
The Environment of Agency & Mutual	Social Context, Safety, Empowerment,
Respect	Language
Access to Information to Trauma	Trauma-Specific Interventions, Safety,
	Empowerment, Language
Opportunities for Connection	Empowerment, Trauma-Specific
	Interventions, Social Context

Emphasis on Strengths

293

294

295

296

Interventions, Language, Safety

Social Context, Trauma-Specific

Cultural Responsiveness & Inclusivity All

Support for Parenting

Trauma-Specific Interventions, Social

Context, Safety

297

298	At the same meeting, trauma leads were asked to develop up to 10 items per domain.
299	Trauma leads worked through a slideshow that presented definitions of each domain.
300	Four items were generated for the domain safety. Six items were generated for the
301	domain language. Four items were generated for the domain social. Six items were
302	generated for the domain Trauma-specific Interventions. Eleven items were generated
303	for the domain empowerment. Seven items were generated for the domain Whole
304	System. Eight items were generated for the domain compassionate leadership. Four
305	items were generated for the domain staff wellbeing.
206	
300	The results of this meeting were shared with the trauma-informed care programme
307	team. The items underwent translation to accommodate common language used by
308	staff for purposes of familiarity and accessibility. This discussion contributed to the
309	removal of some items and the addition of others. The requested domain, staff
310	wellbeing, was removed as it was voted unnecessary in respect of it being represented
311	across the other domains. As two forms were needed, one for staff and another for
312	service users, the translated items were then sent to an experts by experience group
313	working from the Recovery College in Durham, United Kingdom. The experts by
314	experience group worked through the items and translated them for service user
315	accessibility.
316	Both forms, existing only as items, were then placed into two separate SurveyMonkey
317	surveys. Convenience, criterion, and snowball sampling methods were used for the
318	recruitment of participants. Staff and service-user samples were recruited through
319	trauma leads who distributed information on the study. When participants registered

320	interest, the trauma leads then distributed further information through a participant
321	information sheet and a web link to the survey. The surveys asked participants to
322	prioritise the items and leave feedback on articulation. The addition of a RAG rating
323	added a quantifiable indicator to the survey. The staff form was distributed to staff
324	members, and the service-user form was distributed to service users. The inclusion
325	criteria were as follows: staff are identified as being available, engaged in the TEWV
326	NHS trust, and chosen or recommended by those aware of the study. Service users are
327	identified as being available, involved in the NHS trust, and selected by staff. The
328	exclusion criteria were as follows: no affiliation with the NHS TEWV Foundation
329	Trust, under the age of 18 or over the age of 65, or at imminent risk of harming
330	themselves or others. Recruitment was difficult as national restrictions were imposed
331	as a result of the Covid-19 pandemic. The results were analysed using Microsoft Excel
332	to determine priority. A trauma leads meeting was held to discuss these results and it
333	was voted that all items should remain in the final framework, mainly due to high
334	applicability. Although the survey results contradicted this for a few items, the
335	presentation of the survey was questioned for confusion.
336	A quality assurance process took place and involved confirming meaning across items
337	in both staff and service user forms with the research team. The framework was
338	drafted as a table, in which there are four columns: practice point for consideration
339	(item), applicable to service (reason), implementation status (RAG rating), and
340	example (justification). The set of practice points for reflection by the group
341	challenges thinking and enables discussion. The applicable to service column asks
342	why this item needs to be applied in service to facilitate trauma-informed care.
343	However, it is recognised that each item is indicative and may not be applicable in
344	every setting. The implementation column is the RAG rating and questions the user on
345	how trauma-informed they believe their service is with regards to the item in question.
346	The example column asks for examples as to why the service may or may not be

347	delivering trauma-informed care. Reflecting on each practice point can stimulate
348	positive or negative examples and provide meaningful information. The act of
349	assigning a colour can allow the individual or service to reflect on their current
350	standing with trauma-informed service delivery. This can also prompt and motivate
351	individuals and services to improve delivery. Providing examples can be useful for
352	clarity and comparison.
353	Pilot tests were held with both forms with staff and service user sample groups. Five
354	full-length evaluation exercises were held with staff, and one rudimentary exercise
355	was held with service users. Staff focus groups were evaluating the framework and the
356	exercise, whilst the service user group was evaluating the framework. Service users
357	felt empowered when discussing the items of the framework and agreed that the
358	majority of items were essential. Staff found the exercise helpful, and all services
359	requested the completed framework to begin actioning items.
360	A user manual was created to facilitate instructions and to contain the Roots
361	framework. This is now published open access on the Future NHS community
362	platform for practitioners and researchers as long as they have an NHS or public
363	health England email address.
364	6. The Roots Framework Learning Model
365	The Roots framework uses insights from complexity theory. This is recognising that
366	human change is complex because there is rarely one right way of doing something.
367	The service will need to adapt to meet each individual's needs and remain responsive
368	over time. These complexity principles guide the use of Roots towards change and
369	evaluation: change in individuals or organisations is rarely linear. A reflexive
370	approach that evolves is of benefit. There is not a one size fits all interpretation of
371	trauma-informed implementation. Different settings need to define what is needed for
372	them through methods of co-production. Different teams within an organisation,

373	different individuals within teams, and different service users may all display or
374	perceive different strengths concerning the implementation of trauma-informed care.
375	Roots allows for the bringing together of different narratives towards a wider picture.
376	The items emerged from previous examples and are created to be both generic and
377	specific but not exhaustive. New ones can be created as long as they are tangible and
378	observable. Roots needs to be embedded within a learning organisation framework
379	accompanied by an attitude of respect to ensure that progress can be made.

380 7. Study Limitations

381 In the UK, there are many restrictions placed upon social and healthcare research. 382 Research studies must navigate strict ethical standards. In many cases, these standards 383 can form boundaries for what is possible in research. As trauma-informed care is a 384 relatively new concept in the UK, pioneering research must first take the first steps. 385 These initial studies must take place so that more elaborate studies can follow. As 386 Roots is the first trauma-informed evaluation framework to emerge out of the UK, 387 many aspects of the study were impeded. The Tees, Esk and Wear Valleys NHS 388 Foundation Trust has a workforce of over 6700 staff operating from around 100 sites 389 in Durham, Teesside, North Yorkshire and York and Shelby (22). This study is unable 390 to claim to be fully representative as the samples used were often limited to 391 convenience, and the actual number of participants was low. To achieve a confidence 392 level of 99% for a population size of 6700 with a 5% margin of error, 604 staff 393 members must have been involved in the study. For service users, an ideal sample size 394 with the same parameters would be inclusive of 663 participants. In combination, a 395 repeated study would need to consult at least 1200 participants to be statistically 396 significant. The use of qualitative data helps mitigate this through practicality and 397 transparency. The research team had ultimate control over the end-product, meaning 398 that key stakeholders at both staff and service user levels should have been involved in 399 signing off the product. The service user focus group is a good example of this, there

400	were questions raised over some items, but this was not regarded in the finished
401	version. The development of Roots also took place during the Covid-19 pandemic and
402	data collection began when the UK was put under lockdown - this caused delays to
403	the study as all non-covid related research received suspensions from the NHS. There
404	is also a distinct lack of any form of psychometric testing – delays to research and
405	resource expiration made it impossible to continue working on Roots.
10.4	

406 Further Work

407 Roots is the United Kingdom's first foray into the evaluation of trauma-informed 408 services. Much of the precursory work is already done in the United States of America 409 (USA). However, UK healthcare services are delivered very differently than they are 410 in the USA. The development of Roots provides perfect foundations for future work, 411 either using or building on Roots. One of the fundamentals of UK-based development 412 is the practicalities of set-up. Establishing long-term commitment and navigating strict 413 ethical standards are two of the arrangements that must be considered. Acquiring 414 health research authority approval through a completed Integrated Research 415 Application System (IRAS) should be completed early, and applications should be 416 thorough to ensure acceptance. Before use, a thorough psychometric assessment of the 417 tool should be prioritised to ensure reliability and validity. Self-assessment 418 maintenance should be carried out with exceptional regard to the language used. A 419 dynamic self-assessment should be co-produced; the contents of which change with 420 time to ensure consistent reliability with the people they serve.

421 Conclusions

425

422The co-production of the Roots framework involved staff and service-users from the423Tees, Esk and Wear Valleys Foundation NHS Trust and also utilised secondary data424from a national trauma summit titled: *Creating a Narrative for Trauma-informed*

Service Transformation. Relevant secondary data, trauma-informed care programme

426	lead meetings, surveys, focus groups, and consultations with staff and service users
427	constituted the research design. The development of Roots was an experimental
428	process and the research evolved alongside the progression of the trauma-informed
429	care programme at the Tees, Esk and Wear Valleys Foundation NHS Trust. Roots is
430	published open-access and is supported by a user manual.
431	
432	
433	
434	DECLARATIONS:
435	• Ethics approval and consent to participate – The United Kingdom's Research
436	Ethics Committee of the National Health Service's Health Research
437	Authority gave ethical approval for this work. Informed consent was obtained
438	from all individuals included in the study where necessary.
439	• Consent for publication – The authors affirm that human research
440	participants provided informed consent where necessary for publication of
441	data collected.
442	• Availability of data and material – Not Applicable
443	• Competing interests – All authors certify that they have no affiliations with
444	or involvement in any organization or entity with any financial interest or
445	non-financial interest in the subject matter or materials discussed in this
446	manuscript.
447	• Funding – This work was made possible by support from the Tees, Esk and
448	Wear Valleys Foundation NHS Trust and Northumbria University. Funding
449	bodies had no influence on the study.
450	Authors' Contributions

451	• CONCEPTION: ST, PS and AK
452	• METHODOLOGY: ST, PS, AK and PP
453	• DATA COLLECTION: ST
454	• INTERPRETATION OR ANALYSIS OF DATA: ST
455	• PREPARATION OF THE MANUSCRIPT: ST
456	• REVISION FOR IMPORTANT INTELLECTUAL CONTENT: ST,
457	PS, AK, and PP
458	• SUPERVISION: PS, AK, and PP
459	• Acknowledgements: This work was made possible by support from the
460	Tees, Esk and Wear Valleys Foundation NHS Trust and Northumbria
461	University
462	
463	
464	
465	
466	
467	
468	
469	
470	
471	References
472	

473 474 475 476	1.	Baker CN, Brown SM, Wilcox PD, Overstreet S, Arora P. Development and psychometric evaluation of the Attitudes Related to Trauma-Informed Care (ARTIC) scale. School Mental Health. 2016;8(1):61–76.
477 478 479 480	2.	Bassuk EL, Unick GJ, Paquette K, Richard MK. Developing an instrument to measure organizational trauma-informed care in human services: The TICOMETER. Psychology of Violence. 2017;7(1):150–7.
481 482 483 484	3.	Richardson MM, Coryn CLS, Henry J, Black-Pond C, Unrau Y. Development and evaluation of the trauma-informed system change instrument: Factorial validity and implications for use. Child and Adolescent Social Work Journal. 2012;29(3):167–84.
485 486	4.	Fallot RD, Harris M. Trauma-informed services: A self-assessment and planning protocol. Washington, DC: Community Connections. 2006;
487 488	5.	Sweeney K, Griffiths F. Complexity and healthcare: an introduction. Radcliffe Publishing; 2002.
489 490 491 492 493 494	6.	Sweeney A, Clement S, Filson B, Kennedy A. Trauma-informed mental healthcare in the UK: what is it and how can we further its development? Mental Health Review Journal [Internet]. 2016;21(3):174–92. Available from: https://www.emeraldinsight.com/doi/abs/10.1108/MHRJ-01-2015- 0006
495 496 497	7.	Sweeney A, Filson B, Kennedy A, Collinson L, Gillard S. A paradigm shift: relationships in trauma-informed mental health services. BJPsych Advances. 2018;24(5):319–33.
498 499 500	8.	Thirkle SA, Kennedy A, Sice P. A Case for TIC: A Complex Adaptive Systems Enquiry for Trauma Informed Care. International Journal of Systems and Society (IJSS). 2018;5(2):1–12.
501 502 503	9.	Thirkle SA, Kennedy A, Sice P. Instruments for Exploring Trauma- Informed Care. Journal of Health & Human Services Administration. 2021;44(1).
504 505 506	10.	Yatchmenoff DK, Sundborg SA, Davis MA. Implementing Trauma- Informed Care: Recommendations on the Process. Advances in Social Work. 2017;18(1):167–85.
507 508 509 510	11.	Jennings A. Models for developing trauma-informed behavioral health systems and trauma-specific services. Alexandria, VA: National Association of State Mental Health Program Directors, National Technical Assistance Center for State Mental Health Planning. 2004;

511 512 513 514	12.	Johnson A, Nguyen H, Groth M, Wang K, Ng JL. Time to change: A review of organisational culture change in health care organisations. Journal of Organizational Effectiveness: People and Performance. 2016;
515 516 517	13.	Whelan C. Organisational culture and cultural change: A network perspective. Australian & New Zealand journal of criminology. 2016;49(4):583–99.
518 519 520 521	14.	Parmelli E, Flodgren G, Beyer F, Baillie N, Schaafsma ME, Eccles MP. The effectiveness of strategies to change organisational culture to improve healthcare performance: a systematic review. Implementation Science. 2011;6(1):33.
522 523 524	15.	Stacey R. The emergence of knowledge in organization. Emergence, A Journal of Complexity Issues in Organizations and Management. 2000;2(4):23–39.
525 526	16.	Seel R. Emergence in organisations. Retrieved November. 2003;29:2006.
527 528 529	17.	Jung T, Scott T, Davies HTO, Bower P, Whalley D, McNally R, et al. Instruments for exploring organizational culture: A review of the literature. Public administration review. 2009;69(6):1087–96.
530 531	18.	Snowden DJ, Boone ME. A leader's framework for decision making. Harvard business review. 2007;85(11):68.
532 533 534	19.	Goodman LA, Sullivan CM, Serrata J, Perilla J, Wilson JM, Fauci JE, et al. Development and validation of the Trauma Informed Practice Scales. Journal of Community Psychology. 2016;44(6):747–64.
535 536	20.	Evans J. Person-centered care and culture change. Caring for the Ages. 2017;18(8):6.
537 538	21.	Kennedy A. Developing Real World System Capability in Trauma- informed Care: Learning from Good Practice. 2020;
539 540 541	22.	TEWV NHS. NHS Jobs - Employer Information [Internet]. 2021 [cited 2021 Oct 7]. Available from: https://www.jobs.nhs.uk/xi/agency_info/?agency_id=122252
542		

