

1 **Title page**

2 Title: The time is ripe to harmonize global acute febrile illness etiologic investigations

3 Running title: Harmonization of acute febrile illness investigations methods

4 Article category: Brief Report

5 Author names & institutional affiliations: José Moreira^{1,2}, B. Leticia Fernando-Carballo³,

6 Camille Escadafal³, Sabine Dittrich^{3,4}, Patrícia Brasil¹, André M Siqueira¹

7 Affiliations:

8 1.Laboratório de Pesquisa Clínica em Doenças Febris Agudas, Instituto Nacional de
9 Infectologia Evandro Chagas, Fundação Oswaldo Cruz, Rio de Janeiro, Brazil

10 2. Instituto Nacional de Cardiologia, Departamento de Ensino e Pesquisa, Rio de Janeiro,
11 Brazil

12 3. Foundation for Innovative New Diagnostics (FIND), Geneva, Switzerland

13 4. Nuffield Department of Medicine, University of Oxford, Oxford, United Kingdom

14 Corresponding author: José Moreira MD Ph.D; e-mail: jose.moreira@ini.fiocruz.br;

15 telephone number: +55 (21) 97035-8843; Laboratório de Pesquisa Clínica em Doenças Febris

16 Agudas, Instituto Nacional de Infectologia Evandro Chagas, Fundação Oswaldo Cruz, Av.

17 Brasil, 4365 – Manguinhos, Rio de Janeiro, Brazil, CEP: 21040-360; Twitter: @j_moreira_

18

19

20

21

22

23

24

25

26 **Abstract**

27 We investigated the impact of considering different case definitions of fever as inclusion criteria
28 among participants enrolling in an acute febrile illness investigation and observed that adopting a
29 subjective assessment of fever regardless of specific fever cut-off value accounted for the
30 diversity of clinical fever phenotypes and did not mischaracterize the febrile population.

31 **Keywords:** Antipyretic therapy; Fever; Acute febrile illness; Dipyrone; Methodology

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50 **Text**

51 **Introduction**

52 Acute febrile illness (AFI) is a common reason for people seeking care and represents a range
53 of infectious and non-infectious diseases, with wide variation by region of the globe [1]. AFI
54 investigations aim to provide national and subregional epidemiology and infectious diseases
55 surveillance and to inform treatment guidelines about empirical and pathogen-specific
56 management algorithms to prioritize interventions and funding. Several AFI investigations
57 have been conducted primarily in Southeast Asia, with few global initiatives in resource-
58 constrained countries [2]. However, there is no standardized approach to conducting such
59 studies, limiting the interpretation and comparability of study findings.

60 We were involved in a multinational project around fever host-biomarkers [3] and aimed to
61 conduct a pilot study to inform the best criteria for our study site selection. We hypothesized
62 that adopting a subjective or self-reported fever assessment as part of our inclusion criteria
63 would increase the sensitivity threshold for a broad array of febrile clinical phenotypes rather
64 than specifying a cut-off value for fever at presentation. Another hypothesis was that the
65 recent antipyretic intake – largely available over-the-counter- would lower body temperature
66 at arrival at the Emergency Departments (EDs).

67 **Methods**

68 We did a cross-sectional, study of patients presenting to two urban EDs (Unidade de Pronto
69 Atendimento Manguinhos & Unidade de Pronto Atendimento Rocha Miranda) in Rio de
70 Janeiro, Brazil, between October 28, 2018, and March 29, 2019. The study was conducted
71 following the Declaration of Helsinki and the Brazilian National Ethics Research Committee
72 and was approved in August 2018 (IRB approval no:70984617.9.0000.5262). We screened
73 all consecutive non-severe patients who presented to the EDs over the study period and
74 included those who reported fever ≤ 7 days regardless of temperature measured at the EDs.

Research assistants prospectively collected detailed information and interviewed participants or the primary caregivers to obtain data about the demographics, main chief complaint, vital signs, history of fever, peak measured temperature before arrival at the EDs (temperature maximum), the measured axillary temperature at the EDs, home administration of antipyretics, type/dose/timing of last administration, and tests performed at the EDs. Discharge diagnosis and treatment received after discharge was retrospectively collected by reviewing electronic medical records. The attending physician proposed the discharge diagnosis (not involved in the study), and its appropriateness was not re-evaluated. We used the following definitions: fever was subjectively identified by the participant at arrival at the EDs; antipyretics are mostly over-the-counter drugs that act on reducing body temperature usually with added analgesic effect, and were self-administered by the participants before seeking emergency care; adults those aged ≥ 18 years at recruitment, and non-severe AFI was an illness that was codified as blue or green according to the Manchester Triage System at EDs, which does not require hospital admission. We measured the axillary temperature at triage and estimated the magnitude of defervescence, considering the highest measured self-reported temperature before EDs arrival (Tmax). Suspected infection was defined as the initiation of any antimicrobial after an EDs visit.

Statistical analysis

Descriptive analysis was performed to characterize the distributions of several variables. A Chi-square test was used to compare categorical variables between study groups. Continuous variables were compared between the study groups using analysis of variance or the Kruskal-Wallis test if those were found to be normally or non-normally distributed, respectively. We dealt with missing data using the listwise deletion method (or complete case analysis). A logistic regression model was performed to ascertain the effects of recent antipyretic uptake on the likelihood that participants have an overt fever at EDs admission (i.e., ≥ 37.5

°C). A 2-sided *p-value* less than 0.05 was considered statistically significant. All statistical analyses were performed using SPSS software (version 26, IBM, Chicago, Illinois, USA) and Tableau Desktop (version 2020.4.2, Tableau, Seattle, Washington, USA). The primary outcome was to investigate the impact of considering different case definitions of fever as inclusion criteria among febrile patients attending EDs.

Results

During the study period, we triaged 1551 subjects, and 374 [24.1% (95% CI 22-26)] had a history of fever at arrival at the ED, corresponding to a cumulative incidence of fever at EDs of 24 per 100 triaged patients over the five months. These 374 febrile patients constitute our study population, of whom 248 (66.3%), 115 (30.7%), and 11 (2.9%) had a temperature < 37.5 °C, ≥ 37.5 °C, and no temperature measurement registered at arrival, respectively. The mean age was 30.6 [range: 0-84] years, adults (82.5%) and females (54.8%) predominated (Table). The median peak measured temperature before arrival at the ED was 38.8°C (38.0-39.0), and the median duration of fever was 1 [1-3] days.

Hundred ninety-eight febrile patients had a diagnosis of infection, with upper respiratory tract infection (42.4%) followed by undifferentiated febrile illness (15.7%) and urinary tract infection (14.6%) were the primary febrile syndromes (Supplementary Figure 1). Hundred thirty-one patients were prescribed any antibiotic after EDs visit, which beta-lactams (73/131, 55.7%), quinolones (23/131, 17.5%) and macrolides (16/131, 12.2%) were the classes most commonly prescribed. Patients who had temperature ≥ 37.5 °C at presentation were more likely to be diagnosed with an infection [76 (66%) vs. 116 (47%), crude OR: 2.2 (95% 1.4-3.5) and were prescribed antibiotic most frequently compared to those with temperature < 37.5 °C [54 (65%) vs. 75 (48%), crude OR: 2 (95% 1.1-3.5)].

The site of infection varied by the height of temperature mensurated at arrival at EDs (*p*<0.001). Patients who had a temperature < 37.5 °C at presentation were more likely to be

diagnosed with upper respiratory (31.6%) and urinary infections (15.4%) (Supplementary Figure 3). The most frequently etiology diagnostics recorded in those with temperature < 37.5 °C at presentation were acute pharyngitis (20/248, 8%), followed by community-acquired gastroenteritis (19/248, 7.6%), uncomplicated urinary tract infection (16/248, 6.4%), cutaneous abscess (14/248, 5.6%), and community-acquired pneumonia (9/248, 3.6%).

In total, 249/374 (66.6% (95% 61.5-71.3)) subjects reported self-administering an antipyretic at home within a median of 2h [IQR 2-6] before EDs admission (24.1% of the febrile subjects did not recall prior antipyretics intake). Dipyrene (72.2%) was the antipyretic drug most frequently administered, followed by acetaminophen (14.3%) and ibuprofen (6.3%). Most antipyretic takers consumed one class (85.9%), and the median number of doses taken was 2 [IQR 1-3]. Of these 249 subjects, 167/249 (67%) reported the last dose administered at a mean of 2 [2-5] hours before hospital presentation. Subjects who were treated with an antipyretic drug before arrival did not differ from untreated ones regarding the temperature at arrival at the ED visit (37 ± 0.99 vs. 36.8 ± 0.86 , $p = 0.20$) or the documented Tmax before the ED arrival (38.8 ± 0.79 vs. 38.4 ± 0.60 , $p = 0.29$).

The logistic regression model correctly classified 69.1% of the cases (Supplementary Table 1). Males were 2.2 times more likely to exhibit overt fever at presentation than females [aOR: 2.25, 95% CI: 1.19-4.25]. Recent antipyretic intake did not influence the pattern of fever at presentation [aOR:0.81, 95% CI:0.34-1.97].

Discussion

Our study suggests that considering a subjective/self-reported fever as an inclusion criterion rather than a specific fever cut-off value (i.e., ≥ 37.5 °C) at presentation for AFI etiologic investigation results in a more inclusive study population and account for the diversity of clinical fever phenotypes that are usually seen in clinical practice. Specifically, sixty-six

percent of our cohort would not have been included if we restricted our inclusion criteria to a specific febrile cut-off, potentially impacting study recruitment and patient characterization. Almost half (47.8%) of febrile participants with temperature < 37.5 °C at presentation were prescribed at least one antibiotic, and their infections were mainly upper respiratory, justifying their inclusion in AFI studies. Next, our findings suggest that the ubiquitous administration of antipyretics (66.6%) observed in our setting did not influence the febrile presentation of patients attending the EDs evaluated.

We concur with others [2, 4] and argue that there is an urgent need to standardize AFI study protocols to make sure that we can compare results from different studies and pool their results to develop national, regional, or global burden of fever estimates. Such consensus in AFI research would allow better comparisons, translate evidence to practice, and inform treatment policy. A harmonized definition of fever would enable a more robust and consistent capture of different clinical phenotypes, including those conditions where overt fever occurs inconsistently, such as upper respiratory infections and uncomplicated urinary infections. Others [2] have suggested an adapted STRENGTHENING the reporting of OBSERVATIONAL studies in Epidemiology (STROBE) checklist for conducting and reporting AFI research.

In conclusion, we suggest harmonizing the case definition of fever in AFI research for a history of fever regardless of temperature cut-off would enable a complete picture of the different clinical fever phenotypes, thus improving our understanding of the global burden of febrile illness.

Author contributions

JM: writing original draft, visualization, conceptualization. BC: Review and editing. CE: Review and editing. SD: review, editing. PB: conceptualization, writing, editing, supervision. AS: conceptualization, writing, editing, supervision

Funding

The Biomarker for Fever-Diagnostic (BFF-Dx) study was funded by the Dutch, Australian, and UK governments.

Transparency declaration

The authors have no conflict of interest to declare.

References

1. Moreira J, Bressan CS, Brasil P, Siqueira AM. Epidemiology of acute febrile illness in Latin America. *Clin Microbiol Infect.* 2018; 24(8):827-835.
2. Rhee C, Kharod GA, Schaad N et al. Global knowledge gaps in acute febrile illness etiologic investigations: a scoping review. *PLoS Negl Trop Dis.* 2019; 13(11): e0007792.
3. Escadafal C, Geis S, Siqueira AM, et al. Bacterial versus non-bacterial infections: a methodology to support use-case-driven product development of diagnostics. *BMJ Glob Health.* 2020; 5(10)e:003141.2.
4. Newton PN, Guerin PJ. Febrile illness mapping-much of the world without data and without evidence-based treatments. *BMC Med.* 2010; 18(1):287.

Supplementary Tables

Supplementary Table 1. Prediction model for the likelihood of overt fever at presentation in 374 febrile patients attending emergency departments in Rio de Janeiro, Brazil, October 2018 – March 2019.

Supplementary Table 2. Diagnosis, febrile syndrome, and antibiotic prescription in febrile patients attending Emergency departments in Rio de Janeiro, Brazil, October 2018-March 2019

Supplementary Figures

Supplementary Figure 1. Antibiotic class according to the foci of infection in febrile patients attending Emergency departments in Rio de Janeiro, Brazil, October 2018-March 2019

Supplementary Figure 2. Variability in antibiotic prescription accordingly to the foci of infection in febrile patients attending Emergency departments in Rio de Janeiro, Brazil, October 2018-March 2019

Supplementary Figure 3. Foci of infection according to the height of temperature mensurated at arrival in febrile patients attending Emergency departments in Rio de Janeiro, Brazil, October 2018-March 2019

Supplementary Figure 4. Antibiotic class according to the height of temperature mensurated at arrival in febrile patients attending Emergency departments in Rio de Janeiro, Brazil, October 2018-March 2019

Table. Baseline characteristics according to the height of temperature mensurated at arrival in febrile patients attending emergency departments in Rio de Janeiro, Brazil, October 2018-March 2019

	Total febrile population N= 374	History of fever but temperature < 37.5 °C N= 248	History of fever and temperature ≥ 37.5 °C N= 115	p-value
Characteristics				
Demographics				
Age (years), mean (±SD)	31 (16)	32 (15)	28 (16)	0.02
Missing	3 (0.8)			
Male, n- (%)	169 (45.2)	103 (41.5)	62 (53.9)	0.02
Temp at arrival, (°C), mean (±SD)	37.08 (0.9)	36.55 (0.5)	38.24 (0.6)	<0.01
Missing	12 (3.2)			
Tmax (°C), mean (±SD)	38.7 (0.7)	38.7 (0.69)	38.95(0.9)	0.12
Missing	268 (71.7)			
Days of fever, mean (±SD)	1.58 (17.4)	1.45 (21.2)	1.80 (3.2)	0.86
Missing	81 (21.7)			
Degree of defervescence from Tmax, mean (±SD)	1.70 (1.13)	2.16 (0.86)	0.78 (1.04)	<0.01
Diagnosis and management				
Diagnostic testing – n (%)	76 (30.2)	42 (26.1)	32 (38.1)	0.05
Recent antibiotic use – n (%)	21 (12.2)	11 (9.6)	10 (18.9)	0.09
Missing	202 (54)			
Recent antipyretic use – n (%)	249 (66.6)	169 (87.1)	76 (90.5)	0.42
Missing	90 (24.1)			
Antibiotic prescription – n (%)	131 (53.0)	75 (47.8)	54 (65.1)	0.01

SD means standard deviation; Tmax higher temperature self-recorded at home before emergency department visit; Recent antibiotic use was defined as any antibiotic for systemic use started prior to emergency department visit. We excluded antibiotics for systemic use that could also be used for the treatment of tuberculosis in patients with active disease; Recent antipyretic was the use of any over-the counter antipyretic prior to emergency department visit.

Key messages

- Fever is a common reason to seek care in patients attending urban Emergency Departments in Rio de Janeiro, Brazil
- Suspicion of infection is the underlying cause of fever in 53% of the febrile subjects, and the respiratory source is the main focus of infection
- The use of antipyretics among febrile patients attending urban Emergency Departments in Rio de Janeiro is ubiquitous, and dipyrone is the antipyretic of choice in our population. However, such use did not influence overt fever presentation in Emergency Departments
- Our findings are essential for designing fever etiology and question the value of adopting a specific temperature cut-off for fever at enrollment, as in many fever studies. We believe that a history of fever as part of the inclusion criteria is a good surrogate of unwellness and might be incorporated systematically in future fever studies.

Appendix Table of Contents

This supplementary material has been provided by the authors to give readers additional information about their work.

Supplementary Tables


- Supplementary Table 1. Prediction model for the likelihood of overt fever at presentation in 374 febrile patients attending emergency departments in Rio de Janeiro, Brazil, October 2018 – March 2019.
- Supplementary Table 2. Diagnosis, febrile syndrome, and antibiotic prescription in febrile patients attending Emergency departments in Rio de Janeiro, Brazil, October 2018-March 2019
-

Supplementary Figures

- Supplementary Figure 1. Antibiotic class according to the foci of infection in febrile patients attending Emergency departments in Rio de Janeiro, Brazil, October 2018-March 2019
- Supplementary Figure 2. Variability in antibiotic prescription accordingly to the foci of infection in febrile patients attending Emergency Departments in Rio de Janeiro, Brazil, October 2018-March 2019
- Supplementary Figure 3. Foci of infection according to the height of temperature mensurated at arrival in febrile patients attending Emergency Departments in Rio de Janeiro, Brazil, October 2018-March 2019
- Supplementary Figure 4. Antibiotic class according to the height of temperature mensurated at arrival in febrile patients attending Emergency Departments in Rio de Janeiro, Brazil, October 2018-March 2019

Supplementary Figure 1. Antibiotic class according to the foci of infection in febrile patients attending Emergency departments in Rio de Janeiro, Brazil, October 2018-March 2019

Antibiotic class accordingly to the foci of infection in febrile patients attending ED in Rio de Janeiro, Brazil, October 2018 - March 2019



AUFI stands for acute undifferentiated febrile illness; LRTI lower respiratory tract infection; SSTI skin and soft tissue infection; URTI upper respiratory tract infection; and UTI urinary tract infection


Supplementary Figure 2. Variability in antibiotic prescription according to the foci of infection in febrile patients attending Emergency departments in Rio de Janeiro, Brazil, October 2018-March 2019

Variability in antibiotic prescription accordingly to the foci of infection in febrile patients attending ED in Rio de Janeiro, Brazil, October 2018 - March 2019




AUFI stands for acute undifferentiated febrile illness; LRTI lower respiratory tract infection; SSTI skin and soft tissue infection; URTI upper respiratory tract infection; and UTI urinary tract infection

Supplementary Figure 3. Foci of infection according to the height of temperature mensurated at arrival in febrile patients attending Emergency departments in Rio de Janeiro, Brazil, October 2018-March 2019



AUI stands for acute undifferentiated febrile illness; LRTI lower respiratory tract infection; SSTI skin and soft tissue infection; URTI upper respiratory tract infection; and UTI urinary tract infection

Supplementary Figure 4. Antibiotic class according to the height of temperature mensurated at arrival in febrile patients attending Emergency departments in Rio de Janeiro, Brazil, October 2018-March 2019



Supplementary Table 1. Prediction model for the likelihood of overt fever at presentation in 374 febrile patients attending emergency departments in Rio de Janeiro, Brazil, October 2018 – March 2019.

Variable		Adjusted Odds Ratio (95% CI)	p-value
Sex	Female	-	
	Male	2.25 (1.19-4.25)	0.01
Recent antipyretic use [§]	No	-	
	Yes	0.81 (0.34-1.97)	0.65
Age (years)		0.98 (0.95-1.00)	0.10
Time from fever onset (days)		1.00 (0.98-1.01)	0.97
Baseline Heart rate (beats/min)		1.050 (1.03-1.07)	< 0.05

§Non-prescription antipyretic uptake reported by the participant before emergency department care

Dependent variable: proportion of febrile subjects with measured axillary temperature ≥ 37.5 °C at presentation

Supplementary Table 2. Diagnosis, febrile syndrome, and antibiotic prescription in febrile patients attending Emergency Departments in Rio de Janeiro, Brazil, October 2018-March 2019

Presumed diagnosis by febrile syndrome	Number of presentations n/N (%)	Number of patients receiving an antibiotic prescription n/N (%)
Gastrointestinal		
Gastroenteritis	16/17 (94.11)	7/16 (43.75)
Gastritis	1/17 (5.88)	1/1 (100)
Upper respiratory tract infection		
Acute tonsillitis	34/84 (40.47)	34/34 (100)
Acute pharyngitis	22/84 (26.19)	5/22 (22.72)
Acute sinusitis	8/84 (9.52)	6/8 (75)
Common cold	5/84 (5.95)	0/5 (0)
Other upper respiratory tract infections	15/84 (17.85)	9/15 (60)
Lower respiratory tract infection		
Acute bronchitis and COPD exacerbations	3/17 (17.64)	2/3 (66.66)
Community-acquired pneumonia	9/17 (52.94)	8/9 (88.88)
Influenza	2/17 (11.76)	0/2 (0)
Other lower respiratory tract infections	3/17 (17.64)	3/3 (100)
Skin and soft tissue infection		
Cellulitis and abscesses	14/19 (73.68)	14/14 (100)
Other infection of the skin and soft tissue	4/19 (21.05)	4/4 (100)
Bacterial skin infection	1/19 (5.26)	1/1 (100)
Urinary tract infection		
Uncomplicated cystitis & pyelonephritis	28/29 (96.55)	25/28 (89.28)
Other urinary infection	1/29 (3.44)	1/1 (100)
Acute undifferentiated febrile illness		
Arboviruses	5/30 (16.66)	0/5 (0)
Unspecified febrile illness	6/30 (20)	4/6 (66.66)
Leptospirosis	2/30 (6.66)	2/2 (100)
Unspecified viral infection	15/30 (50)	1/15 (6.66)
Other	1/30 (3.33)	1/1 (100)

COPD stands for chronic obstructive pulmonary disease

