

1 The impact of antenatal balanced plate nutrition education for pregnant women on
2 birth weight: a cluster randomized controlled trial in rural Bangladesh

3 Authors:

4 Morseda Chowdhury^{1, 2}, Camille Raynes-Greenow², Patrick J. Kelly², Ashraful Alam²,
5 Kaosar Afsana³, Michael J. Dibley²

6 Affiliations:

7 ¹Health Nutrition and Population Programme, BRAC, BRAC Centre, 75 Mohakhali, Dhaka
8 1212, Bangladesh

9 ²The University of Sydney, Faculty of Medicine and Health, Sydney School of Public Health,
10 Camperdown, New South Wales 2006, Australia

11 ³James P Grant School of Public Health (JPGSPH), BRAC University, Dhaka, Bangladesh

12 Corresponding author: Michael J Dibley, Room 328 Edward Ford Building (A27) Sydney
13 School of Public Health, the University of Sydney, NSW 2006, Australia.

14 Phone: +61 2 9351 520 Email: michael.dibley@sydney.edu.au

15 Short running head: Balanced plate nutrition education & birth weight

16 Trial registration: The Australian New Zealand Clinical Trials Registry number

17 ACTRN12616000080426

18

19 Abstract

20 **Background:** Low birth weight (LBW) is a global public health problem with the highest
21 prevalence in South Asia. It is strongly associated with maternal under nutrition. In South
22 Asia, intra household food distribution is inequitable with lower dietary adequacy in women,
23 regardless of pregnancy. Evidence that nutrition education improves diet during pregnancy
24 and reduces LBW is weak.

25 **Objective:** We aimed to assess the impact of a nutrition education intervention for pregnant
26 women on infant birth weight in a low-income population.

27 **Design:** We conducted a parallel, two arm, cluster-randomized controlled trial in rural
28 Bangladesh, with 36 clusters randomized to receive the intervention (n=445 participants) or
29 standard care (n=448). Eligible participants were ≤ 12 weeks gestation. Intervention
30 participants received education about eating balanced meals to meet daily dietary
31 requirements with the right proportions of food groups (a balanced plate) and practical
32 demonstration from their first trimester until delivery. Primary outcomes were mean birth
33 weight and the incidence of LBW. We included all participants in analyses.

34 **Results:** The mean birth weight difference between the intervention and control groups was
35 127.5 g (95% confidence interval (CI): 11.1, 243.9; $p=0.032$), and the intervention reduced
36 the risk of LBW by 57% (relative risk (RR): 0.43; 95% CI: 0.25, 0.75; $p=0.003$). Post hoc
37 analyses showed a higher birth weight and a greater reduction in LBW amongst adolescent
38 mothers (mean difference: 299.1 g; 95% CI: 101.6, 496.6; $p=0.003$ and RR: 0.28; 95% CI:
39 0.11 to 0.71; $p=0.007$). The mean number of food groups consumed was significantly higher
40 from the third month of pregnancy onwards in the intervention compared to the control group.

41 **Conclusions:** A community-based balanced plate nutrition education intervention effectively
 42 increased mean birth weight and reduced the proportion of infants with LBW in a rural
 43 population of Bangladesh.

44 **Keywords:** Low birth weight, maternal nutrition education, food demonstrations, dietary
 45 diversity, cluster RCT, Bangladesh

46

47

48 Introduction

49 Birth weight is a powerful predictor of infant survival.¹ Low birth weight, defined as less than
50 2500 grams (g), is a composite measure of restricted fetal growth and short duration of
51 gestation, major determinants of infant mortality, childhood growth and development, and
52 adulthood well-being.^{1,2} The lower the birth weight, the higher the risk of mortality.^{3,4}
53 Globally, in 2010, there were an estimated 18 million LBW babies, of whom 59% were term,
54 but small for gestational age and 41% were preterm.⁵ An estimated 606,500 neonatal deaths
55 occur annually in low and middle-income countries due to small for gestational age, which
56 accounts for 22% of neonatal deaths.⁶ South Asia has the highest prevalence of LBW
57 globally (34%).⁶ Bangladesh is ranked fourth in the global burden of LBW.⁵
58 LBW is strongly associated with maternal under-nutrition⁷, which induces suboptimal
59 placental growth, thereby influencing nutrient delivery, altering fetal hormones, and
60 contributing to restricted development.⁸ A World Health Organization study with 111,000
61 women reported that mothers in the lowest quartile of pre-pregnancy weight carried an
62 elevated risk of IUGR of 2.55 (95% confidence interval (CI): 2.3, 2.7) and LBW of 2.38
63 (95% CI: 2.1, 2.5) compared to the upper quartile.⁹ In India, the odds ratio for LBW was
64 three times greater in severe energy-deficient low BMI groups than normal BMI.¹⁰
65 Consistent with the mechanisms and epidemiological evidence is the high prevalence of
66 LBW in rural Bangladeshi women who have a chronic shortfall of food energy throughout
67 pregnancy.¹¹ In South Asia, intra-household food distribution is inequitable with lower
68 energy share, animal source food, and dietary adequacy in women, regardless of pregnancy.¹²
69 There is little data on nutrition education's effects on birth weight in low or middle-income
70 countries, although four meta-analyses have assessed impacts. The first in 1993 found
71 nutrition education significantly increased the consumption of protein and vitamins, and birth

weight of infants by 300 g. However, the studies lacked statistical power or did not report estimates of effect size.¹³ The second meta-analysis in 2012 found a 105 g (95% CI: 18, 193 g) increase in mean birth weight with nutrition education and counseling in pregnancy. When stratified by the study site, the effect failed to reach significance in low and middle-income countries (152 g; 95% CI: -81, 384 g).¹⁴ The third meta-analysis in 2014 reported that dietary counselling significantly increased birth weight in three studies (149 g; $p=0.039$), but two had a small sample size, and another had methodological limitations.¹⁵ The fourth, a Cochrane review,¹⁶ identified two small trials that reported significantly higher birth weight in the intervention group (mean difference 489.8 g, 95% CI: 427.9, 551.6 g). There are no reports of trials of antenatal nutrition education in rural Bangladesh.

Thus, the evidence for nutrition education to improve diet and maternal and neonatal health indicators, including birth weight, remains weak. Most evidence comes from trials with design or analysis limitations¹⁴ and high-income countries where counselling is by professional dietitians or nutritionists.¹⁵ There is a critical knowledge gap about antenatal nutrition education's efficacy and its potential to reduce LBW in South Asia, including Bangladesh. Filling this gap can potentially contribute to the World Health Assembly's global goal of a 30% reduction in the LBW rate by 2025.¹⁷ Therefore, we investigated the impact of antenatal nutrition education with demonstrations of a nutritionally energy-protein balanced diet on infants' birth weight in a low-income population.

Subjects and Methods

Study design and participants

We conducted a two-arm, parallel, cluster randomized controlled trial (cRCT). The James P. Grant School of Public Health; BRAC University Ethical Review Committee; Dhaka; Bangladesh approved the study protocol (No 53, 25th May 2015).

96 We have described the study design.¹⁸ In brief, the study was in the rural district of Sherpur,
 97 which has a low human development index, a high percentage in the lowest economic
 98 quintile (37.1%) compared to the national average (21.7%)¹⁹ and 48% below the poverty
 99 line.²⁰ The Food Security Atlas of Bangladesh classifies Sherpur as having a high/very high
 100 household food insecurity level.²¹ In 2016 in Sherpur the neonatal mortality (41/1000 live
 101 births) and under-five mortality (59/1000 live births) were higher than the national average
 102 (30 and 47/1000 respectively).²²

103 Sherpur has five sub-districts: Jhenaigati (population 160,452), Nakla (189,685), Nalitabari
 104 (251,361), Sherpur Sadar (497,179) and Sreebardi (259,648). Pregnant women, self-reporting
 105 to a hospital or formal practitioner (doctor, nurse, or Family Welfare Visitor), receive
 106 nutrition education as part of antenatal care (ANC). BRAC (a national NGO; formerly known
 107 as Bangladesh Rural Advancement Committee) implements a community-based Maternal,
 108 Neonatal, and Child Health (MNCH) program characterized by prospective pregnancy
 109 surveillance and basic ANC led by community health workers (CHW) known as *Shasthya*
 110 *Kormi*. One *Shasthya Kormi* serves an area of ~10,000 people, along with ten health
 111 volunteers (*Shasthya Shebika*). The *Shasthya Kormi* are village women who have basic 10-
 112 years schooling and have received training on health, nutrition, and sanitation issues for six
 113 weeks, followed by bi-monthly refreshers.

114 In our study, a cluster was the catchment area of one *Shasthya Kormi*, which provided
 115 logistical convenience and minimized intervention spills into the control group. *Shasthya*
 116 *Kormi*, with at least six months' experience in MNCH, were eligible. Participants were
 117 pregnant women 15-49 years, gestation 12 weeks or less, permanently residing in the area,
 118 and not expecting to move from the usual residence until birth. We used the established
 119 BRAC pregnancy detection system using monthly menstrual surveillance to identify new
 120 pregnancies. Women with delayed menstruation for six weeks or more underwent a urine test.

We excluded pregnant women with self-reported chronic disease. All women excluded from the study continued to receive standard care and counselling as in the control clusters. A *Shasthya Kormi* visited each confirmed pregnant woman, screened for eligibility, and obtained verbal informed consent. Recruitment took place from 1st October- 31st December 2016.

Randomization and masking

We used proportionate stratified sampling to randomly select 36 of 135 *Shasthya Kormi* clusters (4, 5, 7, 13, and 7 from the five sub-districts). The clusters were dispersed across the sub-districts and sufficiently separated to minimize the risk of contamination. We selected 36 as the maximum number of *Shasthya Kormi*, who could be overseen by a single BRAC supervisor, and we randomized with 1:1 allocation. All eligible pregnant women served by one *Shasthya Kormi* received the intervention or standard care as per the random assignment. We could not mask families or the research team to group allocation; however, we blinded researchers assessing primary and secondary outcomes to the study hypotheses. We also used an objective outcome (birth weight), which is less prone to ascertainment bias.

Procedures

Intervention *Shasthya Kormi* received two days of nutrition training, including a balanced diet and counselling techniques. They also practiced community counselling skills under supervision and received one-day training on taking consent and data collection. Enrolled participants were visited monthly to provide ANC that included standard nutrition advice. In intervention clusters, balanced plate intervention replaced the standard nutrition advice using a culturally appropriate diet for Bangladeshi pregnant women.²³ This contained seven-food groups (cereal, lentil, animal protein, vegetables, fruits, milk, and oil), giving a daily 2500 kcal energy and included essential micro and macronutrients. The balanced plate was a

combination of foods in appropriate portion sizes to meet requirements. Before the trial, we conducted focus groups to identify locally preferred foods and dietary restrictions. We pilot tested the menu to assess community acceptability and the feasibility of incorporating it into ANC services.

We delivered the intervention individually. Initially, participants received information on a balanced diet and benefits for maternal and fetal health. Then, they learned to prepare a “balanced plate” by demonstrating the right proportions of food groups using the prescribed menu.¹⁸ The community health worker displayed different combinations of food on the plate, with appropriate alternatives, such as replacing meat (an expensive protein source) with cheaper eggs or farm fishes. We emphasized foods commonly missed in the regular diet, such as milk, fruits, and coloured vegetables. Women were encouraged to increase their frequency of food intake to three meals and two snacks per day. To demonstrate measurements, we used a bowl of 250 ml commonly found in most homes. The community health worker invited family decision-makers to observe the demonstration and assist in making a balanced plate.

Delivering the intervention took 45 minutes: 10 for individual counselling, 10 for household counselling, and 25 for demonstration. To help recall messages, we provided a pictorial food chart and a written menu¹⁸. At the first visit, the community health worker delivered eight nutrition messages with a practical demonstration. In subsequent visits, they emphasized messages that were less familiar or difficult to remember. In the counselling, they excluded messages that were easy to recall, such as drinking eight glasses of water or cooking with oil. They also dropped the messages if it was clear the women practiced the behaviour. As the number of messages decreased, the community health worker reduced the visit time to 25 minutes. The mean number of intervention visits was 3.4.

Home visits continued until birth or the end of the pregnancy; ANC from other sources was not discouraged. Husbands were encouraged to re-allocate family budgets to purchase

nutritious food and mothers-in-law to re-distribute family food to increase their daughter-in-law's share.

In the control group, pregnant women received the same frequency of home and standard nutrition advice except for the balanced plate demonstrations. Standard advice included recommendations to consume all food groups and take iron-folic acid and calcium supplements.

Shasthya Kormi collected demographic, socio-economic, and reproductive data at enrolment plus birth weight measurements within 72 hours of birth. Families notified birth or pregnancy loss (miscarriage, abortion, and stillbirth) immediately to their nearby *Shasthya Shebika*.

One field supervisor oversaw the 36 *Shasthya Kormi*, recorded field activities, and conducted follow-up interviews with 5% of participants to confirm visits and measure protocol adherence.

Outcomes

Primary outcomes were the mean birth weight of live-born infants and the proportion of newborns with birth weight less than 2500 g. The *Shasthya Kormi* had many years of experience measuring birth weight, but we retrained them with their standard equipment, a 5 kg Salter spring balance scale with an accuracy of 100 g. They calibrated the scales with a standard weight every week. We also checked the scales monthly during refresher training. For mothers who birthed in health facilities, we recorded the birth weight measured at the facility.

For secondary outcomes, the *Shasthya Kormi* used structured forms to record fetal loss during pregnancy, birth outcome, delivery information, and neonatal or maternal death. They also used semi-structured questionnaires to collect dietary data, with the food list adapted from Food and Nutrition Technical Assistance II Project (FANTA-2).²⁴ These questionnaires

captured information about foods, beverages, and supplements consumed 24 hours before the interview. From this data we formed ten food groups as follows: (1) grains, white roots and tubers; (2) pulses (beans, peas and lentils); (3) nuts and seeds; (4) dairy; (5) meat, poultry and fish; (6) eggs; (7) dark green leafy vegetables; (8) other vitamin A-rich fruits and vegetables; (9) other vegetables; and (10) other fruits. *Shasthya Kormi* collected dietary data monthly, and we calculated the mean number of food groups consumed to assess diet quality.²⁵ We defined animal source protein as consumption of any combination of dairy, meat, poultry, fish or egg food groups.

Statistical analysis

We required 720 pregnant women to achieve 80% power to detect a difference of 100 g in birth weight between groups. We assumed a mean birth weight of 2531 g (standard deviation (SD) 415 g) in the control group based on the findings from a similar trial in Bangladesh.²⁶ We set a significance level of 0.05 (two-sided), and assumed an intra-cluster correlation coefficient (ICC) of 0.03²⁷, and 36 clusters with an average of 10-15 live births per cluster per month. To adjust for 10% pregnancy loss and 15% loss to follow-up based on similar trials,^{26,28} we inflated the required sample size to 900 pregnant women.

We analysed the data with Stata version 15 and conducted intention to treat analyses. We assigned the intervention randomly at the cluster level, but assessed outcomes at the individual level. We estimated the mean between-group difference in birth weight using linear regression and the relative risk for LBW using binomial regression. For both, we included the study group as a covariate and adjusted for cluster randomization using Generalised Estimating Equations (GEE).²⁹ Similarly, we estimated the mean number of food groups consumed using GEE linear regression with the treatment group as a covariate and adjusting for baseline imbalances by including covariates.

We conducted post hoc subgroup analyses for age, education, family income, and place of delivery. We added an interaction term to assess whether there was a difference in intervention effect for any sub-group, using a significance level of 0.01.

Role of the funding source

The funders had no role in study design, participant selection, data collection, and data analyses. The corresponding author had full access to the study's data and final responsibility for the decision to publish.

Results

We identified 2,154 newly pregnant women from August 2015 to February 2016 in the selected clusters. Of these women, 937 did not meet the eligibility criteria (603 had a gestation of >12 weeks, 147 were not permanent residents, and 187 had planned to move out of the study area for childbirth) and 324 declined to participate. We recruited 893 participants, 445 in 18 intervention, and 448 in 18 control clusters, and we lost 44 participants during follow-up, 25 (5.6%) intervention, and 19 (4.2%) control. Birth weight was missing for 22 participants (Figure 1). There were 42 pregnancy losses (abortion or stillbirth), and six neonatal deaths within 24 hours of birth.

There were no important differences in baseline characteristics between the groups (Table 1). There were small differences for mother's age, mother's and father's education, and parity but not enough to warrant an adjustment in analyses. Slightly more intervention participants received four or more ANC visits, but for antenatal care from health centres, this trend reversed (Table 1). When we combined both ANC data sources, the intervention participants received slightly more ANC (intervention 61.7% vs. control 55.6%). There was a slightly higher percentage of intervention participants delivered in a facility (20.7% vs. 16.6%) but

this variable reflects the small differences in ANC. Mean birth weight was higher in facility deliveries (2,915 g, SD 560 g) than in-home births (2,769 g, SD: 410 g).

Supervisors attended 5% of balanced plate demonstrations to observe the appropriateness of message delivery, the use of the menu and food chart, demonstration, problem-solving, and counselling. There were approximately 6-7 of 8 messages delivered in 80% of the participants. The time taken for the demonstrations gradually decreased with increasing numbers of visits. Counselling family members were largely dependent on availability; husbands were present at 15-20% of visits, other family members at >80% of visits.

We found a statistically significant difference in mean birth weight and LBW (Tables 2 and 3). The mean difference in birth weight between groups was 127.5 g (95% CI: 11.1, 243.9; $p=0.032$) adjusted for clustering. The estimated relative risk (RR) of LBW was 57% lower in the intervention group (RR: 0.43; 95% CI: 0.25, 0.75; $p=0.003$) adjusted for clustering.

We conducted subgroup analyses of mother's age, education, and family income (Tables 2 and 3) and found a significant intervention effect on birth weight in adolescents vs. non-adolescents. Mean difference in birth weight between groups was greater in adolescents (299.1 g; 95% CI: 101.6, 496.6; $p=0.003$; vs 95.6 g; 95% CI: -17.3, 208.4; $p=0.097$) (p-value for interaction 0.009) as was reduction in LBW (RR: 0.28; 95% CI: 0.11, 0.71; $p=0.007$ vs RR: 0.54; 95% CI: 0.29, 0.98; $p=0.044$), although this was not statistically significant (p-value for interaction 0.22). There was no evidence of treatment modification by the mother's education or family income (Tables 2 and 3).

To assess bias in ascertaining birth weight, we stratified by participants birthed in a facility and whose babies were weighed by independent observers and those who birthed at home and found no evidence of treatment modification ($p=0.78$).

After adjustment for ANC, the use of iron and folic acid, and clustering, the reduction in mean birth weight and LBW was smaller in the intervention group (Supplemental Tables S1 and S2). The adjusted difference was 121.8 g (95% CI: 11.7, 231.9; $p=0.030$), which is 5.7 g less than the unadjusted analysis (Table 2). The adjusted relative risk of LBW was 0.53 (95% CI: 0.29, 0.98; $p=0.042$), which was smaller than the unadjusted relative risk in Table 3.

The mean number of food groups consumed increased in the intervention group and was significantly higher from the third month of pregnancy. The largest difference was in the eighth month of pregnancy when intervention participants consumed 5.6 food groups/day (95% CI: 5.19, 6.14) compared to 3.8 (95% CI: 3.40, 4.12) in control participants (Fig 2). Starting from the fifth month of pregnancy, the intervention group's mean number of food groups exceeded five (the number for adequate dietary diversity²⁵). The mean number of animal-source protein foods consumed was significantly higher from the fourth month of pregnancy (Figure 2).

Discussion

We demonstrated that antenatal nutrition education with practical demonstrations of preparing a balanced diet increases infant birth weight and reduces LBW in rural Bangladesh. There was a small birth weight reduction after adjusting for differences in antenatal care and iron/folic acid use. The effect of the intervention was greater in adolescent mothers. The findings provide strong evidence of the effectiveness of a sustainable intervention suitable for implementation in community-based health care systems. They demonstrate the potential impact of nutrition education to improve intrahousehold food distribution as a tool to reduce low birth weight in Bangladesh and across South Asia.

The process evaluation of the trial³⁰ provided insights into why the intervention was so impactful. It identified key elements in the intervention strategy, such as practical

demonstration of portion sizes; addressing local food perceptions; demystifying animal-source foods; engaging husbands and mothers-in-law; leveraging women's social networks; and harnessing community health workers' social role, that played a crucial role in achieving the desired adherence. In particular, the approach of involving the mothers-in-law and husbands changed their attitudes towards a fairer allocation of foods to pregnant women. The husbands adhered to the messages and purchased more and nutritious foods for their wives by adjusting the shopping budget and even working extra hours.

Comparison with other studies

Our intervention appears more effective than interventions using micronutrient or food supplements. A large cluster RCT (22,405 pregnancies) in Bangladesh assessed the impact of antenatal micronutrient supplementation and found a 54 g (95% CI: 41, 66) increase in birth weight and a 12% reduction (RR: 0.88; 95% CI: 0.85, 0.91) in LBW.²⁶ Another cluster RCT of lipid-based micronutrient supplementation showed a 41 g birth weight increment without a significant effect on LBW.²⁸

Providing micronutrient supplements even with additional food energy is unlikely to address the fundamental problems of insufficient food intake in pregnancy related to food taboos and inequitable intra-household food distribution. The earlier MINIMat RCT (3267 singleton newborns)³¹, which tested early and late starts to food supplements with micronutrients in pregnancy, found no significant main-effects on birth weight. However, the combination of multiple micronutrients and early food supplementation did improve infant survival. The lack of a control group without food supplements might explain the lack of birth weight response. In contrast, an observational study in Bangladesh reported a 118 g increase in birth weight with daily 608 kcal food supplements combined with iron-folic acid sustained for more than four months.³² Although this evidence is weak, it supports using food-based solutions to address nutrition problems.

Adolescent pregnancy has an increased risk of adverse birth outcomes, including LBW.³³ If severely undernourished, the growing pregnant adolescent and foetus compete for nutrients.³⁴ A recent study from rural Bangladesh found a cessation of linear growth, weight loss, and fat and lean body mass depletion in pregnant adolescents.³⁴ This greater nutritional susceptibility helps explain the greater effect of the balanced plate education in this group.

The balanced-plate intervention significantly increased maternal dietary diversity (Fig 2). Change in dietary patterns is the likely cause of the higher birth weight in the intervention group. A recent cohort study³⁵ followed 374 women from their first antenatal care visit to assess associations between dietary diversity during pregnancy and LBW in rural Ethiopia. The LBW risk doubled with a dietary diversity score of less than four food groups (ARR: 2.06; 95% CI: 1.03, 4.11). A facility-based case-control study³⁶ from rural Ethiopia, which examined iron, folic acid and nutrition counselling, snack consumption, dietary diversity, and maternal undernutrition, also reported similar findings.

There was a significant six-fold increase in the adjusted odds of LBW with an inadequate minimum dietary diversity score (AOR: 6.65; 95% CI, 2.31, 19.16). A cross-sectional study³⁷ from Ghana also reported a protective effect of higher dietary diversity scores for LBW (AOR: 0.10; 95% CI: 0.04, 0.13 per standard deviation change score, $p < 0.0001$). This evidence supports our findings that improved and more diverse dietary intake leads to a reduction in LBW.

Participants in the intervention group consumed more animal-sourced protein foods (Fig 2), reaching a peak at eight months of pregnancy of two animal-sourced foods/day. A large cross-sectional study of pregnant women in Shaanxi province, northwestern China, reported an average weekly consumption of five animal-sourced foods.³⁸ As the frequency of animal-sourced food increased by one time/week, the average birth weight increased by 3.24 g (95%

CI: 1.09, 5.39). Unlike increasing animal-sourced foods in the diet, the consumption of excess protein supplements (>20% of energy as protein) can harm fetal growth.³⁹

Despite earlier evidence of a positive impact of nutrition education on pregnancy outcomes, the delivery method has received little attention. It is particularly important in low literate and poor self-efficacy populations, such as in rural Bangladesh, where women require dietary counselling to be accessible, feasible, and sanctioned by the community. A review of over 300 studies found that nutrition education is more likely to be effective when focusing on behaviour and action than knowledge alone. And it is also likely to be more effective if grounded in theory.⁴⁰ Fishbein and Yzer note that “any given behaviour is most likely to occur; if one has a strong intention to perform the behaviour; if a person has the necessary skills and abilities required to perform the behaviour; and if no environmental constraints are preventing the behavioural performance.”⁴¹ To positively impact dietary behaviour, we designed our intervention to embrace all aspects of this theory. We adopted an innovative counselling technique, including a practical demonstration of a balanced diet and engaged family decision-makers. Similar nutrition counselling intervention was found effective in increasing age-specific complementary feeding practices in rural Bangladesh.⁴²

In Bangladesh, the main source of nutrition education for pregnant women is through ANC, but the contact time is insufficient to affect nutrition change.⁴³ Thus, WHO. Proposes upskilling CHWs in delivering dietary advice is an appropriate approach to complement health services.⁴⁴ In our study, we integrated the balanced plate intervention into BRAC’s existing ANC package delivered by CHWs. The same staff in a recent intervention demonstrated their successful counselling on breastfeeding and complementary feeding.⁴⁵ Previously, we also found that CHWs can positively influence family decision-makers to increase pregnant women’s iron-folic acid consumption.⁴⁶

Strengths and limitations of this study

The evidence presented from this trial is robust. Key strengths are the cluster, randomized controlled trial design with sufficient statistical power to detect clinically important changes in outcomes. The delivery of the trial treatments through an established home-based antenatal care service led to a balance between the treatment groups in the level of antenatal contacts with community health workers. The unblinded intervention was counterbalanced by objective outcomes and blinding participants and data collectors to the aims and hypotheses. The trial had a very high follow-up (93%). We used intention-to-treat analyses and adjusted for unbalanced baseline factors and clustering. Our intervention, based on well-designed qualitative formative research, was logistically simple and culturally appropriate. The key to our success was conducting the trial within existing ANC frameworks. The delivery strategy resonates with maternal and child nutrition recommendations to achieve adequate coverage of nutrition-specific interventions by reaching populations in need.⁴⁷ However, we acknowledge some limitations. First, the *Shasthya Kormi*, who delivered the intervention, also collected birth weight data. Measuring birth weight and providing nutrition counselling are routine tasks for these staff. By blinding the SK to study outcome, we would expect non-differential bias. Also, we found no difference in the effects between women who had birth weight measured by hospital staff versus those who birthed at home with birth weight measured by the SK. Secondly, using catchment areas as clusters might introduce contamination, with the possibility of *Shasthya Kormi* exchanging information. However, this would have biased results towards a null effect. To prevent this, we trained intervention and control group *Shasthya Kormi* separately at regular refresher sessions. Thirdly, we could not quantify food intake and intra-household food distribution, which might have provided further evidence of pathways to the impact we observed. Finally, we excluded over 30% of pregnant women identified mainly because they were already in the second or third trimester when we detected their pregnancy. There is no information about the characteristics of these excluded

women and thus no way to compare them to the women included in the trial. This limits our assessments of the external validity of the trial findings.

Conclusions and policy implications

Our results indicate that a balanced plate nutrition education during pregnancy increases birth weight and reduces LBW. Our trial offers a pragmatic food-based strategy for resource-poor settings. The CHW-led intervention is simple, feasible, and easy to implement in a country with a shortage of human health resources. The intervention is also scalable through existing public and private community health infrastructure.

396 **References**

- 397 1. Black RE, Allen LH, Bhutta ZA, et al. Maternal and child undernutrition: global and
398 regional exposures and health consequences. *Lancet*. 2008;371:243-260.
- 399 2. Cesar G Victora, Linda Adair, Caroline Fall, et al. Maternal and child undernutrition:
400 consequences for adult health and human capital. *Lancet*. 2008;371:340-357.
- 401 3. Wilcox AJ, Russell IT. Birthweight and Perinatal Mortality: II. On Weight-Specific
402 Mortality. *International Journal of Epidemiology*. 1983;12(3):319-325.
- 403 4. Watkins WJ, Kotecha SJ, Kotecha S. All-cause mortality of low birthweight infants in
404 infancy, childhood, and adolescence: population study of England and Wales. *PLoS*
405 *medicine*. 2016;13(5):e1002018.
- 406 5. Lee AC, Katz J, Blencowe H, et al. National and regional estimates of term and
407 preterm babies born small for gestational age in 138 low-income and middle-income
408 countries in 2010. *Lancet Glob Health* 2013;1:e26-36.
- 409 6. Lee AC, Kozuki N, Cousens S, et al. Estimates of burden and consequences of infants
410 born small for gestational age in low and middle income countries with
411 INTERGROWTH-21st standard: analysis of CHERG datasets. *BMJ*. 2017;358(j3677).
- 412 7. Han Z, Mulla S, Beyene J, Liao G, McDonald SD. Maternal underweight and the risk
413 of preterm birth and low birth weight: a systematic review and meta-analyses.
414 *International journal of epidemiology*. 2010;40(1):65-101.
- 415 8. Triunfo S, Lanzone A. Impact of maternal under nutrition on obstetric outcomes.
416 *Journal of endocrinological investigation*. 2015;38(1):31-38.
- 417 9. World Health Organization. *Maternal anthropometry and pregnancy outcomes: a*
418 *WHO collaborative study*. World Health Organization; 1995.
- 419 10. Naidu A, Rao N. Body mass index: a measure of the nutritional status in Indian
420 populations. *European journal of clinical nutrition*. 1994;48:S131-140.
- 421 11. Islam MZ, Akhtaruzzaman M, Lamberg-Allardt C. Nutritional status of women in
422 Bangladesh: comparison of energy intake and nutritional status of a low income rural
423 group with a high income urban group. *Asia Pacific Journal of Clinical Nutrition*.
424 2004;13(1):61-68.
- 425 12. Harris-Fry H, Shrestha N, Costello A, Saville NM. Determinants of intra-household
426 food allocation between adults in South Asia – a systematic review. *International*
427 *Journal for Equity in Health*. 2017;16(107).
- 428 13. Boyd NR, Jr., Windsor RA. A meta-evaluation of nutrition education intervention
429 research among pregnant women. *Health education quarterly*. 1993;20(3):327-345.
- 430 14. Girard AW, Olude O. Nutrition education and counselling provided during
431 pregnancy: effects on maternal, neonatal and child health outcomes. *Paediatric and*
432 *Perinatal Epidemiology*. 2012;26(s1):191-204.
- 433 15. Gresham E, Collins CE, Mishra GD, Byles JE, Hure AJ. Diet quality before or during
434 pregnancy and the relationship with pregnancy and birth outcomes: the Australian
435 Longitudinal Study on Women's Health. *Public Health Nutrition*. 2016;19(16):2975–
436 2983.
- 437 16. Ota E, Hori H, Mori R, Tobe-Gai R, Farrar D. Antenatal dietary education and
438 supplementation to increase energy and protein intake. *Cochrane Database of*
439 *Systematic Reviews*. 2015(6).
- 440 17. Lopes KdS, Ota E, Shakya P, et al. Effects of nutrition interventions during pregnancy
441 on low birth weight: an overview of systematic reviews. *BMJ Global Health*.
442 2017;2(e000389).

18. Chowdhury M, Raynes-Greenow C, Alam A, Dibley M. Making a balanced plate for pregnant women to improve birthweight of infants: a study protocol for a cluster randomised controlled trial in rural Bangladesh. *BMJ Open*. 2017;7(e015393).
19. Sen B, Ali Z. Spatial Inequality in Social Progress in Bangladesh. *The Bangladesh Development Studies*. 2009;XXXII(2).
20. *Poverty Maps of Bangladesh 2010*. World Bank Bangladesh, Bangladesh Bureau of Statistics, and World Food Programme 2011.
21. *The Food Security Atlas of Bangladesh: Towards a poverty and hunger free Bangladesh*. Planning Commission, Bangladesh Bureau of Statistics and UN World Food Programme. 2004.
22. National Institute of Population Research and Training (NIPORT), International Centre for Diarrhoeal Disease Research Bangladesh (icddr), MEASURE Evaluation. *Bangladesh District Level Socio-demographic and Health Care Utilization Indicators*. Dhaka, Bangladesh, and Chapel Hill, NC, USA: NIPORT, icddr, and MEASURE Evaluation; 2019.
23. Nahar Q, Choudhury S, Faruque MO, Sultana SSS, Siddiquee MA. *Desirable Dietary Pattern for Bangladesh*. National Food Policy Capacity Strengthening Programme; 2013.
24. Arimond M, Torheim LE, Wiesmann D, Joseph M, Carriquiry A. Dietary diversity as a measure of the micronutrient adequacy of women's diets: results from rural Bangladesh site. *Washington, DC: Food and Nutrition Technical Assistance II Project (FANTA-2), FHI*. 2009;360.
25. FAO and FHI. Minimum dietary diversity for women: a guide for measurement. *Rome: FAO*. 2016.
26. West Jr PK, Shamim AA, Mehra S, et al. Effect of Maternal Multiple Micronutrient vs Iron-Folic Acid Supplementation on Infant Mortality and Adverse Birth Outcomes in Rural Bangladesh: The JiVitA-3 Randomized Trial. *JAMA*. 2014;312(24):2649-2658.
27. Zeng L, Cheng Y, Dang S, et al. Impact of micronutrient supplementation during pregnancy on birth weight, duration of gestation, and perinatal mortality in rural western China: double blind cluster randomised controlled trial. *BMJ*. 2008;337(7680):1211 - 1215.
28. Mridha MK, Matias SL, Chaparro CM, et al. Lipid-based nutrient supplements for pregnant women reduce newborn stunting in a cluster-randomized controlled effectiveness trial in Bangladesh. *American Society for Nutrition*. 2016;103:236-249.
29. Rabe-Hesketh S, Skrondal A. *Multilevel and longitudinal modeling using Stata*. STATA press; 2008.
30. Alam A, Chowdhury M, Dibley MJ, Raynes-Greenow C. How can we improve the consumption of a nutritionally balanced maternal diet in rural bangladesh? The key elements of the "balanced plate" intervention. *International Journal of Environmental Research and Public Health*. 2020;17(17):6289.
31. Persson LÅ, Arifeen S, Ekström E-C, Rasmussen KM, Frongillo EA, Team MS. Effects of prenatal micronutrient and early food supplementation on maternal hemoglobin, birth weight, and infant mortality among children in Bangladesh: the MINIMat randomized trial. *Jama*. 2012;307(19):2050-2059.
32. Shaheen R, Francisco Ad, Arifeen SE, Ekström E-C, Persson LÅ. Effect of prenatal food supplementation on birth weight: an observational study from Bangladesh1-. *Am J Clin Nutr*. 2006;83:1355–1361.

33. King JC. The risk of maternal nutritional depletion and poor outcomes increases in early or closely spaced pregnancies. *The Journal of nutrition*. 2003;133(5):1732S-1736S.
34. Rah JH, Christian P, Shamim AA, Arju UT, Labrique AB, Rashid M. Pregnancy and lactation hinder growth and nutritional status of adolescent girls in rural Bangladesh. *The Journal of nutrition*. 2008;138(8):1505-1511.
35. Zerfu TA, Umata M, Baye K. Dietary diversity during pregnancy is associated with reduced risk of maternal anemia, preterm delivery, and low birth weight in a prospective cohort study in rural Ethiopia. *The American journal of clinical nutrition*. 2016;103(6):1482-1488.
36. Girma S, Fikadu T, Agdew E, et al. Factors associated with low birthweight among newborns delivered at public health facilities of Nekemte town, West Ethiopia: a case control study. *BMC pregnancy and childbirth*. 2019;19(1):220.
37. Abubakari A, Jahn A. Maternal dietary patterns and practices and birth weight in Northern Ghana. *PloS one*. 2016;11(9):e0162285.
38. Yan H, Dang S, Mi B, et al. Study on relationship between mother's animal sourced food intake during pregnancy and neonate birth weight. *Zhonghua liu xing bing xue za zhi= Zhonghua liuxingbingxue zazhi*. 2017;38(5):615-620.
39. Liberato SC, Singh G, Mulholland K. Effects of protein energy supplementation during pregnancy on fetal growth: a review of the literature focusing on contextual factors. *Food & nutrition research*. 2013;57(1):20499.
40. Contento IR. Nutrition education: linking research, theory, and practice. *Asia Pac J Clin Nutr*. 2008;17(1):176-179.
41. Fishbein M, Yzer MC. Using Theory to Design Effective Health Behavior Interventions. *Communication Theory*. 2003;13(2):164-183.
42. Sanghvi T, Haque R, Roy S, et al. Achieving behaviour change at scale: Alive & Thrive's infant and young child feeding programme in Bangladesh. *Maternal & Child Nutrition*. 2016;12(Suppl. 1):141-154.
43. Ahmed SM, Hossain MA, Chowdhury AMR, Bhuiya AU. The health workforce crisis in Bangladesh: shortage, inappropriate skill-mix and inequitable distribution. *Human Resources for Health*. 2011;9(3).
44. Lehmann U, Sanders D. *Community health workers: what do we know about them? The state of the evidence on programmes, activities, costs and impact on health outcomes of using community health workers*. Geneva: World Health Organization;2007.
45. Alive & Thrive. *Alive & Thrive's approach and results in Bangladesh: Rapid improvements in infant and young child feeding practices resulting from a large-scale program*. Dhaka, Bangladesh: Alive & Thrive;2015.
46. Alam A, Rasheed S, Khan NU, et al. How can formative research inform the design of an iron-folic acid supplementation intervention starting in first trimester of pregnancy in Bangladesh? *BMC Public Health*. 2015;15(374).
47. Bhutta ZA, Das JK, Rizvi A, et al. Evidence-based interventions for improvement of maternal and child nutrition: what can be done and at what cost? *The Lancet*. 2013;382:452-477.

Acknowledgments

This study is part of research towards a Ph.D. at The University of Sydney funded by Endeavour Postgraduate Scholarship (Ph.D.); Department of Education and Training, Australian Government. The study received partial support from the James P. Grant School of Public Health, BRAC University, Bangladesh, and the Sydney Medical School, The University of Sydney, Australia. We thank the *Shasthya Kormi* and the *Shasthya Shebika* of BRAC, and the pregnant women and their families in Sherpur for their time in this study.

Funding:

The study was funded by James P. Grant School of Public Health; Bangladesh, and Sydney Medical School Foundation; Australia

Author contributions:

MC conceived the overall study, developed the study design and data collection methods, and conducted the data analysis, interpreted data, and wrote the first draft of the manuscript. MJD and CRG provided critical input regarding study design, sample size calculation, and outcome evaluation and the statistical analysis plan. KA provided practical guidance in formulating the implementation strategy. AA provided crucial input on formative research and process evaluation design. PK provided technical support for data analyses. CRG and MC obtained funds for the intervention. CRG provided critical input to all drafts and approved the final version for submission. MJD, KA, PK and AA critically reviewed the manuscript and approved the final version for submission. MC, MJD, CRG, PK, KA and AA agreed to be accountable for all investigations necessary to resolve questions related to accuracy or integrity of all or any part of the work.

Competing interests:

None to declare

561 **Data and materials availability:**

562 The data described in this manuscript, the code book, and the analytic code will be made
563 available upon request pending approval by the investigators of an application to use the data.

564

565 **Supplemental Information**

566 **S1 Table: The mean birth weight and the adjusted mean difference in birth weight of**
 567 **balanced -plate intervention compared to usual programs.**

568 **S2 Table: Percentage of low birth weight and adjusted relative risk of balanced-plate**
 569 **intervention compared to usual programs.**

570 **Table 1: Cluster, baseline maternal and household, and antenatal care and delivery**
571 **characteristics by study arm**

	Intervention (n=382)	Control (n=397)
Cluster characteristics		
Number of clusters	18	18
Mean (SD) population in clusters in 2016	9775 (2007)	9639 (1763)
Mean (SD) number of households in clusters	2315 (288)	2321 (419)
Mean (SD) number of women of reproductive age	2639 (542)	2631 (481)
Mean (SD) number of pregnant women enrolled per clusters	24.7 (1.0)	24.9 (0.5)
Baseline maternal and household characteristics		
Mother's age (years):		
Mean (SD)	23.0 (4.6)	24.0 (4.3)
15-19	80 (21.7)	64 (17.0)
20-34	280 (76.1)	307 (81.4)
>34	8 (2.2)	6 (1.6)
Missing	14	20
Mother's education:		
No education	82 (22.0)	89 (23.1)
Primary	250 (67.2)	242 (62.7)
Secondary	27 (7.3)	35 (9.1)
Higher	13 (3.5)	20 (5.2)
Missing	10	11
Monthly family income (SD) in Bangladesh Taka		
Less than 5000	87 (23.9)	79 (21.2)
5000 to less than 10,000	190 (52.2)	209 (56.2)
10,000 or more	87 (23.9)	84 (22.6)
Missing	18	25
Father's education:		
No education	130 (35.0)	151 (39.7)
Primary	188 (50.7)	169 (44.5)
Secondary	34 (9.2)	39 (10.3)
Higher	19 (5.1)	21 (5.5)
Missing	11	17
Marital Status		
Married	372 (97.4)	386 (97.3)
Not married	10 (2.6)	11 (2.7)
Parity:		

	Intervention (n=382)	Control (n=397)
Mean (SD)	1.8 (1.2)	1.9 (1.3)
0	36 (10.3)	48 (12.6)
1	134 (38.5)	115 (30.3)
≥2	178 (51.2)	217 (57.1)
Missing	34	17
Previous pregnancy loss		
Yes	67 (19.3)	87 (22.9)
No	281 (80.8)	293 (77.1)
Missing	34	17
Gestation at enrolment (weeks) Mean (SD)	10.1 (3.2)	9.6 (3.2)
Missing	27	2
Antenatal care and delivery characteristics		
Iron Folic Acid Supplements		
Mean number consumed (SD)	45.6 (61.8)	29.0 (46.8)
Any use	117 (30.6)	70 (17.6)
None	265 (69.4)	327 (83.4)
Antenatal care visit by SK		
Mean number of visits (SD)	3.8 (2.6)	3.6 (2.2)
<4 visits	158 (41.1)	198 (49.9)
≥4 visits	224 (58.6)	199 (50.1)
Antenatal care in health centers		
Mean number of visits (SD)	0.7 (1.1)	1.0 (1.4)
No Visit	227 (59.4)	246 (51.9)
<4 visits	143 (37.4)	169 (42.6)
≥4 visits	12 (3.1)	22 (5.5)
Place of delivery		
Facility	79 (20.7)	66 (16.6)
Home	280 (73.3)	323 (81.4)
Unknown	23 (6.0)	8 (2.0)
Mode of delivery		
Normal	331 (86.7)	361 (90.9)
C-Section	33 (8.6)	34 (8.6)
Unknown	18 (4.7)	2 (0.5)
Time of birth weight measurement		
Within 24h	266 (69.6)	290 (73.1)
24 to 48h	75 (19.6)	61 (15.4)
48 to 72h	13 (3.4)	19 (4.8)
Unknown	28 (7.3)	27 (6.8)

Numbers are frequencies (and column percentage) unless stated otherwise

Table 2: The mean birth weight and the mean difference in birth weight of balanced-plate intervention compared to usual programs.

	Number of participants		Birth weight (g)		Mean difference				P-value	P-value for interaction
	Intervention	Control	Intervention Mean (SD)	Control Mean (SD)	Estimate	(95 %CI)				
Overall	382	397	2861.0 (444.2)	2736.8 (423.2)	127.5	(11.1, 243.9)		0.032		
Mother's age¹ (years)										0.009
<20	80	64	2916.3 (471.1)	2637.5 (481.0)	299.1	(101.6, 496.6)		0.003		
>=20	288	313	2847.6 (443.4)	2762.6 (401.1)	95.6	(-17.3, 208.4)		0.097		
Mother's education²										0.330
None	82	89	2869.5 (460.6)	2693.3 (339.7)	175.1	(29.9, 320.4)		0.018		
Primary	250	242	2839.2 (443.9)	2761.6 (394.4)	92.9	(-28.0, 213.7)		0.132		
Secondary or higher	40	55	2982.5 (435.5)	2741.8 (588.0)	226.1	(-33.5, 485.8)		0.088		
Family income³										0.236
Low	87	79	2819.5 (397.9)	2663.3 (420.4)	166.8	(10.7, 322.8)		0.036		
Medium	190	209	2806.8 (421.7)	2750.7 (390.8)	51.8	(-70.0, 173.6)		0.404		
High	87	84	2995.4 (514.5)	2791.7 (433.2)	183.3	(-40.9, 407.4)		0.109		

Birth weight was analyzed using linear regression. We adjusted all models for clusters using generalized estimating equation (GEE) models, assuming an exchangeable correlation structure and applied a sandwich estimator to standard errors. ¹ n=745 due to 34 missing maternal age values. ² n=758 due to 21 missing maternal education values. ³ n=736 due to 43 missing family income values.

577 **Table 3: Percentage of low birth weight and relative risk of balanced-plate intervention compared to usual programs.**

	Number of participants		Low birth weight (<2,500g)				Relative risk			
	Intervention	Control	Intervention n (%)		Control n (%)		Estimate	(95% CI)	P-value	P-value for interaction
Overall	382	397	37	(10)	87	(22)	0.43	(0.25, 0.75)	0.003	
Mother's age¹ (years)										0.217
<20	80	64	6	(8)	17	(27)	0.28	(0.11, 0.71)	0.007	
>=20	288	313	31	(11)	62	(20)	0.54	(0.29, 0.98)	0.044	
Mother's education²										0.240
None	82	89	11	(13)	22	(25)	0.54	(0.28, 1.05)	0.069	
Primary	250	242	25	(10)	47	(19)	0.50	(0.26, 0.93)	0.029	
Secondary or higher	40	55	1	(3)	12	(22)	0.11	(0.02, 0.8)	0.029	
Family income³										0.441
Low	87	79	9	(10)	25	(32)	0.31	(0.13, 0.76)	0.010	
Medium	190	209	23	(12)	41	(20)	0.60	(0.32, 1.14)	0.117	
High	87	84	5	(6)	12	(14)	0.45	(0.13, 1.53)	0.199	

578 Low birth weight was analyzed using binomial regression with a log link. We adjusted all models for clusters using generalized estimating equation (GEE) models, assuming an
579 exchangeable correlation structure and applied sandwich estimator to standard errors. ¹ n=745 due to 34 missing maternal age values. ² n=758 due to 21 missing maternal
580 education values. ³ n=736 due to 43 missing family income values

581 **Figure 1: Trial flow chart**





Figure 2: The mean number and 95% confidence intervals* of food groups (A) and animal-source protein foods (B) consumed by women per day by treatment group.



<i>n</i> – Intervention	167	362	369	366	345	318	272	178
<i>n</i> – Control	131	367	364	349	339	302	257	159

* The estimated the mean number of food groups consumed and 95% confidence intervals were calculated using linear regression with the study arm as a covariate using Generalised Estimating Equations (GEE) assuming an exchangeable correlation structure and applying the sandwich estimator to standard errors.