

Epidemiological characteristics and transmission dynamics of the outbreak caused by the SARS-CoV-2 Omicron variant in Shanghai, China: a descriptive study

Zhiyuan Chen^{1*}, Xiaowei Deng^{1*}, Liqun Fang^{2*}, Kaiyuan Sun³, Yanpeng Wu¹, Tianle Che², Junyi Zou¹, Jun Cai¹, Hengcong Liu¹, Yan Wang¹, Tao Wang², Yuyang Tian¹, Nan Zheng¹, Xuemei Yan¹, Ruijia Sun¹, Xiangyanyu Xu¹, Xiaoyu Zhou¹, Shijia Ge⁴, Yuxiang Liang¹, Lan Yi¹, Juan Yang¹, Juanjuan Zhang^{1†}, Marco Ajelli⁵, Hongjie Yu^{1†}

1. School of Public Health, Fudan University, Key Laboratory of Public Health Safety, Ministry of Education, Shanghai, China
2. State Key Laboratory of Pathogen and Biosecurity, Beijing Institute of Microbiology and Epidemiology, Beijing, China
3. Division of International Epidemiology and Population Studies, Fogarty International Center, National Institutes of Health, Bethesda, MD, USA
4. Department of Infectious Diseases, Huashan Hospital, Fudan University, Shanghai, China
5. Laboratory for Computational Epidemiology and Public Health, Department of Epidemiology and Biostatistics, Indiana University School of Public Health, Bloomington, IN, USA

*These authors contributed equally to this work.

†Co-corresponding authors:

Juanjuan Zhang, School of Public Health, Fudan University, Key Laboratory of Public Health Safety, Ministry of Education, Shanghai 200032, China. E-mail: zhangjuan@fudan.edu.cn;

Hongjie Yu, School of Public Health, Fudan University, Key Laboratory of Public Health Safety, Ministry of Education, Shanghai 200032, China. E-mail: yhj@fudan.edu.cn

31

32 Abstract: 370 words

33 Main text: 3,992 words

34 **Summary**

35 **Background**

36 In early March 2022, a major outbreak of the severe acute respiratory syndrome
37 coronavirus 2 (SARS-CoV-2) Omicron variant spread rapidly throughout Shanghai,
38 China. Here we aimed to provide a description of the epidemiological characteristics
39 and spatiotemporal transmission dynamics of the Omicron outbreak under the
40 population-based screening and lockdown policies implemented in Shanghai.

42 **Methods**

43 We extracted individual information on SARS-CoV-2 infections reported between
44 January 1 and May 31, 2022, and on the timeline of the adopted non-pharmacological
45 interventions. The epidemic was divided into three phases: i) sporadic infections
46 (January 1–February 28), ii) local transmission (March 1–March 31), and iii) city-wide
47 lockdown (April 1 to May 31). We described the epidemic spread during these three
48 phases and the subdistrict-level spatiotemporal distribution of the infections. To
49 evaluate the impact on the transmission of SARS-CoV-2 of the adopted targeted
50 interventions in Phase 2 and city-wide lockdown in Phase 3, we estimated the dynamics
51 of the net reproduction number (R_t).

53 **Findings**

54 A surge in imported infections in Phase 1 triggered cryptic local transmission of the
55 Omicron variant in early March, resulting in the largest coronavirus disease 2019
56 (COVID-19) outbreak in mainland China since the original wave. A total of 626,000
57 SARS-CoV-2 infections were reported in 99.5% (215/216) of the subdistricts of
58 Shanghai. The spatial distribution of the infections was highly heterogeneous, with 40%
59 of the subdistricts accounting for 80% of all infections. A clear trend from the city
60 center towards adjacent suburban and rural areas was observed, with a progressive
61 slowdown of the epidemic spread (from 544 to 325 meters/day) prior to the citywide
62 lockdown. During Phase 2, R_t remained well above 1 despite the implementation of
63 multiple targeted interventions. The citywide lockdown imposed on April 1 led to a
64 marked decrease in transmission, bringing R_t below the epidemic threshold in the entire
65 city on April 14 and ultimately leading to containment of the outbreak.

67 **Interpretation**

68 Our results highlight the risk of widespread outbreaks in mainland China, particularly
69 under the heightened pressure of imported infections. The targeted interventions
70 adopted in March 2022 were not capable of halting transmission, and the
71 implementation of a strict, prolonged city-wide lockdown was needed to successfully

72 contain the outbreak, highlighting the challenges for successfully containing Omicron
73 outbreaks.

74

75 **Funding**

76 Key Program of the National Natural Science Foundation of China (82130093).

77

78 **Research in context**

79 **Evidence before this study**

80 On May 24, 2022, we searched PubMed and Europe PMC for papers published or
81 posted on preprint servers after January 1, 2022, using the following query:
82 (“SARS-CoV-2” OR “Omicron” OR “BA.2”) AND (“epidemiology” OR
83 “epidemiological” OR “transmission dynamics”) AND (“Shanghai”). A total of 26
84 studies were identified; among them, two aimed to describe or project the spread of the
85 2022 Omicron outbreak in Shanghai. One preprint described the epidemiological and
86 clinical characteristics of 376 pediatric SARS-CoV-2 infections in March 2022, and the
87 other preprint projected the epidemic progress in Shanghai, without providing an
88 analysis of field data. In sum, none of these studies provided a comprehensive
89 description of the epidemiological characteristics and spatiotemporal transmission
90 dynamics of the outbreak.

91

92 **Added value of this study**

93 We collected individual information on SARS-CoV-2 infection and the timeline of the
94 public health response. Population-based screenings were repeatedly implemented
95 during the outbreak, which allowed us to investigate the spatiotemporal spread of the
96 Omicron BA.2 variant as well as the impact of the implemented interventions, all
97 without enduring significant amounts of underreporting from surveillance systems, as
98 experienced in other areas. This study provides the first comprehensive assessment of
99 the Omicron outbreak in Shanghai, China.

100

101 **Implications of all the available evidence**

102 This descriptive study provides a comprehensive understanding of the epidemiological
103 features and transmission dynamics of the Omicron outbreak in Shanghai, China. The
104 empirical evidence from Shanghai, which was ultimately able to curtail the outbreak,
105 provides invaluable information to policymakers on the impact of the containment
106 strategies adopted by the Shanghai public health officials to prepare for potential
107 outbreaks caused by Omicron or novel variants.

108 Background

109 The first wave of coronavirus disease 2019 (COVID-19) in China subsided quickly
110 with strict containment measures in March 2020.¹ As numerous variants have emerged
111 across the globe, China successfully contained multiple COVID-19 outbreaks by
112 adhering to a containment policy. This policy aimed to curb flare-ups of local
113 transmissions in the shortest possible time, which is done by relying on a set of
114 non-pharmacological interventions (NPIs) with adjusted intensities according to the
115 situation on the ground.^{2,3}

116
117 In November 2021, the severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2)
118 Omicron variant was first reported in South Africa.⁴ Its high transmissibility and
119 immune escape properties enabled it to rapidly replace previous strains and become
120 dominant globally.^{5,6} Although the containment policy implemented in China had been
121 effective in containing pre-Omicron outbreaks, its effectiveness against Omicron was
122 unclear. In fact, a total of 750,000 SARS-CoV-2 infections were reported in mainland
123 China in the first five months of 2022,⁷ the majority in Shanghai.

124
125 Shanghai is one of the most populous and economically advanced metropolises in
126 China, with a population of nearly 25 million⁸ and unique features compared to other
127 areas in mainland China and abroad. The population in Shanghai was mostly
128 vaccinated with domestically-developed inactivated vaccines, which had relatively
129 lower effectiveness in preventing SARS-CoV-2 infections due to their lower
130 antibody-neutralizing responses compared to mRNA vaccines^{9,10}. In addition, the
131 vaccination coverage was highest in the young population (100.0% in individuals aged
132 between 12 and 17 years had completed the primary vaccine schedule as of March
133 2022)¹¹ and lowest in the elderly (only 62.0% in individuals aged 60 years or older had
134 completed the primary schedule as of April 2022).¹² Compared to the rest of the cities
135 in mainland China, Shanghai had the highest risk of SARS-CoV-2 infection
136 importation due to its global connectivity: about 30-40% of all international flights
137 arriving in China since 2020 landed in Shanghai.¹³ Although a quarantine period of two
138 weeks was mandatory for all incoming travelers, lapses in these measures may have led
139 to the repeated seeding of SARS-CoV-2 into the community.^{14,15} Consequently, in
140 early March 2022, a major outbreak of the Omicron BA.2 variant started to spread in
141 Shanghai.¹⁶

142
143 In this study, we provide a detailed description of Shanghai's outbreak response
144 timeline, the spatiotemporal distribution of COVID-19 cases, and Omicron
145 transmission dynamics since 2022. The analysis of the Shanghai 2022 Omicron

outbreak could provide invaluable information to policymakers on the impact of the containment strategies adopted by Shanghai to prepare for potential outbreaks caused by Omicron or future novel variants.

Methods

Definition of infection and case

An asymptomatic infection is defined as a PCR-confirmed individual who i) does not meet any of the following clinical criteria: fever, cough, sore throat, and other self-perceived and clinical-identifiable symptoms or signs; and ii) has no radiographic evidence of pneumonia. Laboratory-confirmed cases are categorized into four types based on clinical severity: mild, moderate, severe, and critical cases, with detailed definitions in the **Supplementary Information**. Detailed surveillance, detection, and management of infections and cases are also presented in the **Supplementary Information**.

Public health response

We summarize the NPIs implemented in Shanghai before and during the Omicron waves. Prior to the initial surge in Omicron, Shanghai maintained a baseline level of intensive NPIs against potential outbreaks, including stringent border control policies, symptom-based surveillance, case isolation, tracing of close contacts (requires quarantine in separate facilities) and contacts of contacts, occupation-based screening, targeted screening of individuals at high risk of infection (e.g., contacts of contacts), and a set of other social distancing measures such as travel restrictions and community confinement (**Figure 1**). After the Omicron variant was introduced in Shanghai, a set of additional NPIs were implemented to halt its transmission. In terms of border control measures, the point of entry for international flights to Shanghai was diverted to other airports. At the local scale, the closure of in-person educational activities at all levels was implemented from March 12 to 15, 2022 (starting from primary and secondary school, then followed by universities). From March 16 to March 27, Shanghai adopted grid management at the subdistrict level, fractioning subdistricts into high-risk areas. The lists of high-risk areas changed dynamically over time based on the following principles: the epidemiological situation measured in terms of number of cases and infections, gathering of crowds, population density, social characteristics, and economic activity.¹⁷ During this period, several rounds of nucleic acid screening were performed within high-risk and non-high-risk areas. In high-risk areas, either one or two rounds of population-based nucleic acid screening were performed within 48 hours, together with a lockdown. In non-high-risk areas, a single round of mass PCR screening

was performed between March 18 and 20. Afterwards, the entire city entered a phased stage of lockdown, where eastern Shanghai (which comprises areas east of the Huangpu River, **Supplementary Figure 1**) entered a population-wide lockdown on March 28, and then the rest of Shanghai entered the lockdown phase on April 1. The lockdown restriction was lifted on June 1 when the daily number of infections first declined to 10..

Data sources and collection

Aggregated data on the number of local and imported infections according to symptomatic status in Shanghai from December 2019 were gathered from the Shanghai Municipal Health Commission through a combined approach of manual downloads and compiled scripts. Individual line lists of all SARS-CoV-2 infections were retrieved from multiple publicly available official data sources (websites of municipal health commissions) and integrated with the supplementary information gathered from the websites of local government media (**Supplementary Table 1**). The resulting line list contains the following variables: residential location (address, district, and subdistrict), symptomatic status (including the clinical outcomes of initially asymptomatic infections), date of official reporting, and means of infection identification (i.e., routine screening of community groups or general screening of the quarantined population) (**Supplementary Table 2**). Specifically, community screening groups referred to those residents living in areas reporting no new infections, people proactively looking for healthcare or nucleic acid tests, and essential workers. The quarantined population corresponded to individuals who were quarantined at home or in designated facilities, lived in areas reporting new infections or other key settings under close management, and were identified through screening of close contacts¹⁸. We obtained the timeline of the adopted NPI strategies from the official announcements of the authorities of Shanghai through search engine queries; the list of high-risk areas during March 16 and March 27 was found by searching on local media pages (**Supplementary Table 3**). More details on data collection are provided in the **Supplementary Information**.

Statistical analyses

Definition of the three phases of the epidemic

Three phases of the epidemic were defined according to the staged NPIs policies adopted and the epidemic situation. The first phase covers the period from January 1 to February 28, 2022, during which the main risk was posed by the importation of infections and only small numbers of sporadic and locally transmitted cases were recorded. The second phase started on March 1, immediately before the outbreak of the BA.2 Omicron variant was confirmed¹⁶. In addition, no epidemiological link between

infections reported before and after March 1, 2022, was identified by the authorities. The third phase covers the period from April 1 to May 31, 2022, when the entire city went into staged lockdowns and the epidemic started to subside.

Inference of delay between sampling and reporting

The date of symptom onset or infection was needed to describe the transmission dynamics, whereas we only had the date of official reporting for each infection. We estimated the delayed days between the sampling dates and reporting dates and then generated the sampling date for each infection by assuming that the sampling dates were close to the dates of symptom onset or infection time.

The mean reporting delay was determined by computing the cross-correlation coefficients between the cumulative number of reported SARS-CoV-2 infections and the cumulative number of sampled specimens under different lengths of lag (**Supplementary Figure 2**), the details of which are presented in the **Supplementary Information**. We applied delays of 2 days, 3 days, and 2 days for the periods before March 15, between March 16 and May 14, and after May 15, respectively. In the sensitivity analysis, we considered a four-day delay for the period from March 16 to May 14.

In addition to reporting delay, we also estimated individual locations and the progression of pre-symptomatic infections for data analysis (see **Supplementary Information** for details).

Spatial trends

Trend surface analysis was used to explore the SARS-CoV-2 spread in Shanghai, China. Thin-plate spline regression was used to interpolate the first date of sampling in each $3 \text{ km} \times 3 \text{ km}$ cell of a gridded population of Shanghai to draw a spatial trend surface plot. The local slope (time/distance) of the trend surface was then measured using a 3×3 moving window filter, and the inverse of the slope value was used to estimate the speed of the spatial spread of SARS-CoV-2 (distance/time).^{20,21}

Estimation of epidemiological parameters

The net reproduction number (R_t) was estimated using the EpiEstim R package.²² Briefly, the estimate is based on a Bayesian approach that relies on the knowledge of the time series of infections (sampling date of positive tests in our analysis) and the distribution of the generation time. In the absence of estimates of the generation time for the analyzed outbreak, we assumed it to be gamma-distributed with a mean of 2.72

days (shape = 3.25, scale = 0.84), corresponding to the estimated serial interval during the Omicron wave in Hong Kong in the presence of strict interventions.¹⁹

We estimated the growth rate (r) and 95% confidence interval (CI) in the second epidemic phase by fitting a linear regression to the logarithm of the incidence of infections. The doubling time in the early stage was then estimated as $\ln(2)/r$, where the 95% CI was calculated using the delta method. All analyses and visualization were performed in ArcGIS 10.7 and R (version 4.0.2).

Role of the funding source

The funder of the study had no role in study design, data collection, data analysis, data interpretation, or writing of the report. The corresponding authors had full access to all the data in the study and had final responsibility for the decision to submit for publication.

This study was approved by the institutional review board of the School of Public Health, Fudan University (IRB# 2022-05-0968). All data were collected from publicly available sources. Data were de-identified, and the need for informed consent was waived.

Results

Following the successful containment of the initial epidemic wave caused by the Wuhan outbreak in early 2020, Shanghai was under constant pressure of potential outbreak introductions due to a high influx of international travelers relative to the rest of the cities in mainland China. Despite ranking first in the cumulative number of imported confirmed SARS-CoV-2 infections before 2022, Shanghai had sustained minimum local transmission prior to the emergence of the Omicron variant (**Figure 2a**), which was attributable to well-managed entry quarantine, screenings, and case isolation programs. Nonetheless, sporadic local transmissions events occurred, and all pre-Omicron outbreaks were swiftly suppressed.

Starting in mid-February 2022, Shanghai faced a significant surge of imported infections, the majority of which were linked to the Omicron wave in Hong Kong (**Supplementary Figure 3**). The number of imported infections reached a peak of 81 per day on February 24—more than 19 times the prior daily average in 2020-2021 (4.1 imported infections per day). Overall, from February 15 to March 15, a total of 1,112 imported infections were reported (**Figure 2b**). Compared to any other time point

during the rest of the pandemic, the test positive rate among inbound travelers was at its highest throughout February 2022, reaching a peak of 96.3 infections per 100,000 travelers (**Supplementary Figure 4**).

The sudden upsurge of infected travelers ultimately led to a breach of the entry control program, which might have resulted in the cryptic transmission of the Omicron variant since early March. A cluster of 14 SARS-CoV-2 positive individuals was detected starting from March 1 among attendees of a square dance event for seniors. Because of the Omicron variant's high transmissibility and immune evasive properties, as well as the delayed discovery of the outbreak, the contact tracing program was quickly overwhelmed, leading to a fast-growing epidemic that resulted in a total of 626,000 SARS-CoV-2 infections as of May 31, 2022 (**Supplementary Figure 5**).

Most infections (96.0%) were identified in individuals under quarantine (**Supplementary Figure 6a**). The extent to which the infections were identified by community screening (e.g., essential workers) gradually declined from 13.6% in Phase 2 to 2.7% in Phase 3. Meanwhile, the proportion of infections presenting with symptoms varied from 15.5% in Phase 1 to 9.8% in Phase 3 (**Supplementary Figure 6b**).

The Omicron outbreak was widespread throughout the entire area of Shanghai, with 99.5% (215/216) of the subdistricts reporting positive individuals. However, the spatial distribution of infections was highly heterogeneous, with 40% of the city area accounting for 80% of all infections; 35.5% of infections were identified in the Pudong New Area alone (**Supplementary Figure 5**). We divided the Omicron wave into three phases: Phase 1 (from January 1 to February 28), characterized by a high risk of importation; Phase 2 (from March 1 to 31), characterized by local community transmission; Phase 3 (after April 1), characterized by high levels of local transmission and the implementation of city-scale lockdowns to contain the outbreak. The total incidence of reported infections increased from 0.02 infections per 1,000 individuals in Phase 1 to 2.9 infections per 1,000 individuals in Phase 2 and reaching 21.9 infections per 1,000 individuals in Phase 3 (**Figure 3a-c**). Overall, an incidence of 24.8 infections per 1,000 individuals was observed (**Figure 3d**).

The spatial spread of the reported infections shows a clear trend from the city center towards adjacent suburban and rural areas (**Figure 4a-c**). Despite the implementation of a citywide lockdown in Phase 3, the outbreak continued to expand towards suburban and rural areas (**Figure 4c**). We estimated that spread speed of the infection throughout the city progressively slowed from an average of 544 meters/day in the first week of the

outbreak (February 27-March 5) to approximately 325 meters/day at the end of March (March 25-31), before the city-scale lockdown (**Figure 4d**). Although this decreasing trend was common to all districts, the central areas (including the districts of Jing'an, Yangpu, Hongkou, Putuo, Changning, Xuhui, and Huangpu) showed the fastest rate of spread, especially in the early stage of the outbreak (**Supplementary Figure 7**).

From March 16 to March 27, prior to the blanket lockdown of the entire city, Shanghai launched a set of targeted interventions and management strategies on a smaller geographical scale. The policy was adopted at the subdistrict level, and subdistricts were divided into high-risk areas (see definition in the **Methods** section), where the interventions were implemented, and non-high-risk areas. We further divided the non-high-risk areas into two categories based on their spatial proximity to the high-risk areas: moderate-risk areas (i.e., areas that were not classified as high-risk but were adjacent to high-risk areas) and low-risk areas (i.e., areas that were neither classified as high-risk nor adjacent to high-risk areas) (**Figure 5a**). From March 16 to March 29, the cumulative number of infections per 1,000 individuals was highest in high-risk areas (2.20, 95% CI, 2.18-2.22), followed by moderate-risk areas (1.51, 95% CI, 1.49-1.54) and low-risk areas (0.53, 95% CI, 0.49-0.57) (**Figure 5b-d**). The implemented targeted intervention strategies prior to the citywide lockdown were not sufficient to prevent infections to rise (**Figure 5e**), and R_t remained well above 1 (the epidemic threshold) across all three area categories for the entire period (**Figure 5f**). Moreover, R_t in moderate-risk and low-risk areas increased slightly from March 20 to March 24. The sensitivity analysis that considered alternative values for the delay between sampling and reporting showed similar patterns (**Supplementary Figure 8**). Overall, the growth rate of the epidemic in Phase 2 was estimated to be 0.216 per day (95% CI, 0.210 to 0.222), with infections doubling every 3.21 days (95% CI, 3.12 to 3.29) (**Figure 5g**).

Three days after the implementation of a citywide lockdown on April 1 (Phase 3), R_t started to decrease and fell below the epidemic threshold on April 14 (**Figure 6a-b**). Despite the marked differences in R_t observed between eastern and western Shanghai when the targeted interventions were adopted (Phase 2), immediately after the implementation of the citywide lockdown, the same level of SARS-CoV-2 transmission (R_t) was observed in all of Shanghai (**Figure 6b** and **Supplementary Figure 9**, where a sensitivity analysis confirming this finding is presented).

Discussion

This study provides a comprehensive picture of the spatiotemporal dynamics of the Omicron outbreak in Shanghai in 2022. Our findings highlight that, given the

Shanghai population's susceptibility to the SARS-CoV-2 Omicron BA.2 variant and its high transmissibility, the heightened importation risk of infected individuals from areas with widespread viral circulation was capable of triggering a major epidemic wave. The targeted interventions implemented in March were not enough to halt transmission. The citywide lockdown imposed on April 1 led to a decrease in transmission, bringing the reproduction number below the epidemic threshold in the entire city. Overall, the Omicron outbreak in Shanghai was successfully contained through a combination of stringent measures.

Our findings suggest that when facing a high influx volume of travelers from high-risk areas, breaches in the screening and quarantine of travelers may have the potential to trigger major outbreaks. It is also possible that the wave was simultaneously triggered by domestic travelers, as suggested by previous field investigations by local authorities.²³ We cannot exclude that the outbreak might have originated from multiple sources that led to simultaneous transmission chains prior to the identification of the first local transmission event.

Compared to previous local SARS-CoV-2 outbreaks that emerged in China after the original COVID-19 wave, the rapid spread of the Omicron outbreak in Shanghai was likely driven by the high transmissibility and/or immune-escape properties of Omicron BA.2.²⁴ The immunological landscape of Shanghai at the onset of this outbreak was fragile, with essentially no prior natural immunity (less than 0.01% of the population was infected prior to the outbreak²⁵). In addition, although the coverage of the primary vaccination series exceeded 90% of the total population, the vaccination coverage for individuals aged ≥ 60 years was relatively low (62% for the primary series and 38% for the booster dose²⁶). Moreover, the inactivated vaccines used in Shanghai provided very low protection against Omicron variant infection (approximately 17.0% after receiving a booster dose¹¹, based on the level of neutralizing titer), which quickly wanes over time.²⁷ As a result, the Shanghai population was particularly vulnerable to the Omicron variant. Nonetheless, inactivated vaccines could provide high protection against severe outcomes.^{9,28} These considerations further emphasize the importance of administering vaccines, especially to the most vulnerable segments of the population, and deploying immunization strategies with improved effectiveness against the current circulating variants through procuring currently available mRNA vaccines, approve other domestically developed vaccine candidates with broader protection against Omicron and future variants, and heterologous boosting strategies that have demonstrated improved antibody response.

Another factor that led to the major outbreak might have been the large number of asymptomatic carriers that contributed to widespread silent transmission.²⁹ Here, we found a high proportion (more than 90%) of asymptomatic infections, larger than that estimated for the ancestral lineage (69%³⁰), but in line with other estimates for other SARS-CoV-2 variants in the presence of vaccination (85%³¹). Lower proportions of asymptomatic infections were reported in previous studies,³² including for the Omicron BA.2 outbreak in Hong Kong¹⁹; however, those estimates were obtained in the absence of repeated screenings of the population. Moreover, the criteria adopted for the definition of asymptomatic infection may vary across locations and the study period.

Between March 1 and March 31, a series of public health measures were adopted to reduce transmission between individuals living in areas at different risk levels, and the estimated doubling time in this phase of the epidemic was comparable to that of the Omicron wave in Hong Kong (3.2 days as compared to 3.4 days estimated for Hong Kong¹⁹). However, the implemented interventions were insufficient to curtail the epidemic. In fact, the repeated two-day lockdowns and population screenings implemented in high-risk areas were not sufficient to halt SARS-CoV-2 transmission, and these failed to identify all infected individuals (e.g., those in the early course of the infection, when the viral load was below the detection limit³³). Once the two-day local lockdowns were lifted, unidentified infected individuals were allowed to freely move and transmit the infection to other areas, as shown by the rise in new daily infections identified in areas adjacent to high-risk areas.

The implementation of the citywide lockdown prevented further growth of the epidemic and successfully contained the outbreak. However, given the number of infectious individuals at the time of implementation (over 10,000 new infections reported per day), it took 13 days for R_t to fall below the threshold. It is also possible that it was difficult to transport and isolate all positive individuals in a timely manner into dedicated facilities during the lockdown. Moreover, residents may still have been exposed to the virus inside their buildings.³⁴ It is thus important to carry out quantitative evaluations of the unavoidable risks posed by the adopted strategies. Empirical evidence from Shanghai shows that it is possible to maintain containment policies against the highly transmissible Omicron BA.2 variant, although this required unprecedented efforts to achieve. However, it is important to stress that the interruption of onward transmission in Shanghai might have prevented the infection from spreading into other cities, potentially preventing a major public health crisis in mainland China.¹¹ Nonetheless, it is important to stress that the measures adopted during the Omicron outbreak in Shanghai, including a strict and prolonged city-scale lockdown, would be

socially costly and impractical in the long term, emphasizing the importance refining NPI strategies to be less disruptive of daily lives and boosting population immunity level to significantly reduce the morbidity and mortality burden of SARS-CoV-2.¹¹

A strength of our study is the unique features of the analyzed dataset. In fact, Shanghai's population-based screening policy, which allowed infections to be identified, likely resulted in a high infection ascertainment rate and the identification of the majority of infections throughout the outbreak. This allowed us to investigate the spatiotemporal spread of the Omicron BA.2 variant as well as the impact of the implemented interventions, all without significant amounts of underreporting from surveillance systems, as experienced in other areas. However, our study suffers from the limitations rooted in the uncertainty and fragmentary nature of publicly available sources. First, key variables, such as the date of symptom onset and the addresses of infected individuals, suffered from a high level of missing data. Second, the sparse information on the demographic characteristics of positive individuals limits the scope of our study. Third, the final clinical outcome of infected individuals was only partially available (e.g., we had no information about patients requiring intensive care treatment). Moreover, given the real-time nature of this study and the censoring of the final outcome data, an analysis of the COVID-19 burden is not possible at this point. Finally, this study does not provide quantitative estimates of the impact of specific interventions on transmission dynamics, but providing an overall assessment of the synergetic effect of the adopted interventions.

In conclusion, this study provides a quantitative description of the spatiotemporal spread of the Omicron BA.2 variant in Shanghai and explores the impact of the multifaceted implemented response measures. Our findings highlight the risk of widespread outbreaks in mainland China, particularly under the large pressure of imported infections. The successful containment of the Shanghai outbreak through the implementation of a strict and prolonged citywide lockdown shows that it is possible to successfully contain an Omicron outbreak in China, although it requires the implementation of a set of very stringent interventions. Disentangling the effects of each of the performed interventions might provide further insights into key public health priorities when faced with the emergence of future novel variants.

483 **Contributors**

484 H.Y. conceived and designed the study. H.Y. and JJ.Z. supervised the study. X.D., JJ.Z.,
485 Z.C., JY.Z., J.C., Y.W., Y.T., N.Z., X.Y., R.S., X.X., X.Z., S.G., Y.L., and L.Y.
486 collected and checked data. Z.C., JJ.Z., L.F., T.C., YP.W., JY.Z., J.C., H.L., K.S., Y.W.,
487 T.W., and Y.T. analyzed the data. Z.C., JJ.Z. and H.Y. wrote the first draft of the
488 manuscript. Z.C. JJ.Z, K.S., M.A., L.F., J.Y., and H.Y. interpreted the results and
489 revised the content critically. All authors approved the final version for submission and
490 agreed to be accountable for all aspects of the work.

492 **Declaration of interests**

493 H.Y. has received research funding from Sanofi Pasteur, GlaxoSmithKline, Yichang
494 HEC Changjiang Pharmaceutical Company, Shanghai Roche Pharmaceutical
495 Company, and SINOVAC Biotech Ltd. M.A. has received research funding from
496 Seqirus. None of those research funding is related to this work. All other authors report
497 no competing interests.

499 **Data sharing**

500 The data and code that support the findings of this study will be made available in
501 GitHub upon manuscript acceptance.

503 **Acknowledgments**

504 The findings and conclusions in this report are those of the authors and do not
505 necessarily represent the official position of the NIH. This study was supported by
506 grants from the Key Program of the National Natural Science Foundation of China
507 (grant 82130093 to H.Y.). The funders had no role in study design, data collection, data
508 analysis, data interpretation or writing of the report.

509

510

511 **Figures legends**

512 **Figure 1. Timeline of the public health response in Shanghai by epidemic phase.**

513

514 **Figure 2. Temporal dynamics of local and imported SARS-CoV-2 infections in**
 515 **Shanghai since early 2020. (a)** Number of reported SARS-CoV-2 infections in
 516 Shanghai between 2020-2022, stratified by local and imported infections. **(b)** The
 517 same as in (a), but for the period from January 1 to May 31, 2022.

518

519 **Figure 3. Geographical distribution of SARS-CoV-2 infections. (a-d)** Cumulative
 520 number of new SARS-CoV-2 infections per 1,000 individuals in each phase and
 521 overall.

522

523 **Figure 4. Spatial trends and speed of spread of the epidemic in the three phases.**

524 **(a)** Spatial location of the reported infections during the first phase of the epidemic.
 525 **(b-c)** Estimated arrival time of the epidemic in the different areas of Shanghai.
 526 Estimates are based on the thin spline regression of the interval between the time of
 527 the detection of the first infection in each 3 km × 3 km grid and February 27, 2022.
 528 Triangles indicate the potential source of the outbreak. **(d)** Estimated speed of spread
 529 of SARS-CoV-2 (left axis) and cumulative fraction of affected areas of Shanghai
 530 (right axis). Red dots indicate the speed of spread over time in each cell. The blue line
 531 indicates the average speed per day as obtained using a polynomial regression. Central
 532 areas contain the districts of Jing'an, Yangpu, Hongkou, Putuo, Changning, Xuhui,
 533 and Huangpu.

534

535 **Figure 5. Characterization of the epidemic dynamics between March 16 and**

536 **March 29, 2022. (a)** Location of high-risk, moderate-risk, and low-risk areas. For
 537 each area, its highest risk classification was used. **(b-d)** Number of reported infections
 538 between March 16 and March 29 by area type. **(e)** Number of new reported infection
 539 per 1,000 individuals by area type. **(f)** Estimated R_t between March 16 and March 29
 540 by area type. **(g)** Estimated epidemic growth rate and doubling time (days).

541

542 **Figure 6. Epidemic dynamics under the effect of interventions. (a)** Number of new
 543 SARS-CoV-2 infections by date of sample collection for means of identification. **(b)**
 544 Estimated R_t (mean and 50% confidence interval) in eastern, western, and all
 545 Shanghai areas.

546 **Reference**

- 547 1. Pan A, Liu L, Wang C, et al. Association of Public Health Interventions
548 With the Epidemiology of the COVID-19 Outbreak in Wuhan, China. *Jama*
549 2020; **323**(19): 1915-23.
- 550 2. Luo L, Yang Z, Liang J, et al. Crucial control measures to contain China's
551 first Delta variant outbreak. *Natl Sci Rev* 2022; **9**(4): nwac004.
- 552 3. Li L, Han ZG, Qin PZ, et al. Transmission and containment of the
553 SARS-CoV-2 Delta variant of concern in Guangzhou, China: A
554 population-based study. *PLoS Negl Trop Dis* 2022; **16**(1): e0010048.
- 555 4. World Health Organization. Tracking SARS-CoV-2 variants. 2022.
- 556 5. Viana R, Moyo S, Amoako DG, et al. Rapid epidemic expansion of the
557 SARS-CoV-2 Omicron variant in southern Africa. *Nature* 2022; **603**(7902):
558 679-86.
- 559 6. Arora P, Zhang L, Rocha C, et al. Comparable neutralisation evasion of
560 SARS-CoV-2 omicron subvariants BA.1, BA.2, and BA.3. *The Lancet*
561 *Infectious Diseases*.
- 562 7. National Health Commission of the People's Republic of China. Update on
563 COVID-19. 2022.
- 564 8. Tu WJ, Zeng X, Liu Q. Aging tsunami coming: the main finding from
565 China's seventh national population census. *Aging Clin Exp Res* 2021.

- 566 9. McMenamin ME, Nealon J, Lin Y, et al. Vaccine effectiveness of two and
567 three doses of BNT162b2 and CoronaVac against COVID-19 in Hong Kong.
568 *medRxiv* 2022: 2022.03.22.22272769.
- 569 10. Suah JL, Tng BH, Tok PSK, et al. Real-world effectiveness of homologous
570 and heterologous BNT162b2, CoronaVac, and AZD1222 booster vaccination
571 against Delta and Omicron SARS-CoV-2 infection. *Emerg Microbes Infect*
572 2022; 11(1): 1343-5.
- 573 11. Cai J, Deng X, Yang J, et al. Modeling transmission of SARS-CoV-2
574 Omicron in China. *Nature Medicine* 2022.
- 575 12. Shanghai Municipal Health Commission. The 182th press conference of
576 COVID-19 in Shanghai. 2022.
- 577 13. 2020 Annual Report of Shanghai International Airport 2021.
- 578 14. Siu GK, Lee LK, Leung KS, et al. Will a new clade of SARS-CoV-2
579 imported into the community spark a fourth wave of the COVID-19 outbreak in
580 Hong Kong? *Emerg Microbes Infect* 2020; 9(1): 2497-500.
- 581 15. Gu H, Krishnan P, Ng DYM, et al. Probable Transmission of SARS-CoV-2
582 Omicron Variant in Quarantine Hotel, Hong Kong, China, November 2021.
583 *Emerg Infect Dis* 2022; 28(2): 460-2.
- 584 16. Zhang X, Zhang W, Chen S. Shanghai's life-saving efforts against the
585 current omicron wave of the COVID-19 pandemic. *The Lancet*.

- 586 17. Shanghai Municipal Health Commission. The 125th press conference of
587 COVID-19 in Shanghai. 2022.
- 588 18. Shanghai Municipal Health Commission. The 170th press conference of
589 COVID-19 in Shanghai. 2022.
- 590 19. Mefsin Y, Chen D, Bond HS, et al. Epidemiology of infections with
591 SARS-CoV-2 Omicron BA.2 variant in Hong Kong, January-March 2022.
592 *medRxiv* 2022: 2022.04.07.22273595.
- 593 20. Kraemer MUG, Reiner RC, Jr., Brady OJ, et al. Past and future spread of
594 the arbovirus vectors *Aedes aegypti* and *Aedes albopictus*. *Nat Microbiol* 2019;
595 4(5): 854-63.
- 596 21. Tisseuil C, Gryspeirt A, Lancelot R, Pioz M, Liebhold A, Gilbert M.
597 Evaluating methods to quantify spatial variation in the velocity of biological
598 invasions. *Ecography* 2015; 39.
- 599 22. Cori A, Ferguson NM, Fraser C, Cauchemez S. A new framework and
600 software to estimate time-varying reproduction numbers during epidemics. *Am*
601 *J Epidemiol* 2013; 178(9): 1505-12.
- 602 23. Shanghai Municipal Health Commission. The 123th press conference of
603 COVID-19 in Shanghai. 2022.
- 604 24. Yamasoba D, Kimura I, Nasser H, et al. Virological characteristics of the
605 SARS-CoV-2 Omicron BA.2 spike. *Cell*.

606 25. Li Z, Guan X, Mao N, et al. Antibody seroprevalence in the epicenter
607 Wuhan, Hubei, and six selected provinces after containment of the first
608 epidemic wave of COVID-19 in China. *Lancet Reg Health West Pac* 2021; 8:
609 100094.

610 26. Shanghai Municipal Health Commission. The 167th press conference of
611 COVID-19 in Shanghai. 2022.

612 27. Ranzani OT, Hitchings MDT, de Melo RL, et al. Effectiveness of an
613 Inactivated Covid-19 Vaccine with Homologous and Heterologous Boosters
614 against the Omicron (B.1.1.529) Variant. *medRxiv* 2022:
615 2022.03.30.22273193.

616 28. Jara A, Undurraga EA, Zubizarreta JR, et al. Effectiveness of CoronaVac
617 in children 3 to 5 years during the SARS-CoV-2 Omicron outbreak in Chile. *Nat*
618 *Med* 2022.

619 29. Garrett N, Tapley A, Andriesen J, et al. High Rate of Asymptomatic
620 Carriage Associated with Variant Strain Omicron. *medRxiv* 2022:
621 2021.12.20.21268130.


622 30. Poletti P, Tirani M, Cereda D, et al. Association of Age With Likelihood of
623 Developing Symptoms and Critical Disease Among Close Contacts Exposed
624 to Patients With Confirmed SARS-CoV-2 Infection in Italy. *JAMA Netw Open*
625 2021; 4(3): e211085.

- 626 31. Cohen C, Kleynhans J, von Gottberg A, et al. SARS-CoV-2 incidence,
627 transmission, and reinfection in a rural and an urban setting: results of the
628 PHIRST-C cohort study, South Africa, 2020-21. *Lancet Infect Dis* 2022.
- 629 32. Buitrago-Garcia D, Egli-Gany D, Counotte MJ, et al. Occurrence and
630 transmission potential of asymptomatic and presymptomatic SARS-CoV-2
631 infections: A living systematic review and meta-analysis. *PLoS Med* 2020;
632 17(9): e1003346.
- 633 33. Hay JA, Kissler SM, Fauver JR, et al. Viral dynamics and duration of PCR
634 positivity of the SARS-CoV-2 Omicron variant. *medRxiv* 2022:
635 2022.01.13.22269257.
- 636 34. Zhang X, Wu J, Smith LM, et al. Monitoring SARS-CoV-2 in air and on
637 surfaces and estimating infection risk in buildings and buses on a university
638 campus. *J Expo Sci Environ Epidemiol* 2022: 1-8.
639

Phase 1: January 1 to February 28


Baseline non-pharmaceutical interventions (NPIs)

Border control policy




January 1, 2022 February 28, 2022

Local control policy




Phase 2: March 1 to March 31

Additional NPIs





Phase 3: April 1 to May 31

Additional NPIs



Border control policy
 Policy of case isolation
 Mass screening policy
 Social distancing policy




Symptomatic status and sources

- Local asymptomatic infections
- Local symptomatic cases
- Imported asymptomatic infections
- Imported symptomatic cases


a




b




c





d




Cumulative number
of infections per
1,000 individuals






a


Types of subdistrict

**b**


High-risk areas

**c**




Moderate-risk areas

**d**

Low-risk areas



No. of total infections per 1,000 individuals between March 16 and 29

**e****f****g**