

# Roelfs et al. | Genetics of the brain functional connectome and mental health profiles

## Shared genetic architecture between mental health and the brain functional connectome in the UK Biobank

Daniel Roelfs<sup>1,\*</sup>, Oleksandr Frei<sup>1</sup>, Dennis van der Meer<sup>1,2</sup>, Elleke Tisink<sup>3</sup>, Alexey Shadrin<sup>1</sup>, Dag Alnæs<sup>1,4</sup>, Ole A. Andreassen<sup>1</sup>, Lars T. Westlye<sup>1,5</sup>, Tobias Kaufmann<sup>1,6\*</sup>

<sup>1</sup> NORMENT, KG Jebsen Centre for Neurodevelopmental Disorders, Division of Mental Health and Addiction, Oslo University Hospital & Institute of Clinical Medicine, University of Oslo, Oslo, Norway

<sup>2</sup> School of Mental Health and Neuroscience, Faculty of Health, Medicine, and Life Sciences, Maastricht University, Maastricht, the Netherlands

<sup>3</sup> Department of Complex Trait Genetics, Center for Neurogenomics and Cognitive Research, Vrije Universiteit Amsterdam, Amsterdam Neuroscience, 1081 HV Amsterdam, The Netherlands

<sup>4</sup> Bjørknes College, Oslo, Norway

<sup>5</sup> Department of Psychology, University of Oslo, Oslo, Norway

<sup>6</sup> Department of Psychiatry and Psychotherapy, Tübingen Center for Mental Health, University of Tübingen, Germany

\* Correspondence: Daniel Roelfs & Tobias Kaufmann, PhD.  
Email: [daniel.roelfs@medisin.uio.no](mailto:daniel.roelfs@medisin.uio.no), [tobias.kaufmann@medisin.uio.no](mailto:tobias.kaufmann@medisin.uio.no)  
Postal address: OUS, PO Box 4956 Nydalen, 0424 Oslo, Norway  
Telephone: +47 23 02 73 50, Fax: +47 23 02 73 33

Counts: Main: 2458 words | Abstract: 138 words | Figures: 2 | References: 43

## Roelfs et al. | Genetics of the brain functional connectome and mental health profiles

### **Abstract**

Psychiatric disorders are complex clinical conditions with large heterogeneity and overlap in symptoms, genetic liability and brain imaging abnormalities. Building on a dimensional conceptualization of mental health, previous studies have reported genetic overlap between psychiatric disorders and population-level mental health, and between psychiatric disorders and brain functional connectivity. Here, in 30,701 participants aged 45-82 from the UK Biobank we map the genetic associations between self-reported mental health and resting-state fMRI-based measures of brain network function. Multivariate Omnibus Statistical Test revealed 10 genetic loci associated with population-level mental symptoms. Next, conjunctive FDR identified 23 shared genetic variants between these symptom profiles and fMRI-based brain network measures. Functional annotation implicated genes involved in brain structure and function, in particular synaptic processes. These findings provide further genetic evidence of an association between brain function and mental health traits in the population.

## Introduction

Psychiatric disorders are complex, with a polygenic architecture, and large degree of overlapping symptoms and risk factors. Both imaging and genetics studies have shown numerous but small associations between brain phenotypes, psychiatric disorders, and genetics, such as schizophrenia (SCZ)<sup>1-3</sup>, bipolar disorder (BIP)<sup>4,5</sup>, major depressive disorder (MDD)<sup>6,7</sup>, and anxiety disorder (ANX)<sup>8-10</sup>. Interactions between various brain phenotypes and genetics have been reported across structural<sup>11,12</sup> and functional<sup>2,7,10</sup> imaging modalities.

We have recently deployed a multivariate analysis to study the genetic architecture of brain functional connectivity, revealing genetic variants associated with functional brain connectivity as well as variance in brain activity over time<sup>13</sup>. The results showed meaningful overlap with psychiatric disorders, pointing at synapse-related pathways among the biological processes shared between disorders and brain function<sup>13</sup>.

Previous studies have shown widespread phenotypic and genetic overlap between psychiatric disorders<sup>14-18</sup>. In addition, patients within a diagnostic category can display a wide variety of symptoms. This heterogeneity complicates both diagnosis and therapeutic response due to overlapping symptoms and generally low specificity of diagnostic features<sup>19,20</sup>. While the mental health of any individual in the population varies over the course of a lifetime, most will not meet diagnostic criteria for a psychiatric disorder<sup>21,22</sup>. In order to capture the variance encompassing psychiatric symptoms that is lacking in traditional case-control studies, one can use population-level mental health questionnaires as implemented in the UK Biobank<sup>23</sup>. This facilitates analyses using the continuous scales which enable data-driven clustering methods to extract different profiles each capturing a separate domain relevant to mental health in a sample without individuals diagnosed with a psychiatric disorder, taking advantage of larger sample sizes. Using independent component analysis (ICA), we have previously derived 13 mental health profiles from UK Biobank data, and showed that, although phenotypically independent (by design) they nonetheless share genetic underpinnings<sup>24</sup>.

Here, we aimed to uncover the genetic architecture of mental symptoms and identify shared genetic loci with neurobiological processes related to brain function. Using multivariate analysis<sup>25</sup>, we generated multivariate genome-wide association statistics across our previously identified 13 population-level mental health profiles<sup>24</sup>. This allowed us to identify new gene variants associated with mental health symptoms and traits such as psychosis, depression, and anxiety in the UK Biobank sample not captured in a univariate analysis. Further, we combined this multivariate genetic profile of mental health with GWAS summary statistics of 7 psychiatric disorders and with our previously identified multivariate profiles of

## Roelfs et al. | Genetics of the brain functional connectome and mental health profiles

functional brain connectivity and variance in brain activity over time<sup>24</sup>. This research aims to provide insight into the biological underpinnings of mental health symptoms.

### Methods

#### *Sample and exclusion criteria*

We utilized data from the UK Biobank<sup>26</sup> with permission no. 27412. All participants provided signed informed consent before inclusion in the study. The UK Biobank was approved by the National Health Service National Research Ethics Service (ref. 11/NW/0382). We previously used data from the online follow-up questionnaire on mental health to define 13 phenotypically independent profiles relevant for mental health<sup>24</sup>. In addition, we also utilized imaging data provided by the UK Biobank, the procedure of processing and analyzing the imaging data is described a previous paper<sup>13</sup>. For this study we deployed the summary statistics from these previous studies.

#### *Image acquisition and pre-processing*

The processing pipeline for imaging data used for the multivariate GWAS is described in Roelfs *et al.*<sup>13</sup>. In short, images were acquired using 3T Siemens Magnetom Skyra scanners with a 32 channel head coil (Siemens Healthcare GmbH, Erlangen, Germany) at four different sites in the UK. The fMRI data was recorded using a gradient-echo echo planar imaging sequence with x8 multislice acceleration (TR: 0.735s, TE: 39ms, FOV: 88×88×64 matrix, FA: 52°) with a voxel size of 2.4×2.4×2.4mm. Data is processed by the UK Biobank team following the protocol described in Alfaro-Almagro *et al.*<sup>27</sup>.

#### *Multivariate Genome-Wide Analysis*

In this study we applied the Multivariate Omnibus Statistical Test (MOSTest) to the phenotypic data from the ICA decomposition described in Roelfs *et al.*<sup>24</sup>. MOSTest deploys the univariate test-statistics for each SNP and computes a multivariate test statistic through single random permutations of the genotype vector. This method is described in Van der Meer *et al.*<sup>25</sup>. We performed positional gene mapping using Functional Mapping and Annotation (FUMA)<sup>28</sup>. We also used a built-in tool to follow up these gene mapping analyses using MAGMA to connect the identified genes with tissue types<sup>29</sup>. We analyzed gene sets using the reactome toolbox<sup>30</sup> to identify biological processes associated with the genes associated with the summary statistics identified by FUMA.


#### *Pleiotropy-informed conjunctive false discovery rate*

In order to quantify the degree of genetic overlap and identify shared genetic loci between the mental health profiles and the imaging features we deployed the pleiotropy-informed conjunctive false discovery rate

(conjFDR) through the pleioFDR toolbox<sup>31</sup>. One of the advantages of conjFDR is that it can identify shared genetic loci regardless of effect direction and effect size, a feature that is useful when working with multivariate measures where effect direction might be lacking.

## Results

MOSTest revealed 10 significant loci across the 13 previously identified profiles of mental health (Figure 1). FUMA and its positional mapping tool revealed 48 genes associated with these loci (See Suppl. Table 1), that were linked by MAGMA to a number of brain structures such as the cerebellum and amygdala (Suppl. Figure 1). Among the identified genes, those mapped from the strongest GWAS loci were *ADH1B* and *ADH5* (chromosome 4) and *CRHR1* (chromosome 17).



**Figure 1. Manhattan plot of multivariate genome-wide association statistics for mental health.** Manhattan plot showing the multivariate genome-wide association of our multivariate measure of mental health. We identified 10 loci associated with the multivariate genome-wide association statistics for mental health.

In order to identify shared genomic loci between the mental health profiles and psychiatric disorders, we used GWAS summary statistics from prior case-control studies including schizophrenia (SCZ)<sup>1</sup>, bipolar disorders (BIP)<sup>4</sup>, major depression (MD)<sup>6</sup>, attention-deficit hyperactivity disorder (ADHD)<sup>32</sup>, autism spectrum disorder (ASD)<sup>33</sup>, post-traumatic stress disorder (PTSD)<sup>34</sup>, and anxiety (ANX)<sup>8</sup>, see also Suppl. Table 2. First, we compared the gene set from the multivariate mental health genome-wide association statistics with the gene set from each of the psychiatric disorders. Here we found 35 overlapping genes, 29

# Roelfs et al. | Genetics of the brain functional connectome and mental health profiles


with SCZ, 7 with BIP, and 1 with ADHD (see Suppl. Table 3). It is important to note that the 7 overlapping genes with BIP were mapped from only 2 separate loci. Next, we extracted the loci from each case-control GWAS (202 in total) and assessed whether each locus was significant in the multivariate genome-wide association statistics for mental health profiles as well. Of the 202 loci significant in any of the disorders, one showed genome-wide significance at  $P < 5e-8$  and 122 showed nominal significance at  $P < 0.05$  only in the multivariate genome-wide association statistics for mental health profiles, potentially indicating some shared but small effects.

Next, we explored the genetic overlap between the mental health profiles and the psychiatric disorders through the conjunctive false discovery rate (conjFDR)<sup>31,35</sup> which leverages pleiotropy between two phenotypes to estimate shared genetic determinants. ConjFDR allows for the discovery of shared genetic determinants even when those loci are not genome-wide significant in either of the traits in the analysis. Through conjFDR we identified 35 overlapping loci in total between the multivariate genome-wide association statistics for mental health profiles and psychiatric disorders. We found 10 overlapping loci between the multivariate measure of mental health profiles and BIP, 8 overlapping loci with both MD and ADHD, 5 overlapping loci with SCZ, and 4 overlapping loci with autism (see Suppl. Figure 2). FUMA identified 89 genes associated with these loci (See Suppl. Table 3). We found no overlapping loci between the mental health profiles and ANX or PTSD, which may be related to the limited power in these GWASs (see Suppl. Table 2).

We then calculated the number of shared genetic loci between the multivariate genome-wide association statistics for mental health profiles and the two multivariate measures of the brain functional connectome using conjFDR. We used the GWAS summary statistics from our previous study of brain function<sup>13</sup>. In contrast to the prior study in which we identified genetic overlap between brain function and psychiatric disorders, we here investigated overlaps with the multivariate genome-wide association statistics for mental health to investigate if this approach captures associations not revealed through case-control GWAS approaches. Figure 2 shows two Manhattan plots of the conjunctive FDR analyses between both functional connectivity and node variance with the multivariate GWAS on the multivariate genome-wide association statistics for mental health profiles. Genetic signal was adequate (See Suppl. Figure 3). The multivariate genome-wide association statistics for mental health profiles shared 18 loci with functional connectivity and 5 with node variance. A full list of genes associated with the (in total) 23 unique shared loci between the multivariate summary statistics and FC is presented in Suppl. Table 5. The number of overlapping loci between the brain functional connectome and the multivariate genome-wide association statistics for mental health profiles (18 for FC, 5 for node variance) was generally larger than the number of shared loci between the brain functional connectome and the psychiatric disorders (with the exception of SCZ) identified in our

## Roelfs et al. | Genetics of the brain functional connectome and mental health profiles

previous study<sup>13</sup>. When we mapped the genes from these loci using FUMA and tested for enrichment in gene-sets using the reactome toolbox<sup>30</sup> we found that the genes associated with these shared loci are involved in a number of neurobiologically relevant processes such as axonal growth regulation (NGFR and RHOA) and regulation of transcription factors related through MECP2 (MEF2C, see Suppl. Table 6).



## Discussion

In this study we identified a number of loci associated with multivariate genome-wide association statistics for mental health profiles and found overlapping loci with the measures of brain function and psychiatric disorders. Using MOSTest we were able to leverage the phenotypic overlap between different mental health profiles to identify new loci associated with a multivariate measure of mental health. Genes associated with these loci showed regional expression in different parts of the brain (e.g. cerebellum, amygdala).

Our analysis using conjFDR revealed a number of shared loci and genes between the multivariate genome-wide association statistics and the psychiatric disorders. This demonstrates the shared genetics between psychiatric symptoms regardless of clinical diagnosis, emphasizes the utility of using population-level phenotypes to investigate variance in mental health profiles, and highlights the advantage in



# Roelfs et al. | Genetics of the brain functional connectome and mental health profiles

leveraging pleiotropy between complex phenotypes to boost discovery. We found shared genes with all but two case-control GWAS (ANX, PTSD), which also had the two smallest sample sizes, which may reflect insufficient power to detect an effect<sup>36</sup>, or may indicate the absence of an effect with those disorders. The largest overlap was with SCZ, which shared 29 genes in the geneset with the multivariate measure. Future sample increases in the case-control GWAS may reveal shared genetics with other complex traits, including population based mental health phenotypes and brain imaging features.

We also identified a number of overlapping loci between mental health profiles and fMRI measures of brain function, including 18 shared loci with functional connectivity and 5 shared loci with node variance. The higher number of shared loci for functional connectivity might be partially explained by the number of phenotypes in each composite measure. While the functional connectivity GWAS comprises 210 measures, i.e. partial correlations between 21 brain nodes, the node variance GWAS encompasses only the temporal variance in each node. It is possible that the number of phenotypes included in the multivariate analysis can affect the discovery<sup>25</sup>. Both the functional connectivity and node variance summary statistics had the same sample size (N=30.701). For our analyses this means that the difference in their overlap with the multivariate genome-wide association statistics for mental health is due to either the discrepancy in the number of features contained within the composite measure, or alternatively because of different biological processes underlying both measures. The measures differ in that functional connectivity refers to the correlation between brain networks (edge strength), which is possibly governed by different processes than the temporal variance in activity within brain networks. Overall, we found a number of genes associated with the shared loci that are involved in biologically relevant processes such as axonal growth and energy transport (See Suppl. Table 6). Although more thorough functional analysis is necessary, this could suggest that axonal growth processes is a shared feature between brain connectivity and mental disorders, which would be in line with previous evidence linking axonal growth with both processes independently<sup>37-39</sup>.

The two conjunctive analyses with fMRI measures and the multivariate genome-wide association statistics for mental health each showed a number of overlapping loci. Not all shared loci were unique, this can be partially explained by the definition of the brain networks in our analyses. The functional connectivity and the node variance measures use the same 21 nodes, and, ultimately, the two measures reflect different properties of the same time series. We found that the number of overlapping loci between the multivariate genome-wide association statistics for mental health and the brain functional connectome was generally larger than findings from our previous study highlighting shared genetic loci between psychiatric disorders and the brain functional connectome. This may partly be due to the larger sample size of the multivariate measure of mental health, but it could also reflect that the multivariate genome-



## Roelfs et al. | Genetics of the brain functional connectome and mental health profiles

wide association statistics capture genetic variance more generally related to the brain functional connectome. We found little direct overlap between loci of these two GWAS' separately, which highlights the discovery boost advantage of using conjFDR in phenotypes with generally low heritability.

The main implication from our findings is that we can identify shared genetic variants between a multifactorial measure of mental health in an undiagnosed population sample and fMRI-based measures of brain functional connectivity. Several limitations should be considered. First, the data were obtained from a middle-aged and older White British population, which limits the generalizability of the findings. Further, the mental health questionnaires are self-administered, so the data is vulnerable to various response and self-selection biases<sup>40</sup>. We excluded individuals with a psychiatric diagnosis in our independent component analysis in order to maximize the population variance and to mitigate the influence of a smaller number of individuals with a (diagnosed) psychiatric condition. This results in a healthier sample, lower variance on a number of severe symptom domains and possible survivor biases<sup>24</sup>. This bias also extends to the imaging data, where participants are reportedly healthier than the general population<sup>41</sup>. Further, MOSTest currently lacks effect direction. This complicates further analyses such as genetic correlations that require reliable effect directions. Nonetheless, FUMA and MAGMA revealed brain structures associated with these mapped genes, such as the cerebellum, amygdala, and various parts of the cortex, which have been linked to psychiatric disorders and symptoms. To what degree these shared genes can explain shared clinical characteristics such as symptoms is an important and relevant issue that needs to be answered in future studies.

In conclusion, our multivariate GWAS on 13 mental health symptom profiles showed a number of shared genetic loci with two fMRI-measures reflecting brain function and connectivity. This provides further genetic evidence of an association between brain function and mental health traits in the population.

## References

1. Pardiñas, A. F. *et al.* Common schizophrenia alleles are enriched in mutation-intolerant genes and in regions under strong background selection. *Nat. Genet.* **50**, 381–389 (2018).
2. Pettersson-Yeo, W., Allen, P., Benetti, S., McGuire, P. & Mechelli, A. Dysconnectivity in schizophrenia: where are we now? *Neurosci. Biobehav. Rev.* **35**, 1110–1124 (2011).
3. Trubetskoy, V. *et al.* Mapping genomic loci implicates genes and synaptic biology in schizophrenia. *Nature* **604**, 502–508 (2022).

# Roelfs et al. | Genetics of the brain functional connectome and mental health profiles

4. Mullins, N. *et al.* Genome-wide association study of more than 40,000 bipolar disorder cases provides new insights into the underlying biology. *Nat. Genet.* **53**, 817–829 (2021).
5. Vai, B., Bertocchi, C. & Benedetti, F. Cortico-limbic connectivity as a possible biomarker for bipolar disorder: where are we now? *Expert Rev. Neurother.* **19**, 159–172 (2019).
6. Wray, N. R. *et al.* Genome-wide association analyses identify 44 risk variants and refine the genetic architecture of major depression. *Nat. Genet.* **50**, 668–681 (2018).
7. Zhuo, C. *et al.* The rise and fall of MRI studies in major depressive disorder. *Transl. Psychiatry* **9**, 335 (2019).
8. Otowa, T. *et al.* Meta-analysis of genome-wide association studies of anxiety disorders. *Mol. Psychiatry* **21**, 1391–1399 (2016).
9. Stein, M. B. Neurobiology of generalized anxiety disorder. *J. Clin. Psychiatry* **70 Suppl 2**, 15–19 (2009).
10. Thompson, P. M. *et al.* ENIGMA and global neuroscience: A decade of large-scale studies of the brain in health and disease across more than 40 countries. *Transl. Psychiatry* **10**, 100 (2020).
11. Cheng, W. *et al.* Genetic Association Between Schizophrenia and Cortical Brain Surface Area and Thickness. *JAMA Psychiatry* **78**, 1020–1030 (2021).
12. Smeland, O. B. *et al.* Genetic Overlap Between Schizophrenia and Volumes of Hippocampus, Putamen, and Intracranial Volume Indicates Shared Molecular Genetic Mechanisms. *Schizophr. Bull.* **44**, 854–864 (2018).
13. Roelfs, D. *et al.* Genetic overlap between multivariate measures of human functional brain connectivity and psychiatric disorders. *medRxiv* 2021.06.15.21258954 (2021) doi:10.1101/2021.06.15.21258954.
14. Anttila, V. *et al.* Analysis of shared heritability in common disorders of the brain. *Science* **360**, eaap8757 (2018).
15. Bulik-Sullivan, B. *et al.* An atlas of genetic correlations across human diseases and traits. *Nat Genet* **47**, 1236–41 (2015).

# Roelfs et al. | Genetics of the brain functional connectome and mental health profiles

16. Cross-Disorder Group of the Psychiatric Genomics Consortium. Identification of risk loci with shared effects on five major psychiatric disorders: a genome-wide analysis. *The Lancet* **381**, 1371–1379 (2013).
17. Lee, P. H. *et al.* Genomic Relationships, Novel Loci, and Pleiotropic Mechanisms across Eight Psychiatric Disorders. *Cell* **179**, 1469–1482.e11 (2019).
18. Romero, C. *et al.* Exploring the genetic overlap between 12 psychiatric disorders. *medRxiv* 2022.04.12.22273763 (2022) doi:10.1101/2022.04.12.22273763.
19. Wardenaar, K. J. & de Jonge, P. Diagnostic heterogeneity in psychiatry: towards an empirical solution. *BMC Med.* **11**, 201 (2013).
20. Widiger, T. A. & Clark, L. A. Toward DSM—V and the classification of psychopathology. *Psychol. Bull.* **126**, 946–963 (2000).
21. Kessler, R. C. *et al.* Lifetime Prevalence and Age-of-Onset Distributions of DSM-IV Disorders in the National Comorbidity Survey Replication. *Arch. Gen. Psychiatry* **62**, 593–602 (2005).
22. McGrath, J. J. *et al.* Psychotic Experiences in the General Population: A Cross-National Analysis Based on 31,261 Respondents From 18 Countries. *JAMA Psychiatry* **72**, 697–705 (2015).
23. Doherty, J. L. & Owen, M. J. Genomic insights into the overlap between psychiatric disorders: implications for research and clinical practice. *Genome Med.* **6**, 29–29 (2014).
24. Roelfs, D. *et al.* Phenotypically independent profiles relevant to mental health are genetically correlated. *Transl. Psychiatry* **11**, 202 (2021).
25. van der Meer, D. *et al.* Understanding the genetic determinants of the brain with MOSTest. *Nat. Commun.* **11**, 3512 (2020).
26. Sudlow, C. *et al.* UK biobank: an open access resource for identifying the causes of a wide range of complex diseases of middle and old age. *PLoS Med* **12**, e1001779 (2015).
27. Alfaro-Almagro, F. *et al.* Image processing and Quality Control for the first 10,000 brain imaging datasets from UK Biobank. *NeuroImage* **166**, 400–424 (2018).

# Roelfs et al. | Genetics of the brain functional connectome and mental health profiles

28. Watanabe, K., Taskesen, E., van Bochoven, A. & Posthuma, D. Functional mapping and annotation of genetic associations with FUMA. *Nat. Commun.* **8**, 1826 (2017).
29. de Leeuw, C. A., Mooij, J. M., Heskes, T. & Posthuma, D. MAGMA: Generalized Gene-Set Analysis of GWAS Data. *PLOS Comput. Biol.* **11**, e1004219 (2015).
30. Jassal, B. *et al.* The reactome pathway knowledgebase. *Nucleic Acids Res.* **48**, D498–D503 (2020).
31. Andreassen, O. A. *et al.* Improved detection of common variants associated with schizophrenia and bipolar disorder using pleiotropy-informed conditional false discovery rate. *PLoS Genet* **9**, e1003455 (2013).
32. Demontis, D. *et al.* Discovery of the first genome-wide significant risk loci for attention deficit/hyperactivity disorder. *Nat. Genet.* **51**, 63–75 (2019).
33. Grove, J. *et al.* Identification of common genetic risk variants for autism spectrum disorder. *Nat. Genet.* **51**, 431–444 (2019).
34. Duncan, L. E. *et al.* Largest GWAS of PTSD (N=20 070) yields genetic overlap with schizophrenia and sex differences in heritability. *Mol. Psychiatry* **23**, 666–673 (2018).
35. Smeland, O. B. *et al.* Discovery of shared genomic loci using the conditional false discovery rate approach. *Hum. Genet.* **139**, 85–94 (2020).
36. Visscher, P. M. *et al.* 10 Years of GWAS Discovery: Biology, Function, and Translation. *Am. J. Hum. Genet.* **101**, 5–22 (2017).
37. Devor, A. *et al.* Genetic evidence for role of integration of fast and slow neurotransmission in schizophrenia. *Mol. Psychiatry* **22**, 792–801 (2017).
38. Hsu, W.-C. J., Nilsson, C. L. & Laezza, F. Role of the axonal initial segment in psychiatric disorders: function, dysfunction, and intervention. *Front. Psychiatry* **5**, 109–109 (2014).
39. Mukai, J. *et al.* Molecular substrates of altered axonal growth and brain connectivity in a mouse model of schizophrenia. *Neuron* **86**, 680–695 (2015).
40. Dutt, R. K. *et al.* Mental health in the UK Biobank: A roadmap to self-report measures and neuroimaging correlates. *Hum. Brain Mapp.* **43**, 816–832 (2022).

# Roelfs et al. | Genetics of the brain functional connectome and mental health profiles

41. Lyall, D. M. *et al.* Quantifying bias in psychological and physical health in the UK Biobank imaging sub-sample. *Brain Commun.* **4**, fcac119 (2022).
42. Stahl, E. A. *et al.* Genome-wide association study identifies 30 loci associated with bipolar disorder. *Nat. Genet.* **51**, 793–803 (2019).
43. Duncan, L. E. *et al.* Largest GWAS of PTSD (N=20 070) yields genetic overlap with schizophrenia and sex differences in heritability. *Mol. Psychiatry* **23**, 666–673 (2018).

## Acknowledgements

The authors were funded by the Research Council of Norway (#276082 LifespanHealth, #223273 NORMENT, #283798 ERA-NET Neuron SYNSCHIZ, #249795), the South-East Norway Regional Health Authority (2019101, 2019107, and 2020086), and the European Research Council under the European Union’s Horizon2020 Research and Innovation program (ERC Starting Grant #802998), as well as the Horizon2020 Research and Innovation Action Grant CoMorMent (#847776). This research has been conducted using the UK Biobank Resource (access code 27412, <https://www.ukbiobank.ac.uk/>). E.T. has been supported by the Foundation “De Drie Lichten” and The Simons Foundation Fund in The Netherlands. This work was performed on the TSD (Tjenester for Sensitive Data) facilities, owned by the University of Oslo, operated and developed by the TSD service group at the University of Oslo, IT-Department (USIT). Computations were also performed on resources provided by UNINETT Sigma2 - the National Infrastructure for High Performance Computing and Data Storage in Norway.

## Conflicts of interest

D.R., D.A., O.F., D.vd.M., O.B.S., L.T.W. and T.K. declare no conflicts of interest. O.A.A. is a consultant to HealthLytix and received speakers honorarium from Lundbeck.

## Author contributions

D.R. and T.K. conceived the study; D.R. analyzed the data with contributions from T.K.; All authors contributed with conceptual input on methods and/or interpretation of results; D.R. and T.K. wrote the first draft of the paper and all authors contributed to the final manuscript.

## Data availability

All data used in this study are part of the publicly available UK Biobank initiative (<https://www.ukbiobank.ac.uk/>). Summary statistics for the disorders are publicly available through their


## Roelfs et al. | Genetics of the brain functional connectome and mental health profiles

respective consortia. The summary statistics for the multivariate analyses will be shared on GitHub upon acceptance.


### *Code availability*

Code will be made publicly available via GitHub (<https://www.github.com/norment/open-science>) upon acceptance of the manuscript.

Supplementary Figures








# Roelfs et al. | Genetics of the brain functional connectome and mental health profiles

the largest number of shared loci between the multivariate summary statistics and SCZ (29 loci), BIP (10 loci), followed by ADHD and MDD (both 8 loci).



**Suppl. Figure 3. QQ plots of the genetic signal of the conjFDR between the multivariate genome-wide association statistics for mental health and the brain functional connectome** Figure showing no inflation of signal in the conjFDR at different thresholds. Genetic signal overall was not very strong, but sufficient for the analyses in this pipeline.

## Supplementary Tables

<b>Suppl. Table 1. Mapped genes from the multivariate genome-wide association statistics for mental health</b> Genes mapped by FUMA on the summary statistics from the multivariate summary statistics. We found in total 48 genes associated with the genetic signal of the multivariate measure.				
<i>No</i>	<i>Gene</i>	<i>Chr</i>	<i>Pmin</i>	<i>Ind. Sig. SNP</i>
1	CRB1	1	2.128e-07	rs17641524
2	DENND1B	1	1.060e-08	rs17641524
3	BSN	3	2.759e-04	rs3749237
4	MST1	3	2.871e-08	rs3749237
5	RNF123	3	2.871e-08	rs3749237
6	AMIGO3	3	1.213e-08	rs3749237
7	GMPPB	3	1.213e-08	rs3749237
8	IP6K1	3	1.213e-08	rs3749237
9	CDHR4	3	2.898e-08	rs3749237
10	FAM212A	3	4.738e-08	rs3749237
11	UBA7	3	4.738e-08	rs3749237
12	TRAIIP	3	2.802e-08	rs3749237, rs4688686
13	CAMKV	3	2.802e-08	rs3749237, rs4688686
14	MST1R	3	6.717e-08	rs4688686
15	CTD-2330K9.3	3	6.717e-08	rs4688686
16	MON1A	3	1.294e-07	rs4688686
17	RBM6	3	8.472e-08	rs4688686
18	RBM5	3	6.544e-08	rs4688686
19	SEMA3F	3	5.478e-07	rs4688686
20	KLB	4	4.007e-10	rs35538052
21	METAP1	4	2.364e-08	rs146788033
22	ADH5	4	2.680e-19	rs551676206
23	ADH4	4	3.073e-03	rs146788033
24	ADH1B	4	5.678e-41	rs1229984, rs551676206
25	C4orf17	4		rs551676206
26	TRMT10A	4		rs551676206

<b>Suppl. Table 1. Mapped genes from the multivariate genome-wide association statistics for mental health</b>				
Genes mapped by FUMA on the summary statistics from the multivariate summary statistics. We found in total 48 genes associated with the genetic signal of the multivariate measure.				
<i>No</i>	<i>Gene</i>	<i>Chr</i>	<i>P<sub>min</sub></i>	<i>Ind. Sig. SNP</i>
27	BANK1	4	5.882e-09	rs13107325
28	SLC39A8	4	7.204e-11	rs13107325, rs13105581
29	IRF4	6	2.298e-08	rs12203592
30	PRRX2	9	2.093e-09	rs28428946
31	DCAF5	14	2.136e-07	rs35538328
32	EXD2	14	2.447e-06	rs35538328
33	ARHGAP27	17	1.158e-09	rs62055718
34	PLEKHM1	17	1.534e-13	rs62055718
35	CRHR1	17	2.897e-14	rs62055718, rs199440, rs55938136, rs7224296, rs12938031
36	SPPL2C	17	4.195e-13	rs62055718, rs199440, rs7224296, rs12938031
37	MAPT	17	1.256e-13	rs62055718, rs199440, rs7224296
38	STH	17	1.437e-13	rs62055718, rs199440, rs7224296
39	KANSL1	17	7.431e-14	rs62055718, rs199440, rs7224296
40	ARL17B	17	3.191e-14	rs62055718, rs199440, rs7224296
41	LRRC37A	17	5.397e-13	rs62055718, rs199440, rs7224296
42	LRRC37A2	17		rs62055718, rs199440, rs7224296
43	ARL17A	17		rs62055718, rs199440, rs7224296
44	NSF	17	3.552e-12	rs62055718, rs199440, rs7224296, rs199470
45	WNT3	17	7.193e-12	rs199440, rs7224296
46	PSMG1	21	3.415e-07	rs7281463
47	BRWD1	21	2.851e-07	rs7281463
48	HMG1	21	2.425e-06	rs7281463

372

<b>Suppl. Table 2. Overview of the genetic association analyses on psychiatric conditions used in this study</b>								
List of GWASs we included in the genetic analyses in this paper and their meta data. Sample size in the table and in the summary statistics used in the final analyses may differ since we excluded individuals already included in the UK Biobank and included only individuals with White European ancestry.								
<i>Pheno-type</i>	<i>Consortium</i>	<i>Sample</i>	<i>Citation</i>	<i>N<sub>case</sub></i>	<i>N<sub>control</sub></i>	<i>Number of loci</i>	<i>Lambda<sub>GC</sub></i>	<i>h<sup>2</sup> (se)</i>

Roelfs et al. | Genetics of the brain functional connectome and mental health profiles

SCZ	PGC	Meta analysis of European individuals in PGC and CLOZUK samples	Pardiñas et al., 2018	40,675	64,643	135	1.6831	0.41 (0.01)
BIP	PGC	European individuals	Mullins et al., 2021	41,917	371,549	44	1.4316	0.0853 (<0.01)
MDD	PGC	European individuals	Wray et al., 2018	69,576	161,613	2	1.1973	0.0717 (<0.01)
ADHD	PGC	European individuals	Demontis et al., 2019	19,099	34,194	12	1.2531	0.2354 (0.02)
ASD	PGC, iPSYCH	Meta analysis of European individuals	Grove et al., 2019	18,381	27,969	2	1.1715	0.1941 (0.02)
PTSD	PGC	European individuals	Duncan et al., 2018	2,424	7,113	0	1.0165	0.101 (0.04)
ANX	ANGST	Meta analysis of European individuals	Otowa et al., 2016	7,016	14,745	1	1.0345	0.0763 (0.03)

373

**Suppl. Table 3. Number of overlapping genes in gene sets with the multivariate genome-wide association statistics for mental health (MMMH)** Table showing the number of overlapping genes identified through either overlap in gene sets, the number of genes from the disorder GWAS alone, and the number of shared genes identified through conjFDR.

<i>Disorder</i>	<i>Measure</i>	<i>Number of overlapping genes in gene sets</i>	<i>Number of genes in disorder GWAS alone</i>	<i>Number of shared genes identified through cFDR</i>
ADHD	MMMH	1	22	37
ANX	MMMH	0	0	0
ASD	MMMH	0	6	16
BIP	MMMH	7	184	28
MDD	MMMH	0	1	29
PTSD	MMMH	0	0	0
SCZ	MMMH	27	600	72

374

375

**Suppl. Table 4. List of genes associated with the shared genetic determinants between the psychiatric disorders and multivariate genome-wide association statistics for mental health** Genes associated with conjFDR output from diagnosis and multivariate genome-wide association statistics for mental health. This list only includes psychiatric disorders for which we identified significant loci. The number of mapped genes can be greater than the number of discovered loci since loci can be associated with more than one gene.

<i>Diagnosis</i>	<i>Measure</i>	<i>No</i>	<i>Gene</i>	<i>Chr</i>	<i>Pmin</i>	<i>Ind. Sig. SNP</i>
SCZ	MMMH	1	BARHL2	1	2.568e-07	rs12046000
SCZ	MMMH	2	CRB1	1	4.704e-07	rs10922218
SCZ	MMMH	3	DENND1B	1	1.907e-06	rs10922218
SCZ	MMMH	4	FANCL	2	1.892e-06	rs2125924
SCZ	MMMH	5	BCL11A	2	4.375e-07	rs13019832
SCZ	MMMH	6	XYLB	3	3.220e-07	rs151618
SCZ	MMMH	7	EXOG	3	3.220e-07	rs151618
SCZ	MMMH	8	CCDC71	3	1.795e-07	rs2624847
SCZ	MMMH	9	GPX1	3	1.795e-07	rs2624847
SCZ	MMMH	10	AMT	3	1.795e-07	rs2624847
SCZ	MMMH	11	FAM212A	3	1.795e-07	rs2624847
SCZ	MMMH	12	MST1R	3	1.795e-07	rs2624847
SCZ	MMMH	13	RBM6	3	1.795e-07	rs2624847
SCZ	MMMH	14	FOXP1	3	2.086e-07	rs60756930
SCZ	MMMH	15	BANK1	4	1.651e-10	rs13107325
SCZ	MMMH	16	SLC39A8	4	3.472e-12	rs13107325
SCZ	MMMH	17	HCN1	5	1.601e-06	rs10941692
SCZ	MMMH	18	ZCCHC7	9	4.373e-07	rs308523
SCZ	MMMH	19	RTKN2	10	2.944e-07	rs2306970
SCZ	MMMH	20	ZNF365	10	9.392e-08	rs2306970
SCZ	MMMH	21	TMEM180	10	1.628e-06	rs7909286
SCZ	MMMH	22	ARL3	10	4.597e-08	rs7909286
SCZ	MMMH	23	C10orf32	10	6.281e-07	rs7909286
SCZ	MMMH	24	C10orf32-ASMT	10	4.597e-08	rs7909286
SCZ	MMMH	25	AS3MT	10	4.597e-08	rs7909286

**Suppl. Table 4. List of genes associated with the shared genetic determinants between the psychiatric disorders and multivariate genome-wide association statistics for mental health** Genes associated with conjFDR output from diagnosis and multivariate genome-wide association statistics for mental health. This list only includes psychiatric disorders for which we identified significant loci. The number of mapped genes can be greater than the number of discovered loci since loci can be associated with more than one gene.

<i>Diagnosis</i>	<i>Measure</i>	<i>No</i>	<i>Gene</i>	<i>Chr</i>	<i>Pmin</i>	<i>Ind. Sig. SNP</i>
SCZ	MMMH	26	CNNM2	10	4.597e-08	rs7909286
SCZ	MMMH	27	NT5C2	10	6.005e-06	rs7909286
SCZ	MMMH	28	TAC3	12	1.549e-07	rs703848
SCZ	MMMH	29	MYO1A	12	1.549e-07	rs703848
SCZ	MMMH	30	TMEM194A	12	2.860e-07	rs703848
SCZ	MMMH	31	NAB2	12	2.860e-07	rs703848
SCZ	MMMH	32	STAT6	12	2.860e-07	rs703848
SCZ	MMMH	33	SYT1	12	1.021e-07	rs6539344
SCZ	MMMH	34	PRKD1	14	3.714e-07	rs10134319
SCZ	MMMH	35	NRXN3	14	2.761e-07	rs17764956
SCZ	MMMH	36	APOPT1	14	2.421e-07	rs2295146
SCZ	MMMH	37	XRCC3	14	1.022e-06	rs2295146
SCZ	MMMH	38	AL049840.1	14	3.317e-06	rs2295146
SCZ	MMMH	39	ZFYVE21	14	2.421e-07	rs2295146
SCZ	MMMH	40	PPP1R13B	14	2.421e-07	rs2295146
SCZ	MMMH	41	KIF26A	14	3.317e-06	rs2295146
SCZ	MMMH	42	GOLGA6L9	15	9.837e-08	rs4778687
SCZ	MMMH	43	CPEB1	15	9.837e-08	rs4778687
SCZ	MMMH	44	AP3B2	15	9.837e-08	rs4778687
SCZ	MMMH	45	FURIN	15	4.469e-08	rs4932178
SCZ	MMMH	46	FES	15	5.858e-08	rs4932178
SCZ	MMMH	47	MAN2A2	15		rs4932178
SCZ	MMMH	48	RCCD1	15	7.716e-08	rs4932178
SCZ	MMMH	49	RHBDL3	17	4.403e-07	rs9916114
SCZ	MMMH	50	C17orf75	17	9.641e-07	rs9916114
SCZ	MMMH	51	ZNF207	17	7.052e-07	rs9916114



**Suppl. Table 4. List of genes associated with the shared genetic determinants between the psychiatric disorders and multivariate genome-wide association statistics for mental health** Genes associated with conjFDR output from diagnosis and multivariate genome-wide association statistics for mental health. This list only includes psychiatric disorders for which we identified significant loci. The number of mapped genes can be greater than the number of discovered loci since loci can be associated with more than one gene.

<i>Diagnosis</i>	<i>Measure</i>	<i>No</i>	<i>Gene</i>	<i>Chr</i>	<i>Pmin</i>	<i>Ind. Sig. SNP</i>
SCZ	MMMH	52	PSMD11	17	4.403e-07	rs9916114
SCZ	MMMH	53	CDK5R1	17	5.273e-07	rs9916114
SCZ	MMMH	54	MYO1D	17	5.273e-07	rs9916114
SCZ	MMMH	55	NMT1	17	6.944e-08	rs2532392
SCZ	MMMH	56	FMNL1	17	5.389e-08	rs2532392
SCZ	MMMH	57	ARHGAP27	17	5.389e-08	rs2532392
SCZ	MMMH	58	PLEKHM1	17	5.389e-08	rs2532392
SCZ	MMMH	59	CRHR1	17	5.389e-08	rs2532392
SCZ	MMMH	60	SPPL2C	17	5.389e-08	rs2532392
SCZ	MMMH	61	MAPT	17	5.389e-08	rs2532392
SCZ	MMMH	62	STH	17	1.145e-07	rs2532392
SCZ	MMMH	63	KANSL1	17	5.090e-08	rs2532392
SCZ	MMMH	64	ARL17B	17	6.864e-08	rs2532392
SCZ	MMMH	65	LRRC37A	17	5.389e-08	rs2532392
SCZ	MMMH	66	LRRC37A2	17	5.389e-08	rs2532392
SCZ	MMMH	67	ARL17A	17	5.389e-08	rs2532392
SCZ	MMMH	68	NSF	17	5.389e-08	rs2532392
SCZ	MMMH	69	WNT3	17	1.057e-07	rs2532392
SCZ	MMMH	70	EFCAB13	17	5.879e-07	rs2532392
SCZ	MMMH	71	DCC	18	1.048e-07	rs10502971
SCZ	MMMH	72	TCF4	18	4.085e-08	rs7240986
MDD	MMMH	1	NCKIPSD	3	3.142e-07	rs2029591
MDD	MMMH	2	P4HTM	3	3.142e-07	rs2029591
MDD	MMMH	3	WDR6	3	3.142e-07	rs2029591
MDD	MMMH	4	DALRD3	3	3.142e-07	rs2029591
MDD	MMMH	5	QRICH1	3	3.142e-07	rs2029591

**Suppl. Table 4. List of genes associated with the shared genetic determinants between the psychiatric disorders and multivariate genome-wide association statistics for mental health** Genes associated with conjFDR output from diagnosis and multivariate genome-wide association statistics for mental health. This list only includes psychiatric disorders for which we identified significant loci. The number of mapped genes can be greater than the number of discovered loci since loci can be associated with more than one gene.

<i>Diagnosis</i>	<i>Measure</i>	<i>No</i>	<i>Gene</i>	<i>Chr</i>	<i>Pmin</i>	<i>Ind. Sig. SNP</i>
MDD	MMMH	6	CCDC71	3	3.142e-07	rs2029591
MDD	MMMH	7	GPX1	3	3.142e-07	rs2029591
MDD	MMMH	8	RHOA	3		rs2029591
MDD	MMMH	9	AMT	3	3.142e-07	rs2029591
MDD	MMMH	10	BSN	3	3.142e-07	rs2029591
MDD	MMMH	11	APEH	3	1.386e-06	rs2029591
MDD	MMMH	12	MST1	3	1.386e-06	rs2029591
MDD	MMMH	13	RNF123	3	3.142e-07	rs2029591
MDD	MMMH	14	AMIGO3	3	1.088e-06	rs2029591
MDD	MMMH	15	GMPPB	3	3.142e-07	rs2029591
MDD	MMMH	16	IP6K1	3	8.869e-07	rs2029591
MDD	MMMH	17	CDHR4	3	1.215e-06	rs2029591
MDD	MMMH	18	FAM212A	3	8.869e-07	rs2029591
MDD	MMMH	19	UBA7	3	1.532e-06	rs2029591
MDD	MMMH	20	TRAIP	3	1.290e-06	rs2029591
MDD	MMMH	21	CAMKV	3	1.487e-06	rs2029591
MDD	MMMH	22	RBM6	3	4.871e-07	rs2029591
MDD	MMMH	23	HYAL3	3	3.142e-07	rs2029591
MDD	MMMH	24	ZNF462	9	3.141e-07	rs11789013
MDD	MMMH	25	RP11-508N12.4	9	3.141e-07	rs11789013
MDD	MMMH	26	SEMA6D	15	3.546e-07	rs4338738
MDD	MMMH	27	DCC	18	4.743e-07	rs8084351
MDD	MMMH	28	TCF4	18	4.594e-07	rs2924336
MDD	MMMH	29	ZMYND8	20	1.821e-07	rs6094628
BIP	MMMH	1	TRIM54	2	4.702e-07	rs1260326
BIP	MMMH	2	EIF2B4	2	3.833e-07	rs1260326

**Suppl. Table 4. List of genes associated with the shared genetic determinants between the psychiatric disorders and multivariate genome-wide association statistics for mental health** Genes associated with conjFDR output from diagnosis and multivariate genome-wide association statistics for mental health. This list only includes psychiatric disorders for which we identified significant loci. The number of mapped genes can be greater than the number of discovered loci since loci can be associated with more than one gene.

<i>Diagnosis</i>	<i>Measure</i>	<i>No</i>	<i>Gene</i>	<i>Chr</i>	<i>Pmin</i>	<i>Ind. Sig. SNP</i>
BIP	MMMH	3	SNX17	2	3.833e-07	rs1260326
BIP	MMMH	4	ZNF513	2	3.833e-07	rs1260326
BIP	MMMH	5	PPM1G	2	3.833e-07	rs1260326
BIP	MMMH	6	KRTCAP3	2	3.253e-07	rs1260326
BIP	MMMH	7	GCKR	2	3.253e-07	rs1260326
BIP	MMMH	8	AC109829.1	2	1.823e-06	rs1260326
BIP	MMMH	9	CADM2	3	7.779e-08	rs1872552
BIP	MMMH	10	CNTN5	11	4.121e-07	rs671129
BIP	MMMH	11	TAC3	12	2.144e-07	rs703848
BIP	MMMH	12	MYO1A	12	2.144e-07	rs703848
BIP	MMMH	13	TMEM194A	12	3.734e-07	rs703848
BIP	MMMH	14	NAB2	12	3.734e-07	rs703848
BIP	MMMH	15	STAT6	12	3.734e-07	rs703848
BIP	MMMH	16	CABP1	12	4.644e-07	rs56366632
BIP	MMMH	17	MLEC	12	1.524e-06	rs56366632
BIP	MMMH	18	UNC119B	12	1.373e-06	rs56366632
BIP	MMMH	19	ACADS	12	1.060e-06	rs56366632
BIP	MMMH	20	SPPL3	12	2.124e-06	rs56366632
BIP	MMMH	21	APOPT1	14	3.221e-07	rs2295146
BIP	MMMH	22	XRCC3	14	1.311e-06	rs2295146
BIP	MMMH	23	AL049840.1	14	3.618e-06	rs2295146
BIP	MMMH	24	ZFYVE21	14	3.221e-07	rs2295146
BIP	MMMH	25	PPP1R13B	14	3.221e-07	rs2295146
BIP	MMMH	26	KIF26A	14	3.618e-06	rs2295146
BIP	MMMH	27	FURIN	15	6.946e-08	rs1894401
BIP	MMMH	28	FES	15	6.946e-08	rs1894401

**Suppl. Table 4. List of genes associated with the shared genetic determinants between the psychiatric disorders and multivariate genome-wide association statistics for mental health** Genes associated with conjFDR output from diagnosis and multivariate genome-wide association statistics for mental health. This list only includes psychiatric disorders for which we identified significant loci. The number of mapped genes can be greater than the number of discovered loci since loci can be associated with more than one gene.

<i>Diagnosis</i>	<i>Measure</i>	<i>No</i>	<i>Gene</i>	<i>Chr</i>	<i>Pmin</i>	<i>Ind. Sig. SNP</i>
ASD	MMMH	1	SERPINA1	14	6.175e-08	rs28929474
ASD	MMMH	2	NMT1	17	7.554e-09	rs71375338
ASD	MMMH	3	FMNL1	17	6.107e-09	rs71375338
ASD	MMMH	4	ARHGAP27	17	5.686e-09	rs71375338
ASD	MMMH	5	PLEKHM1	17	5.686e-09	rs71375338
ASD	MMMH	6	CRHR1	17	6.107e-09	rs71375338
ASD	MMMH	7	SPPL2C	17	5.686e-09	rs71375338
ASD	MMMH	8	MAPT	17	6.107e-09	rs71375338
ASD	MMMH	9	STH	17	1.158e-08	rs71375338
ASD	MMMH	10	KANSL1	17	6.107e-09	rs71375338
ASD	MMMH	11	ARL17B	17	6.282e-09	rs71375338
ASD	MMMH	12	LRRC37A	17	5.686e-09	rs71375338
ASD	MMMH	13	LRRC37A2	17	5.686e-09	rs71375338
ASD	MMMH	14	ARL17A	17	5.686e-09	rs71375338
ASD	MMMH	15	NSF	17	6.107e-09	rs71375338
ASD	MMMH	16	WNT3	17	1.160e-08	rs71375338
ADHD	MMMH	1	NCKIPSD	3	1.069e-07	rs373797031
ADHD	MMMH	2	P4HTM	3	1.069e-07	rs373797031
ADHD	MMMH	3	WDR6	3	1.069e-07	rs373797031
ADHD	MMMH	4	DALRD3	3	1.069e-07	rs373797031
ADHD	MMMH	5	QRICH1	3	1.102e-07	rs373797031
ADHD	MMMH	6	CCDC71	3	1.069e-07	rs373797031
ADHD	MMMH	7	GPX1	3	1.069e-07	rs373797031
ADHD	MMMH	8	AMT	3	1.069e-07	rs373797031
ADHD	MMMH	9	BSN	3	1.069e-07	rs373797031
ADHD	MMMH	10	APEH	3	5.050e-07	rs373797031

**Suppl. Table 4. List of genes associated with the shared genetic determinants between the psychiatric disorders and multivariate genome-wide association statistics for mental health** Genes associated with conjFDR output from diagnosis and multivariate genome-wide association statistics for mental health. This list only includes psychiatric disorders for which we identified significant loci. The number of mapped genes can be greater than the number of discovered loci since loci can be associated with more than one gene.

<i>Diagnosis</i>	<i>Measure</i>	<i>No</i>	<i>Gene</i>	<i>Chr</i>	<i>Pmin</i>	<i>Ind. Sig. SNP</i>
ADHD	MMMH	11	MST1	3	5.050e-07	rs373797031
ADHD	MMMH	12	RNF123	3	1.069e-07	rs373797031
ADHD	MMMH	13	AMIGO3	3	5.137e-07	rs373797031
ADHD	MMMH	14	GMPPB	3	1.069e-07	rs373797031
ADHD	MMMH	15	IP6K1	3	5.901e-08	rs373797031
ADHD	MMMH	16	CDHR4	3	4.367e-07	rs373797031
ADHD	MMMH	17	FAM212A	3	2.557e-07	rs373797031
ADHD	MMMH	18	UBA7	3	4.367e-07	rs373797031
ADHD	MMMH	19	TRAIP	3	5.619e-07	rs373797031
ADHD	MMMH	20	CAMKV	3	9.356e-07	rs373797031
ADHD	MMMH	21	RBM6	3	1.998e-07	rs373797031
ADHD	MMMH	22	HYAL3	3	1.069e-07	rs373797031
ADHD	MMMH	23	MEF2C	5	1.695e-07	rs304136
ADHD	MMMH	24	SPATS2	12	8.567e-06	rs933738
ADHD	MMMH	25	KCNH3	12	4.095e-07	rs933738
ADHD	MMMH	26	MCRS1	12	4.095e-07	rs933738
ADHD	MMMH	27	PRPF40B	12	1.716e-06	rs933738
ADHD	MMMH	28	FAM186B	12	1.716e-06	rs933738
ADHD	MMMH	29	FMNL3	12	7.508e-06	rs933738
ADHD	MMMH	30	TMBIM6	12	7.091e-06	rs933738
ADHD	MMMH	31	SH2B3	12	8.013e-07	rs10849943
ADHD	MMMH	32	ATXN2	12	3.011e-06	rs10849943
ADHD	MMMH	33	ALDH2	12	4.842e-07	rs10849943
ADHD	MMMH	34	TMEM116	12	9.193e-07	rs10849943
ADHD	MMMH	35	NAA25	12	4.842e-07	rs10849943
ADHD	MMMH	36	SERPINA1	14	4.065e-07	rs112635299

**Suppl. Table 4. List of genes associated with the shared genetic determinants between the psychiatric disorders and multivariate genome-wide association statistics for mental health**

Genes associated with conjFDR output from diagnosis and multivariate genome-wide association statistics for mental health. This list only includes psychiatric disorders for which we identified significant loci. The number of mapped genes can be greater than the number of discovered loci since loci can be associated with more than one gene.

<i>Diagnosis</i>	<i>Measure</i>	<i>No</i>	<i>Gene</i>	<i>Chr</i>	<i>Pmin</i>	<i>Ind. Sig. SNP</i>
ADHD	MMMH	37	SEMA6D	15	3.759e-07	rs281287

376

**Suppl. Table 5. Gene list of conjFDR between the brain functional connectome and multivariate genome-wide association statistics for mental health**

Genes identified through FUMA associated with the shared loci between the multivariate genome-wide association statistics for mental health and FC or node variance. The number of genes in this gene set can be larger than the number of loci since a locus can be associated more than one gene.

<i>Feature</i>	<i>Measure</i>	<i>No</i>	<i>Gene</i>	<i>Chr</i>	<i>Pmin</i>	<i>Ind. Sig. SNP</i>
FC	MMMH	1	LRPPRC	2	3.806e-06	rs62132319
FC	MMMH	2	CAMKMT	2	3.807e-07	rs62132319
FC	MMMH	3	FANCL	2	1.683e-06	rs2125924
FC	MMMH	4	BCL11A	2	1.977e-07	rs13019832
FC	MMMH	5	ARL8B	3	4.172e-07	rs74677825
FC	MMMH	6	ZNF589	3	1.032e-06	rs6769821
FC	MMMH	7	TMA7	3	4.737e-07	rs6769821
FC	MMMH	8	ATRIP	3	1.134e-06	rs6769821
FC	MMMH	9	TREX1	3	4.737e-07	rs6769821
FC	MMMH	10	NCKIPSD	3	4.737e-07	rs6769821
FC	MMMH	11	IP6K2	3	1.738e-06	rs6769821
FC	MMMH	12	PRKAR2A	3	4.737e-07	rs6769821
FC	MMMH	13	SLC25A20	3	1.499e-06	rs6769821
FC	MMMH	14	ARIH2OS	3	1.532e-06	rs6769821
FC	MMMH	15	ARIH2	3	1.303e-06	rs6769821
FC	MMMH	16	P4HTM	3	4.737e-07	rs6769821
FC	MMMH	17	WDR6	3	4.737e-07	rs6769821
FC	MMMH	18	DALRD3	3	4.737e-07	rs6769821

**Suppl. Table 5. Gene list of conjFDR between the brain functional connectome and multivariate genome-wide association statistics for mental health** Genes identified through FUMA associated with the shared loci between the multivariate genome-wide association statistics for mental health and FC or node variance. The number of genes in this gene set can be larger than the number of loci since a locus can be associated more than one gene.

<i>Feature</i>	<i>Measure</i>	<i>No</i>	<i>Gene</i>	<i>Chr</i>	<i>Pmin</i>	<i>Ind. Sig. SNP</i>
FC	MMMH	19	NDUFAF3	3	1.205e-06	rs6769821
FC	MMMH	20	IMPDH2	3	1.205e-06	rs6769821
FC	MMMH	21	QRICH1	3	4.737e-07	rs6769821
FC	MMMH	22	QARS	3	8.188e-07	rs6769821
FC	MMMH	23	USP19	3	8.188e-07	rs6769821
FC	MMMH	24	LAMB2	3	9.061e-07	rs6769821
FC	MMMH	25	CCDC71	3	9.878e-08	rs6769821; rs2624847
FC	MMMH	26	KLHDC8B	3	4.898e-07	rs6769821
FC	MMMH	27	C3orf84	3	4.898e-07	rs6769821
FC	MMMH	28	CCDC36	3	4.737e-07	rs6769821
FC	MMMH	29	RP11-3B7.1	3	7.060e-07	rs6769821
FC	MMMH	30	C3orf62	3	7.060e-07	rs6769821
FC	MMMH	31	USP4	3	4.737e-07	rs6769821
FC	MMMH	32	GPX1	3	9.878e-08	rs6769821; rs2624847
FC	MMMH	33	RHOA	3	1.320e-06	rs6769821
FC	MMMH	34	TCTA	3	1.339e-06	rs6769821
FC	MMMH	35	AMT	3	9.878e-08	rs6769821; rs2624847
FC	MMMH	36	NICN1	3	4.737e-07	rs6769821
FC	MMMH	37	DAG1	3	1.143e-06	rs6769821
FC	MMMH	38	BSN	3	1.246e-06	rs6769821
FC	MMMH	39	MST1	3	6.078e-07	rs6769821
FC	MMMH	40	RNF123	3		rs6769821
FC	MMMH	41	GMPPB	3	4.737e-07	rs6769821
FC	MMMH	42	FAM212A	3	9.878e-08	rs2624847
FC	MMMH	43	MST1R	3	9.878e-08	rs2624847
FC	MMMH	44	RBM6	3	9.878e-08	rs2624847
FC	MMMH	45	HYAL3	3	1.039e-06	rs6769821



**Suppl. Table 5. Gene list of conjFDR between the brain functional connectome and multivariate genome-wide association statistics for mental health** Genes identified through FUMA associated with the shared loci between the multivariate genome-wide association statistics for mental health and FC or node variance. The number of genes in this gene set can be larger than the number of loci since a locus can be associated more than one gene.

<i>Feature</i>	<i>Measure</i>	<i>No</i>	<i>Gene</i>	<i>Chr</i>	<i>Pmin</i>	<i>Ind. Sig. SNP</i>
FC	MMMH	46	CADM2	3	2.305e-08	rs4856600
FC	MMMH	47	EPHA3	3	2.806e-07	rs73153293
FC	MMMH	48	BANK1	4	2.807e-06	rs13107325
FC	MMMH	49	SLC39A8	4	9.802e-10	rs13107325
FC	MMMH	50	MEF2C	5	2.866e-07	rs618741
FC	MMMH	51	SND1	7	3.498e-07	rs58406876
FC	MMMH	52	LRRC4	7	6.509e-07	rs58406876
FC	MMMH	53	ARL3	10	3.174e-07	rs12765002
FC	MMMH	54	C10orf32	10	3.174e-07	rs12765002
FC	MMMH	55	C10orf32-ASMT	10	3.174e-07	rs12765002
FC	MMMH	56	AS3MT	10	3.174e-07	rs12765002
FC	MMMH	57	CNNM2	10	3.174e-07	rs12765002
FC	MMMH	58	NT5C2	10	3.147e-06	rs12765002
FC	MMMH	59	SH2B3	12	1.732e-07	rs12810456
FC	MMMH	60	ATXN2	12	1.035e-07	rs12810456
FC	MMMH	61	BRAP	12	2.655e-06	rs12810456
FC	MMMH	62	ACAD10	12	2.474e-06	rs12810456
FC	MMMH	63	RP11-162P23.2	12	2.887e-06	rs12810456
FC	MMMH	64	ALDH2	12	1.035e-07	rs12810456
FC	MMMH	65	MAPKAPK5	12	5.899e-07	rs12810456
FC	MMMH	66	TMEM116	12	1.035e-07	rs12810456
FC	MMMH	67	NAA25	12	1.035e-07	rs12810456
FC	MMMH	68	NMT1	17	9.948e-14	rs56328224; rs55938136
FC	MMMH	69	FMNL1	17	9.948e-14	rs56328224; rs55938136
FC	MMMH	70	ARHGAP27	17	9.948e-14	rs56328224; rs55938136
FC	MMMH	71	PLEKHM1	17	9.948e-14	rs56328224; rs55938136
FC	MMMH	72	CRHR1	17	9.948e-14	rs56328224; rs55938136

**Suppl. Table 5. Gene list of conjFDR between the brain functional connectome and multivariate genome-wide association statistics for mental health** Genes identified through FUMA associated with the shared loci between the multivariate genome-wide association statistics for mental health and FC or node variance. The number of genes in this gene set can be larger than the number of loci since a locus can be associated more than one gene.

<i>Feature</i>	<i>Measure</i>	<i>No</i>	<i>Gene</i>	<i>Chr</i>	<i>Pmin</i>	<i>Ind. Sig. SNP</i>
FC	MMMH	73	SPPL2C	17	9.948e-14	rs56328224; rs55938136
FC	MMMH	74	MAPT	17	9.948e-14	rs56328224; rs55938136
FC	MMMH	75	STH	17	9.948e-14	rs56328224
FC	MMMH	76	KANSL1	17	9.948e-14	rs56328224; rs55938136
FC	MMMH	77	ARL17B	17	9.948e-14	rs56328224
FC	MMMH	78	LRRC37A	17	9.948e-14	rs56328224; rs55938136
FC	MMMH	79	LRRC37A2	17	9.948e-14	rs56328224; rs55938136
FC	MMMH	80	ARL17A	17	9.948e-14	rs56328224; rs55938136
FC	MMMH	81	NSF	17	9.948e-14	rs56328224; rs55938136
FC	MMMH	82	WNT3	17	9.948e-14	rs56328224
FC	MMMH	83	ATP5G1	17	4.306e-07	rs11079849
FC	MMMH	84	UBE2Z	17	2.101e-06	rs11079849
FC	MMMH	85	SNF8	17	4.306e-07	rs11079849
FC	MMMH	86	IGF2BP1	17	4.306e-07	rs11079849
FC	MMMH	87	NGFR	17	6.727e-06	rs11079849
node variance	MMMH	1	SIX3	2	4.062e-07	rs490340
node variance	MMMH	2	BANK1	4	6.972e-08	rs13109404
node variance	MMMH	3	EXOC4	7	7.507e-08	rs6467507
node variance	MMMH	4	NMT1	17	4.007e-12	rs71375338; rs55938136
node variance	MMMH	5	FMNL1	17	4.007e-12	rs71375338; rs55938136
node variance	MMMH	6	ARHGAP27	17	2.850e-12	rs71375338; rs55938136

**Suppl. Table 5. Gene list of conjFDR between the brain functional connectome and multivariate genome-wide association statistics for mental health** Genes identified through FUMA associated with the shared loci between the multivariate genome-wide association statistics for mental health and FC or node variance. The number of genes in this gene set can be larger than the number of loci since a locus can be associated more than one gene.

<i>Feature</i>	<i>Measure</i>	<i>No</i>	<i>Gene</i>	<i>Chr</i>	<i>Pmin</i>	<i>Ind. Sig. SNP</i>
node variance	MMMH	7	PLEKHM1	17	2.850e-12	rs71375338; rs55938136
node variance	MMMH	8	CRHR1	17	4.007e-12	rs71375338; rs55938136
node variance	MMMH	9	SPPL2C	17	2.850e-12	rs71375338; rs55938136
node variance	MMMH	10	MAPT	17	4.007e-12	rs71375338; rs55938136
node variance	MMMH	11	STH	17	7.119e-12	rs71375338
node variance	MMMH	12	KANSL1	17	4.007e-12	rs71375338; rs76527351; rs55938136
node variance	MMMH	13	ARL17B	17	3.084e-11	rs71375338
node variance	MMMH	14	LRRC37A	17	2.850e-12	rs71375338; rs55938136
node variance	MMMH	15	LRRC37A2	17	2.850e-12	rs71375338; rs55938136
node variance	MMMH	16	ARL17A	17	2.850e-12	rs71375338; rs55938136
node variance	MMMH	17	NSF	17	4.405e-12	rs71375338; rs55938136
node variance	MMMH	18	WNT3	17	1.450e-11	rs71375338

377

378

<b>Suppl. Table 6. Biological processes mapped by reactome</b> Biological processes associated with the shared genetic determinants between the multivariate genome-wide association statistics for mental health and the brain functional connectome as identified by the reactome toolbox.				
<i>Pathway name</i>	<i>No of genes involved</i>	<i>p-value</i>	<i>p-value (FDR)</i>	<i>Genes involved</i>
Axonal growth stimulation	2	0.0005	0.2235	NGFR; RHOA
Signaling by MST1	2	0.0014	0.2491	MST1; MST1R
MECP2 regulates transcription factors	2	0.0021	0.2491	MEF2C
Axonal growth inhibition (RHOA activation)	2	0.0026	0.2491	NGFR; RHOA
p75NTR regulates axonogenesis	2	0.0030	0.2491	NGFR; RHOA
Respiratory electron transport, ATP synthesis by chemiosmotic coupling, and heat production by uncoupling proteins.	5	0.0038	0.2595	NDUFAF3; QRICH1; ATP5G1; LRPPRC
Inactivation, recovery and regulation of the phototransduction cascade	3	0.0059	0.3361	NMT1; CAMKMT
The phototransduction cascade	3	0.0075	0.3361	NMT1; CAMKMT
Non-integrin membrane-ECM interactions	3	0.0082	0.3361	LAMB2; DAG1
Formation of ATP by chemiosmotic coupling	2	0.0106	0.3361	ATP5G1
ECM proteoglycans	3	0.0164	0.3361	LAMB2; DAG1
Laminin interactions	2	0.0187	0.3361	LAMB2
Cristae formation	2	0.0187	0.3361	ATP5G1
MET activates PTK2 signaling	2	0.0198	0.3361	LAMB2
NFG and proNGF binds to p75NTR	1	0.0199	0.3361	NGFR
Defective POMGNT1 causes MDDGA3, MDDGB3 and MDDGC3	1	0.0199	0.3361	DAG1
EPHA-mediated growth cone collapse	2	0.0210	0.3361	EPHA3; RHOA
The citric acid (TCA) cycle and respiratory electron transport	5	0.0224	0.3361	NDUFAF3; QRICH1; ATP5G1; LRPPRC

**Suppl. Table 6. Biological processes mapped by reactome** Biological processes associated with the shared genetic determinants between the multivariate genome-wide association statistics for mental health and the brain functional connectome as identified by the reactome toolbox.

<i>Pathway name</i>	<i>No of genes involved</i>	<i>p-value</i>	<i>p-value (FDR)</i>	<i>Genes involved</i>
Defective POMT2 causes MDDGA2, MDDGB2 and MDDGC2	1	0.0265	0.3361	DAG1
Defective POMT1 causes MDDGA1, MDDGB1 and MDDGC1	1	0.0265	0.3361	DAG1
Activation, myristoylation of BID and translocation to mitochondria	1	0.0265	0.3361	NMT1
EGR2 and SOX10-mediated initiation of Schwann cell myelination	2	0.0286	0.3361	LAMB2; DAG1
MAPK6/MAPK4 signaling	3	0.0349	0.3361	IGF2BP1; MAPKAPK5
MET promotes cell motility	2	0.0371	0.3361	LAMB2
NADE modulates death signalling	1	0.0394	0.3361	NGFR
p75NTR negatively regulates cell cycle via SC1	1	0.0394	0.3361	NGFR
Cytosolic tRNA aminoacylation	2	0.0448	0.3361	QARS
Respiratory electron transport	3	0.0454	0.3361	NDUFAF3; QRICH1; LRPPRC
Regulation by TREX1	1	0.0458	0.3361	TREX1
Trafficking of myristoylated proteins to the cilium	1	0.0458	0.3361	ARL3
Fanconi Anemia Pathway	2	0.0497	0.3361	FANCL; ATRIP