Research Article

Telomere length among Chinese oldest old

Suey S.Y. Yeung^a, Xingyan Wang^{b,c,d,e}, Suk Ling Ma^f, Yangchao Chen^{g,h}, Stephen Kwok Wing Tsui^{g,i,j}, Nelson Leung Sang Tang^{b,c,d,e,k}, Jean Woo^{a,i}

^a Department of Medicine & Therapeutics, Faculty of Medicine, The Chinese University of Hong Kong, Hong Kong, China

^b Department of Chemical Pathology and Li Ka Shing Institute of Health Sciences, Faculty of Medicine, The Chinese University of Hong Kong, Hong Kong, China

^c Hong Kong Branch of CAS Center for Excellence in Animal Evolution and Genetics, Hong Kong, China

^d KIZ/CUHK Joint Laboratory of Bioresources and Molecular Research in Common Diseases, Hong Kong, China

^e Functional Genomics and Biostatistical Computing Laboratory, CUHK Shenzhen Research Institute, Shenzhen 518172, China

^f Department of Psychiatry, The Chinese University of Hong Kong, Hong Kong, China

^g School of Biomedical Sciences, The Chinese University of Hong Kong, Hong Kong, China.

^h Shenzhen Research Institute, The Chinese University of Hong Kong, Shenzhen, China.

¹ Hong Kong Bioinformatics Centre, The Chinese University of Hong Kong, Hong Kong, China.

^j Centre for Microbial Genomics and Proteomics, The Chinese University of Hong Kong, Hong Kong, China.

^k Cytomics Limited, Hong Kong Science Park, Hong Kong

¹Centre for Nutritional Studies, Faculty of Medicine, The Chinese University of Hong Kong, Hong Kong, China

Short title: Telomere length in oldest old

Corresponding author:

Suey S.Y. Yeung, Rm 124021, 10/F Lui Che Woo Clinical Sciences Building, Prince of Wales Hospital,

Shatin, N.T., Hong Kong, Tel: +852 3505 2190, Email: sueyyeung@cuhk.edu.hk

Number of tables: 2

Number of figures: 2

Word count: 4001

Keywords: Aging, telomere, biomarkers, oldest old, morbidity

1 Abstract

2 Introduction: Telomere length (TL) is generally regarded as a biomarker of aging. TL, which is

3 influenced by sociodemographic factors, has been shown to be inversely associated with morbidity.

4 However, most studies were examined in the youngest old and whether the findings can be extended

5 to the oldest old is less clear. This study examined TL and its associated factors in Chinese oldest old

6 in Hong Kong.

7 Methods: Follow-up data collected after 14 years from the baseline of the Mr. and Ms. Osteoporosis

8 cohort were analyzed. A structured interview on sociodemographic factors and physical

9 measurement was conducted. Frailty and sarcopenia status were respectively determined by Fried's

10 criteria and Asian Working Group for Sarcopenia definition. TL was measured by a molecular

11 inversion probe - quantitative PCR (MIP-qPCR) assay and expressed as telomere / a single copy

12 reference gene (T/S) ratio. Adjusted binary logistic regressions were used to examine the associations

13 between TL and the presence of multimorbidity, age-related diseases, frailty and sarcopenia.

14 **Results:** Among 555 participants (mean age 83.6±3.8 years, 41.3% women), the mean T/S ratio was

15 1.01±0.20. Males had lower T/S ratio (0.97±0.20) compared with females (1.07±0.18) (p<0.001). A

16 lower education level was related to a longer TL (p=0.016). Being a current smoker was related to a

shorter TL (p=0.007). TL was not significantly different across categories of age, subjective

socioeconomic status, drinking status, physical activity level and body mass index (p>0.05). There

19 were no associations between TL and the presence of multimorbidity, diabetes, stroke,

20 cardiovascular diseases, cognitive impairment, frailty and sarcopenia.

21 **Conclusion:** Among Chinese oldest old, males had shorter TL compared with females. TL was not

associated with age-related diseases, frailty and sarcopenia in this age group. TL may not be a

23 biological marker of aging among the oldest old.

24 Introduction

Over the last several decades, life expectancy has increased worldwide. Hong Kong has sustained the world's highest life expectancy at birth since 2013. The latest data reported a life expectancy of 82.9 years for males and 88.0 years for females in Hong Kong [1]. Telomere length (TL), which has been extensively studied in relation to aging and longevity, may play a role in explaining extended longevity among many other factors.

Telomeres are repetitive DNA sequences at the end of chromosomes and serve as caps to preserve genomic integrity and stability by protecting linear DNA molecules from being recognized as double-stranded DNA breaks. TL is influenced by various genetic and non-genetic determinants such as demographic (age, sex), social and environmental factors (diet, smoking, alcohol use, body mass index, socioeconomic status and physical activity) [2-6]. However, most of the studies were conducted in the youngest old (below 75 years old). Studies related to TL beyond the age of 75 years were limited.

37 Telomeres shorten with each cell cycle, therefore, it has been suggested that telomere 38 shortening is one of the hallmarks of aging [7]. Shorter TL was associated with poor survival and age-39 related diseases including cardiovascular diseases, neurodegenerative diseases and diabetes in the 40 youngest old [8,9]. However, these findings were inconsistent among the oldest old. The Leiden 85+ 41 study showed that TL was not associated with age-related diseases [10]. Similarly, the Newcastle 85+ 42 study showed that TL was not associated with multimorbidity, cognitive impairment and disability 43 [11]. While TL was not associated with geriatric syndromes such as frailty and sarcopenia in the 44 youngest old [12], whether this finding can be extended to the oldest old is less clear.

45 Currently, gPCR method was commonly used to assess the TL in epidemiology and population studies [13]. It is also widely known as the relative TL measurement in which TL is expressed as a T / S 46 47 (telomere / a single copy reference gene) ratio. However, many problems have been described for this method. They include the stringent requirements of high quality DNA template, poor 48 49 reproducibility, batch effect, etc [14-17]. Some research groups even suggested that previous 50 reported association between TL and social factors might only represent a systemic bias instead of a 51 real association [18]. A recent paper used the qPCR methods in the UK Biobank samples and found 52 that variance due to technical factors was huge. Technical factors have a strong influence on the 53 results include batch of primer, batch of enzyme, operator among others [19]. Some of the assay 54 problems could be attributed to the use of a multiple-bases mismatch primer (for example the Tel-1b 55 primer) in gPCR for amplification of the telomere repeat motif. Here, we designed a new relative TL 56 measurement method which completely eliminated the use of all mismatch primers by using two 57 molecular inversion probes (MIPs) and an additional ligation step. Instead of using the native

telomere fragments of the sample DNA as the direct template of qPCR, ligation of exogenous MIPs

59 targeting the telomere and another reference repeat motif are used as the template of qPCR.

60 Therefore, this new relative method for TL which could be applied in epidemiology may be helpful to

61 enrich the availability of methods for studying this important topic.

62 Molecular inversion probe (MIP) has been extensively used in molecular biology for almost 63 30 years and was first used as a localized DNA detection method and were called padlock probe. In 64 that version of MIP, there was no gap between the 5' and 3' end of the MIP after hybridization to the 65 targets [20]. Later, it was used as a method of genotyping single nucleotide polymorphisms (SNPs) 66 with a single-base gap between the 5' and 3' end of the MIP [21]. In the recent next generation 67 sequence era, MIP has been further developed to as capturing probe to enrich target templates for 68 library preparation of high throughput sequencing. It is sometimes referred as connector inversion 69 probe (CIP) if gap filling was used as a mean of target enrichment for sequencing. Here, we describe 70 a novel application of MIP in measurement of length (or abundance) of a repeat motif, specifically we 71 measured TL by quantification of ligated (circularized) MIP that had been hybridized to the target 72 motif. And the longer the telomere, more MIP are accommodated and after ligation (circularization) 73 they become templates for quantification. The relative TL is quantified in a reaction using 2 MIPs, one 74 MIP for telomere motif and another MIP for a reference genomic motif. Efficiency corrected delta-75 delta CT values correlate with TL.

To the best of our knowledge, data on the TL measurements in the oldest old and especially in Asia is limited. It is important to fill this research gap given the rapid growth of the aging population. Using samples from the Osteoporotic Fractures in Men and Women (Mr. and Ms. Osteoporosis) study, we examined TL expressed as a T/S ratio and its associated factors among the Chinese oldest old in Hong Kong.

81

82 Materials and Methods

83 Study population

This study is based on the follow-up data collected 14 years after the baseline of the Mr. and Ms. Osteoporosis cohort. A detailed description of the study has been previously reported [22]. In brief, 4000 community-dwelling Chinese men and women aged ≥65 years participated in a health check between 2001 and 2003. Participants were recruited using a stratified sampling method to equally assign them into 65-69, 70-74 and 75+ age groups. They should have been able to walk or take public transport to the study site. Those who had had bilateral hip replacements or were not competent to give informed consent were excluded. Participants were invited to re-attend for a repeated health

91 check between 2015 and 2017. For the present analysis, participants who discontinued the 14-year

92 follow-up assessment (n=2939) and those who attended the assessment but without a valid TL

93 measurement (n=506) were excluded. The final sample included in the analysis was 555. All

94 participants gave their written informed consent to participate in the study. The study was conducted

95 in compliance with the Declaration of Helsinki and was approved by the Clinical Research Ethics

96 Committee of the Chinese University of Hong Kong.

97

98 Characteristics of participants

99 A structured interview was conducted to collect information on sex, age, education level (primary or 100 below vs. secondary or above), subjective socioeconomic status, smoking status (never/ex-smoker vs. 101 current smoker), current drinker (no vs yes), physical activity level, body mass index (BMI) and 102 medical history. Subjective socioeconomic status was assessed by asking participants to mark their 103 self-perceived position on a drawing of an upright ladder with 10 rungs. The most undesirable state 104 with respect to their standing in the community (community status ladder) and in Hong Kong (Hong 105 Kong ladder) was indicated by the lowest rung and the most desirable state was indicated by the 106 highest rung [23]. Physical activity level was assessed using the 12-item Physical Activity Scale for the 107 Elderly (PASE) [24]. Participants reported the average number of hours per day spent in leisure, 108 household, and occupational physical activities in the past week. Activity weights for each item were 109 determined based on the amount of energy spent, and each item score was computed by multiplying 110 the activity weight with daily activity frequency reported. A PASE score was computed by summing 111 each item score. A higher PASE score represents a higher physical activity level. BMI was calculated as 112 body weight (kg) divided by height (meters squared). Body weight was measured using the Physician 113 Balance Bean Scale (Healthometer, Bridgeview, IL) with participants wearing light clothing. Height 114 was measured using the Holtain Harpenden Standiometer (Holtain Ltd, Crosswell, UK). Low BMI was defined as $< 20.0 \text{ kg/m}^2$. 115

116 A list of diseases was used to assess the number of chronic diseases, including diabetes, 117 hyperthyroidism, hypothyroidism, stroke, Parkinson's disease, heart attack, congestive heart failure, 118 hypertension, chronic obstructive pulmonary disease and cancer. Participants were asked if a 119 healthcare professional has even told them that they had or have each of the preceding diseases. 120 Multimorbidity was defined as having two or more chronic diseases. Cognitive impairment was 121 assessed by trained research staff using the Chinese version of the Mini-Mental State Examination 122 (MMSE) [25]. The MMSE scores range from 0 to 30, with a higher score reflecting better cognitive 123 function. Cognitive impairment was defined by a MMSE score of <24 points [26]. Frailty status was 124 defined using the 5-item Cardiovascular Health Study (CHS) frailty phenotype: self-rated exhaustion, 125 weakness (grip strength), slow walking speed, low physical activity and unintentional weight loss [27].

126 The corresponding variables used in this study for the construction of the CHS score were a little of 127 time over the past 4 weeks feel a lot of energy, handgrip strength ≤first quintile, 6-meter walking 128 speed \leq first quintile, PASE score \leq first quintile, and weight loss \geq 5% in the past 6 months. The total 129 score ranges from 0 to 5. A score of 1 to 2 represents a pre-frail state, and a score of \geq 3 represents 130 frailty. Sarcopenia was defined using the Asia Working Group for Sarcopenia (AWGS) [28]. 131 Participants who had low muscle mass and low muscle strength or low physical performance were 132 defined as having sarcopenia. Muscle mass was measured using the dual-energy X-ray 133 absorptiometry and total appendicular skeletal muscle mass (ASM) was calculated. Low muscle mass was defined as ASM/height² <7.0 kg/m² for males and <5.4 kg/m² for females. Low muscle strength 134 135 was defined as handgrip strength of <26 kg in males and <18 kg in females. Low physical performance 136 was defined as a gait speed of <0.8 m/s using the 6-meter walking test at the usual pace [28].

137

138 Telomere length measurements by MIP-qPCR

139 Two MIPs are used, one binds to telomere repeat motif (GGGTTA) and another binds to a common 140 genomic repeat motif such as 4 bp or 5 bp repeats. The 4-bp ATGG repeat motif was used as the 141 target of genomic repeat in this study. The details of DNA sequences and protocol are described in 142 Supplementary Material. The length of the telomere determines the extent of availability of binding 143 sites for the telomere specific MIPs, therefore also related to the abundance of circularized telomere 144 MIP after ligase reaction (as shown in Fig. 1). On the other hand, the availability of genomic repeat 145 motif represents the amount of template DNA added into the ligase reaction as amount of genomic 146 repeat motif (e.g. ATGG) is fairly stable per copy of eukaryotic genome [29]. After a ligase reaction, 147 the two types of circularized MIP are quantified. Here, we used quantitative real-time PCR. However, 148 other quantification methods like digital PCR could be used as well. The ratio of the amount of circularized telomere MIP to that of genomic repeat MIP is proportional to telomere length. Like the 149 150 original relative TL qPCR method, telomere length expressed as T/S ratio were obtained by efficiency 151 corrected delta-delta method. In addition, calibration with samples of given telomere length could also be carried out to convert the T/S ratio to telomere length in kbp unit. Two calibrators with given 152 153 TL (6.9 and 8.1 kbp) measured by Southern blot were used as calibrators which enable conversion of 154 the ratio to telomere length in term of the unit of kbp.

DNA was extracted from blood samples using commercial DNA extraction kit according to the manufacturer's instruction. The MIP for telomere consists of the telomere repeat sequence (GGGTTA) at both 5' and 3' ends connected by linker sequences. Another MIP is used to target a common 4-bp tandem repeat sequence (ATGG) for normalization as roughly the same abundance of such 4-bp motifs are present in each eukaryotic cell. Both MIPs for telomere and 4-bp tandem repeat sequence were 5'-phosphorylated and they were incubated overnight at 50°C with the DNA for hybridization.

Gap filling and ligation was performed by the addition of Ampligase (EpiCentre), T4 polymerase
(Roche) in the presence of 10mM of only dATP and dGTP, and incubated at 37°C for 30 minutes and
inactivated at 95°C for 10 minutes. This ligase reaction circularizes those MIPs that have been
hybridized to their respective target motifs.

165 To quantify the total number of circularized MIPs of the 2 types of repeat motifs, duplex 166 quantitative PCR using specific primers to each type of MIP were performed. These primer pairs were 167 designed specific to PCR amplify circularized telomere and 4-bp tandem repeat MIPs respectively and 168 PCR does not proceed with non-circularized (open) MIPs. A 5' GC rich tail was added to all primers. In 169 addition, two Tagman probes containing the respective repeat sequence (GGGTTA or ATGG) were 170 used in gPCR to differentiate the circularized template of the 2 types of MIP probes in a duplex gPCR 171 performed in the same tube. Delta-delta CT with correction of PCR efficiency were calculated. That is 172 delta Ct of each sample subtracted that of the reference sample to calculate the delta-delta Ct value. 173 Based on the observed qPCR efficiencies of the 2 reactions, the ratio of telomere motif ligation points to genomic 4bp motif ligation points could be calculated as the efficiency corrected delta-delta Ct 174 175 method. This efficiency corrected delta-delta Ct ratio of the 2 motifs of each sample is used as a 176 biomarker of telomere length (T/S ratio). And average values was obtained from duplicated reactions

177 for each sample.

178 In addition, results of TL in kbp unit can be obtained by calibration of T/S ratio against known 179 TL of the calibrator samples. Efficiency corrected delta-delta Ct ratios of individual study samples 180 were regressed onto the calibration line to provide the telomere length in the unit of kbp.

181 Statistical analysis

182 Data are presented as numbers (percentages) and mean±standard deviation (SD). The differences in

183 T/S ratio across categories of sociodemographic factors were examined using analysis of variance

184 (ANOVA) or independent t-test, where appropriate. The associations between TL and each health

185 outcome (the presence of multimorbidity, diabetes, stroke, cardiovascular diseases, cognitive

186 impairment, frailty and sarcopenia) were examined using binary logistic regression models adjusting

187 for covariates. Data are presented as odds ratio (OR) and 95% confidence intervals (CI).

All the analyses were stratified by sex. The analyses were performed using the Statistical Package for the Social Science (version 26.0, Chicago, IL, USA). Results were considered statistically significant when *p*<0.05.

191 **Results**

192 Characteristics of participants

Table 1 shows the descriptive characteristics of the 555 participants. The mean age was 83.6±3.8
years, and 41.3% were females. Most of the participants completed primary school or below (63.4%),
were non-current smokers (97.7%) and had normal cognitive function (75.7%). The mean T/S ratio of
Chinese oldest old was 1.01±0.2.

197 In Chinese oldest old, males had significantly shorter TL (T/S ratio 0.97±0.20) compared with 198 females (T/S ratio 1.07±0.18) (p<0.001). Those who completed primary school or below (T/S ratio 199 1.03±0.19) had higher TL compared with those who completed secondary education or above (T/S 200 ratio 0.98±0.21) (p=0.016). Being a non-current smoker (T/S ratio 1.01±0.20) had significantly higher 201 TL compared with current smoker (T/S ratio 0.86±0.14) (p=0.004). TL was not significantly different 202 across categories of age, subjective socioeconomic status, drinking status, physical activity level and 203 BMI (p>0.05). There were no significant differences in TL between those with or without 204 multimorbidity, diabetes, stroke, cardiovascular diseases, cognitive impairment, frailty and 205 sarcopenia (p>0.05). Figure 2 shows the graphic representation of T/S ratio by age group for males 206 and females.

207Table 2 shows the associations between TL and prevalent frailty and sarcopenia status. In208crude and fully adjusted models, TL was not associated with prevalent frailty (adjusted OR for males2090.55, 95% Cl 0.10-3.06, p=0.490; adjusted OR for females 0.19, 95% Cl 0.02-1.51, p=0.117) and210sarcopenia (adjusted OR for males 4.07, 95% Cl 0.96-17.3, p=0.057; adjusted OR for females 0.74,21195% Cl 0.10-5.62, p=0.769). No associations between TL and the presence of multimorbidity, diabetes,212stroke, cardiovascular diseases and cognitive impairment were found in crude and adjusted models213(p>0.05, data not shown).

214

215 Discussion

The present study found a mean T/S ratio of 1.01 among Chinese oldest old (mean age of 83.6 years),

217 with higher TL observed in males compared with females. TL was longer for those with lower

education levels and among non-current smokers. There were no differences in TL across categories

of other factors being investigated including age, subjective socioeconomic status, smoking status,

220 physical activity level and BMI. TL was not associated with multimorbidity, diabetes, stroke,

221 cardiovascular diseases, cognitive impairment, frailty and sarcopenia in Chinese oldest old.

To the best of our knowledge, this was the first study to investigate TL and its associated factors among the Chinese oldest old. The mean T/S ratio in our participants was higher compared with older adults in China (mean age 61 years, T/S ratio 0.67), US (mean age 71 years, T/S ratio 0.90)

and Finland (mean age 71 years, T/S ratio 0.86) [30-32], but lower than Australian older adults (mean

226 age 77 years, T/S ratio 1.18) [33]. These discrepancies might be explained by the different mean age, 227 methods to measure TL or other sociodemographic factors. Consistent with the literature and the 228 baseline analysis of the same cohort, we found that males had a shorter TL compared with females 229 [5,34]. The sex-difference in TL could potentially be explained by the effect of estrogens. Estrogen 230 level is associated with telomerase activity via telomerase reverse transcriptase (hTERT) gene 231 expression, while and rogen appears to be a negative regulator of telomerase in normal prostate 232 tissue [35]. We found that oldest old with lower education levels had longer TL. A possible 233 explanation is that those with lower educational levels worked for lower-level and physically 234 demanding occupations, such that they were more physically healthy and experienced less work 235 stress compared with those with higher education levels and worked for higher-level occupations. In 236 line with the findings of a meta-analysis on smoking status and TL, we found that current smoker had 237 a shorter TL than non- or former smokers [2]. Harmful chemicals in cigarettes may induce oxidative 238 stress and inflammation, resulting in damage to the telomeric DNA and shortened telomeres. Most of 239 the sociodemographic factors of TL in the younger age group do not appear in the oldest old. This 240 suggests that other factors like genetics may have more influence on TL in the oldest old. Data of 287 241 twin pairs aged 73 to 95 years from the Danish Twin Registry showed that TL was moderately 242 heritable, suggesting genetic control of individual TL [36].

243 In contrast with some studies which found that shorter TL was associated with diabetes, 244 stroke, cardiovascular diseases, cognitive impairment, frailty and sarcopenia in the younger age 245 group (up to a mean age of 75 years) [9,37-39], our study did not find such associations in the oldest 246 old. Our findings were in line with the Leiden 85-plus study which showed that TL was not associated 247 with a history of cardiovascular diseases, hypertension, diabetes, cognitive impairment and dementia 248 in the oldest old. TL in this advanced age group was found to either shorten or lengthen by a 249 considerate amount within a few years. The authors suggested that TL is not a predictive indicator for 250 age-related morbidity, especially among the oldest old, due to a high degree of TL instability in this 251 age group [10]. Similarly, the Newcastle 85+ study showed that TL was not associated with the 252 presence of multimorbidity, cognitive impairment and disability [11]. In a large cohort of the oldest 253 old in Switzerland (mean age 85 years), no significant difference in TL was found between cognitively 254 normal individuals and those with dementia or mild cognitive impairment [40]. No study has 255 examined the association between TL and frailty and sarcopenia in the oldest old. However, the lack 256 of such association in the present study is compatible with a systematic review and meta-analysis in 257 the younger age group [12]. Findings from our study appear to support that TL should not be used unconditionally as a biomarker of the human aging process in older populations [10]. Prospective 258 259 studies are warranted to provide further information on TL attrition and its role in aging in the oldest 260 old.

261 The strengths of this study include the use of data from a relatively large sample of Chinese 262 oldest old and a series of socioeconomic and demographic factors and health outcomes. Several 263 limitations of the present study should be noted. The cross-sectional design restricts us from drawing 264 conclusions about causal relationships. Those with diseases may have passed away earlier, therefore, 265 fewer associations with diseases may be observed. The reliance on self-report of health conditions 266 may be another limitation in the current study. Furthermore, factors accumulated over the life 267 course, which may influence TL, were not captured. Dietary data at 14-year follow-up was not 268 available. However, we found that baseline dietary intake in terms of food groups and dietary 269 patterns were not associated with TL at 14-year follow-up (data not shown). In addition, our results 270 may not be generalizable to other populations as our participants were volunteers who were more 271 likely to be health-conscious and have better health status than the general population. 272 To conclude, among Chinese oldest old, males had lower TL compared with females. A lower 273

- education level and being a non-current smoker was related to higher TL. TL was not associated with
- the presence of multimorbidity, age-related diseases, frailty and sarcopenia in this age group. It
- appears that TL should not be considered a biological marker of aging among the oldest old.

276 Statements of Ethics

- 277 The subjects have given their written informed consent and that the study protocol was approved by
- the Clinical Research Ethics Committee of the Chinese University of Hong Kong (CRE 2003.102).

279 **Conflict of Interest Statement**

280 The method of telomere length measurement is pending patent proceedings.

281 Funding Sources

- 282 This work was supported by the Hong Kong Jockey Club Charities Trust. We thank the generous
- 283 donation of Ms. Therese Pei Fong Chow. The funders had no role in the design, methods, subject
- 284 recruitment, data collections, analysis or preparation of this article.

285 Author Contributions

- Authors contributed to this work are as follows: S.S.Y.Y. performed the statistical analysis,
- 287 interpreted the results, and wrote the manuscript; X.W. performed the experiments; S.L.M. prepared
- 288 the manuscript; Y.C. and S.K.W.T. performed statistical analysis; N.L.S.T. contributed to discussions
- about the results and critically revised the manuscript; J.W. designed the research, contributed to
- 290 discussions about the results and critically revised the manuscript. All the authors read and approved
- the final version submitted.

292 Data Availability Statement

- 293 The datasets generated and/or analyzed during the current study are available from the
- 294 corresponding author on reasonable request.

295 **References**

296	1	Centre for Health Protection: Life Expectancy at Birth (Male and Female), 1971 - 2020. Hong
297		Kong, Department of Health, 2021 [cited 2022 Apr 29]. Available from:
298		https://www.chp.gov.hk/en/statistics/data/10/27/111.html.
299	2	Astuti Y, Wardhana A, Watkins J, Wulaningsih W, Network PR. Cigarette smoking and
300		telomere length: A systematic review of 84 studies and meta-analysis. Environ Res.
301		2017;158:480-489.
302	3	Lin X, Zhou J, Dong B. Effect of different levels of exercise on telomere length: A systematic
303		review and meta-analysis. J Rehabil Med. 2019;51:473-478.
304	4	Robertson T, Batty GD, Der G, Fenton C, Shiels PG, Benzeval M. Is socioeconomic status
305		associated with biological aging as measured by telomere length? Epidemiol Rev. 2013;35:98-
306		111.
307	5	Gardner M, Bann D, Wiley L, Cooper R, Hardy R, Nitsch D, et al. Gender and telomere length:
308		systematic review and meta-analysis. Exp Gerontol. 2014;51:15-27.
309	6	Muezzinler A, Zaineddin AK, Brenner H. Body mass index and leukocyte telomere length in
310		adults: a systematic review and meta-analysis. Obes Rev. 2014;15:192-201.
311	7	Lopez-Otin C, Blasco MA, Partridge L, Serrano M, Kroemer G. The hallmarks of aging. Cell.
312		2013;153:1194-1217.
313	8	Wang Q, Zhan Y, Pedersen NL, Fang F, Hagg S. Telomere length and all-cause mortality: A
314		meta-analysis. Ageing Res Rev. 2018;48:11-20.
315	9	Smith L, Luchini C, Demurtas J, Soysal P, Stubbs B, Hamer M, et al. Telomere length and
316		health outcomes: An umbrella review of systematic reviews and meta-analyses of
317		observational studies. Ageing Res Rev. 2019;51:1-10.
318	10	Martin-Ruiz CM, Gussekloo J, van Heemst D, von Zglinicki T, Westendorp RG. Telomere length
319		in white blood cells is not associated with morbidity or mortality in the oldest old: a
320		population-based study. Aging Cell. 2005;4:287-290.
321	11	Martin-Ruiz C, Jagger C, Kingston A, Collerton J, Catt M, Davies K, et al. Assessment of a large
322		panel of candidate biomarkers of ageing in the Newcastle 85+ study. Mech Ageing Dev.
323		2011;132:496-502.
324	12	Lorenzi M, Bonassi S, Lorenzi T, Giovannini S, Bernabei R, Onder G. A review of telomere
325		length in sarcopenia and frailty. Biogerontology. 2018;19:209-221.
326	13	Cawthon RM. Telomere measurement by quantitative PCR. Nucleic Acids Res. 2002;30:e47.
327	14	Dagnall CL, Hicks B, Teshome K, Hutchinson AA, Gadalla SM, Khincha PP, et al. Effect of pre-
328		analytic variables on the reproducibility of qPCR relative telomere length measurement. PLOS
329		ONE. 2017;12:e0184098.

330	15	Mensà E, Latini S, Ramini D, Storci G, Bonafè M, Olivieri F. The telomere world and aging:
331		Analytical challenges and future perspectives. Ageing Research Reviews. 2019;50:27-42.
332	16	Eisenberg DTA, Kuzawa CW, Hayes MG. Improving qPCR telomere length assays: Controlling
333		for well position effects increases statistical power. Am J Hum Biol. 2015;27:570-575.
334	17	Goldman EA, Eick GN, Compton D, Kowal P, Snodgrass JJ, Eisenberg DTA, et al. Evaluating
335		minimally invasive sample collection methods for telomere length measurement. Am J Hum
336		Biol. 2018;30
337	18	Geronimus AT, Bound J, Mitchell C, Martinez-Cardoso A, Evans L, Hughes L, et al. Coming up
338		short: Comparing venous blood, dried blood spots & saliva samples for measuring telomere
339		length in health equity research. PLOS ONE. 2021;16:e0255237.
340	19	Codd V, Denniff M, Swinfield C, Warner SC, Papakonstantinou M, Sheth S, et al.
341		Measurement and initial characterization of leukocyte telomere length in 474,074
342		participants in UK Biobank. Nat Aging. 2022;2:170-179.
343	20	Nilsson M, Malmgren H, Samiotaki M, Kwiatkowski M, Chowdhary BP, Landegren U. Padlock
344		probes: circularizing oligonucleotides for localized DNA detection. Science (New York, NY).
345		1994;265:2085-2088.
346	21	Hardenbol P, Banér J, Jain M, Nilsson M, Namsaraev EA, Karlin-Neumann GA, et al.
347		Multiplexed genotyping with sequence-tagged molecular inversion probes. Nat Biotechnol.
348		2003;21:673-678.
349	22	Wong SY, Kwok T, Woo J, Lynn H, Griffith JF, Leung J, et al. Bone mineral density and the risk
350		of peripheral arterial disease in men and women: results from Mr. and Ms Os, Hong Kong.
351		Osteoporos Int. 2005;16:1933-1938.
352	23	Adler NE, Epel ES, Castellazzo G, Ickovics JR. Relationship of subjective and objective social
353		status with psychological and physiological functioning: preliminary data in healthy white
354		women. Health Psychol. 2000;19:586-592.
355	24	Washburn RA, Smith KW, Jette AM, Janney CA. The Physical Activity Scale for the Elderly
356		(PASE): development and evaluation. J Clin Epidemiol. 1993;46:153-162.
357	25	Folstein MF, Folstein SE, McHugh PR. "Mini-mental state". A practical method for grading the
358		cognitive state of patients for the clinician. J Psychiatr Res. 1975;12:189-198.
359	26	Creavin ST, Wisniewski S, Noel-Storr AH, Trevelyan CM, Hampton T, Rayment D, et al. Mini-
360		Mental State Examination (MMSE) for the detection of dementia in clinically unevaluated
361		people aged 65 and over in community and primary care populations. Cochrane Database
362		Syst Rev. 2016:CD011145.
363	27	Fried LP, Tangen CM, Walston J, Newman AB, Hirsch C, Gottdiener J, et al. Frailty in older
364		adults: evidence for a phenotype. J Gerontol A Biol Sci Med Sci. 2001;56:M146-156.

365	28	Chen LK, Liu LK, Woo J, Assantachai P, Auyeung TW, Bahyah KS, et al. Sarcopenia in Asia:
366		consensus report of the Asian Working Group for Sarcopenia. J Am Med Dir Assoc.
367		2014;15:95-101.
368	29	Gymrek M, Willems T, Guilmatre A, Zeng H, Markus B, Georgiev S, et al. Abundant
369		contribution of short tandem repeats to gene expression variation in humans. Nat Genet.
370		2016;48:22-29.
371	30	Haapanen MJ, Perala MM, Salonen MK, Guzzardi MA, Iozzo P, Kajantie E, et al. Telomere
372		length and frailty: The Helsinki Birth Cohort Study. J Am Med Dir Assoc. 2018;19:658-662.
373	31	Huang Z, Liu C, Ruan Y, Guo Y, Sun S, Shi Y, et al. Dynamics of leukocyte telomere length in
374		adults aged 50 and older: a longitudinal population-based cohort study. Geroscience.
375		2021;43:645-654.
376	32	Bello GA, Chiu Y-HM, Dumancas GG. Association of a biomarker-based frailty index with
377		telomere length in older US adults: Findings from NHANES 1999-2002. bioRxiv. 2017:191023.
378	33	Yeap BB, Hui J, Knuiman MW, Flicker L, Divitini ML, Arscott GM, et al. U-shaped relationship
379		of leukocyte telomere length with all-cause and cancer-related mortality in older men. J
380		Gerontol A Biol Sci Med Sci. 2021;76:164-171.
381	34	Woo J, Tang NL, Suen E, Leung JC, Leung PC. Telomeres and frailty. Mech Ageing Dev.
382		2008;129:642-648.
383	35	Bayne S, Liu JP. Hormones and growth factors regulate telomerase activity in ageing and
384		cancer. Mol Cell Endocrinol. 2005;240:11-22.
385	36	Bischoff C, Graakjaer J, Petersen HC, Hjelmborg J, Vaupel JW, Bohr V, et al. The heritability of
386		telomere length among the elderly and oldest-old. Twin Res Hum Genet. 2005;8:433-439.
387	37	Martinez-Ezquerro JD, Rodriguez-Castaneda A, Ortiz-Ramirez M, Sanchez-Garcia S, Rosas-
388		Vargas H, Sanchez-Arenas R, et al. Oxidative stress, telomere length, and frailty in an old age
389		population. Rev Invest Clin. 2019;71:393-401.
390	38	Haycock PC, Heydon EE, Kaptoge S, Butterworth AS, Thompson A, Willeit P. Leucocyte
391		telomere length and risk of cardiovascular disease: systematic review and meta-analysis. BMJ.
392		2014;349:g4227.
393	39	Marzetti E, Lorenzi M, Antocicco M, Bonassi S, Celi M, Mastropaolo S, et al. Shorter
394		telomeres in peripheral blood mononuclear cells from older persons with sarcopenia: results
395		from an exploratory study. Front Aging Neurosci. 2014;6:233.
396	40	Zekry D, Herrmann FR, Irminger-Finger I, Ortolan L, Genet C, Vitale AM, et al. Telomere length
397		is not predictive of dementia or MCI conversion in the oldest old. Neurobiol Aging.
398		2010;31:719-720.
399		

400 Figure Legends

- 401 Fig. 1. (A) It represents a sample with a long telomere and (B) represents another sample with a short
- 402 telomere. Template DNA of the sample showing the region with telomere repeats are shown as the
- 403 horizontal line. The length of the horizontal line represents telomere length which is longer in (A).
- 404 Open molecular inversion probes (MIPs) are added to the samples. Both 5' and 3' ends of MIP
- 405 hybridise to the target motif. Then hybridised MIPs are circularised by ligation reaction. These
- 406 circularised MIPs are then used as template for qPCR to quantify the abundance of circularised MIP.
- 407 Longer telomere (A) circularises more MIP.
- 408 Fig. 2. Telomere length (T/S ratio) by age group for males and females.



more circularized MIP. e.g. 6 MIP are circularized and serve as template for qPCR





age group (years)