## Cost and cost-effectiveness of Hepatitis C virus self-testing in four settings: an economic evaluation

Josephine G. Walker PhD<sup>1</sup>, Elena Ivanova PhD<sup>3</sup>, Muhammad S. Jamil PhD<sup>2</sup>, Associate Prof Jason J. Ong PhD<sup>4</sup>, Prof Philippa Easterbrook MD<sup>2</sup>, Emmanuel Fajardo<sup>2,3</sup>, Cheryl Case Johnson MA<sup>2</sup>, Niklas Luhmann MScPH<sup>2</sup>, Prof Fern Terris-Prestholt PhD<sup>5</sup>, Prof Peter Vickerman DPhil<sup>1</sup>, Sonjelle Shilton MPH<sup>3</sup>

- 1. Population Health Sciences, Bristol Medical School, University of Bristol, Bristol, UK
- 2. Global HIV, Hepatitis and STI Programmes, World Health Organization, Geneva, Switzerland
- 3. FIND, Geneva, Switzerland
- 4. London School of Hygiene & Tropical Medicine, London, UK
- 5. Joint UN programme on HIV and AIDS, Geneva, Switzerland

Word count: 3458

3 Figures, 4 Tables, 31 references

Funding: Unitaid and the Netherlands Government funded this study as part of the HEAD-Start project led by FIND. PV also acknowledges support from the NIHR Health Protection Research Unit in Behavioural Science and Evaluation at the University of Bristol.

Competing interests: EI and SJ are current employees of FIND, EF is a former employee of FIND. JGW and PV have received unrestricted research funding from Gilead Sciences unrelated to this research. All authors declare no other conflicts of interest. The findings and conclusions in this report are those of the authors and do not necessarily represent the official position of the World Health Organization.

1

#### Abstract (327 words)

**Introduction**: Globally, there are approximately 58 million people with chronic hepatitis C virus infection (HCV) but only 20% have been diagnosed. HCV self-testing (HCVST) could reach those who have never been tested and increase access and uptake of HCV testing services. We compared cost per HCV diagnosis or cure for HCVST versus facility-based HCV testing services.

**Methods:** We used a decision analysis model with a one-year time horizon to examine the key drivers of economic cost per diagnosis (viraemic infection) or cure following the introduction of HCVST in China (men who have sex with men), Georgia (men 40-49 years), Viet Nam (people who inject drugs, PWID), and Kenya (PWID). HCV antibody (HCVAb) prevalence ranged from 1%-60% across settings. Model parameters in each setting were informed by HCV testing and treatment programmes, HIV self-testing programmes, and expert opinion. In the base case, we assume a reactive HCVST is followed by a facility-based rapid diagnostic test (RDT) and then nucleic acid testing (NAT). We assumed oral-fluid HCVST costs of \$5.63/unit (\$0.87-\$21.43 for facility-based RDT), 62% increase in testing following HCVST introduction, 65% linkage following HCVST, and 10% replacement of facility-based testing with HCVST based on HIV studies. Parameters were varied in sensitivity analysis. Outcomes are reported in 2019 US dollars (\$).

**Results:** Cost per HCV viraemic diagnosis without HCVST ranged from \$35 (Viet Nam) to \$361 (Kenya). With HCVST, diagnosis increased resulting in incremental cost per diagnosis of \$104 in Viet Nam, \$163 in Georgia, \$587 in Kenya, and \$2,647 in China. Differences were driven by HCVAb prevalence. Switching to blood-based HCVST (\$2.25/test), increasing uptake of HCVST and linkage to facility-based care and NAT testing, or proceeding directly to NAT testing following HCVST, reduced the cost per diagnosis. The baseline incremental cost per cure was lowest in Georgia (\$1,418), similar in Viet Nam (\$2,033), and Kenya (\$2,566), and highest in China (\$4,956).

**Conclusions:** HCVST increased the number of people tested, diagnosed, and cured, but at higher cost. Introducing HCVST is more cost-effective in populations with high prevalence.

#### Key messages

#### What is already known on this topic

A recent systematic review of the impact and cost of Hepatitis C virus (HCV) self-testing (HCVST) found no prior evaluations of the cost or cost-effectiveness of HCVST. Indirect evidence can be inferred from studies of HIV self-testing (HIVST), which support the affordability and cost-effectiveness of self-testing in low and middle-income (LMIC) settings in Asia and sub-Saharan Africa. Other recent studies related to HCVST demonstrate the feasibility and usability of HCVST in various LMIC settings.

#### What this study adds

This study is the first to evaluate the cost or cost-effectiveness of HCVST. With qualityassured HCVST coming onto the market in the near future, we build on the best available evidence on potential costs for HCV testing and treatment from four LMIC settings in general and key population groups with HCVAb prevalence ranging from 1% - 60%, and show it is likely to be cost-effective (depending on willingness to pay threshold) particularly where prevalence is high.

#### How this study might affect research, practice or policy

The results of this study indicate that introducing HCVST will help countries diagnose and treat more people with HCV and contribute to achieving elimination goals faster. However, such gains will require additional investment on top of the standard of care. Targeting HCVST toward populations and settings with high HCV burden will be the most affordable option for countries. Efforts to reduce HCV treatment costs, optimize HCVST delivery and linkage services are needed to achieve greatest public health impact of offering this new approach.

#### Introduction

In 2016 the World Health Organization (WHO) launched the Global Health Sector Strategy for Viral Hepatitis with the goal to eliminate viral hepatitis B and C as a public health problem by 2030 [1]. To achieve these elimination targets, the strategy outlined the need to diagnose 90% of infected persons and treat 80% of those diagnosed, alongside scaling up prevention interventions. In 2019, there were approximately 58 million people with chronic HCV globally, of which only 21% had been diagnosed [2].

Reaching individuals not yet tested and aware of their HCV status, including those who are hesitant or unable to access facility-based services, is critical for achieving elimination. Self-testing (a process by which an individual collects his or her own specimen, performs a rapid diagnostic test, and interprets the result) is an additional testing approach to reach those who have never been tested and increase access and uptake of HCV testing services more broadly. During the COVID-19 pandemic in 2020, healthcare systems have been strained, and healthcare resources diverted to tackle the pandemic, resulting in reported disruptions to hepatitis diagnosis and treatment programmes in 43% of countries [2, 3]. Self-testing for HCV (HCVST) may help sustain or increase HCV testing rates despite these challenges, with application to both the general population and key populations, such as people who inject drugs (PWID) and men who have sex with men (MSM).

The WHO-recommended diagnostic pathway of testing someone for HCV starts with a single serological HCV test. Those who have a positive (reactive) result are then tested using HCV RNA nucleic acid testing (NAT), or core antigen, to confirm if viraemic HCV infection is present [4, 5]. While HCVST could replace initial provider-administered serology testing, facility-based confirmation of viraemic infection by NAT or core antigen test will still be necessary.

HCVST uses rapid diagnostic tests (RDT), which detect antibodies in fingerprick/capillary whole blood and/or oral fluid. With evidence emerging and quality-assured HCVST products coming to market in the near future, primarily with those adapting professional use tests for self-testing, it is important to consider the potential impact and affordability of introducing HCVST alongside existing services. In 2020, the World Health Organization began developing normative guidance on HCVST introduction[6]. A systematic review conducted as part of the guidance found no previous studies on the cost or cost-effectiveness of using HCVST[7]. To support the development of these guidelines, we conducted a cost-effectiveness analysis on the introduction of HCVST, with self-testing to be performed by an individual with or without direct support. This analysis projects the short-term costs and outcomes of introducing HCVST alongside standard diagnostic pathways, in four low and middle-income settings, including amongst key populations.

#### **Methods**

We developed a decision tree model representing the path from HCV testing to diagnosis, treatment and cure (Figures 1 and 2). We modelled four scenarios among different countries and populations with varying HCV Antibody (HCVAb) prevalence: PWID (Viet Nam and Kenya), MSM (China), and men 40-49 years (Georgia), where the national approach to HCV testing and treatment varies (Supplementary Material). The costs of distributing self-tests and standard of care testing and treatment, and the care cascade differ for each setting based on local data (Table 1, Supplementary Material). Here we report on the cost per diagnosis of viraemic infection and cost per cure, of HCVST compared to standard facility-based HCV testing pathways alone, and do not model long term outcomes such as infections or disability

adjusted life years averted. Parameter estimates were gathered from real-world examples using literature on HCV testing and treatment programmes and HIV self-testing programmes in each setting, expert opinion, and manufacturers (self-test unit costs). We incorporated data on HCVST usage from recent HCVST feasibility and acceptability trials in the study settings[8].

We explore alternative models and pathways for using HCVST, including whether HCVST is offered alongside HIVST (Kenya and China), whether confirmatory NAT testing for viraemic infection is done immediately on samples from those with a reactive facility-based serologic test (reflex testing; Kenya and Viet Nam), and whether self-tests are peer-guided (Kenya and Vietnam). In all settings, we examined two pathways of linkage from self-testing to facility-based care –first where a positive self-test leads to the standard pathway of facility-based repeat serologic testing ("repeat serologic testing") or second, directly to confirmatory NAT testing for viraemic infection ("direct to NAT"). Although core antigen testing is a WHO-recommended option for confirmation of viraemic infection, in this study, we only evaluate pathways using NAT testing as it is the main confirmation method for viraemic infection used in all the case study settings.

#### Model structure

The decision tree model (Figure 1) is based on a cross-sectional evaluation of the proportion of the population of interest who do not know their status, using testing and linkage to treatment rates expected within a one year time horizon. The population examined differs by case study setting. The model represents three pathways for the study population (Figure 1A): (1) standard of care testing, in which people receive facility-based testing and do not get self-tests; (2) no testing, for the proportion of the population who remain untested by any method; and (3) self-testing, which provides testing for a subset of the group who otherwise would not access testing, as well as replacing some of the standard of care tests with self-tests.

Those in the no testing pathway are assumed not to be diagnosed or access HCV care in the modelled year. Standard of care testing consists of facility-based anti-HCV antibody testing, with anti-HCV positive individuals receiving confirmatory HCV RNA nucleic acid testing (NAT). Those with confirmed infection are referred for pre-treatment clinical assessment, treated, and then evaluated for a sustained virological response using NAT (typically 12 weeks after end of treatment). Self-testing follows a similar pattern, but self-test results are separated into five possible outcomes depending on the modelled scenario (Figure 1A): 1) self-test result conducted but result not reported, 2) positive self-test result which links directly to confirmatory NAT testing for viraemic infection (direct to NAT scenario), 3) positive self-test result with retesting using standard of care anti-HCV antibody testing (repeat serologic testing scenario), 4) negative self-test, which is assumed not to lead to follow up testing, or 5) an invalid test result, such as if the result is not readable by the individual, in which case they would receive repeat serology testing. Details of how parameters and costs are incorporated within the model are given in Supplementary Tables 1 and 2.

#### Analysis

For each setting, we compared the introduction of HCVST in terms of numbers of individuals diagnosed with viraemic infection or cured, total cost, cost per diagnosis, and cost per cure, to a counterfactual standard of care scenario in which no self-testing occurs. Cost per diagnosis, cost per cure and the incremental cost-effectiveness ratio (ICER) were calculated. The ICER divides the difference in total cost between the HCVST and the counterfactual scenario

(incremental cost) by the difference in the number of people diagnosed/cured between the HCVST scenario and the no self-testing scenario (incremental effect). This provides a measure of the extra cost per extra person diagnosed or cured with the introduction of HCVST.

#### Base case assumptions

We use the repeat serologic testing pathway as the base case, which assumes that individuals with a reactive self-test presenting to a healthcare facility (which could include "community" testing sites such as harm reduction centres) are tested by the standard of care pathway starting from a facility-based serologic test. This facility-based serologic test is assumed to be the SD Bioline HCV rapid test (Abbott Diagnostics, IL, USA) which has an overall sensitivity of 95% (95% CI 93-96%) and specificity of 100% (95% CI 99-100%) [9].

In the base case HCVST analysis for each setting, we present the cost-effectiveness of using oral-fluid based self-tests (OraQuick® HCV Rapid Antibody Test, OraSure Technologies, PA, USA), as these were evaluated in the HCVST usability study [8]. The sensitivity and specificity of the OraQuick® test on an oral sample are reported to be 98% (95% CI 97%-99%) and 100% (95% CI 90%-100%), respectively [10]. We adjust the sensitivity and specificity to account for misinterpretation of results during self-testing, as observed by interreader agreement in the usability studies in each setting (88-98%, Table 1) [8]. In addition, we make the assumption that 3% of self-tests are used incorrectly so that no result can be reported, but which still lead to the individual linking to facility-based testing (invalid test result). The cost of the oral-fluid based HCVST was estimated to be \$4.50 plus 25% overheads to account for human resources and infrastructure (\$5.63 total), based on the authors' expert opinion, drawing on experience from HIVST and knowledge of current HCV diagnostic test pricing.

The uptake of self-testing and linkage to care parameters were determined based on randomised controlled trials of HIV self-testing, with 65% of reactive or invalid self-tests linking to facility-based testing, and self-testing leading to a 62% increase in the number of people tested [11]. Therefore, the number of people undertaking self-tests is calculated as a function of the number accessing standard of care testing in each setting and assuming 10% of people that otherwise would access standard of care testing, use self-tests instead (substitution) as in HIVST models [12]. In addition, following linkage to care, we assume no difference in treatment initiation or success parameters between the standard of care vs self-testing scenarios, as a systematic review of HIV self-testing trials showed no difference in treatment initiation for those who were self-tested (risk ratio 0.98, 95% CI 0.86-1.11) [11]. Setting-based parameter assumptions, such as the standard of care costs, HCV prevalence, and cascades of care, are presented in Table 1.

#### Sensitivity analysis

Base case assumptions were varied in one-way sensitivity analysis, to reflect uncertainty in parameters, details are presented in Table 2.

#### Costing

We gathered previously published costing data including from research studies and programme reports, identified through literature searches, previous research by the authors in each setting, and consultation with WHO focal points in each country (Table 1). Costs were identified for HCV testing and treatment, and/or HIV self-testing in each setting, accounting for overheads, staff time, training, outreach, facilities, and start-up costs where available,

from the healthcare providers' perspective. In two settings (Kenya and China), HCVST costs were assumed to be incremental adding to existing HIV self-testing programmes (Table 1).

Most cost data were identified in United States Dollars (USD) from between 2017-2019, with these being adjusted for inflation as necessary to present all costs in 2019 USD by using the consumer price index (CPI) for the study country [13]. The CPI value for Kenya was not available for 2019, so it was assumed to grow in the same ratio from 2018 as seen from 2017 to 2018. Some cost data from China were received in Chinese Yuan (RMB) [Chen, personal communication; Ong, personal communication], these were assumed to represent prices in 2019 and were converted to USD using the 2019 average exchange rate per USD (6.91 RMB per USD) from the International Monetary Fund's International Financial Statistics.

#### Results

#### Cost and outcomes of HCVST in the base case

Tables 3 and 4 show the ICERs for the base case implementation of HCVST compared to the counterfactual of no HCVST. In addition to increasing the number of individuals diagnosed, introducing HCVST increases the cost per diagnosis in all settings. In the base case, it is assumed that introducing HCVST will increase the number of individuals tested by 62%, which increases the numbers diagnosed and cured by 30.6% in Viet Nam, 34.6% in Kenya, 35.0% in Georgia, and 34.6% in China (Figure 2), due to slight differences in the cascade of care in each setting (see Table 1).

The ICER per additional person diagnosed with the introduction of HCVST is lowest in Viet Nam (\$104), \$163 in Georgia, \$587 in Kenya, and \$2,647 in China (Table 5). The variations in the cost per diagnosis by setting relate to the differences in HCV prevalence in each study setting, with cheaper costs in the settings with higher prevalence. The ICER per person cured ranges from \$1,418 in Georgia, to \$2,030 in Viet Nam, \$2,566 in Kenya, and \$4,956 in China (Table 6). The HCVST cost per cure is driven by treatment costs in each setting.

#### Costs and outcomes of the standard of care HCV testing

In contrast to the intervention, the cost per HCV diagnosis with no HCVST (excluding treatment-related costs) is estimated to be \$35 in Viet Nam, \$55 in Georgia, \$162 in China, and \$361 in Kenya. The cost per cure is more comparable across settings - \$1,238 in Georgia, \$1,839 in China, \$1,943 in Viet Nam, and \$2,284 in Kenya, as the cost of treatment is similar across most settings (between \$1,415-1,543 in Kenya, Viet Nam, and China, but approximately half at \$784, in Georgia). Of note, there is a marked difference in the cost per person receiving facility-based testing, varying 25-fold from \$0.87 in China to \$21.43 in Kenya (Table 1), due to differences in test type and variation in consumable costs.

The absolute numbers of people diagnosed (Table 3) or cured (Table 4) in one year in the absence of HCVST are dependent on the population size and prevalence in each setting. In China, this is equivalent to 53 diagnosed and 41 cured per 100,000 MSM (antibody prevalence 1.0%); in Viet Nam 18,440 diagnosed and 14,440 cured per 100,000 PWID (antibody prevalence 66.0%); in Georgia 1,333 diagnosed and 801 cured per 100,000 men aged 40-49 (antibody prevalence 22.7%); and in Kenya 3,353 diagnosed and 2,691 cured per 100,000 PWID (antibody prevalence 12.9%).

#### Sensitivity analysis

The differences in the ICER per diagnosis and cure for HCVST under the sensitivity analysis scenarios are shown in Figure 3 and Supplementary Figure 3, compared to the base case. Similar patterns are seen across all settings.

The cost per diagnosis (viraemic infection) (Figure 3) is highly sensitive to the cost of HCVST. The largest decrease in the ICER per diagnosis is seen when the HCVST price is matched to the standard of care RDT test cost, except for Kenya where this increases the ICER slightly, due to the high standard of care test cost. The ICER per diagnosis increases the most when the cost of the HCVST increases, there is low uptake of HCVST, when linkage is low, there is low performance of the HCVST or high substitution of standard of care tests with self-tests. This pattern holds across all countries, with the EIA as standard of care scenario also increasing the ICER per diagnosis in Georgia and Kenya. Low self-test success has little impact on the ICER. Reductions are seen in the ICER with use of the blood-based HCVST, even at double its usual price, as well as when there is high uptake of HCVST, high linkage to facility-based testing, high inter-reader agreement, and low substitution of standard of care tests with self-tests. Proceeding from a positive HCVST direct to NAT testing also reduces the ICER in all settings.

Differences in the ICERs in sensitivity analysis are driven by changes in the number of individuals diagnosed and the total cost. The number diagnosed is influenced most by changes in the uptake of HCVST and linkage to healthcare facilities after self-testing. The sensitivity of the ICER per cure to changes in parameters within each setting is impacted by the same factors as the cost per diagnosis. Although the relative magnitudes differ slightly, the pattern is the same as seen in the cost per diagnosis (Supplementary Figure 3).

#### Discussion

We evaluated the cost-effectiveness of HCVST compared to facility-based testing across four settings and populations with a wide range of HCVAb prevalence. Based on the assumptions that introduction of HCVST can increase the number of people who know their status, are diagnosed with chronic HCV, and are successfully treated, we found that incremental cost per HCV diagnosis for adding HCVST to the facility-based testing standard of care varied widely by setting, from \$104 in Viet Nam to \$2,647 in China, while cost per cure was lowest in Georgia (\$1,418), and highest in China (\$4,956). In all settings, HCVST resulted in more people diagnosed or cured compared to the standard of care, at a higher cost, meaning it is not cost-saving. Variation in cost per diagnosis between settings is due to differences in prevalence and test costs, while differences in the cost per cure were driven primarily by treatment costs rather than the cost of diagnosis.

The ICER (cost per diagnosis or cost per cure) of HCVST is impacted strongly by the price of the HCVST itself, and the uptake of the tests, with higher uptake leading to a reduction in the ICER. Conversely, with greater substitution of standard of care tests by self-tests, the ICER increases due to the higher cost of HCVST compared to facility-based RDT testing. The tests' performance and usability in terms of inter-reader agreement had little impact on the ICER due to generally high values for these parameters. If we do not undertake confirmatory HCV antibody testing at the facility but instead go direct to NAT, then the ICER decreases slightly because of decreases in test costs and increases in the number of people diagnosed from avoiding some false negatives resulting from the facility-based RDT.

This study is the first to evaluate the potential cost and impact of HCVST in terms of increasing access to HCV diagnosis and cure, and so by necessity, we had to make assumptions about some parameters. However, our study is strengthened because we used real-world examples for the case study settings, using locally observed costs, HCV prevalence, HCVST feasibility trials, and cascades of care for the standard of care pathway. Where local estimates were not available for undertaking HCVST, we drew on work for HIVST in the same populations, including adapting the costs of implementation of HIVST in Kenya and China.

This study has several limitations, particularly in uncertainty around parameters regarding uptake of self-testing, linkage to confirmatory testing, and in the lack of information about how HCVST will be implemented. We used data from HIV self-testing where possible, but differences between the diseases could affect the relevance of this data for HCV self-testing. In addition, the four specific case studies may not be broadly generalisable. However, they were selected to represent different populations (PWID, MSM, general population), a wide range of HCV prevalence, and different testing regimes. In addition, the study only uses a one-year time horizon, focusing on the outcomes of the number of people diagnosed and cured. We do not capture the long-term benefits of diagnosis and curing people of HCV, which will lead to reduced morbidity and mortality from end-stage liver disease and reduce onward transmission, as well as avert costs of liver disease care. In this preliminary study, it was not feasible to incorporate a model of disease progression and HCV transmission.

Although there is currently limited data and experience of using HCVST, there are key transferable lessons from work on HIV. For example, with both HIV and HCV there may be an impact of stigma on uptake of testing, multi-step testing is required, both HIV and HCV affect similar key populations, and a significant proportion maybe co-infected. However, at present there is a much higher proportion of people living with HCV who do not know their status (80%) compared to around 20% for HIV. Self-testing is useful to target those who are unlikely to otherwise access care, and this may become increasingly important as countries get closer to reaching HCV elimination targets. The possibility of a cure for HCV compared to lifelong treatment for HIV, and the largely asymptomatic nature of HCV infection will also impact testing uptake.

In the early days of HIVST research, modelling studies predicted that HIVST would be costeffective [12] and subsequent studies confirmed this. Cost-effectiveness studies of HIVST in LMIC also found the introduction of self-testing led to increased diagnosis and linkage to care, but with an additional cost to identify each case [17,26]. The first HCVST real world implementation studies are currently underway, piloting different models of HCVST distribution in Malaysia and Pakistan[14, 15]. Costing and cost-effectiveness analysis based on these and future studies will allow policymakers to make an informed decision on optimal approaches to implement HCVST in each setting. Future work on HCVST should aim to present outcomes in terms of cost per quality adjusted life year or disability adjusted life year to allow decision-makers to compare value for money across different types of interventions. Our results indicate that the introduction of HCVST may increase the overall numbers of HCV-infected people diagnosed and cured, but will require additional investment compared to the current standard of care facility-based testing and treatment pathway. These additional costs need to be minimised through ensuring that test costs are kept low and linkage to care rates are high.

#### References

- 1. World Health Organization, *Global health sector strategy on viral hepatitis 2016-*2021. Towards ending viral hepatitis. 2016, World Health Organization: Geneva.
- 2. World Health Organization, Second round of the national pulse survey on continuity of essential health services during the COVID-19 pandemic. Interim report. 2021. p. 91.
- 3. Blach, S., et al., *Impact of COVID-19 on global HCV elimination efforts*. J Hepatol, 2021. **74**(1): p. 31-36.
- 4. World Health Organization, *Guidelines on hepatitis B and C testing*. 2017, World Health Organization: Geneva.
- 5. Easterbrook, P.J. and W.H.O.G.D. Group, *Who to test and how to test for chronic hepatitis C infection 2016 WHO testing guidance for low- and middle-income countries.* J Hepatol, 2016. **65**(1 Suppl): p. S46-S66.
- 6. World Health Organization, *Recommendations and guidance on hepatitis C virus selftesting*. 2021, World Health Organization: Geneva.
- 7. Fonner, V.A., et al., *Web Annex C: Hepatitis C virus self-testing: systematic review report.*, in *Recommendations and guidance on hepatitis C virus self-testing.* 2021, World Health Organization: Geneva.
- 8. Reipold, E.I., et al. *Self-testing for HCV: multi-country evidence on usability and acceptability.* in *CROI.* 2021.
- 9. Vetter, B.N., et al., Sensitivity and specificity of rapid diagnostic tests for hepatitis C virus with or without HIV coinfection: a multicentre laboratory evaluation study. J Infect Dis, 2020.
- 10. Tang, W., et al., *Diagnostic accuracy of tests to detect Hepatitis C antibody: a metaanalysis and review of the literature.* BMC Infect Dis, 2017. **17**(Suppl 1): p. 695.
- 11. Jamil, M., et al., ANNEX 3: Should HIV self-testing be offered as an additional approach to delivering HIV testing services? A GRADE systematic review and values and preferences. 2019, World Health Organization: Geneva.
- 12. Cambiano, V., et al., Assessment of the Potential Impact and Cost-effectiveness of Self-Testing for HIV in Low-Income Countries. J Infect Dis, 2015. **212**(4): p. 570-7.
- 13. World Bank, *Consumer price index* (2010=100). 2020.
- Shilton, S., et al., Protocol for a cluster randomised controlled trial of secondary distribution of hepatitis C self-testing within the context of a house-to-house hepatitis C micro-elimination programme in Karachi, Pakistan. BMC Public Health, 2022.
  22(1): p. 696.
- 15. Shilton, S., et al., A quasi-randomised controlled trial of online distribution of homebased hepatitis C self-testing for key populations in Malaysia: a study protocol. Trials, 2022. **23**(1): p. 304.
- 16. National Statistics Office of Georgia, *Demographic Situation in Georgia (2019)*. 2020: Tbilisi. p. 185.
- 17. Stone, J., et al., *Modelling the Impact of Prevention and Treatment Interventions on HIV and Hepatitis C Virus Transmission Among People Who Inject Drugs in Kenya*. medRxiv, 2021.
- 18. Ong, J., et al., *Cost-effectiveness of community-based organization led HIV selftesting vs. facility-based HIV rapid-diagnostic testing among men who have sex with men in China.* Unpublished.
- 19. Des Jarlais, D., et al., Using dual capture/recapture studies to estimate the population size of persons who inject drugs (PWID) in the city of Hai Phong, Vietnam. Drug Alcohol Depend, 2018. **185**: p. 106-111.

- 20. Hagan, L.M., et al., *Hepatitis C prevalence and risk factors in Georgia*, 2015: setting *a baseline for elimination*. BMC Public Health, 2019. **19**(Suppl 3): p. 480.
- 21. Akiyama, M.J., et al., *Prevalence, estimated incidence, risk behaviours, and genotypic distribution of hepatitis C virus among people who inject drugs accessing harm-reduction services in Kenya: a retrospective cohort study.* Lancet Infect Dis, 2019. **19**(11): p. 1255-1263.
- 22. Jin, F., et al., *Prevalence and incidence of hepatitis C virus infection in men who have sex with men: a systematic review and meta-analysis.* Lancet Gastroenterol Hepatol, 2021. **6**(1): p. 39-56.
- Moles, J.P., et al., *HIV control programs reduce HIV incidence but not HCV incidence among people who inject drugs in HaiPhong, Vietnam.* Sci Rep, 2020. 10(1): p. 6999.
- 24. Rapoud, D., et al., *Towards HCV elimination among people who inject drugs in Hai Phong, Vietnam: study protocol for an effectiveness-implementation trial evaluating an integrated model of HCV care (DRIVE-C: DRug use & Infections in ViEtnamhepatitis C).* BMJ Open, 2020. **10**(11): p. e039234.
- 25. Averhoff, F., et al., *Progress and challenges of a pioneering hepatitis C elimination program in the country of Georgia.* J Hepatol, 2020. **72**(4): p. 680-687.
- 26. Mafirakureva, N., et al., An intensive model of care for hepatitis C virus screening and treatment with direct-acting antivirals in people who inject drugs in Nairobi, Kenya: a model-based cost-effectiveness analysis. Addiction, Unpublished/In press.
- 27. Mangenah, C., et al., *Economic cost analysis of door-to-door community-based distribution of HIV self-test kits in Malawi, Zambia and Zimbabwe*. J Int AIDS Soc, 2019. **22 Suppl 1**: p. e25255.
- 28. Tskhomelidze, I., et al., *Economic evaluation of the Hepatitis C elimination program in Georgia*. Unpublished.
- 29. Due, O.T., et al., *Cost-Utility Analysis of Direct-Acting Antivirals for Treatment of Chronic Hepatitis C Genotype 1 and 6 in Vietnam.* Value Health, 2020. **23**(9): p. 1180-1190.
- 30. Shilton, S., et al., *Feasibility and effectiveness of models of HCV viraemia testing at harm reduction sites in Georgia: a prospective 3 arm study.* Unpublished, 2019.
- 31. The Global Fund to Fight AIDS Tuberculosis and Malaria. *Price & Quality Reporting Price Reference Report*. 2021 25 February 2021]; Available from: https://www.theglobalfund.org/en/sourcing-management/price-quality-reporting/.

#### **Figures**

Figure 1: Flow charts representing scenarios analysed for the introduction of selftesting (A) and representing the pathway of care from receiving a positive HCV antibody test through to treatment and cure (B). In A, the dashed lines represent the two possible self-testing scenarios leading to either repeat serologic testing or direct to NAT, which are examined separately. Blue boxes show where B links to A. Abbreviations: HCV = Hepatitis C virus; HCVAb = HCV antibody; NAT = nucleic acid test; SVR12 = sustained virologic response (cure).



Figure 2: Cascade of care of patients tested, antibody positive, diagnosed viraemic, treated, and cured in each setting for the standard of care with no HCVST compared to the introduction of HCVST (Base case analysis). Values above each bar show the percent of previous step in cascade of care within standard of care or base case cascades (eg. percent antibody positive out of all tested, percent diagnosed viraemic out of antibody positive).



Figure 3: Tornado plot showing sensitivity analyses of how changes in parameter assumptions effects the incremental cost per diagnosis for each country. The sensitivity analyses are described in detail in Table 2. Note that the x-axis scale is different for each country, but order of y-axis is the same.



\* The vertical line represents the base case incremental cost per diagnosis, as shown in Table 5. The end of each bar represents the incremental cost per diagnosis in each modelled scenario, with the length of the bar representing the magnitude of the difference from the base case. Numbers at end of bars are the alternative values for those parameters in that sensitivity analysis.

#### Tables

# Table 1: Assumptions and parameters used in the analysis that vary by country. Parameters which do not vary by country are presented in Supplementary Table 3. Sources marked with \* represent programme data re-analysed for this study

	Kenya PWID	Georgia men 40-49	Vietnam PWID	China MSM	Source*
Transition parameters					
Size of study population HCV Antibody prevalence	13,450 13% (11-15%)	234,200 23% (18-29%)	5,000 66% (46-87%)	17,000 1% (0.6- 1.5%)	[16-19] [20-24]
Undiagnosed with HCV	71%	76%	70%	80%	[21, 25]*
Chronic hepatitis C Prevalence among those Ab+	77%	80%	84%	75%	[24-26]*
Standard of care test uptake among unknown status per year	50%	13%	50%	10%	[18, 24, 25]*
Uptake of self-tests among otherwise untested (to achieve 62% increase in testing)	31%	7.8%	31%	6.2%	[11]
Inter-reader agreement of HCVST	97%	98%	88%	97%	[8]
Receive NAT test after facility-based serologic test (assume reflex testing in Kenya and Vietnam)	100%	81%	100%	90%	[25, 26]
Link to Care if NAT positive	92%	90%	89%	90%	[21, 24]*
Start Treatment if linked to care	92%	90%	96%	90%	[21, 24]*
Cured if start treatment	95%	74%	92%	98%	[24-26]*
Not tested for SVR	0%	25%	5%	0%	[24-26]*
Cost parameters (2019 USD)					
Cost of undertaking self-test (excluding test kit)	15.46	3.00	10.00	2.52	[18, 27]
Facility-based RDT cost	21.43	2.79	2.22	0.87	
NAT test cost (diagnosis or SVR12) Pre-treatment costs – blood tests, liver	103.56	27.82	26.00	20.26	[26, 28]
disease staging, etc. [For Vietnam and China calculate as 10% of total treatment costs]	123.29	40.89	171.40	157.00	
Average treatment cost	1501.49	784.37	1542.60	1414.64	[26, 28, 29]

Scenario name	Description	Base case
		assumption
Direct to NAT	Patients are tested using NAT testing following a reactive	Repeat serologic
	self-test, without facility-based serology testing	testing at facility
EIA standard of	Standard of care antibody testing (including in the no self-	Standard of care
care	testing counterfactual for this scenario) and repeat	antibody testing by
	serologic testing are by enzyme immunoassays (EIA),	RDT (95%
	which are more expensive than RDT (assumed double	sensitivity and
	RDT test cost in Kenya and Vietnam, \$32.76 in Georgia,	100% specificity)
	and \$5.07 in China [24, 26, 30]) and have sensitivity and	
	which the PDTs are compared	
Blood-based	Using a blood-based self-test based on the PMC First	HCV/ST by oral-fluid
HCVST	Response HCV Card Test (Premier Medical Corporation	hased test with
110001	Mumbai India) which has an overall sensitivity of 96%	98% sensitivity and
	(94-97%) and specificity of 99% (99-100%) [9]. The	100% specificity
	professional use version of this tests costs an average of	and costs \$5.63
	\$0.90 in the Global Fund pricing list [31]. We assume the	··· · · · · · · · · · · · · ·
	self-test cost will be double this cost to account for	
	additional costs of packaging, plus 25% overheads	
	(\$2.25)	
High cost blood-	Use blood-based self-test but assume the total cost is	
based HCVST	doubled to \$4.50 due to uncertainty in the market price of	
	the HCVST in each setting.	
High cost oral fluid	Double the cost of oral fluid-based HCVST to \$11.25	Oral fluid-based
HCVST		HCVST is \$5.63
Equal cost oral fluid	Set the cost of HCVST including distribution costs to be	Distribution costs
HUVSI	equal to the standard of care RDT test cost in each	Vary by setting
	Setting. Reduced HCV/ST performance to 0.0% consitivity and 0.7%	
performance	specificity, reflecting reduced test performance observed	sensitivity and
performance	in samples from people co-infected with HIV [9]	100% specificity
High inter-reader	Increase the inter-reader agreement to be 100% to reflect	Inter-reader
agreement	maximum successful usage of the self-tests	agreement varies
		by setting (88-98%)
High or low linkage	Assume 50% or 80% of positive self-tests link to facility-	65% of positive
	based repeat serological testing. Higher linkage to care is	self-tests link to
	possible, particularly in Vietnam and Kenya, where testing	facility
	is assumed to be peer-led.	000/1
High or low HCVST	Increase the uptake of self-testing to reach an 80%	62% increase in
иртаке	increase in overall testing or reduce the uptake of self-	overall testing
High or low	Vary the properties of these using self-testing instead of	10% substitution
substitution	facility-based testing to be 20% or 5% while keeping the	
3003010001	increase in overall testing at 62%	
Low or high self-	Vary the proportion of invalid self-test results to be 5% or	3% of self-test
test success	1%	results invalid

## Table 2: Scenarios explored in one-way sensitivity analysis. See main text for full details of base case assumptions.

Setting	Scenario	Total cost	Total diagnosed	Incremental cost	Incremental diagnosed	Incremental cost per diagnosis (ICER)
Kenya	No HCVST	\$162,685	451	-	-	-
	Base case HCVST	\$254,194	607	\$91,509	156	\$587
Georgia	No HCVST	\$171,037	3,123	-	-	-
_	Base case HCVST	\$349,430	4,216	\$178,393	1,093	\$163
Viet	No HCVST	\$32,413	922	-	-	-
Nam	Base case HCVST	\$61,566	1,203	\$29,152	282	\$104
China	No HCVST	\$1,416	9	-	-	-
	Base case HCVST	\$9,393	12	\$7,977	3.0	\$2,647

## Table 3: Incremental cost per diagnosis of implementing HCVST (base-case analysis). Note cost per diagnosis excludes treatment costs.

#### Table 4: Incremental cost per cure of implementing HCVST (base-case analysis).

Setting	Scenario	Total cost	Total cured	Incremental cost	Incremental cured	Incremental cost per cure
Kenva	No HCVST	\$826 740	362	-	-	
Ronya	Base case HCVST	\$1.147.729	487	\$320.989	125	\$2.566
Georgia	No HCVST	\$2,322,642	1,876	-	-	-
5	Base case HCVST	\$3,254,115	2,533	\$931,473	657	\$1,418
Viet	No HCVST	\$1,402,615	722	-	-	-
Nam	Base case HCVST	\$1,850,412	943	\$447,797	221	\$2,030
China	No HCVST	\$12,785	7	-	-	-
	Base case HCVST	\$26,690	9	\$11,905	2.4	\$4,956