

Periodic epidemic outbursts explained by local saturation of clusters

Louis Gostiaux,¹ Wouter J. T. Bos,¹ and Jean-Pierre Bertoglio¹

Univ Lyon, École Centrale de Lyon, INSA Lyon, Université Claude Bernard Lyon 1, CNRS, Laboratoire de Mécanique des Fluides et d'Acoustique, UMR 5509, 36 Avenue Guy de Collongue, F-69134, ECULLY, France

(Dated: 31 August 2022)

Adding the notion of spatial locality to the susceptible-infected-removed (or SIR) model, allows to capture local saturation of an epidemic. The resulting minimum model of an epidemic, consisting of five ordinary differential equations with constant model coefficients, reproduces slowly decaying periodic outbursts, as observed in the COVID-19 or Spanish flu epidemic. It is shown that if immunity decays, even slowly, the model yields a fully periodic dynamics.

Mathematical models of epidemics remain indispensable in the fight against diseases such as the COVID-19 crisis. These models help to take political decisions such as mobility restrictions. An example of such a model is the compartmental SIR¹ model, which is the paradigm-model to illustrate the elementary dynamics of epidemic spreading in a well-mixed community.

The SIR dynamics describe the exponential increase of infected individuals upto a level where the epidemic saturates through herd-immunity. After this saturation, the number of infected individuals declines exponentially towards zero. In most epidemics, this global immunity is not attained after the first peak, or wave of infections. Indeed, another feature of epidemics is quasi-cyclic behavior with several waves of infection spaced by a period of several months. This behavior is for instance observed during the COVID-19 pandemia², the 2014 Ebola outbreak in Guinea³ as well as during the influenza epidemic in the early 20th century⁴.

The main contribution of the present work is the insight that these two features (rapid saturation and quasi-cyclic behavior) are caused by the same underlying physical phenomenon: we propose to model the global effect of local epidemic saturation. To understand how exactly this local herd immunity leads to epidemic waves, we derive and analyse a novel variant of the SIR model, introducing the minimum amount of additional complexity required to obtain successive epidemic waves.

The SIR model is too simple to describe the dynamics of a realistic epidemic, in particular since no distinction is made between susceptible individuals which are in contact, or on the contrary, far away from contagious individuals. This well-mixedness assumption can be leveraged in various ways. One possible way is to radically change the approach and consider agent-based descriptions^{5,6}. If one wants to keep a compartmental approach, diffusion can be added to the system^{7,8}, memory-effects⁹, or one can consider SIR-type models on lattices or using small-world networks¹⁰⁻¹². A minimal refinement was proposed to introduce the notion of space-dependence in the compartmental approach, dividing the community into only two groups: individuals close to, and those far away from the infected¹³. It is this approach, which successfully introduced the effect of local herd-immunity on the first wave of the epidemics, that we will improve in order to model successive outbursts.

In the domain of mathematical modeling of epidemics, various ways are explored to reproduce quasi-cyclic behavior¹⁴.

Examples are to adapt predator-prey descriptions^{15,16} or to introduce time-dependent model constants¹⁷⁻¹⁹. This latter approach needs to arbitrarily set the period and the beginning of the pandemic. However, the emergence of pandemics is in general not correlated to typical seasons, and can occur at any time²⁰. What is new here is that we do not explicitly try to model the quasi-cyclic behavior; instead we demonstrate that it is a direct consequence of the spatial dynamics of an epidemic. The present investigation suggests that saturation is attained well before global collective immunity is reached, due to local collective immunity in clusters, and that after a phase of decline, the epidemic revives when local outbreaks have decayed sufficiently to become mathematically equivalent to new small-size clusters.

a. SBIGR, the global dynamics of an epidemic in a non-fully-mixed community. The SBIGR approach can be understood as follows. The total population consists of N individuals. Within well-defined clusters the local dynamics is given by a SIR like approach. Indeed, we define a sub-ensemble N^* of individuals within local clusters around the infected. These individuals are either susceptible B , infected I or recovered/removed G . For the individuals far away from the infected, we have in our description two types: susceptibles S and removed R . It is important to understand that this sub-ensemble, further named the blob, represents multiple independent clusters with little or no connectivity, that will share similar constant physiological and social parameters, leading to similar dynamical equations.

Within the blob, corresponding to the ensemble of clusters, we have a SIR type of dynamics,

$$\frac{dB}{dt} = -\beta \frac{BI}{N^*} + \phi_{SB} \quad (1)$$

$$\frac{dI}{dt} = \beta \frac{BI}{N^*} - \gamma I \quad (2)$$

$$\frac{dG}{dt} = \gamma I - \phi_{GR} \quad (3)$$

$$(4)$$

with $N^* = B + I + G$. For a fixed total population size, we have then

$$\frac{dS}{dt} = -\phi_{SB} + \chi R \quad (5)$$

$$\frac{dR}{dt} = \phi_{GR} - \chi R \quad (6)$$

$$(7)$$




FIG. 1. Representation of the SBIGR model. The susceptible part of the population is subdivided into individuals B inside the blob, and S , outside the blob. The recovered or removed are also subdivided into individuals inside R and individuals outside Q the blob. The blob population is constituted by $N^* = B + I + R$. Model coefficients with an asterisk are normalized by N^* , e.g. $\kappa^* \equiv \kappa/N^*$. The dashed arrow from the R to the S compartment reflects the decrease of natural immunity.




FIG. 2. Graphical representation of the evolution of an (a) acceleration, or (b) deceleration epidemic. The "blob" around the infected I represents all people that are in close contact with contagious individuals. The blob expands when the local concentration of I is large. The blob will shrink if little infected are present. S and B are susceptibles, G, R are removed/recovered. Change in surface, δA during a time-interval is indicated as a shaded area.

where we have added an exchange term χR from the R to the S compartment. This term represents the decrease of immunity after infection, that we will consider null ($\chi = 0$) for the moment. Without the fluxes ϕ_{SB} and ϕ_{GR} we have a local SIR system, with B, G taking the place of S, R , respectively. Schematically, the compartments are shown in Fig. 1, where the fluxes ϕ_{SB} and ϕ_{GR} are indicated by arrows in both directions, corresponding to a growing or shrinking epidemic. The novelty of the SBIGR model is to estimate these fluxes with a physical model for the spatial evolution of the clusters, introducing a diffusion approach for the blob evolution²¹.

In Fig. 2 we have represented a visual representation of the model. We now consider that the blob evolves but that the individuals remain fixed in space and that they are uniformly spaced. The surface of the blob is A^* and the total surface containing all individuals N is denoted A . We have then that $N^*/N = A^*/A$.

On the left of Fig. 2 we illustrate the blob expansion, governed by the proximity of infected people (inside the blob, close to the border) with susceptibles (outside the blob). In a time-interval δt , it is observed that the number of susceptible individuals which were outside the blob but who will become part of it is given by

$$[\delta B]^+ = \delta A^+ \frac{S}{A}, \quad (8)$$

where S/A is the average concentration of susceptibles outside the blob. In the schematic, δA^+ corresponds to the increase of surface of the blob. This increase of surface should be, at least dimensionally, proportional to $\delta A^+ \sim \tilde{\kappa} \delta t$, where $\tilde{\kappa}$ is a diffusion coefficient. The unknown at this point is how $\tilde{\kappa}$ depends on the other parameters of the problem. What we do know is that the blob will expand when the concentration of infected individuals is large enough. We should therefore have a dependency on the local proportion of infected in the blob, I/N^* , and we have therefore in the simplest form $\tilde{\kappa} = \kappa I/N^*$ and thereby

$$\phi_{SB}^+ \approx \left[\frac{\delta B}{\delta t} \right]^+ = \frac{\kappa SI}{A N^*}. \quad (9)$$

Let us now consider the case when the blob shrinks. Similarly, the amount of individuals that were in the blob, but will find themselves outside it, are given by the decrease of surface, multiplied by the local concentration of the B . We have therefore, if the blob shrinks, that

$$[\delta S]^+ = \delta A^- \frac{B}{A^*}. \quad (10)$$

If we now model the shrinking by a negative diffusion process, independent of the presence or not of infected people, and we introduce ζ the diffusion coefficient associated with this process, we have

$$\phi_{SB}^- \approx \left[\frac{\delta S}{\delta t} \right]^+ = \zeta \frac{B}{A^*} = \frac{\zeta BN}{A N^*}. \quad (11)$$

Thereby we have modeled both the positive and the negative contributions to the flux between S and B ,

$$\phi_{SB} = \frac{\kappa IS}{A N^*} - \frac{\zeta BN}{A N^*}. \quad (12)$$

Exactly the same derivation, considering now the G, R individuals, leads to the expression

$$\phi_{GR} = -\frac{\kappa IR}{A N^*} + \frac{\zeta GN}{A N^*}. \quad (13)$$

In order to obtain these expressions we assumed that $N \gg N^*$, which is an assumption which is well satisfied in most applications. In the following, since A is fixed, we will replace the quantities κ/A and ζ/A by κ and ζ so that κ^{-1} and ζ^{-1} can be associated with the typical timescales of diffusion during expansion and shrinking, respectively.




FIG. 3. (a) Numerical integration of the SBIGR model [Eqs. (14)]. The figure shows a stackplot of the quantities B, I, G . Slowly decaying oscillations are observed for this set of parameters. In the inset it is illustrated that the time-interval T between two peaks of the number of infected individuals (indicated by dots in the main plot) is set by the parameter κ/ζ .

In the present description all terms are expressed as dimensional quantities. In order to simplify the analysis, we introduce dimensionless variables, where all compartments are normalized by N . For instance we note $\tilde{S} = S/N$, and introduce the same normalization for B, I, G, R . This naturally implies $\tilde{N}^* = N^*/N$. Omitting in the following the tildes, since we will only use normalized quantities, we have

$$\begin{aligned} \frac{dS}{dt} &= -\kappa^* SI + \zeta^* B + \chi R \\ \frac{dB}{dt} &= -\beta^* BI + \kappa^* SI - \zeta^* B \\ \frac{dI}{dt} &= \beta^* BI - \gamma I \\ \frac{dG}{dt} &= \gamma I + \kappa^* IR - \zeta^* G \\ \frac{dR}{dt} &= -\kappa^* IR + \zeta^* G - \chi R, \end{aligned} \quad (14)$$

where we introduced for the coefficients β, κ and ζ the notation $x^* \equiv x/N^* = x/(B + I + G)$.

Before numerically integrating the model, we first interpret the physical meaning of the 5 model-parameters.

The value of γ^{-1} represents the typical duration of the contagious period of an individual and β will determine the contagiousness, i.e., the rate at which an infected individual contaminates susceptible individuals. These two quantities determine to a large extent the initial, exponential phase of an epidemic. In particular, their ratio β/γ determines the reproduction number in the beginning of an epidemic. This number $\beta/\gamma = \mathcal{R}_0(0) \approx 2$, at least in the beginning of the COVID19 pandemic²². The new quantities in the SBIGR model are the growing and shrinking time-scales $1/\kappa$ and $1/\zeta$, as well as $1/\chi$, if decrease of immunity is taken into account.

The κ parameter, associated with the expansion of the blob sets the height of the first epidemic peak. Indeed, the SBIGR

dynamics allow to reproduce the saturation of an outbreak before the total population has attained herd-immunity. It is possible to estimate the peak-value of the number of infected individuals analytically (see appendix). The peak is given by

$$I_{max} \approx \frac{\kappa}{\gamma} \left(\frac{\beta - \gamma}{\beta} \right)^2 S. \quad (15)$$

The parameter ζ does not appear in this expression, which governs the short-time behavior of the epidemic wave, but is key in determining the long-time dynamics. Indeed, the blob deflates after local herd immunity has been attained, and we show in the appendix that at long times the blob evolves as

$$\frac{dN^*}{dt} \approx -\zeta, \quad (16)$$

which allows to show that for given β and γ the typical decay time scales like $T \sim \kappa/(\zeta\gamma)$.

These ideas are further assessed by numerical integration of the model. Thereto we need the definition of initial conditions and values for the control parameters. For the initial conditions we consider the case of the very beginning of an epidemic where $B(0) = G(0) = R(0) = 0$, $I(0) = I_0 \ll 1$, and $S(0) = 1 - I_0$. The parameters β and γ are determined such that $\beta/\gamma = 2$. We take $\gamma = 0.18$ which gives an order of magnitude of the infectious period of approximately 1 week (72% of the individuals have been cured or removed from the I compartment after 7 days). We set for the moment $\chi = 0$ (persistent immunity).

We illustrate the model with $\kappa = 0.002$ and $\zeta = 0.00015$ in Fig. 3. The most salient feature of this stackplot is the cyclic nature of the epidemic: even in the absence of decay of immunity ($\chi = 0$), the SBIGR model, representing the epidemic as an inflating and deflating blob, exhibits well-defined waves, characteristic for pandemics such as COVID19.

In order to understand the influence of the parameters on the period of the waves, we have systematically varied ζ for different values of κ (keeping $\kappa > \zeta$). It is shown in the inset of Fig. 3, as demonstrated in the supplementary materials, that the effective period between successive local maxima of the number of infected is proportional to κ/ζ .

A standing question is obviously whether this very simple model shows more than qualitative agreement with a realistic pandemic. However, quantitative comparison over long time-periods is not straightforward. Clearly, in the current pandemic, different countries have used different control strategies such as strict lockdowns. These would to some extent influence the parameters β and κ of the model, and possibly ζ , so that these parameters cannot be chosen strictly constant anymore. This will possibly influence the periodic character of the pandemic, and more certainly modify the relative heights of successive peaks.

Varying the model parameters as a function of time is a common way to reproduce a posteriori the evolution of an epidemic²³. This will however severely complicate the analysis of the model. It is already remarkable that the present model with constant coefficients reproduces saturation of the first wave followed by a close to periodic slowly decaying

South Africa




FIG. 4. (a) Comparison of the results of the SBIGR model [Eqs. (14)] with data of newly reported cases for South Africa. Results for different values of κ are shown for a fixed value of κ/ζ .




FIG. 5. Phase-space plot of the SBIGR model [Eqs. (14)] with decrease of immunity, $\chi > 0$. For the current parameters, the value $\chi > 0.0007$, corresponding to a typical immunity decrease time $t_{1/2} \approx 1000\text{days}$ leads to a limit cycle, representing a non-vanishing periodic epidemic. For $\chi < 0.0007$, the amplitude of the peaks decreases in time.

dynamics. We have investigated different countries and have observed that some countries show a more periodic behavior than others. One of the countries where the data of the number of daily new infected cases is most periodic is South-Africa. We keep $\beta = 0.36$, $\gamma = 0.18$ and the linear relation in the inset of Fig. 3 allows to determine the value for κ/ζ . Subsequently results for different values of κ are shown in Fig. 4.

The most important observation is that the SBIGR model allows to reproduce, for these model-parameters, the correct period of the waves and a good estimate of the order of magnitude of the number of infected individuals. Indeed the exact number is most probably underestimated significantly in the beginning of the epidemic since tests were less available. Furthermore, what our model does obviously not take into account in its simplest form, is the genetic evolution of the virus leading to variants which have different properties, thereby affecting the values of β, γ . Nevertheless, even without taking these effects into account, the results are strikingly similar, and we hope that more sophisticated models based on this framework, beyond the scope of the present work, will signif-

icantly improve our understanding of the epidemics.

b. The existence of a limit-cycle. The epidemic waves in Fig. 4 are very pronounced. Their amplitude decays and clearly the number of individuals will eventually tend to zero and stay there. Indeed, since the population-size N is fixed and lasting immunity is obtained, the epidemic will eventually die out. If we model, as in Fig. 1 the transfer from the R to the S compartment by a linear transfer term $\pm\chi R$, the natural decay of immunity can be taken into account. This can correspond both to genetic evolution of the virus, or to the evolution of the immune-system of the individuals in the R -compartment. We have added, for the same parameters as used to model the evolution of the COVID19 pandemic in South-Africa, the transfer term and we have varied the value of χ .

We illustrate using a phase-space plot in BIG space in Fig. 5 that the dynamics of our model tends to a limit-cycle for $\chi > 7 \cdot 10^{-4}$, corresponding to a $t_{1/2} \approx 10^3$, the time at which the immunity has decayed for an isolated person to 50%. This value of $t_{1/2}$ is an order of magnitude larger than estimated for the COVID19 disease, where $t_{1/2} = \mathcal{O}(10^2)^{24}$. Considering that in the current pandemic the natural immunity acquired after infection decays with a typical timescale approximately 3 months, we can conclude that, within the scope of our model, the Covid19 virus will evolve on such a limit-cycle.

Discussion & Conclusion. The main conclusion that can be drawn from the current work is that epidemic waves can be caused by the spatial nature of the spreading of the disease which will, at long times, often be slower than the local saturation. This saturation allows to decrease the local concentration of infected so that the spread slows down. However, this saturation does not, by any means indicate the end of an epidemic, since the deflation of the blob, representing the ensemble of clusters around infected individuals will eventually lead to a situation where a new spread of the disease is possible. The resulting epidemic waves can therefore not be eradicated by a lock-down which is shorter than at least several times ζ^{-1} . Obviously, the model needs to be refined before precise quantitative predictions can be formulated. The power of our approach is that, to reproduce the main features (saturation and cyclic dynamics), the SBIGR model does not need to model the effect of social-distancing, finite incubation time, demographic evolution, lock-down, seasonal fluctuations, vaccination, evolution of the virus etc. The complexity of the model remains therefore limited to 5 quantities, evolving according to 5 ODEs with constant parameters, opening a way to efficiently construct more complex models to assess the influence of different factors on epidemic waves.

Data Availability Statements The data that support the findings of this study are openly available from the Johns Hopkins University data-base².

ACKNOWLEDGMENTS

The authors thank Benjamin Pillot (UMR Espace-Dev 228) for developing the Pycovid-19 package²⁵ used to access data of the JHU-database and for useful advices. We also thank Simon Bos Sanz for suggesting to add the model for the decay

of immunity.

APPENDIX

Short time behavior: height of the first peak. In order to better understand the influence of the different parameters, we analytically estimate the influence of the peak as a function of κ assuming $\zeta = 0$. The resulting equation for B becomes then,

$$\frac{dB}{dt} = -\beta^*BI + \kappa^*SI. \quad (17)$$

A maximum for B is obtained by setting the time-derivative to zero leading to the solution for the maximum of B at time τ ,

$$B(\tau) = \frac{\kappa}{\beta}S(\tau). \quad (18)$$

In order to estimate the height of the first peak, we assume exponential increase of both I and G in this phase. The exponential of I is determined by the exponent $\beta - \gamma$ and G will increase with the same exponential time-dependence. Indeed, during this phase, we have

$$\frac{dG}{dt} \approx \gamma I, \quad (19)$$

so that, if $I \sim \exp(\beta - \gamma) \sim G$, we have

$$G = \frac{\gamma}{\beta - \gamma}I \quad (20)$$

and we assume that this approximation holds until the number of infected individuals peaks. We now set the time-derivative in the equation for I equal zero to obtain at this peak,

$$\gamma I(B + I + G) = \beta BI \quad (21)$$

since $I > 0$, and using Eqs. (20) and (18), we have at $t = \tau$,

$$I(\tau) = \frac{(\beta - \gamma)^2}{\gamma\beta}B(\tau) = \frac{(\beta - \gamma)^2}{\gamma\beta} \frac{\kappa}{\beta}S(\tau). \quad (22)$$

If, finally, we assume that the number of infected peaks approximately at the same time as B , and that $S(\tau) \approx S(0) = 1$, we obtain that the first peak of the number of infected is given by

$$I_{\max} = \frac{(\beta - \gamma)^2}{\beta^2} \frac{\kappa}{\gamma}. \quad (23)$$

This expression shows thus that the height of the first peak of the epidemic is directly proportional to the value of κ .

Time-scale of the epidemic waves. An estimate of the period of the waves can be obtained by considering the dynamics of the blob. The blob is defined as $N^* = B + I + G$. Summing the equations for B, I, G yields

$$\frac{dN^*}{dt} = \kappa I \frac{S+R}{N^*} - \zeta \frac{B+G}{N^*}. \quad (24)$$

We can assess the long-time evolution of an epidemic wave, when $I \downarrow 0$. In this limit, the second term becomes dominant, and $N^* \approx B + G$. We obtain then the expression,

$$\frac{dN^*}{dt} \approx -\zeta. \quad (25)$$

The decay of the blob is therefore linear, to that we can estimate the order of magnitude of the time T from the peak of a wave to the beginning of the next wave by

$$T \approx \frac{N_{\max}^*}{\zeta}. \quad (26)$$

In order to estimate the maximum blob size, we combine the expressions for the maximum of I, G and B (relations (20), (18) and (22)) to find the expression,

$$N_{\max}^* \approx \frac{\kappa}{\gamma}S, \quad (27)$$

so that the typical decay-time can be estimated, to be

$$T \approx \frac{\kappa}{\zeta\gamma}. \quad (28)$$

This expression is thus an order of magnitude estimate of the typical decay-time of an epidemic wave.

REFERENCES

- ¹William Ogilvy Kermack and Anderson G McKendrick. A contribution to the mathematical theory of epidemics. *Proceedings of the royal society of london. Series A, Containing papers of a mathematical and physical character*, 115(772):700–721, 1927.
- ²Ensheng Dong, Hongru Du, and Lauren Gardner. An interactive web-based dashboard to track covid-19 in real time. *The Lancet infectious diseases*, 20(5):533–534, 2020.
- ³Beth P Bell. Overview, control strategies, and lessons learned in the cdc response to the 2014–2016 ebola epidemic. *MMWR supplements*, 65, 2016.
- ⁴Jeffery K Taubenberger and David M Morens. 1918 influenza: the mother of all pandemics. *Revista Biomedica*, 17(1):69–79, 2006.
- ⁵Charles M Macal and Michael J North. Tutorial on agent-based modeling and simulation. In *Proceedings of the Winter Simulation Conference*, 2005., pages 14–pp. IEEE, 2005.
- ⁶M. Owliya, Mozafar Saadat, Rachid Anane, and M. Goharian. A new agents-based model for dynamic job allocation in manufacturing shopfloors. *IEEE Systems Journal*, 6(2):353–361, 2012.
- ⁷Julia V Noble. Geographic and temporal development of plagues. *Nature*, 250(5469):726–729, 1974.
- ⁸Alessandro De Rosis. Modeling epidemics by the lattice boltzmann method. *Phys. Rev. E*, 102:023301, Aug 2020.
- ⁹Michael Bestehorn, Thomas M Michelitsch, Bernard A Collet, Alejandro P Riascos, and Andrzej F Nowakowski. Simple model of epidemic dynamics with memory effects. *Physical Review E*, 105(2):024205, 2022.
- ¹⁰Duncan J Watts and Steven H Strogatz. Collective dynamics of ‘small-world’ networks. *nature*, 393(6684):440–442, 1998.
- ¹¹Mark EJ Newman, I Jensen, and RM Ziff. Percolation and epidemics in a two-dimensional small world. *Physical Review E*, 65(2):021904, 2002.
- ¹²Alexander A Alemi, Matthew Bierbaum, Christopher R Myers, and James P Sethna. You can run, you can hide: the epidemiology and statistical mechanics of zombies. *Physical Review E*, 92(5):052801, 2015.
- ¹³Wouter Bos, Jean-Pierre Bertoglio, and Louis Gostiaux. Modeling the role of clusters and diffusion on the evolution of covid19 infections during lock-down. *medRxiv*, 2020.

- ¹⁴Herbert W Hethcote and Simon A Levin. Periodicity in epidemiological models. In *Applied mathematical ecology*, pages 193–211. Springer, 1989.
- ¹⁵Alfred J Lotka. Analytical note on certain rhythmic relations in organic systems. *Proceedings of the National Academy of Sciences*, 6(7):410–415, 1920.
- ¹⁶Vito Volterra. Fluctuations in the abundance of a species considered mathematically. *Nature*, 118(2972):558–560, 1926.
- ¹⁷Herbert W Hethcote. Asymptotic behavior in a deterministic epidemic model. *Bulletin of Mathematical Biology*, 35:607–614, 1973.
- ¹⁸Kenneth L Cooke and James L Kaplan. A periodicity threshold theorem for epidemics and population growth. *Mathematical Biosciences*, 31(1-2):87–104, 1976.
- ¹⁹Mircea T. Sofonea, Bastien Reyn  , Baptiste Elie, Rams  s Djidjou-Demasse, Christian Selinger, Yannis Michalakakis, and Samuel Alizon. Epidemiological monitoring and control perspectives: application of a parsimonious modelling framework to the covid-19 dynamics in france. *medRxiv*, 2020.
- ²⁰Patrick R Saunders-Hastings and Daniel Krewski. Reviewing the history of pandemic influenza: understanding patterns of emergence and transmission. *Pathogens*, 5(4):66, 2016.
- ²¹Axel Brandenburg. Piecewise quadratic growth during the 2019 novel coronavirus epidemic. *Infectious Disease Modelling*, 5:681–690, 2020.
- ²²Roy M Anderson, Hans Heesterbeek, Don Klinkenberg, and T D  irdre Hollingsworth. How will country-based mitigation measures influence the course of the covid-19 epidemic? *The lancet*, 395(10228):931–934, 2020.
- ²³Kevin Linka, Mathias Peirlinck, and Ellen Kuhl. The reproduction number of COVID-19 and its correlation with public health interventions. *Computational Mechanics*, 66(4):1035–1050, October 2020.
- ²⁴Jennifer M Dan, Jose Mateus, Yu Kato, Kathryn M Hastie, Esther Dawen Yu, Caterina E Faliti, Alba Grifoni, Sydney I Ramirez, Sonya Haupt, April Frazier, et al. Immunological memory to sars-cov-2 assessed for up to 8 months after infection. *Science*, 371(6529):eabf4063, 2021.
- ²⁵Covid-19 python library by b. pillot and l. gostiaux. <https://framagit.org/benjaminpillot/covid-19>. Accessed: 2020-07-03.