

Long-Term Opioid Use and Emotional, Cognitive, and Sensorimotor Function in Chronic Back Pain

Gaelle Rached MD MSc^{1,2}, Andrew D. Vigotsky MS^{1,2,3}, Paulo Branco PhD^{1,2}, Rami Jabakhanji PhD^{1,2}, Thomas J. Schnitzer MD PhD^{1,4}, A. Vania Apkarian PhD^{1,2,4,5}, and Marwan N. Baliki PhD^{1,4,6}

¹ Center for Translational Pain Research, and Center of Excellence for Chronic Pain and Drug Abuse Research, Northwestern University Feinberg School of Medicine, Chicago, IL, USA

² Department of Neuroscience, Northwestern University Feinberg School of Medicine, Chicago, IL, USA

³ Departments of Biomedical Engineering and Statistics, Northwestern University, Evanston, IL, USA

⁴ Department of Physical Medicine and Rehabilitation, Northwestern University Feinberg School of Medicine, Chicago, IL, USA

⁵ Department of Anesthesiology, Northwestern University Feinberg School of Medicine, Chicago, IL, USA

⁶ Shirley Ryan Ability Lab, Chicago, IL, USA

Correspondence to: Dr A.V. Apkarian

E-mail : a-apkarian@northwestern.edu

Abstract

This study examines cognitive, emotional, sensory, and motor function in patients with chronic back pain (CBP) with and without long-term opioid exposure (CBP+O vs. CBP–O) and the influence of race and sex on these outcomes. We recruited 64 CBP+O (mean 7.4 years of opioid use) and 64 matched CBP–O. We collected pain intensity, pain duration, NIH Toolbox outcomes, and measures of opioid use: dosage, duration, blood levels, and withdrawal symptoms. We compared both groups to a normative sample, examined sex and race influences separately, and related CBP+O’s NIH Toolbox outcomes to their opioid use measures.

CBP+O reported greater pain interference and poorer emotional function. Sex moderated opioid exposure’s effects on cognitive performance and social satisfaction, while race moderated motor dexterity and taste outcomes. In CBP+O, 1) opioid dosage was associated with cognitive speed and sensory performance; 2) blood levels of opioids corresponded with poorer attention, motor endurance, and more severe stress and negative affect; 3) opioid use duration was related to improved working memory; while 4) higher withdrawal symptoms were associated with poorer cognitive performance, worsened negative emotions, and decreased grip strength.

The study demonstrates both positive and negative outcomes associated with long-term opioid use in CBP, highlighting the need to consider sex- and race-related factors when assessing function in this population.

Introduction

The opioid epidemic is a major public health crisis affecting the United States, with devastating health, economic, and societal costs ¹. Despite controversial evidence of efficacy, opioids are widely used for treating chronic pain and thus contribute to the ongoing epidemic ²⁻⁶. Chronic pain and opioid use have largely been studied independently compared to healthy controls, although they are intertwined. Chronic pain is associated with impairments in memory, attention, processing speed, and executive functioning ^{7, 8}. Similarly, long-term opioid prescription use is associated with declines in global cognitive performance, particularly in memory, language, and attention ⁹. In addition, prescription opioid misuse and chronic pain worsen negative affect and psychological well-being ¹⁰.

Despite overwhelming evidence that chronic pain and long-term opioid use separately impact multiple psychological domains, insufficient research characterizes both conditions concurrently. Consequently, the characteristics of long-term opioid use in chronic pain patients compared to chronic pain without opioids are poorly understood. Here, we investigated the effect of opioids on patients with chronic back pain (CBP) by comparing CBP on long-term opioids (CBP+O) to matched CBP without opioids (CBP-O) using a standard multidimensional assessment tool, the NIH toolbox, which measures cognitive, emotional, sensory, and motor function ¹¹, providing a holistic picture of neurological and behavioral health. As the patients studied were 58% female and 30% black, we also interrogated sex and race influences on NIH toolbox outcomes. Finally, we examined how opioid dosage, opioid blood levels, opioid duration, and withdrawal symptoms related to the NIH toolbox scores in CBP+O.

61

62 **Methods**

63 ***Recruitment***

64 We recruited 64 CBP+O patients (pain duration > 6 months, mean = 16.5 years) on opioid therapy
 65 for at least three months. We selected 64 CBP−O patients (mean pain duration = 16.7 years) from
 66 a previously recruited sample (Figure 1). Patients were matched on age, sex, pain intensity, and
 67 pain duration using optimal matching ¹² (Table 1). All participants were fluent in English, over 18
 68 years of age, and could understand instructions and questionnaires. Recruitment was from the
 69 Northwestern Medicine (NM) healthcare system and Shirley Ryan Ability Lab. Our exclusion
 70 criteria included (1) treatment with a spinal cord stimulator; (2) diagnosis of rheumatoid arthritis,
 71 ankylosing spondylitis, acute vertebral fractures, fibromyalgia, low back/spine oncologic history,
 72 and other comorbid neurological disorders, including major depression and psychiatric disorder
 73 requiring treatment; (3) involvement in litigation regarding their back pain, having a disability
 74 claim, or receiving workman's compensation; (4) significant comorbid diseases such as
 75 uncontrolled hypertension, unstable diabetes mellitus, renal insufficiency, congestive heart failure,
 76 coronary or peripheral vascular disease, chronic obstructive lung disease, or malignancy; (5)
 77 pregnancy; (6) Beck Depression Inventory (BDI) > 28. The protocols were approved by
 78 Northwestern University IRB (STU00207384, STU00205398).

79

80 ***Measures***

81 We collected demographic information from participants, including age, sex, race, and pain
 82 intensity (0–10 numeric rating scale, NRS) and duration. Participants completed the four
 83 neuropsychological domains of the NIH toolbox: cognition, emotion, motor, and sensation ¹¹.

Tests requiring personal interaction between the experimenter and participants were skipped during the height of the COVID-19 pandemic.

For the CBP+O group, we collected opioid duration and dosage, which we converted to morphine milligram equivalent (MME) using CDC's conversion factors ¹³. We also collected plasma to quantify systemic opioid levels. Plasma opioid concentrations were converted to a relative opiate equivalent (ROE), taking into consideration the affinity of each opioid to the mu-opioid receptor (MOR) ¹⁴ and respective molar weights ¹⁵⁻²². Withdrawal was assessed by the Subjective Opiate Withdrawal Scale (SOWS), a self-report questionnaire to rate common signs and symptoms of opiate withdrawal ²³.

Blood sample collection, analysis, and ROE calculations

We also collected blood samples from participants to quantify systemic levels of opioids. The concentrations of eight analytes (oxymorphone, hydromorphone, oxycodone, hydrocodone, fentanyl, buprenorphine, methadone and tramadol) and the presence of three others (morphine, codeine, and heroin) in plasma were determined by liquid chromatography-tandem mass spectrometry after sample preparation by solid-phase extraction, using deuterated analogs (oxymorphone-D3, hydromorphone-D6, oxycodone-D6, hydrocodone-D6, buprenorphine-D4, methadone-D3, and tramadol-C13D3) as internal standards, based on the method reported by Langman et al ²⁴. One hundred microliters of the plasma sample were diluted with 300 uL of 1% acetic acid, vortexed and loaded onto a Phenomenex Strata-X-Drug B 96-well plate. After washing with 100 mM sodium acetate buffer and methanol, the analytes were eluted with an ethyl acetate/isopropanol/ammonium hydroxide (7:2:1 v/v/v) mixture and evaporated to dryness under a stream of nitrogen. The samples were reconstituted in 15% methanol in water and analyzed by a

Sciex 6500+ Qtrap LC-MS/MS system equipped with a Shimadzu Nexera2 UPLC system. Samples were eluted from a Phenomenex Kinetex 2.6 μ m Biphenyl (50 x 2.1 mm) column with a mobile phase consisting of A: water with 0.1 % formic acid and B: methanol with 0.1 % formic acid at a flow rate of 0.6 mL/min with a gradient from 10% B to 85% B in 4.5 min. Samples were run in duplicate, and the average opioid concentration was used. The opioid concentrations were then converted to a relative opiate equivalent (ROE), taking into consideration the affinity of each opioid to the mu-opioid receptor (MOR)¹⁴, and the respective molar weights¹⁵⁻²². We realize that limiting tramadol to its MOR affinity excludes other aspects of the tramadol effect; however, we are specifically interested in the opiate activity on MOR. Samples with peak presence of opioids below the lower limit of quantitation were imputed as the halfway point from zero to the measurable lower quantifiable threshold.

Statistical Analysis

NIH toolbox differences between CBP+O and CBP-O were analyzed using multiple regression with age, gender, pain intensity, and pain duration included as covariates—the same covariates on which the participants were matched. Because we used one-to-one matching, we calculated cluster-robust variances. For each outcome, each group's score was also compared to normative data using a one-sample *t*-test, with a mean T-score of 50 and a standard deviation of 10 in the normative sample. To investigate the effects of sex and race, we performed separate multiple regressions: one with a sex-by-group interaction and another with a race-by-group interaction.

To map NIH toolbox scores onto CBP-O's opioid parameters, we calculated rank-based partial correlations (95% CI) between the NIH toolbox outcomes and measures of opioid use,

adjusted for age, sex, pain intensity, and pain duration. To do so, (1) all the measures were converted to ranks, (2) opioid use metrics and NIH toolbox metrics were residualized with respect to the covariates, and (3) we calculated Pearson correlation coefficients using the residuals. Measures of opioid use were quantified in terms of opioid dosage (MME), opioid blood level (ROE), opioid duration (years), and opioid withdrawal (SOWS). MME and ROE were log-transformed due to the skewed distributions.

Results

We independently evaluated the groups' NIH toolbox outcomes and contrasted the groups with and without sex and race interactions (figure 2). In the cognition domain, Flanker's attentional inhibitory control test was most perturbed in both groups compared to normative values, while picture vocabulary was worse in CBP+O. In the emotion domain, CBP+O was worse than CBP-O on all five tests, especially regarding social satisfaction. All six motor tests were abnormal but with divergent differences: balance and walking endurance were similarly diminished in both groups; grip strength was worse in CBP-O; and dexterity was worse in CBP+O. In the sensation domain, taste was similarly worse than normative values across both groups and pain interference was worse in both groups, but more so in CBP+O.

Sex moderated the effects of opioid exposure on social satisfaction, grip strength, and walking endurance. Race moderated the group effect on grip strength, dexterity, stress, working memory, and flanker inhibitory control.

Adjusted partial correlations within the CBP+O group examined the influence of opioid use measures on NIH toolbox outcomes (Figure 3): 1) Higher opioid use dosages were related to better processing speed, salt and odor identification, and worse dexterity. 2) Greater opioid blood

levels were related to higher stress, decreased well-being, increased negative affect, and diminished walking ability. 3) Longer durations of opioid use were related to better memory performance. 4) More severe opioid withdrawal signs (SOWS) were related to worse vocabulary and memory, worse outcomes for all five emotion measures, lower grip strength and quinine taste, and increased pain interference.

Discussion

Our study examined the association of chronic pain and long-term opioid use on cognitive, sensory, emotional, and motor functioning. Both CBP+O and CBP-O experienced more pain interference than the normative sample. Although our study groups were matched on back pain intensity and duration, CBP+O exhibited more pain interference, which we surmise may be related to opioid-induced hyperalgesia^{25, 26}; these effects were not moderated by race and sex. Within CBP+O, pain interference was most strongly related to withdrawal signs, weakly related to opioid dosage (consistent with earlier data²⁷) and opioid blood levels, and appeared to be unrelated to the duration of opioid use (contrary to an earlier report²⁸).

The emotional domain of the NIH toolbox was the dimension most perturbed by opioid exposure. All five emotion-related tests exhibited poorer outcomes in CBP+O, especially regarding social satisfaction, which was moderated by sex. Like pain interference, all five measures were associated with withdrawal and less consistently with blood levels of opioids. Given that this is a cross-sectional observational study, causal relationships remain unclear. However, given that pain interference and emotion domain were all related to state measures (blood opioids, withdrawal), it is suggested that the worsening of these outcomes is a consequence of opioid exposure.

In the motor domain, some measures were improved, and others were exacerbated in CBP+O. Moreover, these outcomes were differentially moderated by race. In the sensory taste outcomes, both CBP–O and CBP+O showed similar abnormalities moderated by race. In the cognitive domain, CBP+O and CBP–O performed poorly on attentional tasks, consistent with previous studies demonstrating that chronic pain is associated with impaired general cognitive functions^{8, 29, 30}. These results have broad implications for the daily functioning and rehabilitation of individuals with chronic pain. Previous results regarding long-term opioid use and cognition remain contradictory and inconclusive³¹⁻³⁶.

Disparities in cognitive performance among individuals with chronic pain relative to race highlight the need for further investigation and consideration of race-related factors when assessing and treating cognitive impairments in this population. Interestingly, sex effects were less prominent than race and included specific outcomes in cognition, emotion, and motor domains. Both factors remain to be studied more thoroughly in the future.

This study has three noteworthy limitations. First, our cross-sectional design makes the causality unclear. However, this is partially overcome by our matching and relating outcomes to opioid measures. Second, we excluded patients who have a clinical diagnosis of depression and have a BDI > 28, decreasing the generalizability of our results while allowing us to characterize behavior and functioning in patients with minimal psychiatric confounds. Third, our matched sample comes from a previously recruited sample. Since the data were collected by different people at different time points, potentially affecting our measured values.

Conclusion

Our study provides important insights into the impact of chronic pain and long-term opioid use on various domains of functioning. Chronic pain was associated with cognitive, sensory, emotional, and motor impairments, with attention deficits being a prominent feature. Our results suggest that long-term opioid use may not significantly contribute to cognitive impairments beyond chronic pain itself. However, long-term opioid use is associated with worse emotional functioning. Additionally, our study highlights potential race and sex disparities in cognitive performance and social satisfaction, respectively, among individuals with chronic pain. Overall, our study furthers our understanding of the complex interactions between chronic pain and long-term opioid use in different aspects of patients' lives, which can inform more effective treatment strategies for this patient population.

Table 1: Demographic characteristics of patients with chronic back pain not consuming opioids (CBP – O) and on long-term opioids (CBP + O)

	CBP - O (N=64)	CBP + O (N=64)	P-value ^a
Age (years)			
Mean (SD)	59.3 (12.6)	59.6 (11.2)	0.90
Sex			
Male	28 (43.8%)	24 (37.5%)	0.59
Female	36 (56.3%)	40 (62.5%)	
Race			0.01
Black/African American	20 (31.3%)	20 (31.3%)	
White	33 (51.6%)	43 (67.2%)	
Other	10 (15.6%)	1 (1.6%)	
Missing	1 (1.6%)	0 (0%)	
Pain Intensity (0-10)			
Mean (SD)	5.22 (2.20)	5.33 (1.98)	0.77

	CBP - O (N=64)	CBP + O (N=64)	P-value ^c
Pain Duration (years)			
Mean (SD)	16.5 (19.3)	16.7 (12.2)	0.94
Opioid Dosage (MME)			
Mean (SD)	NA	66.5 (196)	NA
Median [Min, Max] ^a	NA	20.0 [5.00, 1440]	
Missing	NA	5 (7.8%)	
Opioid Blood Level (ROE)			
Mean (SD)	NA	0.0146 (0.0736)	NA
Median [Min, Max] ^a	NA	0.000400 [0, 0.552]	
Missing	NA	7 (10.9%)	
Opioid Withdrawal (SOWS)			
Mean (SD)	NA	8.69 (8.42)	NA
Opioid Use Duration (years)			
Mean (SD)	NA	7.36 (6.14)	NA
Opioid Type (N)^b			
Buprenorphine	0	1	NA
Codeine	0	2	NA
Fentanyl Patch	0	3	NA
Hydrocodone	0	25	NA
Hydromorphone	0	1	NA
Methadone	0	2	NA
Morphine	0	3	NA
Oxycodone	0	11	NA
Oxymorphone	0	1	NA
Tapentadol	0	1	NA
Tramadol	0	25	NA

^a MME and ROE were log-transformed due to their respective skewed distribution; thus, median, and interquartile range (IQR) is also reported.

^b Eleven patients were taking two types of opioids thus the total is higher than 100%.

^c P values were calculated using a *t*-test for continuous variables and chi-square test for categorical variables. NA = not available.

Figures & Legends




Figure 1: Recruitment flowchart of patients with chronic back pain not consuming opioids (CBP – O) and on long-term opioids (CBP + O).





Figure 2: The Emotional, Cognitive, and Sensorimotor Scores in Patients with Chronic Back Pain with (CBP+O) or without Long-Term Opioid Use (CBP-O) and Their Relationships with Sex and Race.

(A) Estimated marginal means (95% CI) of the NIH toolbox components for patients with chronic back pain with (green, $n=64$) and without (pink, $n=64$) long-term opioids. Point estimates and their CIs were obtained using a linear regression adjusted for age, sex, pain intensity, and pain duration with cluster-robust variances. The grey line indicates the

mean in a normative sample (normative T-score = 50 ± 10 , $n = 1000$). Each group can thus be directly compared to the normative sample using their estimate and 95% CI. The number of participants per group per measure is shown. The second column shows group differences (CBP+O minus CBP-O, 95% CI) for each subscale. Data points with CIs not crossing the line on 0 indicate $P < 0.05$. The third column shows opioid group by sex interactions (95% CI). Positive values indicate that the CBP+O minus CBP-O is greater in females, and negative values indicate the opposite (CBP+O minus CBP-O is greater in males). The fourth column shows the opioid group by race interaction (95% CI); positive values indicate that CBP+O minus CBP-O is greater in blacks, and negative values indicate the opposite (CBP+O minus CBP-O is greater in whites). We did not include the “Other” category for race in the analysis. **(B)** Group differences in Grip-Strength Dominant, Social Satisfaction, 2min Walk Endurance stratified by sex. **(C)** Group differences in Grip-Strength Dominant, Social Satisfaction, 2min Walk Endurance stratified by race. Box plots show the upper and lower quartiles and medians, and point estimates show the mean.

A



B




Figure 3: The Relationship between Emotional, Cognitive, and Sensorimotor Scores and Opioid Dosage, Opioid Blood Level, Opioid Use Duration, and Withdrawal Signs in Patients with Chronic Back Pain with Long-Term Opioid Use (CBP+O).

(A) Partial rank correlations (95% CI) between the NIH Toolbox items and measures of opioid use adjusted for age, sex, pain intensity, and pain duration. Measures of opioid use are quantified in terms of opioid dosage (log(MME)), opioid blood level (log(ROE)), opioid duration (years) and opioid withdrawal (SOWS). The numbers of participants per NIH Toolbox test and per opioid use measure are represented on the plot. (B) Scatter plots showing the partial rank correlations between some NIH Toolbox items and measures of opioid use. Nine-Hole Pegboard Dexterity-Dominant was correlated with

opioid dosage ($\log(\text{MME})$), 2-Minute Walk Endurance with opioid blood level ($\log(\text{ROE})$),
 List Sorting Working Memory with opioid duration, and Negative Affect and Picture
 Vocabulary with Withdrawal Severity (SOWS). Solid lines in the scatter plots show fitted
 lines and shaded areas are 95% confidence intervals. Since we used partial rank
 correlations, the axes in (B) are based on residualized ranks rather than original units.

Acknowledgments:

This work made use of the Mary Beth Donnelley Clinical Pharmacology Core at Northwestern University, which has received support from the NIH (1S10OD012016-01 / 1S10RR019071-01A1), Soft and Hybrid Nanotechnology Experimental (SHyNE) Resource (NSF ECCS-1542205), the State of Illinois, and the International Institute for Nanotechnology (IIN).

Meghan Ford, data collection

Elizabeth Yan, data collection

Maryam Abdallah, data collection

Byron K Yip, data collection

Santiago Espinosa, data collection

Rahia Parveen Shuaib, data collection

References

1. Volkow ND, Blanco C. The changing opioid crisis: development, challenges and opportunities. *Mol Psychiatry*. 2021;26(1):218-33. Epub 20200204. doi: 10.1038/s41380-020-0661-4. PubMed PMID: 32020048; PMCID: PMC7398847.

2. Nury E, Schmucker C, Nagavci B, Motschall E, Nitschke K, Schulte E, Wegwarth O, Meerpohl JJ. Efficacy and safety of strong opioids for chronic noncancer pain and chronic low back pain: a systematic review and meta-analyses. *Pain*. 2022;163(4):610-36. Epub 2021/07/31. doi: 10.1097/j.pain.0000000000002423. PubMed PMID: 34326292.

3. Volkow ND, McLellan AT. Opioid Abuse in Chronic Pain — Misconceptions and Mitigation Strategies. *New England Journal of Medicine*. 2016;374(13):1253-63. doi: 10.1056/NEJMr1507771. PubMed PMID: 27028915.

- 325 4. Nagpal AS, Eckmann M, Weinstein SM. Maintaining Opioid Prescription for Chronic
326 Back Pain: Pro Versus Con. PM R. 2018;10(1):83-9. doi: 10.1016/j.pmrj.2017.12.002. PubMed
327 PMID: 29413121.
- 328 5. Voon P, Karamouzian M, Kerr T. Chronic pain and opioid misuse: a review of reviews.
329 Subst Abuse Treat Prev Policy. 2017;12(1):36. Epub 20170815. doi: 10.1186/s13011-017-0120-
330 7. PubMed PMID: 28810899; PMCID: PMC5558770.
- 331 6. Dahlhamer JM, Connor EM, Bose J, Lucas JL, Zelaya CE. Prescription Opioid Use Among
332 Adults With Chronic Pain: United States, 2019. Natl Health Stat Report. 2021(162):1-9. Epub
333 2021/09/16. PubMed PMID: 34524076.
- 334 7. Higgins DM, Martin AM, Baker DG, Vasterling JJ, Risbrough V. The Relationship
335 Between Chronic Pain and Neurocognitive Function: A Systematic Review. Clin J Pain.
336 2018;34(3):262-75. doi: 10.1097/ajp.0000000000000536. PubMed PMID: 28719507; PMCID:
337 PMC5771985.
- 338 8. Moriarty O, McGuire BE, Finn DP. The effect of pain on cognitive function: A review of
339 clinical and preclinical research. Progress in Neurobiology. 2011;93(3):385-404. doi:
340 <https://doi.org/10.1016/j.pneurobio.2011.01.002>.
- 341 9. Warner NS, Hanson AC, Schulte PJ, Habermann EB, Warner DO, Mielke MM.
342 Prescription opioids and longitudinal changes in cognitive function in older adults: A population-
343 based observational study. Journal of the American Geriatrics Society. 2022;70(12):3526-37. doi:
344 <https://doi.org/10.1111/jgs.18030>.
- 345 10. Martel MO, Dolman AJ, Edwards RR, Jamison RN, Wasan AD. The Association Between
346 Negative Affect and Prescription Opioid Misuse in Patients With Chronic Pain: The Mediating

- 347 Role of Opioid Craving. The Journal of Pain. 2014;15(1):90-100. doi:
348 <https://doi.org/10.1016/j.jpain.2013.09.014>.
- 349 11. NIH. NIH Toolbox® Scoring and Interpretation Guide for the iPad: National Institutes of
350 Health and Northwestern University
351 ; 2021 [updated 5/25/2021; cited 2022 9/8/2022].
- 352 12. Ho D, Imai K, King G, Stuart EA. MatchIt: Nonparametric Preprocessing for Parametric
353 Causal Inference. Journal of Statistical Software. 2011;42(8):1 - 28. doi: 10.18637/jss.v042.i08.
- 354 13. Prevention CfDCA. Analyzing prescription data and morphine milligram equivalents
355 (MME)2021.
- 356 14. Volpe DA, Tobin GAM, Mellon RD, Katki AG, Parker RJ, Colatsky T, Kropp TJ, Verbois
357 SL. Uniform assessment and ranking of opioid Mu receptor binding constants for selected opioid
358 drugs. Regulatory Toxicology and Pharmacology. 2011;59(3):385-90. doi:
359 <https://doi.org/10.1016/j.yrtph.2010.12.007>.
- 360 15. Information NCfB. PubChem Compound Summary for CID 5284604, Oxymorphone.
361 2022.
- 362 16. Information NCfB. PubChem Compound Summary for CID 5284603, Oxycodone2022.
- 363 17. Information NCfB. PubChem Compound Summary for CID 3345, Fentanyl2022.
- 364 18. Information NCfB. PubChem Compound Summary for CID 5284570,
365 Hydromorphone2022.
- 366 19. Information NCfB. PubChem Compound Summary for CID 5284569, Hydrocodone2022.
- 367 20. Information NCfB. PubChem Compound Summary for CID 5284371, Codeine2022.
- 368 21. Information NCfB. PubChem Compound Summary for CID 33741, Tramadol2022.
- 369 22. Information NCfB. PubChem Compound Summary for CID 4095, Methadone2022.

23. Handelsman L, Cochrane KJ, Aronson MJ, Ness R, Rubinstein KJ, Kanof PD. Two new rating scales for opiate withdrawal. *Am J Drug Alcohol Abuse*. 1987;13(3):293-308. doi: 10.3109/00952998709001515. PubMed PMID: 3687892.
24. Langman LJ, Korman E, Stauble ME, Boswell MV, Baumgartner RN, Jortani SA. Therapeutic monitoring of opioids: a sensitive LC-MS/MS method for quantitation of several opioids including hydrocodone and its metabolites. *Ther Drug Monit*. 2013;35(3):352-9. doi: 10.1097/FTD.0b013e318283e29a. PubMed PMID: 23666565.
25. Lee M, Silverman SM, Hansen H, Patel VB, Manchikanti L. A comprehensive review of opioid-induced hyperalgesia. *Pain Physician*. 2011;14(2):145-61. PubMed PMID: 21412369.
26. Wilson SH, Hellman KM, James D, Adler AC, Chandrakantan A. Mechanisms, diagnosis, prevention and management of perioperative opioid-induced hyperalgesia. *Pain Manag*. 2021;11(4):405-17. Epub 20210329. doi: 10.2217/pmt-2020-0105. PubMed PMID: 33779215; PMCID: PMC8023328.
27. Denawa Y, Kurtz W, Conermann T. The Social and Functional Implications of High-Versus Low-Dose Opioids on Chronic Non-Cancer Pain. *Pain Physician*. 2019;22(4):401-11. PubMed PMID: 31337176.
28. Morasco BJ, Yarborough BJ, Smith NX, Dobscha SK, Deyo RA, Perrin NA, Green CA. Higher Prescription Opioid Dose is Associated With Worse Patient-Reported Pain Outcomes and More Health Care Utilization. *The Journal of Pain*. 2017;18(4):437-45. doi: <https://doi.org/10.1016/j.jpain.2016.12.004>.
29. Prefontaine K, Rochette A. A Literature Review on Chronic Pain: The Daily Overcoming of a Complex Problem. *British Journal of Occupational Therapy*. 2013;76(6):280-6. doi: 10.4276/030802213x13706169932905.

30. Sjøgren P, Christrup LL, Petersen MA, Højsted J. Neuropsychological assessment of chronic non-malignant pain patients treated in a multidisciplinary pain centre. *Eur J Pain*. 2005;9(4):453-62. Epub 20041105. doi: 10.1016/j.ejpain.2004.10.005. PubMed PMID: 15979026.
31. Kendall SE, Sjøgren P, Pimenta CAM, Højsted J, Kurita GP. The cognitive effects of opioids in chronic non-cancer pain. *Pain*. 2010;150(2):225-30. doi: 10.1016/j.pain.2010.05.012. PubMed PMID: 20554115.
32. Jamison RN, Schein JR, Vallow S, Ascher S, Vorsanger GJ, Katz NP. Neuropsychological effects of long-term opioid use in chronic pain patients. *J Pain Symptom Manage*. 2003;26(4):913-21. doi: 10.1016/s0885-3924(03)00310-5. PubMed PMID: 14527760.
33. Rowbotham MC, Twilling L, Davies PS, Reisner L, Taylor K, Mohr D. Oral opioid therapy for chronic peripheral and central neuropathic pain. *N Engl J Med*. 2003;348(13):1223-32. doi: 10.1056/NEJMoA021420. PubMed PMID: 12660386.
34. Tassain V, Attal N, Fletcher D, Brasseur L, Dégieux P, Chauvin M, Bouhassira D. Long term effects of oral sustained release morphine on neuropsychological performance in patients with chronic non-cancer pain. *Pain*. 2003;104(1-2):389-400. doi: 10.1016/s0304-3959(03)00047-2. PubMed PMID: 12855350.
35. Schiltenwolf M, Akbar M, Hug A, Pfuller U, Gantz S, Neubauer E, Flor H, Wang H. Evidence of Specific Cognitive Deficits in Patients with Chronic Low Back Pain under Long-Term Substitution Treatment of Opioids. *Pain physician*. 2014;17:9-20. doi: 10.36076/ppj.2014/17/9.
36. Lee DM, Pendleton N, Tajar A, O'Neill TW, O'Connor DB, Bartfai G, Boonen S, Casanueva FF, Finn JD, Forti G, Giwercman A, Han TS, Huhtaniemi IT, Kula K, Lean MEJ, Punab M, Silman AJ, Vanderschueren D, Moseley CM, Wu FCW, McBeth J. Chronic widespread pain is associated with slower cognitive processing speed in middle-aged and older European men.

416 Pain. 2010;151(1):30-6. Epub 20100619. doi: 10.1016/j.pain.2010.04.024. PubMed PMID:
417 20646831.
418