

Knowledge Mapping of Nigeria's Scientific Contribution to Antimicrobial Resistance Research: A visualized investigation using VOS viewer and Cite Space

Adeiza Shuaibu Suleiman^{1*}, Shuaibu Mohammed Gazali², Shuaibu Bello Abdulmalik³

^{*1} Department of Pharmaceutical Microbiology, Faculty of Pharmaceutical Sciences Ahmadu Bello University Zaria.

²Department of Microbiology, Faculty of Chemical and Life sciences, Usmanu Danfodiyo University, Sokoto, Nigeria

³Department of Veterinary Microbiology, Faculty of Veterinary Medicine, Usmanu Danfodiyo University, Sokoto, Nigeria

Adeiza Shuaibu Suleiman^{1*} = Corresponding author

email = suleykestler2@gmail.com

ABSTRACT

Objective: A comprehensive overview of the research output, including conceptual and temporal trends, is required to address the growing AMR burden in Nigeria, which was previously lacking. This study uses bibliometric analysis to show this.

Methods: Data from a query of AMR articles in the Dimensions database (PubMed included) were subjected to bibliometric analysis (1972–2022). Bibliometric indicators, topic networks were established and topic trends were analysed by frequency, relevance scores, and trends over time.

Results: A total of 0.2% of the papers on AMR published worldwide were written by authors or institutions from Nigeria. In 2021, publications grew by 13.6%, highlighting the field's increased vigour. The vast majority of publications (57.18%) were in the field of Medical and Health Sciences, with 2428 papers. The top journal for AMR papers with at least one Nigerian institution affiliation was African Journal of Clinical and Experimental Microbiology, which published 1.8% of these papers. Antimicrobial stewardship, Clinical and laboratory practices on AMR, Public health implications, Traditional and molecular methods, and Phytomedicine and drug discovery were prominent areas of focus for Nigerian researchers.

Conclusion: This study provides a comprehensive overview of the AMR research output, highlighting the need of increased research capacity to address the burden of AMR in Nigeria. Because Nigeria is overly reliant on conventional methods of discovery and development, innovation is essential for the future of antibiotics.

Keywords: Antimicrobial Resistance, Bibliometrics, antimicrobial stewardship, drug discovery, Nigeria

Knowledge Mapping of Nigeria's Scientific Contribution to Antimicrobial Resistance Research. A visualized investigation using VOS viewer and Cite Space

Introduction

Microbes can become resistant to antimicrobials through mutation or adaptation, which renders treatments ineffective. This phenomenon is known as antimicrobial resistance (AMR)[1]. The World Health Organization (WHO) estimates that Antimicrobial Resistance kills over 700,000 people across the globe yearly, an estimate that may well soar to 10 million by the year 2050[2]. According to a needs assessment on antimicrobial use and resistance in Nigeria published in 2017 by the federal ministries of health, agriculture, and the environment, certain antimicrobial agents have increased resistance rates of 70% to 100% in certain states[3]. In a country where infectious diseases are a major cause of morbidity and mortality, Nigeria considers AMR a priority on the national public health agenda[3]. The Nigerian response to antimicrobial resistance led by the Nigeria Centre for Disease Control (NCDC) began in 2016 following a situational analysis that investigated common antimicrobial-resistant pathogens recovered from hospitals, animal sources, agricultural and environmental sources[4]. Overall, the National Action Plan (NAP) strives to address five key pillars in accordance with the WHO Global Action Plan on Antimicrobial Resistance[5]. The fundamentals entail boosting knowledge and awareness of AMR among health personnel and the general public; developing a One Health surveillance system; strengthening prevention efforts, control, and biosecurity; promoting rational use of antimicrobials and access to quality medications; and conducting research into antimicrobial alternatives, new diagnostics, and drug discovery[4]. The problem of AMR in Nigeria is multisectoral.

67 The ease with which antibiotics are available and can be purchased without the need
68 for a prescription thrives in an environment where the antibiotic market is poorly
69 regulated. The problem of improper disposal of expired antibiotics and left-over
70 medicines by the general public, which mostly end up in conventional waste bins and
71 dumps due to lack of awareness on the part of the users and weak enforcement on
72 the part from regulatory authorities[6]. The problem of indiscriminate use impacts
73 human health and animal welfare alike [7]. In the human health sector, many
74 incidences of fever are addressed empirically with antibiotics[8]. Although a few
75 antimicrobials are prescribed by a doctor, patients and their families frequently press
76 healthcare professionals for antimicrobials and, in most cases, self-medicate with
77 antimicrobials even for mild viral infections[9]. Even with the Nigerian Veterinary
78 Council's warnings against excessive use of antimicrobials in the animal health
79 sector, several farmers habitually add antimicrobials into animal feed or water to
80 promote growth and, in the case of poultry farming, to boost egg size to ensure
81 better profit. Several farmers in the country use antimicrobials in post-harvest
82 storage to preserve the yield[10]. As a result, human and animal waste, particles
83 from crop sprays, discharge from health facilities, pharmaceutical companies, and
84 farmlands carrying antimicrobial-resistant pathogenic microbes and antimicrobial
85 residues find their way into the ecosystem[11]. It thus results in successions of
86 negative impact on food production, bodies of water, as well as fish farms[11].
87 Despite increased awareness of antimicrobial resistance, the general situation is
88 deteriorating, and the scientific community and Institutions have endured in research
89 and publications to fashion newer antimicrobial peptides (AMPs), antibiotic
90 combinations, and monitoring systems to effectively control AMR[12]. The amount of
91 reliance on data increases with the growth of scientific publications and information

technologies in healthcare. These big data raise various issues to be resolved by innovative big data analytics, like Knowledge mapping via bibliometrics.

Knowledge mapping is an assessment tool that blends concepts and techniques from mathematics, visuals, and other disciplines with bibliometric methods[13]. Cite Space, and VOS viewer are the most popular bibliometric analysis tool for Knowledge mapping via bibliometrics[14]. Knowledge mapping has been applied in many research fields[13,15,16]. To date, a knowledge mapping analysis of AMR research contributions and inclinations in Nigeria has remained hidden. This study aimed at filling this gap. Understanding Nigeria's contribution to AMR research requires quantifying and mapping AMR-related publications. This study examines the research conducted by institutions in Nigeria using the Dimensions database as a foundation. We focus on bibliometric elements like international collaboration, subject area classification, active authors and institutions, journals and highly cited papers. In this study, we generated visualized knowledge maps of AMR research in Nigeria and analysed scientific hotspots, emerging/future trends, prolific authors and key research institutions of momentous research activities in AMR, which we hope will provide clarity about the nation's priority for AMR research; and serve as reference for future collaborative networks.

METHODS

Data acquisition and query criteria

Bibliographic data on antimicrobial resistance publications in Nigeria were retrieved from the Dimensions™ database on August 20th 2022. The database is one of the most comprehensive for bibliometric and patentometric analysis. It contains millions of publications cited over 1.6 billion times linked with Patents, clinical trials, policy

documents, datasets and grants [15,17]. This database is completely inclusive of PubMed, making it highly reliable. We used the search criteria: ‘antibiotics OR antifungal OR Antibacterial OR antimicrobial resistance AND Nigeria’ to spotlight research articles published between 1972 and 2022. The search was not limited by language, and it included all accessible data sets. Both title and abstracts were search to ensure subject-specific outcome for the benefit of accuracy. Publications indexed in the database that featured a minimum of one author connected to a Nigerian institution, including a university or organization, qualified for inclusion.

Data processing and parsing

The dplyr and ggplot2 packages of R- studio v.4.1.2 (2022-02-16) was used to examine, parse and visualize bibliographic data. Imported raw files were cured of false positives and duplicates. Top 100 documents were arbitrarily selected per year and checked for presence of false positive results. A key term was ascribed to noun-phrases with similar meanings (e.g., "Antimicrobial sensitivity test ", and “AST” were merged to “Antimicrobial susceptibility test “). Multiple or similar keyword occurrences in a publication were consolidated as a single count. Network mapping and visualizations were rendered with Visualization of Similarities (VOS) Viewer software (Leiden University, Netherlands)[18]. This method adopts diverse clusters and Color spectrum ; the bubble size symbolizes the total frequency, the colours indicate the cluster type, and the lines of conjunct (straight or curved) illustrates connectedness [19]. Indices like, scientific production, mean citations, authors productivity, affiliation and countries, preferred journals and keywords, were used to assess the present and likely future trends of AMR research in Nigeria [20,21]. The terms used in keyword clustering analysis were extracted from title and abstract

fields of included bibliographic documents. copyright statements and Structured abstract labels were ignored in the analysis. Full counting method with minimum occurrence threshold of 20 ($T \geq 20$) was used. The top 60% of calculated relevance scores were used in the analysis.

In order to determine the burstness and centrality index of the bibliographic components, Cite Space (Version 6.1.R3 Basic) was used[22]. The following limits were set: time span (1972 to 2022), years per slice (one), pruning (pathfinder and Pruning Sliced Networks), selection criteria (g-index), link strength (cosine) and scope (within slices) and other default settings.

Furthermore, the top cited articles were also extracted and sorted by field citation ratio. The Field Citation Ratio (FCR) and Relative Citation Ratio (RCR) are citation-based measures of an articles' scholarly impact[23]. FCR was determined by dividing the number of citations received by an article by average number of citations received by articles published within the same year and Fields of Research (FoR) class. While the RCR is measured as the number of citations received by a paper adjusted to the citation count garnered by National Institute of Health (NIH) funded publications in the same research field and year[23].

RESULTS

Nigeria's annual contribution in AMR Research

Nigerian authors and institutions have played a key role in production of a total of 4,246 papers on antimicrobial resistance. As noted in Figure 1, AMR-related research in Nigeria have shown an annual upward tendency that peaked in 2021(13.6%). Based on our research, there are 2,176,583 publications on AMR globally and Nigeria have contributed at least 0.2% of that number. Table S1 shows

the distribution of scientific output by field of research; Medical and Health Sciences (n = 2428, 57.18%), Biological Sciences (n = 806, 18.98%), Agricultural and Veterinary Sciences (n = 409, 9.63%).

Influential Contributor Analyses

1. Authors and Co-Cited Authors contributing to AMR research

A total of 14391 authors were involved in AMR research. OKEKE IN published the most papers indexed in Dimension database (n = 31), followed by LAMIKANRA A (n = 28), and OLAYINKA BO (n = 27) (Figure 1). The top ranked author (cited) by centrality is BAUER AW (1985) with centrality of 0.13 followed by Okeke IN (2000), with centrality of 0.07 (Figure 3). The co-authorship network map was constructed authors (n = 1000) who had published at least five papers on AMR (Figure 4). Authors that share the same colour space represent the same cluster. There were active collaborations in AMR research in Nigeria, especially among authors in the same cluster, such as OKEKE IN AND LAMIKANRA A (green), SOWUNMI A and ODUOLA AMJ (coffee brown), ILIYASU G and HABIB AG (lime green), OLAYINKA BO AND ONAOLAPO JA (yellow) etc.

Authors with citation bursts are defined as those that are cited frequently over a while. Through detecting bursts, Table 1 lists the top Authors of publications on AMR with the strongest citation bursts. COKER A on the top of the list with the maximum burst duration of 15 years and strength of 4.54, followed by OGUNSOLA F with a 14-year citation burst with 4.45 burst strength. NGWAI Y had a minimum burst duration of 1 year and the highest strength of 8.7. Recently published authors of AMR research with the strongest burst were NGWAI Y (8.7), OLAYINKA B (8.53) and OKEKE IN (7.27). This shows that their work may have formed a hot and leading

topic. The table also show a list the top cited authors of publications in AMR with the strongest citation bursts. AKO-NAI AK on the top of the list with the maximum burst duration of 18 years and strength of 6.5, followed by TENOVER FC A with an 8-year citation burst with 5.86 burst strength. OKEKE IN had the strongest burst 11.6 among cited authors.

2. Contributions and Cooperation Network Among Institutions/Organizations

The top ranked institutions/organizations in AMR research collaboration by centrality are University of Ibadan (1972), with centrality of 0.33. The second one is Ahmadu Bello University (1972), with centrality of 0.14. The third is University of Lagos (1985), with centrality of 0.13. The 4th is International Institute of Tropical Agriculture (1975), with centrality of 0.13. The 5th is Obafemi Awolowo University (1982), with centrality of 0.10 (Figure 3). Figure 5, depicts the cooperation network among Nigerian institutions/organizations on AMR research, where the nodes represent the institutions/organizations, and the sizes denote number of publications. The lines between two institutions/organizations denote that they cooperate. The density of the links of an institutions/organization indicates the cooperation degree. The denser of the lines, the more collaboration for one institutions/organization with other institutions/organization. From the figure, University of Ibadan (n =366, links = 1586), University of Lagos (n =218, links = 1079), University of Nigeria (n =172, links = 430), Ahmadu Bello University (n =152, links = 647), Obafemi Awolowo University (n =141, links = 566) and Nigerian Institute of Medical Research (n =132, links = 883) have more lines. Therefore, they have more cooperation with other institutions/organizations.

3. Analysis of Top Journals publishing Nigeria's contribution to AMR research

By analysing the distribution of journals accepting Nigeria's contribution in this field, we could accurately identify the main part of the academic research, and papers published in such journals that could be supported by academic research. Figure 2 shows the top journals that published Nigeria's contributions in AMR research. The number of publications in these journals was about 21.83% of the total number of contributions in this field. The number of articles published in African Journal of Clinical and Experimental Microbiology was the largest, with 75 articles, accounting for 1.8% of the total. Followed by PLoS One and African Journal of Medicine and Medical Sciences, with 50 articles each (1.2%). In terms of mean citation, American Journal of Tropical Medicine and Hygiene had the highest (33.3%).

4. AMR Research themes and trends in Nigeria

Keywords are the condensation and reaction to the main content of the article, which can reflect the hot topic and the development trend related to the research field. We ran keyword co-occurrence network analysis with VOS viewer and arrived with a graph with 953 nodes and 6 Clusters (Figure 6). The keywords "*Escherichia coli*" and "*Staphylococcus aureus*" displayed the highest frequency of 881 and 540, respectively. Followed by the keywords "ESBL" (470), "*Salmonella*" (468), "tetracycline" (465) and "plasmid" (412). Thematic research group of focus by Nigeria authors are; Antimicrobial stewardship (25.50%), Clinical and laboratory (25.39%), public health implication (23.92%), traditional and molecular methods (16.79%) and phytomedicine and drug discovery (8.39%). Additional insights on the clusters are described below;

a) Antimicrobial stewardship theme-based cluster (red, n= 243 items, 25.50%)

Antimicrobial stewardship, antibiotic prescription, antimicrobial resistance, aquatic environment, antimicrobial usage, attitude, awareness, burden, compliance, combination, evaluation, knowledge, middle income country, misuse, monitoring, mortality, observation, perception, point prevalence, practice, prescription, rational use, self-medication, standard, surveillance, survey, Systematic review, world health organization etc.

b) Clinical and laboratory theme-based cluster (green, n= 242 items, 25.39%)

Patients, outpatient, treatment, case, *Acinetobacter baumannii*, 30ug, Adult, age group, *Aeromonas hydrophilia*, amoxicillin, amoxiclav, ampicillin, antibiotic, antenatal clinic, antibiogram, antibiotics resistance profile, antibiotic susceptibility test, antibiotic resistance, Augmentin, azithromycin, bacillus, bacteraemia, bacteria isolate, bacteria pathogen, bacterial infection, bacterial resistance, biofilm, blood agar, cefepime, cefixime, cephalosporin, cerebrospinal fluid, choice, cloxacillin, conventional biochemical test, cotrimoxazole, disk diffusion method, drug resistance, *Enterobacter* spp, first line drugs, gram stain, high resistance, high vaginal swab, erythromycin, intensive care unit, *Klebsiella*, Kirby Bauer disc diffusion method, levofloxacin, microbiology laboratory, multi drug resistant, pattern, otitis medium, *Pseudomonas* spp, *Shigella* spp, standard microbiological techniques, samples, UTI, wound infection, swab etc.

c) Public health implication theme-based cluster (blue, n= 228 items, 23.92%)

Isolate, strain, gene, animal, human, Abattoir environment, antibiotics resistance gene, animal production, antimicrobial residue, aquaculture, biofilm formation, cattle, coliform, environmental sample, faecal sample, ESBL, farm, fluoroquinolones, food handlers, gastroenteritis, indiscriminate use, livestock, *Listeria spp*, meat sample, MLST, market, oxytetracycline, phenotypic characterization, poultry environment, pork, public health implication, raw meat, reservoir, resistance profile, *Salmonella sp.* salmonellosis, seafood, serotyping, whole genome sequencing, standard bacteriological method, *Vibrio spp*, significance, virulence gene, resistance genes etc.

d) Traditional and molecular methods cluster (yellow, n= 160 items, 16.79%)

Molecular characterization, resistance gene, molecular detection, multiple antibiotics resistance gene, amp gene, Betalactamase, *blaZ*, *ctxm*, *bla-shv*, *bla tem*, PCR, carbapenem, cefotaxime, CIP, clinical infection, clinical isolate, clinical sample, detection, plasmid profile, plasmid, *pmqr*, sequencing, dissemination, double disc synergy test, *E. coli*, ESBI, gram negative, integrons, laboratory standard institute, MDR, MARI, *mecA*, gene, pathogens of public health concern, methicillin resistant *Staphylococcus aureus*, nasal swab, vancomycin, carriage, etc.

e) Phytomedicine and drug discovery themed cluster (purple, n= 80 items, 8.39%)

Plant extract, aqueous extract, antibacterial activity, *Candida spp*, concentration, efficacy, ethanol, flavonoid, Leafe, methanol, medicinal plants, microbial isolate, microorganisms, phytochemical, range, spectrum, tannin, discovery, technology, saponin, etc.

Top cited articles with high Field Citation Ratio (FCR) and Relative Citation Ratio (RCR)

This study also revealed the top cited AMR articles indexed in Dimension database and sorted by FCR (Table 2). The FCR of the top 20 articles ranged from 4.09 to 36.54. The RCR ranged from 1.01 to 9.42. Table S2 shows the Nigeria's top scholarly contributions to AMR indexed in Scopus database.

Discussion

AMR is an unfortunate result of antibiotic misuse and this study confirms that Institutions in Nigerian have contributed at least 0.2% of the global research output on AMR. A marked rise in scientific output started in 2016 and peaked in 2021. Although data for 2022 is incomplete, evidence of a continuing rise in research output is apparent. In general, the number of AMR documents in the database has gradually increased, implying a significant scientific research output. A similar upward trend has also been reported in earlier bibliometric studies[24–26]. This parallels the increased burden of AMR in Nigeria. Further, the marked rise in scientific output coincides with Nigeria's response to a national rise in the burden of antimicrobial resistance that led to the development of the NAP (2017–2022) that sought to address crucial risk factors in harmony with the WHO Global Action Plan on AMR. Connecting the gap between research institutions, pharmaceutical industries and legislators is pertinent because the role of research institutions does not end with the production of scientific data. Research should translate to development of actionable and sustainable policies and practices. These findings also highlight the importance of strengthening research capacity across Nigerian institutions.

The majority of the documents (57.18%) were in the field of Medical and Health Sciences.

This result is comparable to (25,26), where the majority of research output was in a related field. There is a need for more research output in all subject areas, as well as more multidisciplinary research, to advance the science of AMR.

The centrality of an institution/organisation reflects its recognition in the field of AMR research[27]. By analysing both the centrality and number of articles, Nigeria's AMR research strength was observed to be primarily focused at universities, research institutions and NGOs. The University of Ibadan was Nigeria's leading research institution in terms of scientific outputs and centrality. International Institute of Tropical Agriculture was the leading nongovernmental organization (NGO) in terms of centrality and Nigerian Institute of Medical Research lead in scholarly output among research institutes.

We found that OKEKE IN ranked first in publication frequency and second to BAUER AW (0.13) in terms of centrality (0.07), indicating that the author was part of an influential core author group in the field of AMR research in Nigeria. Majority of Nigerian authors had a centrality index of less than one. This result indicated that more researchers should participate actively in studies on AMR burden and contribute important scientific input to this field.

African Journal of Clinical and Experimental Microbiology was ranked as the top journal, publishing AMR documents with a minimum of one Nigerian institutional affiliation. The same journal was also the most cited in terms of published documents. Although Nigerian journals were among the top publishers of AMR research featuring at least a Nigerian author, there is a need to invest in

strengthening local journals in Nigeria in order to achieve comparable indexing status and scientific standards to those of the world's top journals.

The frontier in a research area depicts the current growth path of a discipline in bibliometrics, and thus the references in the frontier article represent the field's intellectual core[28]. We described Nigeria's scholarly core in the field of AMR by mapping references. Themes on antimicrobial stewardship, Clinical and laboratory practice, public health implication, traditional/molecular methods, and phytomedicine and drug discovery are major research trends of Nigeria's scientific community. These thematic distributions align with the observations of [29]. It is pertinent to boost research in all thematic areas of AMR research in the country to significantly facilitate processes that will slow down the spread of AMR in the country. Research themes exploring the use of molecular and AI systems for AMR detection, surveillance, and drug discovery must be enhanced to confront rising bacterial resistance by significantly reducing the time taken to find new antimicrobials, improving diagnostic and treatment precision, as well as lowering costs[30]. A hotspot is a scientific topic of discussion in a series of documents which are inextricably connected to a particular time period[31]. The keywords are generalizations of the literature's topic[31]. A high-frequency keyword analysis can be used to identify hotspots in the field of AMR research. From our results "*Escherichia coli*" and "*Staphylococcus aureus*", "ESBL", "*Salmonella*", "tetracycline" and "Plasmid" were the most frequently used keywords. Nigerian researchers could conduct advanced studies based on the frequently discussed topics in this field.

An FCR range of 4.09 to 36.54 reflect the moderate to high scientific influence of Nigerian researchers in AMR research field. The theme of the top-cited documents in both the dimension and Scopus databases discussed are similar. The articles were

focused on key AMR issues and determinants of human and animal importance relevant to the Nigerian and African population. Including; infectious diseases, Clinical and community health. Specifically, they highlighted the Prevalence and/or Emergence of colistin resistance *Klebsiella pneumoniae*, Community-Associated Methicillin-Resistant *Staphylococcus aureus*, carbapenem resistance, Enterobacteriaceae, Multidrug Resistant Non-O1/Non-O139 *Vibrio cholerae*, trimethoprim resistance gene *dfpG* in *Staphylococcus aureus*, Plasmid-mediated quinolone resistance and betalactamases in *Escherichia coli*, CTX-M-15 extended-spectrum Beta-lactamase from *Klebsiella pneumoniae* etc. The identification of top-cited articles is useful as it reflects the diverse scope and health research fields that are of high interest to the readership of AMR, in addition to helping researchers identify the authors and institutions that have highly contributed to such influential studies for future research collaborations.

We are unlikely to witness a return to the heyday of antibiotic discovery, when numerous chemotypes were being regularly discovered and brought into clinical use in a short time frame. What does the antibiotic future hold? An assortment of approaches to infection treatment and prevention will have to be offered in the future. Anti-virulence strategies, Small-molecule antibiotics, peptide-based antibiotics, Bacteriophage Monoclonal antibodies, Probiotic strategies and Vaccines are future trends in AMR and drug discovery.

Increasingly, the traditional Waksman platform returns recognized molecules. with recent breakthroughs in analytical chemistry, synthetic biology, genome mining, and genomics, the spectacular chemical diversity of nature can be taken advantage of by moving beyond the traditional Waksman platform. For instance, many environmental bacteria like actinomycetes harbour genetic programs to produce 20 to 40 natural

products, while fungi can contain even more[32]. Avoiding actinomycetes bacteria is one strategy, for example. Some studies have identified 2 new antibiotics with new mechanisms of action, teixobactin, from the betaproteobacterium *Eleftheria terrae* isolated from a soil specimen[33], and darobactin, from the gamma proteobacterium *Photorhabdus khaini* isolated from the gut of insect-killing nematodes[34].

The Myers laboratory has developed innovative modular pathways for the synthesis of new therapeutic tetracycline, macrolide, and lincosamide antibiotics, significantly expanding the diversity of these compound classes well beyond limits of natural biosynthesis[32]. Artificial intelligence approaches could also mine chemical libraries for therapeutic potential in order to develop new synthetic scaffolds[30].

Despite the difficulties in bringing peptide-based antibiotics to the clinic (most suffer from haemolytic activity or nephrotoxicity), optimism remains, and many AMPs are in various stages of development, targeted toward a range of pathogens[32].

Though naturally narrow in spectrum and must be combined with antibiotics, Antivirulence strategies aimed at abrogating pathogen virulence are a future trend. Sibofimloc, which prevents pathogenic *E. coli* from adhering to host epithelia is entering phase 2 trials[35].

Adjuvants or antibiotic combinations, such as b-lactamase inhibitors, can aid in the fight against resistance. A combo of bedaquiline, pretomanid, and linezolid, for instance, was recently approved to treat of Extensively drug - resistant M. tuberculosis, and more combinations are being studied[36].

In current history, the effectiveness and narrow spectrum of phage have been acknowledged as a valuable resource in the treatment of highly drug-resistant infections. The concept of using bacterial phages to treat drug-resistant infections is

becoming more appealing. Phage therapy is increasingly being viewed as an alternative to conventional antibiotics in medicine, spurring more clinical trials, the majority of which are in phases one and two [32].

There are currently only 3 approved antimicrobial Monoclonal antibodies (mAbs), two of which inhibit the virulence of *Bacillus anthracis* (raxibacumab and obiltoxaximab), and one of which inhibits the virulence of *Clostridium difficile* (bezlotoxumab), reducing the incidence of reoccurrence. Only bezlotoxumab withstood clinical trials in its entirety[37], whereas the raxibacumab and obiltoxaximab were approved under the animal rule for bioterror organisms. But besides their proven effectiveness, few are now in late-stage clinical trials (phase 2, *P. aeruginosa*: panobacumab/Aerumab; phase 2, *S. aureus*: omodenbamab/514G3 and; phase 3, *S. aureus*: suvrattoxumab and tosattoxumab[32].

Limitations

The data of this study were mainly derived from the dimensions database on August 20th 2022, and other literature databases like Web of Science, Open Alex and Semantic scholar were not searched. Therefore, we might not have been able to retrieve all papers on this topic, and selection bias may have been present.

Conclusion and Recommendations

According to this research, Nigerian institutions and organizations contribute moderately to AMR research output. However, there is a pressing need to strengthen the country's research capacity and expedite the use of scientific insights provided by its scientists across all subject areas in order to make a significant contribution to development.

Nigeria is home to more than 206,139,589 people (an annual change of 2.58%)[38,39], and effective research is crucial to the advancement of the country. According to our research, Nigeria's contribution to the world's AMR scientific output was only 0.2%. This is scanty, and it emphasizes that AMR response is not always tied to local data, instead it relies on adapting global policies and those of global health organizations. Developing local data to better understand AMR trends is still an excellent method for Nigeria to respond to health emergencies. Nigeria's potential to respond to health issues needs to be strengthened, and investing in research is one excellent way to do so.

Federal and state academic universities, which form the bulk of academic research institutions in Nigeria, have been on strike for over 65 months since the country's independence in 1999. This illustrates that Nigeria's academic institutions are underrated as significant contributors to the nation's information economy and growth plans. A university ought to be a valuable source of knowledge that helps with innovation and national growth[39]. The combined effect of resistance pressures and a lack of creativity and determination to develop next-generation therapies threatens modern medicine's very existence. Artificial intelligence (AI) frameworks are not being utilized in the country. Whole-genome sequencing for antimicrobial susceptibility testing (WGS-AST) is used to diagnose AMR with high accuracy and consistency by extracting large and high-dimensional datasets. AI systems are used to improve this process. Fundamentally, the antimicrobial resistance (AMR) problem is one of kinetics: While the development of antimicrobials is slow, resistance develops quickly. AI technologies have the potential to ease synthesis of antimicrobial peptides with machine/deep learning, New and cost-effective antibiotics, appropriate antibiotic prescription, prediction of antibiotic resistance,

prediction of the severity of infection (efficiency of distinguishing infectious and non-infectious diseases) and by extension, Mortality reduction. Nigeria should adopt current AI practices regarding antibiotic resistance across the world, with Nigeria as the central focus.

The burden of AMR is Monumental and existential and the Right priority ought to be given towards achieving Sustainable standards in the fight against it. Evidence-driven economy is imperative and national stakeholders must be devoted to backing education by adapting the United Nations Educational, Scientific and Cultural Organizations (UNESCO) recommendations[40]. Investment in improving the student's professor ratio in Nigerian Universities will aid in improving the quality of scientific output. New antibiotics are always welcomed, the big question is, where to find these candidates? Nigerian universities must be revitalized so that they can meet the country's development needs and herald a generation of patriots with impactful knowledge.

Funding: No funding was received

Conflicts of interest: According to the authors, there are no conflicts of interest.

472 References

- 473 1. Prestinaci F, Pezzotti P, Pantosti A. Antimicrobial resistance: a global multifaceted
474 phenomenon. *Pathog Glob Health* [Internet]. 2015 [cited 2022 Sep 17];109:309–18.
475 Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4768623/>
- 476 2. Marie Nyhagen Vikan I. Survey on the activities of the World Health Organization
477 (WHO) against antimicrobial resistance (AMR). 2016 Mar 22; Available from:
478 <http://dx.doi.org/10.26226/morressier.56d6be7cd462b80296c961cc>
- 479 3. Chukwu EE, Oladele DA, Awoderu OB, Afocha EE, Lawal RG, Abdus-salam I, et
480 al. A national survey of public awareness of antimicrobial resistance in Nigeria.
481 *Antimicrob Resist Infect Control* [Internet]. 2020 [cited 2022 Sep 16]; 9:72. Available
482 from: <https://aricjournal.biomedcentral.com/articles/10.1186/s13756-020-00739-0>
- 483 4. Achi CR, Ayobami O, Mark G, Egwuenu A, Ogbolu D, Kabir J. Operationalising
484 One Health in Nigeria: Reflections From a High-Level Expert Panel Discussion
485 Commemorating the 2020 World Antibiotics Awareness Week. *Frontiers in Public*
486 *Health* [Internet]. 2021 May 31;9. Available from:
487 <http://dx.doi.org/10.3389/fpubh.2021.673504>
- 488 5. Harant A. Assessing transparency and accountability of national action plans on
489 antimicrobial resistance in 15 African countries. *Antimicrobial Resistance & Infection*
490 *Control* [Internet]. 2022 [cited 2022 Sep 17]; 11:15. Available from:
491 <https://doi.org/10.1186/s13756-021-01040-4>
- 492 6. Knobler SL, Infections I of M (US) F on E, Lemon SM, Najafi M, Burroughs T.
493 Factors Contributing to the Emergence of Resistance [Internet]. *The Resistance*
494 *Phenomenon in Microbes and Infectious Disease Vectors*. 2003 Mar 26; Available
495 from: <http://dx.doi.org/10.17226/10651>
- 496 7. Experts caution Nigerians against indiscriminate use of antibiotics to reduce future
497 health crisis [Internet]. WHO | Regional Office for Africa. [cited 2022 Sep 17].
498 Available from: [https://www.afro.who.int/news/experts-caution-nigerians-against-](https://www.afro.who.int/news/experts-caution-nigerians-against-indiscriminate-use-antibiotics-reduce-future-health-crisis)
499 [indiscriminate-use-antibiotics-reduce-future-health-crisis](https://www.afro.who.int/news/experts-caution-nigerians-against-indiscriminate-use-antibiotics-reduce-future-health-crisis)
- 500 8. Auta A, Banwat SB, David S, Dangiwa DA, Ogbole E, Tor-anyiin AJ. Antibiotic
501 Use in Some Nigerian Communities: Knowledge and Attitudes of Consumers.
502 *Tropical Journal of Pharmaceutical Research* [Internet]. 2013 [cited 2022 Sep
503 17];12:1087–92. Available from:
504 <https://www.ajol.info/index.php/tjpr/article/view/99922>
- 505 9. Stivers T. Managing Patient Pressure to Prescribe Antibiotics in the Clinic.
506 *Paediatr Drugs* [Internet]. 2021 [cited 2022 Sep 17];23:437–43. Available from:
507 <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC8375467/>
- 508 10. Adesokan HK, Akanbi IO, Akanbi IM, Obaweda RA. Pattern of antimicrobial
509 usage in livestock animals in south-western Nigeria: The need for alternative plans.
510 *Onderstepoort J Vet Res* [Internet]. 2015 [cited 2022 Sep 17];82:816. Available from:
511 <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6238793/>

11. Kotwani A, Joshi J, Kaloni D. Pharmaceutical effluent: a critical link in the interconnected ecosystem promoting antimicrobial resistance. *Environ Sci Pollut Res* [Internet]. 2021 [cited 2022 Sep 17];28:32111–24. Available from: <https://doi.org/10.1007/s11356-021-14178-w>
12. Sinha R, Shukla P. Antimicrobial Peptides: Recent Insights on Biotechnological Interventions and Future Perspectives. *Protein and Peptide Letters*. 2019;26:79–87.
13. Liang C, Luo A, Zhong Z. Knowledge mapping of medication literacy study: A visualized analysis using CiteSpace. *SAGE Open Medicine* [Internet]. 2018 [cited 2022 Sep 16];6:205031211880019. Available from: <http://journals.sagepub.com/doi/10.1177/2050312118800199>
14. Ding X, Yang Z. Knowledge mapping of platform research: a visual analysis using VOSviewer and CiteSpace. *Electron Commer Res* [Internet]. 2022 [cited 2022 Sep 17];22:787–809. Available from: <https://doi.org/10.1007/s10660-020-09410-7>
15. Adeiza S, Shuaibu A. Trends in Monkeypox Research: A Sixty Year Bibliometric Analysis. *Microbes and Infectious Diseases* [Internet]. 2022 Jul 13;0(0):0–0. Available from: <http://dx.doi.org/10.21608/mid.2022.147680.1334>
16. Li Y, He M, Zou Z, Bian X, Huang X, Yang C, et al. Artificial liver research output and citations from 2004 to 2017: a bibliometric analysis. *PeerJ* [Internet]. 2019 [cited 2022 May 14];6:e6178. Available from: <https://peerj.com/articles/6178>
17. Hook DW, Porter SJ, Herzog C. Dimensions: Building Context for Search and Evaluation. *Front Res Metr Anal* [Internet]. 2018 [cited 2022 May 12];3:23. Available from: <https://www.frontiersin.org/article/10.3389/frma.2018.00023/full>
18. Van Eck N, Waltman L. Software survey: VOSviewer, a computer program for bibliometric mapping. *scientometrics*. Akadémiai Kiadó, co-published with Springer Science+ Business Media BV ...; 2010;84:523–38.
19. Perianes-Rodriguez A, Waltman L, van Eck NJ. Constructing bibliometric networks: A comparison between full and fractional counting. *Journal of Informetrics* [Internet]. 2016 [cited 2022 May 12];10:1178–95. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S1751157716302036>
20. Huang T, Wu H, Yang S, Su B, Tang K, Quan Z, et al. Global Trends of Researches on Sacral Fracture Surgery: A Bibliometric Study Based on VOSviewer. *Spine* [Internet]. 2020 [cited 2022 May 12];45:E721. Available from: https://journals.lww.com/spinejournal/Abstract/2020/06150/Global_Trends_of_Researches_on_Sacral_Fracture.18.aspx?context=FeaturedArticles&collectionId=2
21. Suleiman AS, Abdulmalik SB, Ghazali SM. Extended Spectrum Betalactamase Research mapping; a 32-year Temporal and geographical perspective. 2022 Aug 2; Available from: <http://dx.doi.org/10.21203/rs.3.rs-1898374/v1>
22. Chen C. CiteSpace II: Detecting and visualizing emerging trends and transient patterns in scientific literature. *J Am Soc Inf Sci* [Internet]. 2006 [cited 2022 Sep 16];57:359–77. Available from: <https://onlinelibrary.wiley.com/doi/10.1002/asi.20317>

23. What is the FCR? How is it calculated? [Internet]. Dimensions. [cited 2022 Sep 17]. Available from: <https://dimensions.freshdesk.com/support/solutions/articles/23000018848-what-is-the-fcr-how-is-it-calculated->
24. Sweileh WM, Moh'd Mansour A. Bibliometric analysis of global research output on antimicrobial resistance in the environment (2000–2019). *Global Health Research and Policy* [Internet]. 2020 [cited 2022 Sep 17];5:37. Available from: <https://doi.org/10.1186/s41256-020-00165-0>
25. Torres RT, Carvalho J, Cunha MV, Serrano E, Palmeira JD, Fonseca C. Temporal and geographical research trends of antimicrobial resistance in wildlife-a bibliometric analysis. *One Health*. Elsevier; 2020;11:100198.
26. Payumo J, Alocilja E, Boodoo C, Luchini-Colbry K, Ruegg P, McLamore E, et al. Next Generation of AMR Network. *Encyclopedia* [Internet]. Multidisciplinary Digital Publishing Institute; 2021 [cited 2022 Sep 17];1:871–92. Available from: <https://www.mdpi.com/2673-8392/1/3/67>
27. Leydesdorff L. Betweenness centrality as an indicator of the interdisciplinarity of scientific journals. *Journal of the American Society for Information Science and Technology* [Internet]. 2007 [cited 2022 Sep 17];58:1303–19. Available from: <https://onlinelibrary.wiley.com/doi/abs/10.1002/asi.20614>
28. Qi S, Hua F, Xu S, Zhou Z, Liu F. Trends of global health literacy research (1995–2020): Analysis of mapping knowledge domains based on citation data mining. Lahiri A, editor. *PLOS ONE* [Internet]. 2021 Aug 9;16(8):e0254988. Available from: <http://dx.doi.org/10.1371/journal.pone.0254988>.
29. Luz CF, van Niekerk JM, Keizer J, Beerlage-de Jong N, Braakman-Jansen LMA, Stein A, et al. Mapping twenty years of antimicrobial resistance research trends. *Artificial Intelligence in Medicine* [Internet]. 2022 [cited 2022 Sep 16];123:102216. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S0933365721002098>
30. Rabaan AA, Alhumaid S, Mutair AA, Garout M, Abulhamayel Y, Halwani MA, et al. Application of Artificial Intelligence in Combating High Antimicrobial Resistance Rates. *Antibiotics* [Internet]. 2022 [cited 2022 Sep 16];11:784. Available from: <https://www.mdpi.com/2079-6382/11/6/784>
31. Wang Z, Ma D, Pang R, Xie F, Zhang J, Sun D. Research Progress and Development Trend of Social Media Big Data (SMBD): Knowledge Mapping Analysis Based on CiteSpace. *IJGI* [Internet]. 2020 [cited 2022 Sep 16];9:632. Available from: <https://www.mdpi.com/2220-9964/9/11/632>
32. Cook MA, Wright GD. The past, present, and future of antibiotics. *Sci Transl Med* [Internet]. 2022 [cited 2022 Sep 16];14:eabo7793. Available from: <https://www.science.org/doi/10.1126/scitranslmed.abo7793>
33. Ling LL, Schneider T, Peoples AJ, Spoering AL, Engels I, Conlon BP, et al. A new antibiotic kills pathogens without detectable resistance. *Nature*. Nature Publishing Group; 2015;517:455–9.

34. Imai Y, Meyer KJ, Iinishi A, Favre-Godal Q, Green R, Manuse S, et al. A new antibiotic selectively kills Gram-negative pathogens. *Nature* [Internet]. 2019 Nov 20;576(7787):459–64. Available from: <http://dx.doi.org/10.1038/s41586-019-1791-1>
35. Reinisch W, Hébuterne X, Buisson A, Schreiber S, Desreumaux P, Primas C, et al. Safety, pharmacokinetic, and pharmacodynamic study of sibofimloc, a novel FimH blocker in patients with active Crohn's disease. *Journal of Gastroenterology and Hepatology*. Wiley Online Library; 2022; <https://doi.org/10.1111/jgh.15828>
36. Wright GD. Antibiotic Adjuvants: Rescuing Antibiotics from Resistance. *Trends in Microbiology* [Internet]. 2016 Nov;24(11):862–71. Available from: <http://dx.doi.org/10.1016/j.tim.2016.06.009>.
37. Allio T. The FDA Animal Rule and its role in protecting human safety. *Expert Opinion on Drug Safety* [Internet]. 2018 Sep 12;17(10):971–3. Available from: <http://dx.doi.org/10.1080/14740338.2018.1518429>.
38. Worldometer - real time world statistics [Internet]. Worldometer. [cited 2022 Sep 16]. Available from: <http://www.worldometers.info/>
39. Adebisi YA, Pius M. Nigeria's scientific contributions to COVID-19: A bibliometric analysis. *Annals of Medicine and Surgery* [Internet]. 2022 [cited 2022 Sep 16];80:104316. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S2049080122010767>, <https://doi.org/10.1016/j.amsu.2022.104316>
40. A thematic analysis regarding education goals in Nigeria and select UNESCO *Education for All* priorities - ProQuest [Internet]. [cited 2022 Sep 18]. Available from: <https://www.proquest.com/openview/5539ae2be9fff78c286b9ea39ada831f/1?pq-origsite=gscholar&cbl=18750>

It is made available under a CC-BY-NC-ND 4.0 International license.




Figure 2: Nigeria's annual contribution to AMR publication growth.




Figure 3: The top Authors and Journals of AMR research in Nigeria



Figure 3: Centrality of top Universities (Right) and Authors (Left). Notes: each node in the network represents cited Author /University, the number of papers published by the author is represented by the size of the nodes, and the stroked rings represent the centrality index.

It is made available under a [CC-BY-NC-ND 4.0 International license](#).

648
649
650
651
652
653
654
655
656




Figure 4: The co-authorship map of 1000 authors of authors in AMR research domain ($T \geq 5$). Notes: The size of node reflects the author's co-occurrence frequencies, the link specify the co-occurrence association between authors, and the same colour of node represent the same cluster. In the figure there are 74 clusters, 7063 links and a total link strength of 10240.




Figure 5: Institutional collaboration network: A minimum threshold of 20 ($T = 20$) documents per institution was set for inclusion. Notes: The size of the connecting lines, circles and colour vary based on total link strength and number of publications and cluster respectively. Normalization method used was Lin/log modularity and full counting method.

It is made available under a [CC-BY-NC-ND 4.0 International license](#).




Figure 6: keyword co-occurrence network and clusters of AMR research (T \geq 40, contains 1481 items, 14 clusters and 449516 links; max lines = 200, total link strength = 5333437). Notes: The node and word size reflects the co-occurrence frequencies, the link indicate the co-occurrence relationship, and the same colour of node signify the same cluster. Network map of the trend topics according to the keywords used. Indicator shows the current publications from purple to yellow. More studies focused on vaccine, drug treatment and spike protein have been published recently. The size of the circles represents the frequency of appearance as the keywords. The distance between the two circles indicates their correlation.

Table 1: Top25 Authors and Cited Authors in AMR research with the Strongest Citation Bursts (Sorted by Beginning year)

Authors	Strength	Begin	End	1972 - 2022
OLUKOYA D	5.65	1988	1996	
COKER A	4.54	1992	2007	
BAKARE R	6.31	1995	2002	
OGUNSOLA F	4.45	2002	2016	
ADELOWO O	4.81	2008	2020	
LOWE O	5.91	2012	2018	
ABIMIKU A	5.47	2012	2019	
UMOH VJ	4.88	2013	2015	
KWAGA J	4.63	2013	2022	
AKINDUTI P	4.21	2014	2020	
ANYANWU M	5.69	2015	2019	
OJO O	4.78	2015	2022	
ODUYEBO O	6.62	2016	2022	
ILIYASU G	4.23	2016	2022	
MOHAMMED Y	4.23	2016	2022	
EHINMIDU J	4.3	2017	2020	
IDOWU E	4.18	2017	2022	
CHAH K	4.63	2018	2022	
MUKHTAR M	4.02	2018	2022	
NGWAI Y	8.7	2019	2020	
OKEKE I	7.27	2019	2022	
IROHA I	4.8	2019	2022	
OLAYINKA B	8.53	2020	2022	
ODETOYIN B	4.26	2020	2022	
ADEBOWALE O	3.96	2020	2022	
Top Cited Authors				
Cited Authors	Strength	Begin	End	1972 - 2022
AKO-NAI AK	6.5	1999	2017	
MENKIR A	6.92	2007	2012	
OKEKE IN	11.6	2009	2013	
TENOVER FC	5.86	2009	2017	
GHEBREMEDHIN B	7.0	2011	2014	
PATERSON DL	7.63	2012	2017	
LAWSON L	6.82	2013	2019	
SHITTU AO	5.86	2013	2018	
BAUER AW	5.29	2013	2014	
LIVERMORE DM	6.43	2014	2016	
AIBINU I	6.82	2015	2020	
ALTSCHUL SF	5.65	2015	2018	
MAGIORAKOS A □	9.44	2016	2022	
OJO OE	7.07	2017	2020	
ADESOKAN HK	5.75	2018	2022	
ADELOWO OO	5.63	2018	2022	
OLOSNO	7.93	2019	2022	
IBRAHIM SS	6.56	2019	2022	
ALHAJI NB	6.01	2019	2022	
KARIUKI S	5.98	2019	2020	
KRUMPERMAN PH	5.97	2019	2022	
IGBINOSA EO	5.86	2019	2022	
ABUBAKAR U	7.96	2020	2022	
AWORH MK	6.91	2020	2022	
ASHLEY EA	5.84	2020	2022	

Notes: The Blue bars mean the reference had been published; the red bars = Citation burstness (The start and end of the burst)

period is consistent with the length of red segment).

Table 2: Top 20 scholarly contributions to AMR authored by a Nigerian (Sorted by FCR)

Authors	DOI	Year	Times cited	RCR	FCR
Stewardson et al	10.1016/s1473-3099(18)30792-8	2019	92	8.45	36.54
Olaitan et al	10.1016/j.ijantimicag.2014.07.020	2014	203	9.42	24.66
Nurjadi et al	10.1093/jac/dku174	2014	75	3.13	19.42
Shittu et al	10.1186/1471-2180-11-92	2011	123	4.36	17.65
Ghebremedhin et al	10.1128/jcm.00648-09	2009	119	4.11	17.11
Kesah et al	10.1046/j.1469-0691.2003.00531.x	2003	96	2.22	16.92
Adesokan et al	10.4102/ojvr.v82i1.816	2015	61	3.37	14.57
Fashae et al	10.3855/jidc.909	2010	79	2.56	11.58
Aibinu et al	10.1111/j.1469-0691.2011.03730.x	2011	58	2.49	11.44
Marin et al	10.1371/journal.pntd.0002049	2013	82	3.23	9.96
Titilawo et al	10.1016/j.scitotenv.2015.03.095	2015	61	2.67	8.77
Ogbolu et al	10.1089/jmf.2006.1209	2007	77	1.01	8.09
Ogbolu et al	10.1016/j.ijantimicag.2010.08.019	2010	70	2.76	6.92
Titilawo et al	10.1007/s11356-014-3887-3	2015	63	2.73	6.21
Akinyemi et al	10.1016/j.puhe.2004.04.009	2005	53	1.15	5.98
Okeke et al	10.3201/eid0604.009913	2000	148	1.47	5.85
Fortini et al	10.1093/jac/dkr085	2011	71	2.76	5.69
Chigor et al	10.3390/ijerph7103831	2010	53	0.97	5.32
Iwalokun et al	10.1089/jmf.2004.7.327	2004	83	1.41	5.2
Soge et al	10.1093/jac/dki429	2005	64	1.83	4.09

675

676

677

