

1 **Influence of high-energy laser therapy to the patellar** 2 **tendon on its ligamentous microcirculation:** 3 **a quasi-experiment**

4 **Andreas Brandl** ^{1,2}, **Christoph Egner**², **Ursel Reisser**³, **Christian Lingenfelder**³, **Robert Schleip** ^{2,4,*}

5 ¹ Department of Movement and Training Science, Faculty of Psychology and Human Movement Science, University of
6 Hamburg, Hamburg, Germany

7 ² DIPLOMA Hochschule, Bad Sooden-Allendorf, Germany

8 ³ Zimmer Medizinsysteme, 89231 Neu-Ulm, Germany

9 ⁴ Conservative and Rehabilitative Orthopedics, Department of Sport and Health Sciences, Technical University of Munich,
10 Munich, Germany

11 * Corresponding author

12 Email: robert.schleip@tum.de

13 **Abstract**

14 Laser therapeutic applications, such as the use of high energy lasers (HILT), are widely used in physical therapy, but
15 basic studies on the mechanisms of action of HILT on tendinous/ligamentous tissue are largely lacking. The aim of this study
16 was to investigate microcirculatory changes of the patellar tendon by HILT. 21 healthy volunteers were treated with HILT.
17 Before and after the intervention, as well as 10 minutes later, the microcirculation was measured by noninvasive laser
18 spectroscopy (O2C device). Tissue temperature was recorded at the measurement time points using thermography. Blood flow
19 increased significantly by 86.38 AU ($p < 0.001$) after the intervention and by 25.76 AU ($p < 0.001$) at follow-up. Oxygen
20 saturation increased by 20.14% ($p < 0.001$) and 13.48%, respectively ($p < 0.001$), whereas relative hemoglobin decreased by
21 6.67 AU and 7.90 AU, respectively. Tendon temperature increased by 9.45° and 1.94° Celsius, respectively. Acceleration of
22 blood flow by improving the flow properties of erythrocytes and platelets may have caused the results.

23 **Introduction**

24 Laser therapy applications are widely used in physical therapy. However, there is a lack of empirical research on
25 fascial structures, their variability and adaptability [1]. Non-invasive therapy devices are becoming increasingly important in
26 the context of musculoskeletal rehabilitation. High-intensity laser therapy (HILT) has been little studied in previous research
27 as an intervention method for knee osteoarthritis [2]. Basic studies on the mechanisms of action of HILT on dense parallel-
28 fibred connective tissues (such as tendons or ligaments) are scarce.

29 Vascular factors have been identified in previous studies as critical for the development of tendinopathies. Reduced
30 blood flow, especially after activity, represents the greatest risk factor. Studies in the Achilles tendon have shown that structural
31 parameters - such as Achilles tendon thickness, tendon structure as analyzed by ultrasound tendon analysis, or foot posture
32 index – do not significantly contribute to the prediction of Achilles tendinopathy [3], whereas microcirculation in the capillary
33 venous area serves as a significant risk predictor.

34 Furthermore, microcirculation in tendon tissue plays a prominent role in regeneration after injury. In this context,
35 oxygen, blood viscosity, endothelial functionality and blood vessel permeability are important parameters that emphatically
36 influence wound healing [4].

37 Hypoxia in tendon tissue promotes the production of proinflammatory cytokines, which can be observed in the early
38 stage of the development of tendinopathy and is therefore discussed as a trigger for such a condition. These hypoxia-related
39 proteins are key mediators of cellular inflammation and apoptosis, processes that cause a shift in collagen matrix synthesis
40 toward an increased proportion of type III collagen (which tends to form fibrils with lower mechanical strength). A shift in
41 matrix structure from type I collagen, which makes up 70% of the dry mass in healthy tendon tissue, to type III collagen reduces
42 the resistance of the tendon to tensile forces, thereby favoring ruptures [5]. Hypoxia further causes local acidosis, corresponding
43 to increased intratendinous lactate accumulation, which further negatively influences pathogenesis [6].

44 Recent studies indicate that tendinopathies are associated with new vascularization with swelling and pain [7].
45 However, what seems like a paradox could, according to Järvinen [8], also represent persistent preceding hypoxia. Since tissue
46 regeneration requires an adequate supply of oxygen and nutrients, therefore, neovascularization in tendinopathy can be
47 interpreted as a sign of both persistent hypoxia and a failed attempt to repair the tendon. It is known from the fields of cancer
48 research and retinopathy that hypoxia-induced neovessels are hyper-permeable, resulting in vascular leakage and reduced blood
49 flow. This provides insufficient oxygen and nutrients needed for tissue maintenance and possible regeneration [8].

50 HILT decreases erythrocyte deformability and platelet aggregation, resulting in membrane revitalization, viscosity
51 reduction, and erythrocyte stress adaptation. In addition, the lifespan of erythrocytes and platelets is prolonged and their flow
52 properties are improved under HILT [9–11]. Therefore, HILT is also discussed as a promising method for the treatment of
53 various blood diseases [9–13].

54 The aim of this study was to investigate microcirculatory changes of the patellar tendon by HILT.

55 **Methods**

56 **Study design overview**

57 The study was a quasi-experiment with one intervention group. Measurements were taken before and after the
58 intervention and at a 10-minute follow-up according to the SPIRIT guidelines [14]. The study protocol was prospectively
59 registered with the German Clinical Trials Register (DRKS00028155) on 18.02.2022. The study has been reviewed and
60 approved by the ethical committee of the DIPLOMA Hochschule, Germany (Nr. 1021/2021), has been carried out in accordance
61 with the declaration of Helsinki and has obtained written informed consent from the participants [15].

62 **Setting and participants**

63 The study was conducted in a school of physiotherapists, in a medium-sized city in middle Germany. The number of
64 participants was calculated based on the results of a previous study and set at 18 [16,17]. The acquisition was carried out via
65 direct contact, a notice board, and the distribution of information material in the school.

66 Inclusion criteria were: (a) A generally healthy constitution; (b) the persons to be treated must have intact thermal
67 sensitivity and be able to perceive and communicate pain; (c) a BMI between 18 and 29.9; (d) female or male subjects aged 18
68 to 60 years; (e) sitting position for 15 minutes must be pain-free for the subjects.

69 Exclusion criteria were: (a) generally valid contraindications to physiotherapeutic and osteopathic treatments of the
70 patellar tendon (i.e., fractures, tumors, infections, severe cardiovascular and metabolic diseases); (b) pregnancy; (c) rheumatic
71 diseases; (d) taking medication that affects blood circulation, pain or mind; (e) taking muscle relaxants; (f) skin changes (e.g.
72 neurodermatitis, psoriasis, urticaria, decubitus ulcers, hematoma); (g) surgery or other scars in the patellar tendon region; (h)
73 previous mental illness; (i) surgery in the last three months; (j) prosthetics or internal knee arthroplasty; (k) acute inflammation.

74 **Study flow**

75 First, the anthropometric data, age, gender, height, and weight were collected by the investigators (AB, RS). See the
76 flowchart in Fig. 1 for the course of the study.

77 **Fig. 1. Flow diagram of the study. The "Loss to follow-up" has been removed, as it is not relevant for the study**
78 **design.**

79 Before the measurements, the subjects were given information regarding the implementation of the intervention and
80 measurements. They lay on their backs with their heads bent at 160° (cervical spine flexion) on a treatment couch. The knees
81 were placed underneath and a knee flexion of 145° - 150° was preset. The measurements were taken in daylight between 9:00
82 a.m. and 03:00 p.m. and at a room temperature of 22 degrees Celsius. The initial measurement (baseline) was performed first.
83 Then the HILT intervention (OptonPro, Zimmer Medizinsysteme GmbH, Neu-Ulm, Germany, Fig. 2B) was carried out (Fig.
84 2C). A total amount of energy of 800 J (100 J/cm²) was applied to a predefined area of 2 cm x 4 cm on the patellar tendon (Fig.
85 2A).

86 **Fig 2. A show the prefabricated template defining the treatment area of 2 cm x 4 cm. B shows the OptonPro high**
87 **intensity laser therapy device. C shows the laser application to the patellar tendon.**

88 **Outcomes**

89 Measurement of the microcirculation of the patellar tendon was performed using a laser Doppler flowmeter and tissue
90 spectrometer (O2C, LEA Medizintechnik GmbH, Heuchelheim, Germany) (Fig. 3). This makes it possible to simultaneously
91 determine blood flow, oxygen saturation, and relative hemoglobin levels of the measured tissue. The device determines these
92 parameters at the venous end of the capillary and thus provides information about the local microcirculation.

93 **Fig 3. A: Hardware for microcirculation measurement using a laser Doppler flowmeter and a tis-sue spectrometer. B:**
94 **Probe placement.**

95 Temperature measurement using thermography covers temperature ranges from -20° to 400° Celsius (Flir One,
96 Teledyne FLIR LLC, Wilsonville, US). Measurement accuracy is +/- 3° or +/- 5%. This is valid 60 seconds after switching on
97 the instrument in an ambient temperature of 15° to 35° Celsius and the target temperature is in the range of 5° and 120° Celsius.
98 To ensure measurement at a reproducible measuring point, the template (Fig. 2A) was provided with a marker that ensured
99 exact positioning of the image section for temperature determination (Fig. 4).

100 **Fig 4. Thermography of the patellar tendon.**

101 **Statistics**

102 The standard deviation (SD), mean, and 95% confidence interval (95% CI) were determined for all parameters. There
103 were no outliers in the data. The outcome variables were normally distributed as determined by the Shapiro-Wilk test ($p >$
104 0.05). Differences between measurements were tested for significance using a repeated measures ANOVA. For significant
105 main effects, pairwise comparisons were performed with Bonferroni correction. The significance level was set at $p = 0.05$.

106 Libreoffice Calc version 6.4.7.2 (Mozilla Public License v2.0) was used for the descriptive statistics. The inferential
107 statistics were carried out with the software R, version 3.4.1 (R Foundation for Statistical Computing, Vienna, Austria).

108 **Results**

109 The anthropometric data and baseline characteristics are shown in Table 1. Of 28 subjects screened between
110 18/02/2022 and 30/03/2022, 21 met eligibility criteria (Fig. 1). None of the subjects reported any adverse side effects of HILT
111 treatment. The raw data on which the study results are based are in the S1 Data Supporting information.

112 **Table 1. Baseline characteristics.**

Baseline characteristics	Participants (n=21) mean ± SD
Gender (men/woman)	15/6
Age (years)	22.1 ± 4.2
Height(m)	1.75 ± 0.1
Weight (kg)	76.0 ± 11.5
BMI (kg/m ²)	24.9 ± 3.5
Sport activity (hours per week)	6.0 ± 4.2

113 SD standard deviation. n number.

114 A repeated-measures ANOVA revealed that temperature had a statistically significant difference between
115 measurements, $F(2, 40) = 342.46$, $p < 0.001$, partial $\eta^2 = 0.96$. A Bonferroni-adjusted post-hoc analysis revealed significantly
116 ($p < 0.001$) higher temperature after the intervention ($M_{Diff} = 9.45$, 95%-CI[8.73, 10.16]) and 10 minutes later ($M_{Diff} = 1.94$,
117 95%-CI[1.24, 2.65]) compared to pre-intervention.

118 There was also a significant difference for blood flow, $F(1.56, 31.23) = 116.15$, $p < 0.001$, partial $\eta^2 = 0.85$. A
119 Bonferroni-adjusted post-hoc analysis showed significantly ($p < 0.001$) higher flow after intervention ($M_{Diff} = 86.38$, 95%-
120 CI[71.62, 101.14]) and 10 minutes later ($M_{Diff} = 25.76$, 95%-CI[16.65, 34.87]) compared to before the intervention. These
121 values also exceeded the minimum detectable changes of 9.2 arbitrary units (AU) [3].

122 A significant difference was found for oxygen saturation (SO₂), $F(2, 40) = 25.01$, $p < 0.001$, partial $\eta^2 = 0.56$. A
123 Bonferroni-adjusted post-hoc analysis revealed significantly ($p < 0.01$) higher values after the intervention ($M_{Diff} = 20.14$, 95%-
124 CI[13.41, 26.88]) and 10 minutes later ($M_{Diff} = 13.48$, 95%-CI[8.10, 18.85]) compared to before the intervention.

125 A significant difference was found for relative hemoglobin concentration (rHb), $F(2, 40) = 7.24$, $p = 0.002$, partial η^2
126 = 0.27. A Bonferroni-adjusted post-hoc analysis revealed significantly ($p < 0.01$) lower rHb values after the intervention (M_{Diff}
127 = -6.67, 95%-CI[-11.46, -1.88]) and 10 minutes later ($M_{Diff} = -7.90$, 95%-CI[-12.95, -2.86]) compared with the preintervention
128 period.

129 The changes between the baseline measurement, the measurement after treatment, and the measurement 10 minutes
130 later are shown in Table 2 and Fig. 5.

131

132 **Table 2. Descriptive statistics.**

Outcome	t₀ (n=21) mean ± SD	t₁ (n=21) mean ± SD	t₂ (n=21) mean ± SD
Temperature (°C)	28.2 ± 1.1	37.7 ± 1.7***	30.2 ± 1.3***
Flow (AU)	38 ± 19	123 ± 35***	65 ± 24***
SO ₂ (%)	37 ± 16	57 ± 20***	51 ± 17***
rHb (AU)	45 ± 10	38 ± 7*	36 ± 8*

133 SD, standard deviation; n, number; AU, arbitrary units; SO₂, oxygen saturation; rHb, relative hemoglobin
134 concentration. Significant at the level * < 0.05. *** < 0.001.

135 **Fig. 5. Changes between baseline (t₀), after treatment (t₁), and 10 minutes later (t₂). AU, arbitrary units; SO₂, oxygen**
136 **saturation; rHb, relative hemoglobin concentration. Error bars show the standard error of the mean value.**
137 **Significant at the level * < 0.05; *** < 0.001; **** < 0.0001.**

138 Discussion

139 To the authors' knowledge, this is the first study to investigate the influence of HILT on patellar tendon
140 microcirculation. The results here showed significant changes in all measured parameters.

141 The thermal changes, which were recorded by infrared thermography, were as expected. There was an increase of
142 9.45° Celsius immediately after HILT application and a reduction to 1.94° Celsius at follow-up compared to the baseline
143 measurement.

144 Blood flow increased both immediately after intervention and at follow-up (86.38 AU and 25.76 AU, respectively).
145 These values not only showed statistical significance, they also exceeded the minimum detectable changes of 9.2 AU
146 determined by Wezenbeek et al. [3], although the authors determined these values on the Achilles tendon and the
147 transferability to the patellar tendon is debatable. Furthermore, an increase in SO₂ was also observable, which was 20.14%
148 after the intervention and 13.48% at follow-up. This improvement of the capillary blood supply could follow a vasodilatation
149 by thermal influences [18]. However, this should also have caused an increase in rHb. In a study preceding this one, which
150 investigated the influences of purely local thermal changes on the patellar tendon on the microcirculation, an increase in rHb
151 was observed [17]. Nevertheless, the results of this study show a different picture, which is why HILT-specific mechanisms
152 of influence can also be discussed here.

153 In laboratory studies, there is some evidence of changes in erythrocytes and platelets due to laser therapy that
154 significantly affect flow characteristics. Thereby, the blood viscosity decreases and certain stress adaptation of erythrocytes
155 occurs [9–11]. The phenomenon of an apparent decrease in relative hemoglobin observed in the study would also be
156 consistent with this. The rHb value was decreased by 6.67 AU after the intervention and by 7.9 AU at follow-up. To the
157 authors' knowledge, this response has not been observed in other studies and may also indicate improved microcirculatory
158 flow properties that increase blood flow velocity in the venous-capillary vasculature. This would explain that the O2C BF
159 meter detects a faster clearance of erythrocytes, leading to a decrease in rHb [19]. However, it should be critically noted in
160 these considerations that although statistically significant results are available here, the observed value changes are relatively
161 small, and it seems questionable whether the minimum detectable changes were exceeded.

162 Recent studies have also identified red cell changes in long-COVID-19 patients that could affect microcirculation
163 and lead to typical long-COVID-19 symptoms such as fatigue and muscle weakness [20,21]. Therefore, in addition to the
164 scattered work on laser therapy as a method to treat long-COVID-19, work focusing on HILT-specific enhancement of
165 erythrocyte flow properties is of great interest here [12,13].

166 The study had the strengths mentioned above, but some limitations should be taken into account. For example, it
167 was conducted in a quasi-experimental design that was unblinded and not placebo-controlled. The study group was relatively
168 young, physically active, and of normal weight. Therefore, transferability of the study results to other populations is limited.
169 The results should be understood under the premise of a basic scientific study.

170 Significant changes in blood flow could be detected after HILT application, which also exceeded the minimal
171 detectable changes. Here, HILT may have an effect beyond mere vasodilation, improving the flow properties of erythrocytes
172 and platelets. Further research investigating these relationships and a controlled design with a comparison group receiving
173 heat-only application represent promising future study approaches.

174 **Conclusion**

175 This is, to the authors' knowledge, the first study to evaluate the effect of HILT on the microcirculation of the
176 patellar tendon. It found a significant improvement in blood flow and oxygen saturation after HILT intervention, which were
177 still detectable 10 minutes later. On the other hand, the relative amount of hemoglobin decreased significantly, which could
178 indicate an acceleration of blood flow due to improved flow properties of erythrocytes and platelets.

179 Supporting information

180 S1 Data.

181 (XLSX)

182 References


- 183 1. Willard FH, Vleeming A, Schuenke MD, Danneels L, Schleip R. The thoracolumbar fascia: anatomy, function and
184 clinical considerations. *J Anat.* 2012;221: 507–536. doi:10.1111/j.1469-7580.2012.01511.x
- 185 2. Wyszynska J, Bal-Bocheńska M. Efficacy of High-Intensity Laser Therapy in Treating Knee Osteoarthritis: A First
186 Systematic Review. *Photomedicine and Laser Surgery.* 2018;36: 343–353. doi:10.1089/pho.2017.4425
- 187 3. Wezenbeek E, Willems T, Mahieu N, De Muynck M, Vanden Bossche L, Steyaert A, et al. The Role of the Vascular
188 and Structural Response to Activity in the Development of Achilles Tendinopathy: A Prospective Study. *Am J Sports*
189 *Med.* 2018;46: 947–954. doi:10.1177/0363546517750854
- 190 4. Kösters AK, Ganse B, Gueorguiev B, Klos K, Modabber A, Nebelung S, et al. Effects of low-intensity pulsed
191 ultrasound on soft tissue micro-circulation in the foot. *International Orthopaedics (SICOT).* 2017;41: 2067–2074.
192 doi:10.1007/s00264-017-3574-3
- 193 5. Millar NL, Reilly JH, Kerr SC, Campbell AL, Little KJ, Leach WJ, et al. Hypoxia: a critical regulator of early human
194 tendinopathy. *Ann Rheum Dis.* 2012;71: 302–310. doi:10.1136/ard.2011.154229
- 195 6. Sikes KJ, McConnell A, Serkova N, Cole B, Frisbie D. Untargeted metabolomics analysis identifies creatine,
196 myo-inositol, and lipid pathway modulation in a murine model of tendinopathy. *Journal Orthopaedic Research.*
197 2022;40: 965–976. doi:10.1002/jor.25112
- 198 7. Calanni L, Zampella C, Micheletti P, Greco D, Negro M, D’Antona G. Correlation between Patellar Tendon
199 Mechanical Properties and Oxygenation Detection by Near Infrared Spectroscopy in Males. *Muscle Ligaments and*
200 *Tendons J.* 2021;11: 54. doi:10.32098/mltj.01.2021.06
- 201 8. Järvinen TA. Neovascularisation in tendinopathy: from eradication to stabilisation? *Br J Sports Med.* 2020;54: 1–2.
202 doi:10.1136/bjsports-2019-100608
- 203 9. Al-Hasan AKJ. Effects of low-and high-level pulsed Nd: YAG laser irradiation on red blood cells and platelets indices
204 of albino rats in vitro. *Iraq Med J.* 2017;1: 10–19.
- 205 10. Deryugina AV, Ivashchenko MN, Ignatiev PS, Balalaeva IV, Samodelkin AG. Low-level laser therapy as a modifier of
206 erythrocytes morphokinetic parameters in hyperadrenalinemia. *Lasers Med Sci.* 2019;34: 1603–1612.
207 doi:10.1007/s10103-019-02755-y
- 208 11. Pasternak K, Nowacka O, Wróbel D, Pieszyński I, Bryszewska M, Kujawa J. Influence of MLS laser radiation on
209 erythrocyte membrane fluidity and secondary structure of human serum albumin. *Mol Cell Biochem.* 2014;388: 261–
210 267. doi:10.1007/s11010-013-1917-y
- 211 12. Fekrazad R. Photobiomodulation and Antiviral Photodynamic Therapy as a Possible Novel Approach in COVID-19
212 Management. *Photomed Laser Surg.* 2020;38: 255–257. doi:10.1089/photob.2020.4868

- 213 13. Moskvin S, Askhadulin E, Kochetkov A. Low-Level Laser Therapy in Prevention of the Development of Endothelial
214 Dysfunction and Clinical Experience of Treatment and Rehabilitation of COVID-19 Patients. *Rehabil Res Pract*.
215 2021;2021: e6626932. doi:10.1155/2021/6626932
- 216 14. Chan A-W, Tetzlaff JM, Gøtzsche PC, Altman DG, Mann H, Berlin JA, et al. SPIRIT 2013 explanation and
217 elaboration: guidance for protocols of clinical trials. *BMJ*. 2013;346. doi:10.1136/bmj.e7586
- 218 15. World Medical Association. World Medical Association Declaration of Helsinki: ethical principles for medical
219 research involving human subjects. *JAMA*. 2013;310: 2191–4. doi:10.1001/jama.2013.281053
- 220 16. Faul F, Erdfelder E, Lang A-G, Buchner A. G*Power 3: a flexible statistical power analysis program for the social,
221 behavioral, and biomedical sciences. *Behav Res Methods*. 2007;39: 175–191.
- 222 17. Spieß C. Untersuchung des Einflusses von induzierten Temperaturwechseln auf die Durchblutung der Patellar-Sehne.
223 Diploma Hochschule. 2021.
- 224 18. Kellogg DL. In vivo mechanisms of cutaneous vasodilation and vasoconstriction in humans during thermoregulatory
225 challenges. *J Appl Physiol*. 2006;100: 1709–1718. doi:10.1152/jappphysiol.01071.2005
- 226 19. Bender D, Tweer S, Werdin F, Rothenberger J, Daigeler A, Held M. The acute impact of local cooling versus local
227 heating on human skin microcirculation using laser Doppler flowmetry and tissue spectrophotometry. *Burns*. 2020;46:
228 104–109. doi:10.1016/j.burns.2019.03.009
- 229 20. Thomas T, Stefanoni D, Dzieciatkowska M, Issaian A, Nemkov T, Hill RC, et al. Evidence of Structural Protein
230 Damage and Membrane Lipid Remodeling in Red Blood Cells from COVID-19 Patients. *J Proteome Res*. 2020;19:
231 4455–4469. doi:10.1021/acs.jproteome.0c00606
- 232 21. Kubánková M, Hohberger B, Hoffmanns J, Fürst J, Herrmann M, Guck J, et al. Physical phenotype of blood cells is
233 altered in COVID-19. *Biophys J*. 2021;120: 2838–2847. doi:10.1016/j.bpj.2021.05.025

234

medRxiv preprint doi: <https://doi.org/10.1101/2022.09.27.22280424>; this version posted September 27, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.

It is made available under a [CC-BY 4.0 International license](https://creativecommons.org/licenses/by/4.0/).



Figure

Enrollment

Assessed for eligibility (n=28)

Allocation



Allocated to intervention (n=21)
• Received allocated intervention (n=21)
• intervention aborted (n=0)

Analysis



Analysed (n=21)
• Excluded from analysis (n=0)

Excluded (n=7)
• Not meeting the inclusion criteria (n=7)

Figure



Figure



Figure

medRxiv preprint doi: <https://doi.org/10.1101/2022.09.27.22280424>; this version posted September 27, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.

It is made available under a [CC-BY 4.0 International license](https://creativecommons.org/licenses/by/4.0/).



Figure