

Physician burnout and symptom of anxiety and depression: Burnout in

Consultant Doctors in Ireland Study (BICDIS)

Short title: Anxiety and depression in physician burnout: implications for the post-Covid landscape

Genevieve Crudden¹

Fabio Margiotta²

Anne M. Doherty^{3,4}

¹ Department of Psychiatry, Galway University Hospitals, Newcastle Road, Galway, Ireland

² Department of Psychiatry, Townsville University Hospital, Queensland, Australia.

³ Department of Psychiatry, University College Dublin, 63 Eccles Street, Dublin 7, Ireland.

⁴ Department of Liaison Psychiatry, Mater Misericordiae University Hospital, 63 Eccles Street, Dublin 7, Ireland.

Corresponding author

Anne M. Doherty

Department of Psychiatry

University College Dublin

63 Eccles Street,

Dublin 7,

Ireland.

Tel +353871385336

Email: anne.doherty@ucd.ie

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

Abstract

Background: Burnout is a work related syndrome comprising three major dimensions: emotional exhaustion, depersonalisation and reduced professional efficacy. It is recognised that burnout may be associated with psychopathological sequelae, such as symptoms of anxiety and depression.

Aims: This study aimed to examine the relationship between burnout and symptoms of depression and anxiety in senior doctors in Ireland. We examined personality traits as a mediating factor on burnout, in relation to work related stress among the participants.

Methods: This study was a national cross sectional survey of consultants. Data included basic demographic data, work-related data, burnout as measured by the Maslach Burnout Inventory, personality traits, and symptoms of depression and anxiety from the Depressive Anxiety Stress Scale [DASS].

Results: Results were obtained from 477 consultants, yielding a response rate 21.9%. Nearly half (42%) reported high burnout levels. The Depression and Anxiety Stress Scale revealed high levels of stress symptoms, and lower levels of anxiety symptoms. After controlling for gender and age, depressive symptoms were associated with high levels of emotional exhaustion and a higher clinical workload

Conclusions: The this study reported a high level of work related burnout, associated with symptoms of depression and anxiety in senior doctors. This is concerning, and strategies are required that consider the relationship between personality and burnout. Further research to evaluate strategies to ensure physician wellbeing and optimal delivery of patient care are required to address this serious problem.

Keywords: Burnout, depression, anxiety, psychological stress, Ireland

Introduction

Burnout is an occupational phenomenon related to stress, and has been described as “a particular response to stress involving psychological withdrawal. It is a costly coping mechanism seen as the end-point of a process which begins with the stress response”. [1] It is defined by the World Health Organisation (WHO) in ICD-11 as “a syndrome conceptualized as resulting from chronic workplace stress that has not been successfully managed. It is characterized by three dimensions: feelings of energy depletion or exhaustion; increased mental distance from one’s job, or feelings of negativism or cynicism related to one’s job; and reduced professional efficacy” [2]. The WHO classifies it as an occupational condition and not as a medical condition.

It can have a significant effect on the health of the individual worker and can affect how they view and approach their work: “What starts out as an important meaningful and challenging work becomes unpleasant, unfulfilling and meaningless.” [3]

In the decades preceding the Covid-19 pandemic, physician burnout has become an area of increasing concern, both from an occupational medicine standpoint, and from the perspective of patient safety. First described in 1974, the construct of burnout has been developed into a measurable three-dimensional model comprising emotional exhaustion, cynicism (depersonalisation) and ineffectiveness (low personal accomplishment). [4] It is a common problem with some studies reporting rates as high as 50%, and it may have a profound effect on the lives of the physicians affected. [5]

Doctors have been identified as a group who are at increased risk of developing work-related stress. In a large study comparing US physicians with the general population, although higher educational attainment was identified as a protective factor for burnout in the general population, a medical degree conferred an increased risk independent of hours worked. [5] The authors concluded that the occurrence of physician burnout is set apart from the larger societal trend, and that there are factors specific to physicians and their work which predispose this group to poorer outcomes. Riley suggested that burnout in doctors may be the end consequence of the interplay of challenging work environment, personality factors, subjective lack of control and insufficient reward. [6] The author suggests that the

“conscientious and committed personalities” working in medicine are at a loss when faced with certain environmental strains. These strains include the “demand control imbalance” identified by Karasek, [7] along with effort reward imbalance, and inadequate support in the work environment. [8] The extent of the role of personality in relation to burnout remains unclear in general, and has not been studied in the Irish context. Previous research has indicated that personality traits of neuroticism or 'negative affectivity' has been associated with greater stress and burnout in doctors [9]

The impact of burnout on patient care is well established. Burnout is associated with lower patient satisfaction, involvement in adverse events, and failure across domains of professionalism which affect patient care, including empathy, interpersonal communication and adherence to guidelines. [10-12] It has been suggested that burnout may be associated with issues relating to patient safety, as both burnout and patient safety relate to organisational, social and individual factors which depend on environmental resources. [13] There is also evidence that medical errors and their sequelae can in turn cause or worsen practitioner burnout and mental health problems, an effect that may be compounded by organisational factors including an organisational culture of blame, inadequate personal support and suboptimal administrative support [14, 15]. Two key systematic reviews examined the relationship between burnout and patient safety. Hall et al identified a relationship between poor patient safety outcomes such as errors, and noted a lack of prospective studies. [14] Dewa et al reported a strong relationship between burnout and problems with safety, but the relationship between burnout and patient satisfaction with care is less well defined. [15]

Burnout is associated with increased risk of health problems in doctors: a range of difficulties including cardiovascular disease and shorter life expectancy, problematic alcohol use, broken relationships, depression, and suicide. [16] While burnout is an occupational phenomenon rather than a mental illness, there are many areas of overlap with depression, and in particular the experience of hopelessness, helplessness and powerlessness are found in individuals with burnout and with depression. [2, 17] People who are severely affected by burnout are likely to also report symptoms of depressive illness: one Finnish study reported that half of those with severe burnout were also experiencing a depressive episode. [18]

This study aimed to investigate the prevalence of burnout and associated factors and their relationship with the development of psychiatric morbidity in a cohort of senior physicians working in the Irish Healthcare system. We sought to explore the impact of burnout and personality factors on the development of depressive and anxiety symptoms.

Methods

This study utilised a cross-sectional design, to study the prevalence of burnout in a national population of senior hospital doctors (consultants) in Ireland. The methods were previously described by Margiotta et al. [19] We will summarise here for ease of reference: an anonymous survey was distributed to members of the Irish Hospital Consultants Association (IHCA) via their electronic database. The IHCA is a representative organisation for hospital consultants across Ireland with 2160 members. An email was sent to the membership including a hyperlink to the electronic survey, the study information sheet and the consent form. Participation was voluntary, and responses were completely anonymised. Ethical approval was obtained from the Clinical Research Ethics Committee of the Saolta group. Data were collected between 1st September and 30th November 2016.

Participants were asked to provide demographic details and information and details of their current working climate and lifestyle. The questions encompassed work-related characteristics and stressors. These included the person's medical specialty, their type of contract, satisfaction with remuneration, hours worked (weekly) , time spent face-to-face with patients, and any experience of medico-legal or disciplinary action.

Burnout itself was measured by the Maslach Burnout inventory General Survey (MBI-GS) which contains 3 subscales, namely, emotional exhaustion, depersonalization or cynicism, and professional efficacy. High levels of emotional exhaustion and cynicism are associated with burnout, while high levels of professional efficacy are protective. This questionnaire is validated in medical populations of clinicians internationally. Although it is not recommended that the three internal scales of MBI-GS be combined, previous studies have created a model which adds high levels of emotional exhaustion to high levels of cynicism and low professional efficacy. [20]

The Depressive Anxiety Stress Scale (DASS) is a validated self-report instrument which measures symptoms of depression, anxiety and stress. [21] It measures each item on a 4-point Likert scale. Items related to low positive affect, decreased self-esteem, hopelessness, and poor motivation are categorised as depression; while items related to autonomic arousal and fearfulness were categorised as anxiety. Items relating to irritability and over-reactivity are categorised as stress. [22] It has been validated in clinical and non-clinical populations. [23]

The Big Five Inventory (BFI-10) is a 10-item schedule to measure personality on the traits. [24] This is a 10-item scale with questions related to the five personality traits, answered on a 5 point Likert scale from “disagree strongly”, through “disagree a little”, “neither agree nor disagree”, “agree a little” to “agree strongly”. It is an abbreviated version of the Big Five Inventory (John et al, 1991). [25]

Data were analysed using the Statistical Package for the Social Sciences (SPSS) (Version 27.0). To make comparison between groups we examined basic bivariate characteristics. Stepwise linear regression analysis was used to investigate the association of demographic variables, personality and psychiatric morbidity with burnout after controlling for the population characteristics. To investigate these hypotheses we performed further multivariable linear regression analysis with the DASS outcomes (depression, anxiety and stress) as predictors and the three burnout dimensions as the dependent variable.

Results

The characteristics of this group were previously described. [19] Of the 2,160 consultants invited to participate from the database of the Irish Hospital Consultants Association, 472 commenced the survey (response rate = 21.9%), and 433 completed the questionnaire: these were suitable for analysis. The socio-demographic characteristics of the group are outlined in Table 1. Of the respondents, 56.6% were male and 43.3% were female doctors, the mean age was 49.1 years and the majority were married or in a long-term relationship (82.7%). Medical physicians were the most represented specialty (19.4%), followed by psychiatrists (18.7%) and surgeons (15.7%). Just under one-third of respondents reported

having been involved in a work-related legal action (31.9%), and 2.1% reported an experience of formal disciplinary action in the workplace.

Although mean DASS scores for the whole group were below the cut off for caseness, 25.8% of hospital consultants reported levels of depressive symptoms above the cut-off and 13.8% reported anxiety symptoms above the cut-off. Almost 60% of participants reported stress levels above the cut off for caseness (See Table 2). The mean BFI (personality) scores are also shown in Table 2 and are significantly higher than in a normative general population sample where means sit between 3.2 (SD 0.83) for agreeableness and 4.1 (SD 0.69) for conscientiousness. [26] In this study population these means were 7.4 (SD 1.7) and 9.0 (1.3) respectively, more than double the general population scores in the literature.

We performed linear regression to examine the association between the three dimensions of burnout, psychiatric morbidity, and its contributory factors. After controlling for age and gender, higher scores on the emotional exhaustion burnout subscale were significantly associated with elevated symptoms of depression on the DASS depressive symptom subscale ($p < 0.001$). These were also associated with higher rates of face-to-face patient contact ($p = 0.001$). Individual doctors with reduced satisfaction with remuneration ($p = 0.001$) were more likely to develop psychiatric morbidity (table 3).

Elevated burnout scores on the cynical subscale were associated with high DASS stress symptoms ($p < 0.001$) after controlling for age and gender. Greater weekly hours of face-to-face patient contact was associated with higher scores on the cynicism subscale ($p = 0.022$) and poor satisfaction with remuneration was likewise associated with higher scores on the cynicism subscale ($p < 0.001$) as shown in table 3.

Professional efficacy is the third subscale of the MBI-GS and is a protective factor for the development of burnout. In this study we found that after controlling for age and gender there was an inverse relationship with the stress subscale of the DASS ($p < 0.001$), i.e., among those reporting higher professional efficacy, higher levels of stress were found. Professional efficacy is associated with lower levels of burnout. It was also significantly associated with elevated face-to-face time with patients (0.024), and there was no

association in this model with satisfaction with remuneration and professional efficacy (table 3).

Discussion

This study reported a high prevalence of burnout among hospital consultants in Ireland. High proportions screened positive for stress (60%), symptoms of depression (26%) and symptoms of anxiety (14%). Burnout is associated with higher scores of depressive symptoms and stress as measured by DASS. Higher levels of patient contact were associated with increased emotional exhaustion and cynicism, but also with elevated levels of the protective professional efficacy. Low levels of satisfaction with remuneration were associated with increased emotional exhaustion and cynicism, but there was no association with professional efficacy.

Using the composite measure of burnout, higher levels of burnout were found in this sample compared with those reported by O'Dea in a previous study of Irish general practitioners (GPs) (10.8% vs 6.6%). [20] O'Dea's study reported higher levels of emotional exhaustion (52.7% compared with 42.7%) and lower levels of the professional efficacy which is protective for burnout (16.3% compared with 1.2%). [20] O'Dea et al found similar rates of burnout to UK-based GPs according to a European study, where UK GPs scored higher than those based in Greece, Turkey, Italy and Bulgaria. [27]

Compared to interns, Irish consultants appear have lower rates of burnout than more junior hospital doctors. Two-fifths 42% of Irish interns reported a low sense of professional efficacy, and a majority (51.5%) reported high rates of emotional exhaustion [24]. It is unknown whether the working conditions of younger doctors, or more general factors associated with age predispose them to burn-out. For example, there may be an association between age and working hours as junior doctors face greater pressure to increase knowledge and skills through practice. Furthermore, their lack of experience may mean they take longer to complete tasks, thereby resulting in increased working hours. [28]

The response rate in this study compares favourably with previous response rates in similar studies: Al Humadi et al recruited 225 senior doctors, a response rate of 16.3%, and Garcia et al reported a response rate of 17.1% in their US based studies [29, 30].

Al Humadi et al reported rates of 6.2% depression, 6.6% suicidal ideation, and 19.6% burnout during the Covid-19 pandemic, all associated with history of prior depression or anxiety. Suicidal ideation and burnout were also associated with younger age. Female physicians reported reduced work-life balance and higher rates of burnout. In our study gender was not found to be significantly associated with burnout. A more recent study of Irish consultants conducted during the Covid-19 pandemic in 2021, found that three-quarters screened positive for burnout in the context of the pandemic. [31]

The association of burnout with longer time spent face-to-face with patient may be a proxy for longer hours of clinical work (given that the mean amount of patient contact was 5.5 hours per day), or it may perhaps represent the higher emotional toll of patient care, especially in certain specialities. Previous studies have found a similar association between high clinical workload and burnout. [32]

High rates of work-related burnout in consultants are highly concerning. Symptoms of stress, anxiety and depression indicate the high psychological burden of burnout. Personality factors, particularly conscientiousness and agreeableness, appear to have an influence on physician burnout. There is a need for research to identify appropriate strategies (both short-term and long-term) to minimise the impact of burnout and to ensure physician wellbeing. There is an urgent need for evidence-based strategies perhaps modulating the relationship between personality and burnout, to mitigate the impact on symptoms of depression and anxiety and to optimise doctors' health for optimal health care delivery.

Any interventions must include a consideration of both organisational and individual level measures. West et al conducted a systematic review of the evidence for the management of burnout and found strong evidence base for the use of mindfulness, stress management programmes and small group (including reflective) discussions [33]. There was limited

evidence for organisational interventions, mainly due to a lack of studies: it is much more challenging to perform a randomised controlled trial of organisational change than a mindfulness programme [33, 34]. At an individual level, doctors tend to identify organisational and structural factors as the priority areas for improving their working lives and addressing burnout [35].

The limitations of this study include its cross-sectional design and the use of DASS, a screening rather than a diagnostic measure. The small response rate, while not dissimilar to other similar studies, suggests the potential for response bias: people feeling the effects of burnout may be more likely to respond to a survey on this topic, but conversely doctors who are feeling especially overstretched may not feel able to take on another task. It is, however, a large national sample of senior clinicians which includes broad representation across geographical location and speciality.

Given these high rates of burnout and stress that were reported pre-covid, it would be important to consider the changes that have occurred in healthcare since 2020. There is an urgent need to consider the need to consider measures to mitigate the impact of burnout and to consider the causes, especially structural causes that it may be possible to mitigate.

Burnout is common in senior doctors, and it is associated with high levels of stress and symptoms of depression and anxiety. Given that physician mental health and suicide is an internationally growing area of concern, these findings suggest that burnout is associated with poorer mental health. There is an urgent need for more research on the relationship between burnout and depression and anxiety in the post-covid world and for the introduction of measures to mitigate its effect on doctors and patients.

Key points

What is already known about this subject:

- Burnout is a common problem among healthcare professionals especially doctors and includes three domains: emotional exhaustion, cynicism and professional efficacy.
- Burnout is associated with poorer mental and physical health, and people who meet the threshold for burnout frequently endorse symptoms of depression.

What this study adds:

- High-levels of burnout and stress were reported in this study of senior doctors (consultants) in this pre-covid study.
- Emotional exhaustion is associated with depressive symptoms after controlling for age, gender and higher workload.

What impact this may have on practice or policy:

- Addressing factors which contribute to burnout may have a positive impact on physician mental health.
- Work-related factors must be considered in any strategies designed to address physician burnout.

Author contributions

Conceptualisation: GC, FM AMD. Data curation: GC, FM AMD. Formal analysis: GC, AMD. Investigation: GC, FM AMD. Methodology: GC, FM AMD. Project administration: GC, FM. Supervision: AMD. Visualisation : GC, FM AMD. Writing - original draft: GC. Writing – review and editing: GC, FM, AMD.

Acknowledgements

The authors acknowledge the Irish Hospital Consultants Association for allowing access to the membership email, and to its members for participating in this study.

330

331 **Competing interests**

332 The authors have no interests to declare.

333

334 **Funding**

335 This research did not receive any specific grant from funding agencies in the public,
336 commercial, or not-for-profit sectors.

337

338

339

References

1. Cherniss C. Stress, burnout, and the special services provider. *Special Services in the Schools*. 1985;2(1):45-61.
2. WHO. ICD-11 for mortality and morbidity statistics. QD85 Burn-out. Geneva: WHO; 2020.
3. Maslach C, Leiter MP. Understanding the burnout experience: recent research and its implications for psychiatry. *World psychiatry*. 2016;15(2):103-11.
4. Maslach C, Jackson SE. The measurement of experienced burnout. *Journal of organizational behavior*. 1981;2(2):99-113.
5. Shanafelt TD, Boone S, Tan L, Dyrbye LN, Sotile W, Satele D, et al. Burnout and satisfaction with work-life balance among US physicians relative to the general US population. *Arch Intern Med*. 2012;172(18):1377-85.
6. Riley GJ. Understanding the stresses and strains of being a doctor. *Med J Aust*. 2004;181(7):350-3.
7. Karasek Jr RA. Job demands, job decision latitude, and mental strain: Implications for job redesign. *Administrative science quarterly*. 1979;285-308.
8. Johnson JV, Hall EM. Job strain, work place social support, and cardiovascular disease: a cross-sectional study of a random sample of the Swedish working population. *Am J Public Health*. 1988;78(10):1336-42.
9. Deary IJ, Blenkin H, Agius RM, Endler NS, Zealley H, Wood R. Models of job-related stress and personal achievement among consultant doctors. *Br J Psychol*. 1996;87 (Pt 1):3-29.
10. Shanafelt TD, Balch CM, Bechamps G, Russell T, Dyrbye L, Satele D, et al. Burnout and medical errors among American surgeons. *Ann Surg*. 2010;251(6):995-1000.
11. Shanafelt TD, Gradishar WJ, Kosty M, Satele D, Chew H, Horn L, et al. Burnout and career satisfaction among US oncologists. *J Clin Oncol*. 2014;32(7):678-86.
12. Anagnostopoulos F, Liolios E, Persefonis G, Slater J, Kafetsios K, Niakas D. Physician burnout and patient satisfaction with consultation in primary health care settings: evidence of relationships from a one-with-many design. *J Clin Psychol Med Settings*. 2012;19(4):401-10.
13. de Lima Garcia C, Bezerra IMP, Ramos JLS, do Valle J, Bezerra de Oliveira ML, Abreu LC. Association between culture of patient safety and burnout in pediatric hospitals. *PLoS One*. 2019;14(6):e0218756.
14. Hall LH, Johnson J, Watt I, Tsipa A, O'Connor DB. Healthcare Staff Wellbeing, Burnout, and Patient Safety: A Systematic Review. *PLoS One*. 2016;11(7):e0159015.
15. Dewa CS, Loong D, Bonato S, Trojanowski L. The relationship between physician burnout and quality of healthcare in terms of safety and acceptability: a systematic review. *BMJ Open*. 2017;7(6):e015141.
16. Kakiashvili T, Leszek J, Rutkowski K. The medical perspective on burnout. *Int J Occup Med Environ Health*. 2013;26(3):401-12.
17. Wurm W, Vogel K, Holl A, Ebner C, Bayer D, Mörk S, et al. Depression-Burnout Overlap in Physicians. *PLoS One*. 2016;11(3):e0149913.
18. Ahola K, Honkonen T, Isometsä E, Kalimo R, Nykyri E, Aromaa A, et al. The relationship between job-related burnout and depressive disorders--results from the Finnish Health 2000 Study. *J Affect Disord*. 2005;88(1):55-62.
19. Margiotta F, Crudden G, Byrne D, Doherty AM. Prevalence and co-variates of burnout in consultant hospital doctors: burnout in consultants in Ireland Study (BICDIS). *Ir J Med Sci*. 2019;188(2):355-64.

20. O'Dea B, O'Connor P, Lydon S, Murphy AW. Prevalence of burnout among Irish general practitioners: a cross-sectional study. *Ir J Med Sci.* 2017;186(2):447-53.
21. Lovibond PF, Lovibond SH. The structure of negative emotional states: comparison of the Depression Anxiety Stress Scales (DASS) with the Beck Depression and Anxiety Inventories. *Behav Res Ther.* 1995;33(3):335-43.
22. Clark LA, Watson D. Tripartite model of anxiety and depression: psychometric evidence and taxonomic implications. *J Abnorm Psychol.* 1991;100(3):316-36.
23. Antony MM, Bieling PJ, Cox BJ, Enns MW, Swinson RP. Psychometric properties of the 42-item and 21-item versions of the Depression Anxiety Stress Scales in clinical groups and a community sample. *Psychological assessment.* 1998;10(2):176.
24. Rammstedt B, John OP. Measuring personality in one minute or less: A 10-item short version of the Big Five Inventory in English and German. *Journal of research in Personality.* 2007;41(1):203-12.
25. John OP, Donahue EM, Kentle RL. Big five inventory. *Journal of Personality and Social Psychology.* 1991.
26. Rammstedt B. The 10-item Big Five Inventory: Norm values and investigation of sociodemographic effects based on a German population representative sample. *European Journal of Psychological Assessment.* 2007;23(3):193-201.
27. Soler JK, Yaman H, Esteva M, Dobbs F, Asenova RS, Katic M, et al. Burnout in European family doctors: the EGPRN study. *Fam Pract.* 2008;25(4):245-65.
28. Amofo E, Hanbali N, Patel A, Singh P. What are the significant factors associated with burnout in doctors? *Occupational medicine.* 2015;65(2):117-21.
29. Al-Humadi S, Bronson B, Muhrad S, Paulus M, Hong H, Cáceda R. Depression, Suicidal Thoughts, and Burnout Among Physicians During the COVID-19 Pandemic: a Survey-Based Cross-Sectional Study. *Acad Psychiatry.* 2021;45(5):557-65.
30. Garcia LC, Shanafelt TD, West CP, Sinsky CA, Trockel MT, Nedelec L, et al. Burnout, Depression, Career Satisfaction, and Work-Life Integration by Physician Race/Ethnicity. *JAMA Netw Open.* 2020;3(8):e2012762.
31. Doherty AM, Colleran GC, Durcan L, Irvine AD, Barrett E. A pilot study of burnout and long covid in senior specialist doctors. *Ir J Med Sci.* 2021:1-5.
32. Garcia CL, Abreu LC, Ramos JLS, Castro CFD, Smiderle FRN, Santos JAD, et al. Influence of Burnout on Patient Safety: Systematic Review and Meta-Analysis. *Medicina (Kaunas).* 2019;55(9).
33. West CP, Dyrbye LN, Erwin PJ, Shanafelt TD. Interventions to prevent and reduce physician burnout: a systematic review and meta-analysis. *Lancet.* 2016;388(10057):2272-81.
34. De Simone S, Vargas M, Servillo G. Organizational strategies to reduce physician burnout: a systematic review and meta-analysis. *Aging Clin Exp Res.* 2021;33(4):883-94.
35. Dillon EC, Tai-Seale M, Meehan A, Martin V, Nordgren R, Lee T, et al. Frontline Perspectives on Physician Burnout and Strategies to Improve Well-Being: Interviews with Physicians and Health System Leaders. *J Gen Intern Med.* 2020;35(1):261-7.

Table 1: Demographics and work related characteristics

Characteristics		
Personal characteristics		
Age (years), mean (SD)		49.14 (6.9)
Gender, n (%)	Male	253 (56.6)
	Female	194 (43.4)
Region, n (%)	Dublin	177 (39.2)
	Munster	139 (30.6)
	Connacht	63 (14.0)
	Leinster	57 (12.6)
	Ulster	15 (3.3)
Relationship status, n (%)	Married	374 (82.7)
	Long term relationship	16 (3.5)
	Single	33 (7.3)
	Divorced/separated	25 (5.5)
	Widowed	4 (0.9)
Habits, n (%)	Smoking	17 (3.7)
	Alcohol	389 (87.8)
	Exercise	355 (78.9)
Work characteristics		
Specialty, n (%)	Medicine	86 (19.9)
	Psychiatry	81 (18.7)
	Surgery	68 (15.7)
	Anaesthesia	47 (10.9)
	Radiology	46 (10.6)
	Paediatrics	30 (6.9)
	Pathology	29 (6.7)
	Obstetrics & Gynaecology	25 (5.8)
	Emergency Medicine	16 (3.7)
	Ophthalmology	5 (1.2)
Hours spent in “face-to-face” with patients per day, mean (SD)		5.5 (4.3)
Public or private practice, n (%)	Wholly in the public system	206 (46.5)
	Part public/ private	173 (39.1)
	Part academic/ public/ private	28 (6.3)
	Solely private	18 (4.1)
	Part academic/ part public	12 (2.7)
	Retired	4 (0.9)
	Locum	2 (0.5)
Satisfaction with remuneration, n (%)		209 (47.6)
Formal disciplinary action at work, n (%)		9 (2.1)
Work-related lawsuit, n (%)		140 (31.9)
Hours worked/week, mean (SD)	Total	51.8 (11.5)
	Public setting	30.9 (18.2)
	Private	7.25 (11.4)
	Managerial	3.76 (6.5)
	Academic	2.1 (4.6)
	Other	0.83 (3.7)

Table 2: Mean scores on scales measuring depression, anxiety and stress (DASS) and personality traits (BFI)

Scale/ Subscale	Mean (SD)	N (%) above cut-off
DASS total	10.9 (9.7)	
DASS depression subscale	3.3 (4.0)	107 (25.8%)
DASS anxiety subscale	1.6 (2.5)	58 (13.8%)
DASS stress subscale	6.0 (4.4)	245 (59.2%)
BFI - extraversion	6.3 (2.1)	
- agreeableness	7.4 (1.7)	
- conscientiousness	9.0 (1.3)	
- Neuroticism	5.0 (2.2)	
- Openness	7.4 (1.5)	
DASS = Depressive Anxiety Stress Scale BFI – Big Five Inventory		

Table 3: Linear regression models with Emotional exhaustion, Cynicism and Personal Efficacy as the dependent variables

	B	p-value	Confidence Interval
<i>Emotional exhaustion</i>			
Age	-0.046	0.313	-0.136 – 0.044
Gender	0.946	0.136	-0.3 – 2.192
DASS depression	1.032	<0.001	0.882 – 1.182
Face-to-face	0.219	0.001	0.085 – 0.352
Satisfaction with remuneration	2.111	0.001	0.889 – 3.333
<i>Cynicism</i>			
Age	0.014	0.746	2.799 – 13.922
Gender	0.742	0.227	-0.073 – 0.102
DASS stress	1.033	<0.001	0.9 – 1.167
Face-to-face	0.152	0.022	0.022 – 0.281
Satisfaction with remuneration	2.681	<0.001	1.501 – 3.861
<i>Professional Efficacy</i>			
Age	-0.047	0.342	-0.143 – 0.05
Gender	-0.992	0.14	-2.312 – 0.328
DASS stress	-0.459	<0.001	-0.607 – -0.311
Face-to-face	0.196	0.024	0.026 – 0.367
Satisfaction with remuneration	-0.948	0.15	-2.241 – 0.345