| 1  | Is Kinesio taping effective for peripheral tissue perfusion in women with mild to   |
|----|---|
| 2  | moderate chronic venous insufficiency? A randomized controlled trial  |
| 3  |   |
| 4  | Short title: Kinesio taping in chronic venous insufficiency   |
| 5  |   |
| 6  | Maria Luiza Vieira Carvalho <sup>1</sup> ; Viviane De Menezes Caceres <sup>2</sup> ; Isabella de Oliveira                 |
| 7  | Nascimento <sup>1</sup> ; Henrique Silveira Costa <sup>3</sup> ; Pedro Henrique Schedit Figueiredo <sup>3</sup> ; Vanessa |
| 8  | Pereira Lima <sup>3</sup> ; Debora Pantuso Monteiro <sup>1</sup> ; Danielle Aparecida Gomes Pereira <sup>1*</sup>         |
| 9  | <sup>1</sup> Graduate Program in Rehabilitation Sciences, Department of Physical Therapy – Universidade Federal           |
| 10 | de Minas Gerais, Beio Horizonte, Minas Gerais, Brazil.  |
| 11 | <sup>2</sup> Adelaide Medical School, Faculty of Health and Medical Sciences – The University of Adelaide,                |
| 12 | Adelaide, Australia   |
| 13 | <sup>3</sup> Graduate Program in Rehabilitation and Functional Performance, Department of Physical Therapy –              |
| 14 | Universidade Federal Dos Vales Do Jequitinhonha e Mucuri, Diamantina, Minas Gerais, Brazil                                |
| 15 |   |
| 16 | * Corresponding Author:   |
| 17 | E-mail: danielleufmg@gmail.com (DAGP)   |
|    |   |

## 19 Abstract

20 Objective: To evaluate the acute effect of Kinesio Taping<sup>®</sup> on peripheral tissue perfusion in 21 women with chronic venous insufficiency (CVI). Methods: This randomized, double-blind 22 controlled trial included 59 women with mild to moderate CVI. They were randomized to a 23 control group (n = 23; 54.08  $\pm$  9.04 years) and Kinesio Taping<sup>®</sup> group (KT) (n = 36; 55.87  $\pm$ 24 9.97 years). Near-infrared spectroscopy was positioned in the medial gastrocnemius muscle for 25 assessment of resting tissue perfusion 48 h after the first day of evaluation and after placement 26 of the Kinesio Taping<sup>®</sup> tape. The evaluation also consisted of performing movements of the 27 plethysmography examination. To verify the comparisons of pre- and post-Kinesio Taping® 28 data, the variation delta was used for analysis. Mann-Whitney U test was performed an an alpha 29 of 5% was considered statistically significant. Results: There wasn't a significant difference 30 between groups regarding the peripheral tissue perfusion evaluation variables Peripheral 31 Oxygen Saturation Difference - DELTA\_SPO2: (KT Group 3.21 (0.84-3.62); Control 2.21 32 (1.59-4.83), p = 0.219) and Difference in deoxygenate hemoglobin values – DELTA\_HHB 33 (units) KT Group -0.62 (-2.14–0.67); Control Group -0.07 (-2.15–2.62) p = 0.238). Despite the 34 lack of statistical significance, the KT group had a 785.7% greater drop in HHB values than the 35 control group. Conclusions: Acute use of Kinesio Taping® in women with CVI did not 36 significantly alter tissue perfusion of calf muscles. However, it was possible to observe 37 percentage differences in venous retention to be considered from a clinical point of view.

38 Trial registration: Brazilian Registry of Clinical Trials (REBEC) RBR-9bhbrp.

39 Keywords: varicose veins; rehabilitation; bandage; pain; physiotherapy specialty.

## 41 Introduction

Chronic venous insufficiency (CVI) is characterized by persistent lower limb venous 42 43 hypertension as a consequence of valvular incompetence, thrombosis, and a calf muscle 44 pump function failure [1]. CVI has a prevalence of 17% in the U.S. adult population and 45 is especially frequent in postmenopausal women because of a combination of various 46 risk factors [2]. Among other symptoms, prolonged venous hypertension can lead to a 47 perception of tiredness, heaviness, cramps, heat, or burning sensations on the skin or 48 entire limb [3,4]. Individuals with CVI present with venous retention not only at rest but 49 also during and after walking activities secondary to muscle pump dysfunction [5]. The severity of the disease has been related to reduced mobility of the talocrural joint, which 50 51 worsens the gait of affected people and contributes to the emergence of venous ulcers 52 [6]. Tissue damage in chronic venous insufficiency results from perivascular 53 inflammation caused by a variety of cytokine mechanisms that weaken the usual dermal barrier against pathogens and allergens [7]. Lymphatic dysfunction, detected by means 54 55 of nucleotide lymphangiography, is present in up to one third of cases of chronic venous 56 insufficiency and may resolve with correction of the venous abnormalities [7]. The subjective complaints related to vein hypertension and reduced ankle range of ankle 57 58 motion (ROAM) can be alleviated by compression therapy and an outpatient vascular 59 exercise program [8]. However, treatment adherence, especially with compression 60 therapy still low, and the main reasons include pain, discomfort, application difficulties, 61 skin problems, uncomfortable footwear, and inadequate lifestyle advice from health care 62 professional [8,9].

Kinesio Taping® (KT) is a therapeutic tool and has become increasingly popular in
clinical practice for the treatment of orthopedic changes, pain, muscle changes, and
edema [10]. In addition to these conditions, authors have proposed its use to reduce

| 66 | lymphatic and venous edema with promising results for symptoms of lower limb                |
|----|---|
| 67 | sensation, lameness, edema, cramps, pain, venous filling time, calf muscle power, and       |
| 68 | amount of extracellular fluid, the clinical severity of disease and physical function       |
| 69 | [11,12], and electromyographic activity of the gastrocnemius muscles [13]. However,         |
| 70 | little scientific evidence is available on the application of KT in people with vascular    |
| 71 | condition, and its usefulness in CVI has been poorly reported [11].                         |
| 72 | It is known that the homeostasis of interstitial fluid is critically important in the       |
| 73 | physiology and pathogenesis of tissues and is of therapeutic significance with important    |
| 74 | clinical implications [14]. The advocates of KT claim that convolution or mechanical        |
| 75 | deformation of the skin could lift epidermis away from the dermis altering the flow of      |
| 76 | interstitial fluid, blood, and lymph of the superficial tissues, which in turn might reduce |
| 77 | swelling and pressure of pain receptors under the skin [15].                                |
| 78 | However, the precise action mechanism of KT is not fully explained, and to further          |
| 79 | understand its benefits in individuals with CVI, it is essential to evaluate acute tissue   |
| 80 | perfusion responses using this treatment technique. Support in the study of peripheral      |
| 81 | adaptations is offered by the use of near-infrared spectroscopy (NIRS) [16]. This           |
| 82 | technique noninvasively explores tissue microvascular hemodynamics, monitoring the          |
| 83 | local balance between oxygen delivery and consumption [16]. The NIRS technique              |
| 84 | allows exploration of a limited muscle region and offers a rapid study of the changes in    |
| 85 | perfusion, skeletal muscle blood flow, and mitochondrial respiratory function [14-16].      |
| 86 | The present study aimed to evaluate the acute effect of KT on on peripheral tissue          |
| 87 | perfusion in women with CVI and compare with a placebo control group. The main              |
| 88 | hypothesis of the present study was that KT in the calf muscles could acutely improve       |
| 89 | cutaneous microvascular function and subsequent muscle oxidative metabolism in              |
| 90 | women with CVI.   |

#### 5

## 91 Methods

### 92 Study Design

- 93 This study of acute effect of KT is part of a randomized, double-blind clinical trial. It
- 94 was approved by the Research Ethics Committee of the Institution and registered at the
- 95 Brazilian Registry of Clinical Trials (REBEC) -
- 96 http://www.ensaiosclinicos.gov.br/rg/RBR-9bhbrp/.
- 97 The study was conducted between May 2017 and June 2018 and it was performed
- respecting the Declaration of Helsinki. In total, 63 volunteers were recruited; two were
- 99 excluded owing to signs of deep vein thrombosis and heart failure and two owing to
- 100 problems with data capture in NIRS (Fig 1).
- 101
- 102 Fig 1. Flow chart of study.
- 103 Fig 1 Legend: NIRS Near Infrared Spectroscopy
- 104
- 105 Participants were randomized to the control group (GC) or the Kinesio Taping group
- 106 (GKT) according to a list previously generated by the project coordinator using software
- 107 available at <u>www.randomization.com</u>. The random allocation sequence was performed
- in blocks of four participants, allocation 1:1. Participants from both groups and
- 109 evaluators were blinded in relation to GC and GKT allocation. Only the physical
- therapist who applied the KT tapes knew which group the participant belonged to.

6

## 111 **Participants**

| 112 | Sample selection was non-probabilistic, and all participants were informed about the   |
|-----|--|
| 113 | risks and benefits of the study interventions, before signing the written informed   |
| 114 | consent form. The definitive sample size was calculated in GPower $3.1^{^{\ensuremath{^$ |
| 115 | sample number reached 10 participants in each group. Considering the delta of the  |
| 116 | variables described for the study, the effect size found was 0.75, and the calculated $N$  |
| 117 | was 24 participants per group, for an alpha of 5% and a power of 80%.  |
| 118 | From May 2017 to June 2018, women with CVI aged 30 to 79 years old, from Jenny de  |
| 119 | Andrade Faria Institute for Elderly and Women's Health Care at the UFMG - Hospital   |
| 120 | das Clínicas, were enrolled in the study. The participants were included in the study  |
| 121 | according to the following inclusion criteria: being classified in stages C1, C2, or C3 of   |
| 122 | the Clinical Etiology Anatomy Pathophysiology Classification of Chronic Venous   |
| 123 | Disease (CEAP). The CEAP clinical item is classified from C0 to C6: C0 indicates the   |
| 124 | absence of clinical signs, C1 - telangiectasies or reticular veins, C2 - varicose veins, C3  |
| 125 | - edema, C4 - skin changes and subcutaneous tissue, C5 - healed ulcer, and C6 - ulcer  |
| 126 | active [17]. The exclusion criteria on the study were: hormone replacement therapy,  |
| 127 | symptoms of intermittent claudication, lymphedema, thrombosis diagnosis for less than  |
| 128 | three months, contraindication to using KT as open wounds, experiences trauma, had   |
| 129 | edema from heart or kidney disease, had cancer, known KT tape allergy, and pregnancy.  |
| 130 | If the data captured by NIRS did not allow analysis, they were excluded.   |
|     |  |

## 131 **Experimental procedure**

Two evaluations were performed for data collection, with an interval of 48 hours
between assessments. The first day consisted of an initial assessment to collect
demographic and clinical data such as age, sex, weight, height, body mass index, and

7

| 135 | analysis of disease severity according to CEAP of all participants included in the study          |
|-----|---|
| 136 | (Fig 1). NIRS was positioned in the posterior calf region of the dominant leg (medially           |
| 137 | in the medial gastrocnemius muscle) to collect the Oxygen Saturation (StO <sub>2</sub> ) and      |
| 138 | deoxyhemoglobin (HHB) data at rest for 3 min in the supine position. Subsequently, the            |
| 139 | volunteers performed the motion protocol of the plethysmography exam, which                       |
| 140 | consisted of elevation of the lower limbs to a height of 15 cm for 5 min. After that, they        |
| 141 | were asked to move to the orthostatic position without support on the evaluated leg,              |
| 142 | until the stabilization of NIRS StO <sub>2</sub> and HHB data, and were then requested to perform |
| 143 | a bipodal plantar flexion. After plantar flexion, the participants were again instructed to       |
| 144 | keep the evaluated leg at rest and without support. After stabilization of the $StO_2$ data, it   |
| 145 | was requested to perform ten plantar flexions in bipodal support. At the end of this              |
| 146 | procedure, the volunteer remained with the evaluated leg raised to stabilize the NIRS             |
| 147 | StO <sub>2</sub> and HHB data [18].   |
| 148 | On the second assessment day, the same protocol used in the first assessment to collect           |
| 149 | NIRS variables was performed on the second evaluation before and after KT placement.              |
| 150 | In GKT volunteers, the ribbons were placed in the prone position, the gastrocnemius               |
| 151 | muscles were stretched to measure the ribbons, and the "Y" cut was performed. Firstly,            |
| 152 | the leg was positioned in neutral, for fixation of the bases, without tension, in the             |
| 153 | proximal region of the popliteal fossa at the muscular origin, both in the medial and             |

154 lateral regions. Subsequently, passive stretching of the triceps suralis muscle was

performed by means of ankle dorsiflexion and KT placement with approximately 15%

tension. In the end, the muscle stretching was removed, followed by the end of the

157 application of the tensionless anchor. For participants of the CG, the KT was placed in

the same places, but the necessary positions and tensions were not followed. NIRS

variables were measured for 40 min before and after Kinesio Tex® [19] tape placement,

8

after which, participants from both groups were asked to perform the movements of the

161 air plethysmography protocol again.

### **162 Outcome Measures**

- 163 Muscle metabolism and related biomarkers were assessed by a continuous wave NIRS
- 164 system (PortaMon System, ARTINIS MEDICAL SYSTEMS BV<sup>®</sup>). Near-infrared light
- (760 850 nm) propagating through biologic tissue is partly absorbed or scattered by
- the tissues and partly recollected by the detector; therefore, the intensity of the
- recollected light provides information about oxyhemoglobin (HbO<sub>2</sub>) and HHB
- 168 concentrations. The equipment measures change in oxygenation in terms of  $HbO_2$  and
- 169 HHB in real time. In addition, it calculates the StO<sub>2</sub> index as a percentage, which
- reflects the average  $StO_2$  of muscle tissue. NIRS optodes with an interoptode distance of
- 171 3.5 cm, allowing light penetration of approximately 20 mm, were placed on the medial
- side of the calf (gastrocnemius muscles) and secured with tape.
- The parameters considered were those from motion protocol of the plethysmographyexam such as:
- Venous HHB volume (VV\_HHB): the total venous filling volume of HHB after the
- individual has changed from supine to standing position.
- HHB venous filling index (IEV\_HHB): the venous filling rate obtained by the ratio of
- 178 90% of VV\_HHB to 90% of the filling time of VV\_HHB.
- HHB value after plantar flexion (HHB\_1FP)
- HHB ejection fraction (FE\_HHB): calculated by the ratio of the HHB ejection volume
- 181 (VE\_HHB) to VV\_HHB (FE\_HHB = VE\_HHB / VV\_HHB).

9

| 182 | • Venous retention fraction (FRET): the product of the ratio between the lowest HHB |
|-----|---|
|     |   |

- 183 value after ten plantar flexions and the VV\_HHB.
- Peripheral Oxygen Saturation Difference (DELTA\_SPO<sub>2</sub>): result of subtracting SPO<sub>2</sub>
- 185 values before and after KT tape placement.
- Difference in deoxygenate hemoglobin values (DELTA\_HHB): result of subtracting
- 187 HHB values before and after KT tape placement.

### **188 Statistical procedures**

189 Statistical analysis of the data was performed by a researcher who was blinded to which

190 group the sample volunteers belonged to. Data normality distribution was performed

using the Shapiro-Wilk test and histograms. For descriptive analyses, categorical

192 variables were presented as percentages (%), while numerical variables were described

as means with their standard deviation (SD) or median with interquartile range (p25-

194 p75), depending on their distribution. Independent t-test and chi-square test were used

195 to compare baseline characteristics between groups. To verify comparisons of data

196 before and after KT placement, the variation delta was used for analysis. Mann-

197 Whitney U test was performed for peripheral tissue perfusion comparisons between the

intervention and control groups. An alpha of 5% was considered statistically significant.

The IBM Statistical Package for Social Sciences (SPSS) version 19.0 software was usedfor data analysis.

## 201 **Results**

Out of the 63 women recruited for the study, 59 women with a mean age  $54.78 \pm 9.37$ 

203 years met the inclusion criteria and were randomly assigned to the placebo control

group (n: 23) or the experimental KT group (n: 36). The recruitment flowchart and

10

follow-up of participants are summarized in Fig 1. Baseline demographic and clinical

206 characteristics of the study sample were similar between groups for all variables (Table

207 1).

### **Table 1. Baseline demographic and clinical characteristics of the sample.**

|                          | Control Group (N=23) | KT Group (N=36) | р     |
|--------------------------|----------------------|-----------------|-------|
|                          |                      |                 |       |
| Age (years)              | 55.87±9.97           | 54.08±9.04      | 0.480 |
| BMI (Kg/m <sup>2</sup> ) | $29.73 \pm 7.44$     | 30.17±6.64      | 0.813 |
| CVI Symptoms (%)         | 52.2                 | 63.9            | 0.372 |
| Thrombosis history (%)   | 4.3                  | 8.3             | 0.553 |
| Smoking (%)              | 8.7                  | 8.3             | 0.961 |
| CEAP (%)                 |                      |                 |       |
| 1                        | 4.3                  | 2.8             |       |
| 2                        | 39.2                 | 44.4            | 0.918 |
| 3                        | 56.5                 | 52.8            |       |

209 KT - Kinesio Taping; BMI - body mass index; Kg/m<sup>2</sup> - kilograms per square meter; CVI

210 – Chronic Venous Insufficiency; CEAP - Clinical Etiology Anatomy Pathophysiology
 211 Classification of Chronic Venous Disease.

212

213 Regarding the outcome variables of the peripheral tissue perfusion evaluation study,

there were no significant posttreatment differences between groups (Table 2). Despite

the absence of significant difference between groups, the KT group showed a 785.5%

216 greater drop in HHB compared to the control group.

### Table 2. Comparison of the variation of data obtained from NIRS in median and interquartile range (25-75) between KT group and

| Variables                      | Delta Control Group (N=23) | Delta KT Group (N=36)   | Difference between groups (%) | р     |
|--------------------------------|----------------------------|-------------------------|-------------------------------|-------|
| VV_HHB (units)                 | 1.03 (-8.91 – 6.36)        | 0.80 (-4.44 - 0.80)     | 22.8                          | 0.384 |
| IEV_HHB (units /seconds)       | 0.23 (0.06 - 0.69)         | 0.33 (0.14 – 0.52)      | 41.5                          | 0.901 |
| HHB_1FP (units)                | -0.19 (-10.26 - 5.32)      | -0.30 (-5.00 – 5.39)    | 57.9                          | 0.465 |
| FE_HHB (%)                     | 19.59 (12.09 - 71.42)      | 28.15 (9.98 - 63.93)    | 43.7                          | 0.852 |
| FRET (%)                       | 186.75 (97.16 – 318.53)    | 142.79 (60.18 – 248.29) | 23.5                          | 0.384 |
| DELTA_SPO <sub>2</sub> (units) | 2.21 (1.59 - 4.83)         | 3.21 (0.84 - 3.62)      | 31.0                          | 0.219 |
| DELTA_HHB (units)              | -0.07 (-2.15 – 2.62)       | -0.62 (-2.14 – 0.67)    | 785.7                         | 0.238 |

220 Legend: KT - Kinesio Taping; VV\_HHB - Venous deoxyhemoglobin volume; IEV\_HHB - Deoxyhemoglobin venous filling index; HHB\_1FP -

221 Deoxyhemoglobin value after plantar flexion; FE\_HHB - Deoxyhemoglobin ejection fraction; FRET - Venous retention fraction; DELTA\_SPO<sub>2</sub>

- Peripheral Oxygen Saturation Difference; DELTA\_HHB - Difference in deoxyhemoglobin values.

12

| 224 | Fig 2 shows the | variation in HHB | (DELTA HHB | ) according to CE | AP among women |
|-----|-----------------|------------------|------------|-------------------|----------------|
|     |                 |                  | \[         | / 0               | 0              |

- allocated to the control and KT groups. Although there was no statistically significant
- 226 difference in DELTA\_HHB between groups, there was a more evident graphic
- reduction in the median of this variable in women classified as C2 in the KT group. The
- 228 median DELTA\_HHb of women classified as CEAP 2 and allocated to the KT group (-
- 1.45 (-2.33-0.70)) dropped 56% more compared to its counterpart in the control group (-
- 230 0.63 (-2.90-2.14)).

231

### Fig 2. Variation in deoxygenated hemoglobin according to CEAP.

Fig 2 Legend - DELTA\_HHB: variation in deoxygenated hemoglobin; CEAP: Clinical
Etiology Anatomy Pathophysiology Classification of Chronic Venous Disease.

235

## 236 **Discussion**

237 In the present study, no statistically significant differences were found among variables

that evaluated tissue perfusion between GKT and CG, in women with CVI, CEAP < 4

and overweight/obese. Despite this, it is important to highlight, from a clinical point of

view, the differences in percentage values. GKT presented a 22.8% lower venous filling

volume after KT tape placement than CG (placebo), indicating a change in venous

system compliance. We could observe a 785.7% decrease in HHB value and a 31%

increase in GKT SPO<sub>2</sub> compared to CG. These alterations may suggest that clinical

changes have occurred, since changes in the variables infer compliance and

consequently modify venous hypertension and thus favor tissue oxygenation.

246 By analyzing the percentage changes in relation to the venous system filling index,

247 GKT presented a 41.5% higher filling speed. In contrast, this group had HHB emptying

- at 57.9% greater plantar flexion, 43.7% better muscle pump power, and 23.5% lower
- venous retention. That is, although the venous system of GKT was more compliant,

13

when there was muscle contraction, there was emptying of the vascular bed and less
venous retention, indicating that using the tape may favor muscle activation and
positively influence the venous system response.

253 We believe that it was not possible to verify a significant difference owing to the high

variability of NIRS data, as well as the time of use of the tape. Aguilar-Ferrándiz et al

255 (2013) [13] reported a protocol to use KT. A study was directed to responses regarding

the electromyographic activity of the medial and lateral gastrocnemius muscles [13].

257 Throughout this study, the authors detected positive changes with increased muscle

activity in women with CVI who used the tape three times a week for four weeks.

259 Therefore, it can be inferred that chronic changes in muscle activation alter vascular

responses, which in turn indirectly modify tissue perfusion, is directly dependent on the

261 difference in intraluminal and tissue pressures. Therefore, the use of KT may be

important in the treatment of venous stasis and may be used for secondary prevention of

263 CVI.

264 The literature confirms that longer use of KT can also interfere with the symptoms and

clinical variables of women with CVI, as shown in the study by Aguilar-Ferrándiz et al.

266 (2013) [13], with reduced sensation of weight in the legs, lameness, venous vein,

edema, and muscle cramps. In addition, changes in clinical parameters such as increased

venous filling time and calf muscle power, assessed by photoplethysmography, occur.

269 These data demonstrated that stimuli on skin surface caused by the placement of KT

270 may cause changes in the venous system and perfusion [13]. However, the time of

application, permanence of the tape, and form of evaluation appear to be relevant

272 factors to be considered for its use.

In the present study, tissue perfusion was evaluated after 40 min of KT placement, and although there was no statistically significant change, the percentage concentration of

14

| 275 | HHB showed clinically relevant variation, which may contribute to the clinical practice      |
|-----|--|
| 276 | for women with CVI. Tape usage time appears to be an important variable for more             |
| 277 | representative results, but this limitation is inherent in the proposed experimental         |
| 278 | design.  |
| 279 | In the present study, the NIRS tissue perfusion assessment method was used; however,         |
| 280 | the variability of data associated with a small sample was a limiting factor for the results |
| 281 | presented. Additionally, the sample consisted only of women with mild CVI severity.          |
| 282 | Studies should be conducted with a more heterogeneous sample, with greater disease           |
| 283 | severity.  |
|     |  |

284

# 285 Conclusion

From the present study, it was possible to verify that acute use of KT in women with

287 CVI could not alter tissue perfusion of the calf muscles significantly. However, it was

288 possible to observe percentage differences in venous retention to be considered from a

clinical point of view. Therefore, future studies are necessary to verify if long-term KT

use could promote changes in tissue perfusion in women with CVI.

291

**Ethics Committee:** approved by the Research Ethics Committee of the Federal University of Minas Gerais (CAAE 57708416.0.0000.5149).

294

295 **Sources of funding**: Fundação de Amparo à Pesquisa do Estado de Minas

296 Gerais - FAPEMIG [grants: Programa Pesquisador Mineiro PPM-00369-18 and CDS -

15

| 297 | APQ-02820-12]; | Conselho Naciona | l de Desenvolvimento | Científico e T | ecnológico - |
|-----|----------------|------------------|----------------------|----------------|--------------|
|-----|----------------|------------------|----------------------|----------------|--------------|

298 CNPq [grant 307597/2011-3] and PROEX/CAPES [grant 23038.002321/2020-79].

299

# 301 **References**

- 302
- 303 1. Eberhardt RT, Raffetto JD. Chronic venous insufficiency. Circulation. 2014;130:
- 304 333–346. doi:10.1161/CIRCULATIONAHA.113.006898
- 305 2. Bromen K, Pannier-Fischer F, Stang A, Rabe E, Bock E, Jöckel KH. [Should sex
- 306 specific differences in venous diseases be explained by pregnancies and hormone
- 307 intake?]. Gesundheitswesen. 2004;66: 170–174. doi:10.1055/S-2004-813019
- 308 3. Aldunate; JLCB, Isaac C, Ladeira PRS de, Carvalho VF, Ferreira MC. Venous
- 309 ulcer in lower extremities. Rev Med. 2010;89: 158–163.
- doi:10.11606/ISSN.1679-9836.V89I3/4P158-163
- 4. Piazza G. Varicose veins. Circulation. 2014;130: 582–587.
- 312 doi:10.1161/CIRCULATIONAHA.113.008331
- 5. Hosoi Y, Yasuhara H, Shigematsu H, Aramoto H, Komiyama T, Muto T. A new
- 314 method for the assessment of venous insufficiency in primary varicose veins
- using near-infrared spectroscopy. J Vasc Surg. 1997;26: 53–60.
- doi:10.1016/S0741-5214(97)70147-2
- 6. Belczak CEQ, Godoy JMP de, Seidel AC, Silva JA, Junior GC, Belczak SQ.
- 318 Assessing the influence of daily activities in the volumetry of inferior limbs via
- 319 circumference measurement and water displacement volumetry. J Vasc Bras.
- 320 2004;3: 304–310.
- Raju S, Neglén P. Chronic venous insufficiency and varicose veins. N Engl J
  Med. 2009;360: 2319–2327.
- doi:10.1056/NEJMCP0802444/SUPPL\_FILE/NEJM\_RAJU\_2319SA1.PDF

17

- 8. Heinen MM, van der Vleuten C, de Rooij MJM, Uden CJT, Evers AWM, van
- 325 Achterberg T. Physical activity and adherence to compression therapy in patients
- with venous leg ulcers. Arch Dermatol. 2007;143: 1283–1288.
- 327 doi:10.1001/ARCHDERM.143.10.1283
- 328 9. Van Hecke A, Grypdonck M, Defloor T. A review of why patients with leg
- ulcers do not adhere to treatment. J Clin Nurs. 2009;18: 337–349.
- doi:10.1111/J.1365-2702.2008.02575.X
- 10. Gloviczki P, Comerota AJ, Dalsing MC, Eklof BG, Gillespie DL, Gloviczki ML,
- et al. The care of patients with varicose veins and associated chronic venous
- diseases: Clinical practice guidelines of the Society for Vascular Surgery and the

American Venous Forum. J Vasc Surg. 2011;53: 2S–48S.

- doi:10.1016/J.JVS.2011.01.079
- 11. Aguilar-Ferrándiz ME, Castro-Sánchez AM, Matarán-Peñarrocha GA, Guisado-
- 337 Barrilao R, García-Ríos MC, Moreno-Lorenzo C. A randomized controlled trial
- of a mixed Kinesio taping-compression technique on venous symptoms, pain,
- peripheral venous flow, clinical severity and overall health status in
- 340 postmenopausal women with chronic venous insufficiency. Clin Rehabil.
- 341 2014;28: 69–81.
- 342 doi:10.1177/0269215512469120/ASSET/IMAGES/LARGE/10.1177\_026921551
- 343 2469120-FIG1.JPEG
- 344 12. Aguilar-Ferrándiz ME, Moreno-Lorenzo C, Matarán-Peñarrocha GA, García-
- 345 Muro F, García-Ríos MC, Castro-Sánchez AM. Effect of a mixed kinesio taping-
- 346 compression technique on quality of life and clinical and gait parameters in
- 347 postmenopausal women with chronic venous insufficiency: double-blinded,

| 348 randomized controlled trial. Arch Phys | s Med Rehabil. 2014;95: 1229–1239. |
|--|------------------------------------|
|--|------------------------------------|

- doi:10.1016/J.APMR.2014.03.024
- 350 13. Aguilar-Ferrándiz ME, Castro-Sánchez AM, Matarán-Peñarrocha GA, García-
- 351 Muro F, Serge T, Moreno-Lorenzo C. Effects of kinesio taping on venous
- 352 symptoms, bioelectrical activity of the gastrocnemius muscle, range of ankle
- 353 motion, and quality of life in postmenopausal women with chronic venous
- insufficiency: a randomized controlled trial. Arch Phys Med Rehabil. 2013;94:
- 355 2315–2328. doi:10.1016/J.APMR.2013.05.016
- 356 14. Swartz MA, Fleury ME. Interstitial flow and its effects in soft tissues. Annu Rev
  357 Biomed Eng. 2007;9: 229–256.
- 358 doi:10.1146/ANNUREV.BIOENG.9.060906.151850
- Wiig H. Pathophysiology of tissue fluid accumulation in inflammation. J Physiol.
  2011;589: 2945–2953. doi:10.1113/JPHYSIOL.2011.206136
- 361 16. Grassi B, Quaresima V. Near-infrared spectroscopy and skeletal muscle oxidative
- function in vivo in health and disease: a review from an exercise physiology
- 363 perspective. J Biomed Opt. 2016;21: 91313. doi:10.1117/1.JBO.21.9.091313
- 17. Eklöf B, Rutherford RB, Bergan JJ, Carpentier PH, Gloviczki P, Kistner RL, et
- 365al. Revision of the CEAP classification for chronic venous disorders: Consensus
- 366 statement. J Vasc Surg. 2004;40: 1248–1252. doi:10.1016/J.JVS.2004.09.027
- 18. Yamaki T, Nozaki M, Sakurai H, Takeuchi M, Soejima K, Kono T. The utility of
- quantitative calf muscle near-infrared spectroscopy in the follow-up of acute deep
- 369 vein thrombosis. J Thromb Haemost. 2006;4: 800–806. doi:10.1111/J.1538-
- 370 7836.2006.01859.X

- 19. Kase K, Tatsuyuki H, Tomoki O. Development of Kinesio Tape. Kinesio Taping
- Perfect Manual. Kinesio Taping Assoc. 1996;6: 117–188.





