medRxiv preprint doi: https://doi.org/10.1101/2022.11.23.22282583; this version posted November 28, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in

perpetuity. It is made available under a CC-BY-NC 4.0 International license .

Sinonasal disease among patients with primary ciliary dyskinesia – an international study

Authors

Yin Ting Lam¹⁻², Jean-François Papon³⁻⁴, Mihaela Alexandru³⁻⁴, Andreas Anagiotos⁵, Miguel Armengot⁶⁻ ⁷, Mieke Boon⁸, Andrea Burgess⁹, Suzanne Crowley¹⁰, Sinan Ahmed. D. Dheyauldeen¹¹⁻¹², Nagehan Emiralioglu¹³, Ela Erdem Eralp¹⁴, Christine van Gogh¹⁵, Yasemin Gokdemir¹⁴, Onder Gunaydın¹⁶, Eric G. Haarman¹⁷, Amanda Harris¹⁸⁻¹⁹, Isolde Hayn²⁰, Hasnaa Ismail-Koch⁹, Bülent Karadag¹⁴, Céline Kempeneers²¹, Sookyung Kim³, Philipp Latzin²², Natalie Lorent²³, Ugur Ozcelik¹³, Charlotte Pioch²⁴, Anne-Lise ML Poirrier²⁵, Ana Reula²⁶⁻²⁷, Jobst Roehmel²⁴, Panayiotis Yiallouros²⁸⁻²⁹, on behalf of the EPIC-PCD team*, Myrofora Goutaki^{1,22}

Affiliations

- 1. Institute of Social and Preventive Medicine, University of Bern, Switzerland.
- 2. Graduate School for Cellular and Biomedical Sciences, University of Bern, Bern, Switzerland.
- 3. Assistance Publique-Hôpitaux de Paris (AP-HP), Université Paris-Saclay, Hôpital Bicêtre, Service d'ORL, Le Kremlin-Bicêtre, France.
- 4. Faculté de Médecine, Université Paris-Saclay, Le Kremlin-Bicêtre, France.
- 5. Department of Otorhinolaryngology, Nicosia General Hospital, Nicosia, Cyprus.
- 6. Department of Otorhinolaryngology, and Primary Ciliary Dyskinesia Unit, La Fe University and Polytechnic Hospital, Valencia, Spain.
- 7. Medical School, Valencia University, Valencia, Spain.
- 8. Department of Paediatrics, University Hospital, Leuven, Belgium.
- 9. Primary Ciliary Dyskinesia Centre, Southampton Children's Hospital, Southampton NHS Foundation Trust, Southampton, UK.
- 10. Paediatric Department of Allergy and Lung Diseases, Oslo University Hospital, Oslo, Norway.
- 11. Department of Otorhinolaryngology, Head and Neck Surgery, Oslo University Hospital, Oslo, Norway.
- 12. Faculty of Medicine, University of Oslo, Norway.
- 13. Department of Pediatric Pulmonology, Hacettepe University, School of Medicine, Ankara, Turkey.
- 14. Department of Pediatric Pulmonology, Marmara University, School of Medicine, Istanbul, Turkey.
- 15. Department of Otorhinolaryngology Head and Neck Surgery, Amsterdam UMC, Amsterdam, The Netherlands.
- 16. Department of Otorhinolaryngology, Hacettepe University, School of Medicine, Ankara, Turkey.
- 17. Department of pediatric pulmonology, Emma Children's Hospital, Amsterdam UMC, Vrije Universiteit, Amsterdam, The Netherlands.
- 18. Southampton Children's Hospital, University of Southampton, Southampton, UK.
- 19. Primary Ciliary Dyskinesia Centre, NIHR Respiratory Biomedical Research Centre, University of Southampton, Southampton, UK.
- 20. Department of Otorhinolaryngology, Head and Neck Surgery, Charité-Universitätsmedizin Berlin, Berlin, Germany.
- 21. Division of Respirology, Department of Pediatrics, University Hospital Liège, Liège, Belgium.
- 22. Division of Paediatric Respiratory Medicine and Allergology, Department of Paediatrics, Inselspital, University Hospital, University of Bern, Switzerland.
- 23. Department of Respiratory Diseases, University Hospital, Leuven, Belgium.
- 24. Department of Pediatric Pulmonology, Immunology and Critical Care Medicine, Charité-Universitätsmedizin Berlin, Berlin, Germany.

NOTEST. Derearing the arts are constituted by performing and a generative and a generative

- 26. Biomedical Sciences Department, CEU-Cardenal Herrera University, Castellón, Spain.
- 27. Molecular, Cellular and Genomic Biomedicine Group, IIS La Fe, Valencia, Spain.
- 28. Medical School, University of Cyprus, Nicosia, Cyprus.
- 29. Pediatric Pulmonology Unit, Hospital 'Archbishop Makarios III', Nicosia, Cyprus.

* a complete list of group members appears in the acknowledgments

Correspondence: Myrofora Goutaki, Institute of Social and Preventive Medicine, University of Bern, Mittelstrasse 43, 3012 Bern, Switzerland. myrofora.goutaki@ispm.unibe.ch

Tel.: +41 31 684 59 73

Message:

Our study reinforces the importance of regular sinonasal examinations for patients of all ages with

PCD and the need for developing evidence-based sinonasal treatments as part of overall PCD

management.

Keywords:

Primary ciliary dyskinesia, epidemiology, orphan pulmonary diseases, sinusitis, upper airway, QoL

(upper airway system)

Abstract

Background: Although most patients with primary ciliary dyskinesia (PCD) report sinonasal symptoms, little is known about symptom frequency and severity. We describe sinonasal manifestations among PCD patients using data from the Ear, nose, and throat (ENT) Prospective International Cohort of PCD patients.

Methods: We included data from participants with routine clinical ENT examinations and complete FOLLOW-PCD symptoms questionnaires from the same visit or within 2 weeks. We compared the prevalence, reported symptoms, and clinical findings among children and adults and identified potential factors associated with increased risk of sinonasal disease using ordinal regression.

Results: We included 397 (53% male) participants from 12 centres with median age 15 years (IQR 9– 22). Almost all (352; 89%) reported chronic nasal symptoms. More adults (34; 26%) than children (11; 5%) reported anosmia or hyposmia. Among 140 participants who completed SNOT-22 questionnaires, quality of life was moderately affected by sinonasal symptoms (median score 31; IQR 22–45). We observed nasal polyps among 52 (15%) of 352 participants and hypertrophic inferior nasal turbinates among 129 (34%) of 353 participants; facial pain was recorded among 51 (13%) of 353 participants. More adults than children had nasal polyps, hypertrophic turbinates, deviated septum, and facial pain. We found age 10 and older the only factor associated with increased risk of sinonasal disease.

Conclusions: Our study reinforces the importance of regular sinonasal evaluations for patients of all ages with PCD and the need for developing evidence-based guidelines for sinonasal treatments as part of overall PCD management.

Introduction

Sinonasal symptoms among patients with primary ciliary dyskinesia (PCD) are as common as lower respiratory symptoms [1, 2]. Often present from birth, rhinitis is one of the first signs of PCD and usually persists throughout life [3–6]. With impaired respiratory ciliary movement and reduced mucociliary clearance, nasal secretions depend only on gravity and airflow transport [1, 7]. Sinonasal problems may manifest with rhinorrhea or blocked nose, facial pain, and headaches [8, 9]. With PCD, symptoms are part of daily life, often considered normal, and likely underreported during routine consultations. Sinonasal disease is also characterised by recurrent upper respiratory infections, often leading to chronic rhinosinusitis (CRS). Despite the clinical burden, sinonasal manifestations are frequently neglected and in many centres, ear-nose-throat (ENT) assessments are not part of routine multidisciplinary PCD care, particularly for adults [10–16]. Since sinuses may function as bacterial reservoirs for pulmonary infections later leading to lung function impairment, sinus infections are often considered only after unsuccessful treatment of pulmonary infections [17–22]. Encountered among 15–30% patients with PCD compared with 3–4% among the general population, nasal polyps are also common [6, 24]. Other sinonasal manifestations among patients with PCD include hypoplasia or agenesis of paranasal sinuses [8, 23].

The few published studies on sinonasal manifestations in PCD are mostly retrospective, include small numbers of 20–60 participants who are primarily children, and obtain data from chart reviews where symptoms were collected in a nonstandard way [25, 2]. Little is known about progression of sinonasal disease with age or with increased frequency of sinonasal symptoms. We aimed to describe the prevalence of patient-reported sinonasal symptoms and sinonasal examination findings among children and adults with PCD and identify possible risk factors associated with sinonasal disease.

Methods

Study design and population

Our study analyses cross-sectional baseline data from the ENT prospective, international cohort of patients with PCD (EPIC-PCD)—the first PCD cohort focused on upper airway disease manifestations [26]. We set up EPIC-PCD in February 2020 to follow PCD patients at their routine ENT consultations. Participants did not undergo additional testing for our study purposes. EPIC-PCD is hosted at the University of Bern (clinicaltrials.gov identifier: NCT04611516). For our collaborative study, 12 participating centres (Amsterdam, Ankara, Berlin, Bern, Cyprus, Istanbul, Leuven, Liège, Oslo, Paris, Southampton, Valencia) in 10 countries contributed data. For our analysis, we included data entered in the database by 31 July 2022 participants with PCD of all ages who underwent ENT examinations and completed symptoms questionnaires at the same visit or within 2 weeks.

We received ethical approval from all participating centers and human research ethics committees in accordance with local legislation. We obtained informed consent or assent from either participants or parents or caregivers of participants 14 years or younger. Our report conforms with the Strengthening the Reporting of Observational studies in Epidemiology (STROBE) statement [27].

Patient-reported symptoms and quality of life

For collecting patient-reported symptoms, we used the disease-specific FOLLOW-PCD questionnaire (version 1.0), which is part of the FOLLOW-PCD form developed to collect clinical information for research and clinical follow-up in a standardised way [28]. There are age-specific versions of the FOLLOW-PCD questionnaire for adults, adolescents 14–17 years, and parents or caregivers of children with PCD 14 years and younger. The FOLLOW-PCD questionnaire is available in languages of participating centres. Sinonasal symptoms questions ask about frequency and characteristics of symptoms during the past three months, specifically focusing on chronic nasal symptoms, snoring, and headaches, as well as more frequent ENT symptoms during the past 12

months. Symptom frequency options included daily, often, sometimes, rarely, and never (five-point Likert scale). Lifestyle questions asked about smoking exposure and living conditions during the past 12 months. Depending on available response categories, we recoded missing answers as "unknown," "no," or "never."

Based on local protocols, if distributed during the clinic visit we also collected information about quality of life (QoL) using the Sino-Nasal Outcome Test (SNOT-22) [29]. SNOT-22 is a validated CRS health-related QoL outcome measure. Participants score CRS-related items 0–5 each from "no problem" to "problem as bad as it can be". In total, SNOT-22 ranges between 0–110; corresponding to a mild (0–20), moderate (21–50), or severe (\geq 51) effect of CRS on QoL.

Sinonasal examinations

The EPIC-PCD is nested in routine care and follows participants at their usual ENT consultations. Performed by an ENT specialist according to local protocols, routine ENT consultations included clinical sinonasal examinations by nasal endoscopy or anterior rhinoscopy if tolerated by the participant. Examination findings were recorded in a standardised way using the ENT examination module of the FOLLOW-PCD form [28]. We recorded the proportion of the total nasal cavity volume occupied by nasal polyps using a semi-quantitative measure— the Lildholdt score— described as "partially blocking" (Lildholdt scores 1–2) and "fully blocking" (Lildholdt score 3) [30]. We recorded, reported, and present missing information from sinonasal examinations as missing.

Diagnosis and other clinical information from charts

Participants were diagnosed according to European Respiratory Society (ERS) guidelines [31]. Definite PCD diagnosis was confirmed by presence of hallmark ultrastructural defects seen in transmission electron microscopy (TEM) or by identification of bi-allelic pathogenic mutations in PCD genes. Remaining participants were diagnosed by a combination of several tests, including nasal nitric oxide, high-speed video microscopy analysis, or immunofluorescence. We collected data on laterality defects from medical records, and when it was available, past medical history information

particularly about neonatal rhinitis. Lastly, in addition to the basic dataset, some participating centres contributed information on prescribed sinonasal management. We entered all collected data in the study database—which uses the Research Electronic Data Capture (REDCap) software—based on the FOLLOW-PCD form [28].

Statistical analysis

We described characteristics of the population, patient/parent-reported sinonasal symptoms, and sinonasal examination findings, for the total population and separately among age groups 0–6, 7–14, 15–30, 31–50, and 50 years and older. For continuous variables, we used median and interquartile range (IQR); for categorical variables, we used numbers and proportions, calculating Wilson 95% confidence intervals (CI) for proportions. We compared differences between age groups using Pearson's Chi square, Wilcoxon rank-sum, and Kruskal-Wallis rank test. We created a composite outcome variable for sinonasal disease consisting of three variables: patient-reported headache while bending down as a proxy for sinusitis, ENT examination findings of nasal polyps, and facial pain. Each of them scored either 0 (absence) or 1 (presence). Total scores ranged from 0 to 3. We assessed factors possibly associated with sinonasal disease such as age, age of diagnosis, sex, study centre, smoking status of either active or passive smoke exposure, and season when ENT consultations occurred in a multivariable ordinal logistic regression model. We chose factors based on clinical importance and data availability. There was collinearity of age and age of diagnosis so it was not possible to include both in our main model; separate models showed similar results so we included age. After exploring linear and non-linear effects of age as continuous variable, we chose to include age by decades in the final model. We excluded study centre from the full model due to restricted sample size; however, we conducted sensitivity analyses with study centre alone and with age. Lastly, among a subgroup of participants with available TEM results, we repeated the model including age and category of ciliary ultrastructural defect to study if ciliary ultrastructural defect

was associated with risk for sinonasal disease. We performed all analyses with Stata version 15 (StataCorp LLC, Texas, USA).

Results

Study population

By the end of July 2022, 448 (89%) of 505 invited patients with PCD enrolled in the EPIC-PCD cohort (figure 1). Of them, 397 (53% male) participants with median age 15 years (IQR 9–22) entered in the database fulfilled eligibility criteria for ENT consultation and completed FOLLOW-PCD questionnaire at the same visit or within 2 weeks (table 1). Among participants, 257 (65%) were children, 140 (35%) adults, and 142 (36%) had situs inversus totalis. With regards to participant diagnostic status, 252 (63%) had definite PCD diagnoses based on ERS guidelines [31] with a bi-allelic PCD-causing mutation or a hallmark defect identified by TEM (table S1); 125 (32%) participants were diagnosed by a combination of several other tests; and 21 (5%) were newly diagnosed participants with diagnostic results pending at enrolment.

Patient-reported symptoms and quality of life

In total, chronic nasal symptoms were very common; most (352; 89%) participants reported nasal symptoms during the past three months (table 2). Over half of participants (204; 52%) reported chronic nasal symptoms daily or often, which most commonly "persisted all the time" for over one-third (145; 37%) of participants. Rhinorrhoea was the most commonly (167; 47%) reported nasal symptom, although nasal discharge colour varied. Some participants reported anosmia or hyposmia (45; 11%) and nearly half (190; 48%) reported snoring. Of participants reporting snoring, 46 participants snored almost every night (12%), 83 even during periods when they did not have colds (44%). Most participants reported headaches (243; 62%), which for some occurred mainly while bending down (43; 18%). Much fewer participants (27; 7%) suffered from migraines. More ENT symptoms were reported during December when compared with other months. In comparison with children, more adults reported anosmia or hyposmia (26% vs 5%, *p* <0.005), headaches (72% vs 55%,

p <0.110), and migraines (15% vs 2%, p <0.001). We did not find other differences with patientreported sinonasal symptoms by age or sex. Only 26 (7%) participants reported no sinonasal symptoms (figure S1).

In total, 140 (35%) participants who were most commonly adults completed SNOT-22 questionnaires. The median score was 31 (IQR 22–45), reflecting a moderate effect of CRS symptoms on QoL (table 2). Median SNOT-22 scores were higher with age; we observed the most severe effect on QoL (63; IQR 35–79) among participants aged 50 years and older (p <0.001), and among participants with daily nasal symptoms (figure S2).

Sinonasal clinical examinations

We excluded 15 of 397 study participants from our analysis of sinonasal examination findings for remote sinonasal consultations without sinonasal examinations. Among the remaining 382 participants, recording of sinonasal findings from sinonasal examinations was incomplete for some (table 3). For 165 (43%) participants, the nose appeared blocked, while nasal discharge was mainly serous (87; 30%) or sero-mucous (122; 43%). Abnormal nasal mucosa findings were recorded for 169 (44%), specifically mucosal oedema for 108 (26%) participants. Nasal polyps were assessed in 353 participants, and identified among 52 (14%; median age 20 years; IQR 14–36) participants with 19 (37%) located bilaterally. Of the 52 participants, 39 (75%) had nasal polyps either partially (9; 17%) or fully (9; 17%) blocking nasal passages. Nasal turbinates were hypertrophic in 129 participants (34%) and 118 participants had deviated septum (31%). 51 participants had facial pain (13%) at examination. When compared with children, more adults had nasal polyps, hypertrophic turbinates, deviated septum, and facial pain (all *p* <0.003). We did not find differences according to sex.

Information on management of upper airways

At baseline, 76 (19%) participants had hospitalisations since previous consultation, yet it was unattributed to upper respiratory infections (table S2). A small proportion of participants (17; 4%) underwent elective operations, 9 of them for sinonasal complications (53%), during this period.

Nearly one-quarter (83; 24%) of 339 participants were prescribed nasal corticosteroids, most commonly for year-round use (70; 84%). Out of 203 participants, the most common relevant nasal corticosteroid instruction involved regular nose blowing (170; 84%); out of 208 participants, instructions commonly involved nasal rinsing (190; 91%)—both instructions recommended mostly year-round use. Lastly, 50 (19%) of 269 participants (19%) were prescribed upper airway nebulisation prescriptions with isotonic saline (17; 34%), hypertonic saline (23; 46%), or other medication (9; 18%). Most commonly (47; 95%) prescribed using year-round.

Factors associated with sinonasal disease

We found age 10 years and older associated with higher risk of sinonasal disease, more profoundly when comparing participants ages 31–40 years with ages 0–10 years (odds ratio OR: 14.29, 95% CI: 5.17–39.48). Even after accounting for age, risk also differed based on study centre (table S3). We did not find associations with sex, tobacco smoke exposure, or season when consultations took place (figure 2). In the subgroup analyses of 197 participants with available TEM results (table S1), we found no association between ciliary ultrastructural defect class and risk of sinonasal disease (figure S3).

Discussion

Our study benefitted from data from the first prospective, multicentre, international ENT cohort of patients with PCD. Even though we performed our study during the COVID-19 pandemic, with much lower prevalence of viral infections, most participants reported chronic nasal symptoms, most commonly rhinorrhea. Our results showed sinonasal symptoms and clinical examination findings indicated chronic inflammation that is also more common with increasing age. Overall, QoL— measured by SNOT-22—was moderately affected by CRS (median score 31; IQR 22–45). Anterior rhinoscopic or endoscopic findings, such as nasal polyps, hypertrophic turbinates, and deviated septum, as well as facial pain at examination, were more commonly found among adults

than children. We found the risk of sinonasal disease increased with age and associated with study centre.

Strengths and limitations

Our study's main strength includes our use of data from a large, prospective, international cohort with an overall recruitment rate of 89%. We are the first to describe patient-reported sinonasal symptoms and sinonasal examination findings obtained during the same consultation for PCD. Another strength is our use of FOLLOW-PCD—it allowed standardised records of diseasespecific information and comparisons between participating centres. We excluded participants if their data were not yet entered in the study database or they did not meet eligibility criteria. We have no reasons to think exclusions were not random or affected the representation of our study population, yet participants with more sinonasal symptoms might be more willing to join EPIC-PCD when invited. We expect small risks of recall bias for patient-reported symptoms since questionnaires ask about the last three months. However, these symptoms are unspecific and part of participants' daily life, so they might be underreported. Particularly, assessing anosmia or hyposmia among young children is difficult and possibly underreported by parents. Mostly, adult participants completed SNOT-22 questionnaires, which is expected since it is only validated for adults and not used at all participating centres. We still do not know if the score sufficiently capture or underestimate effects from CRS on QoL among children. Another reason might be that the prevalence of CRS increases with age among adults in the general population; in particular CRS without nasal polyps is more prevalent among adults 40 years or younger and CRS with nasal polyps is more evident among adults 40 years or older [24]. Although our cohort was set up at the beginning of the COVID-19 pandemic, based on data from the COVID-PCD study SARS-CoV-2 infections were infrequent and caused generally mild to moderate symptoms among people with PCD [32], probably due to participants' careful shielding behaviours [33]. Generally, people with PCD and their shielding behaviours likely led to fewer infections, resulted in lower prevalence, and underestimated sinonasal problems, yet almost all participants reported nasal symptoms.

Comparison with other studies

Previous studies of upper respiratory symptoms also showed ENT symptoms common among people with PCD; however, ENT symptom definitions varied, making comparisons difficult. For instance, in a prospective study using a nationwide survey based on the FOLLOW-PCD questionnaire in Switzerland, 70 (94%) of 74 participants reported chronic nasal symptoms with rhinorrhoea (65%), blocked nose (55%), or anosmia (38%) [9]. In comparison with this study, an older population or differences in upper airway management, could explain higher prevalence of symptoms. Differences in upper airway management among participating study centres might explain the risk of sinonasal disease differences we found, yet in depth comparisons require more detailed data. Similar to our findings, a prospective study in North America described CRS among 47 children with nasal polyps (3; 6%), snoring (23; 49%), and a mean SNOT-22 score of 36.4 [34]. In a retrospective study in France, 63 of 64 adults reported sinonasal problems along with pathological nasal endoscopic findings [35], which is similar to our adult population's chronic nasal problems. In the same study, there was no correlation of ENT disease severity with ciliary ultrastructural defects. In another study assessing 39 adults with PCD and CRS in Italy, 59% had nasal polyps and more severely affected QoL—measured by SNOT-22 score—than those without nasal polyps [36]. Their findings were more severe than among our population, probably because we included children and more young adults. A study including 67 adults with PCD in Japan supports our finding that nasal polyps were observed more frequently with increasing age [37]. Although we did not observe similarly, higher odds of having CRS have been described for tobacco smoke exposure in the general population. However, our population reported a small number of participants exposed to tobacco, particularly active smoking [38].

Conclusion

We found sinonasal problems persist throughout life among people with PCD. In particular, more adults had nasal polyps and reported anosmia or hyposmia, showing complications of CRS increase with age, possibly due to ongoing chronic inflammation. Although most participants

frequently reported sinonasal symptoms, not all were prescribed sinonasal treatment or management, which could be due to patients underreporting or lack of standardised care and evidence-based PCD management guidelines for upper airways. Our study reinforces the importance of regular sinonasal examinations for PCD patients of all ages and the need to develop evidencebased sinonasal treatments as part of the overall PCD management.

Availability of data and materials

The datasets used and analysed during the current study are available from the study PI Dr Myrofora Goutaki (myrofora.goutaki@ispm.unibe.ch) upon reasonable request.

Author contribution: M Goutaki developed the concept and designed the study. M Goutaki, YT Lam manage the study. YT Lam cleaned and standardised the data and performed statistical analyses supervised by M Goutaki. YT Lam and M Goutaki drafted the manuscript. All authors commented and revised the manuscript. YT Lam and M Goutaki take final responsibility for the content.

Statement on funding sources and conflicts of interest: This study is funded by a Swiss National Science Foundation Ambizione fellowship (PZ00P3_185923). The authors participate in the BEAT-PCD (Better experimental approaches to treat PCD) clinical research collaboration, supported by the European Respiratory Society, and most centres are members of the PCD core of ERN-LUNG (European Reference Network on rare respiratory diseases). P Latzin received grants or honoraria for participation in data safety monitoring boards or advisory boards from Vertex, Vifor, OM Pharma, Polyphor, Santhera (DMC), and Sanofi Aventis within the last 36 months. J Roehmel received grants, clinical study recompensations from Vertex, INSMED, Medical Research Council/UK, BMBF, Mukoviszidose Institut.

Acknowledgements: We thank all people with PCD and their families participating in EPIC-PCD and PCD support organisations (especially PCD Family Support Group UK; Association ADCP France; Kartagener Syndrom und Primäre Ciliäre Dyskinesie e. V. Deutschland/ Deutschschweiz; Asociación Nacional de Pacientes con Discinesia Ciliar Primaria DCP España/PCD Spain) for their close collaboration. We also thank all researchers from the participating centres involved in enrolment, data collection, and data entry who work closely with us (listed below as collaborators of the EPIC-PCD study). We are grateful for everyone who contributed to translations of the FOLLOW-PCD questionnaire in Dutch, Flemish, French, Norwegian, Spanish, and Turkish. We thank Kristin Marie Bivens (ISPM, University of Bern) for her editorial assistance.

Collaborators of the EPIC-PCD study (listed in alphabetical order): Dilber Ademhan (Hacettepe University, Turkey), Mihaela Alexandru (AP-HP, France), Andreas Anagiotos (Nicosia General Hospital, Cyprus), Miguel Armengot (La Fe University and Polytechnic Hospital, Spain), Lionel Benchimol (University Hospital of Liège, Belgium), Achim G Beule (University of Münster, Germany), Irma Bon (Vrije Universiteit, the Netherlands), Mieke Boon (University Hospital Leuven, Belgium), Marina Bullo (University of Bern, Switzerland), Andrea Burgess (University of Southampton, UK), Doriane Calmes (University Hospital of Liège, Belgium), Carmen Casaulta (University of Bern, Switzerland), Marco Caversaccio (University of Bern, Switzerland), Nathalie Caversaccio (University of Bern, Switzerland), Bruno Crestani (RESPIRARE, France), Suzanne Crowley (University of Oslo, Norway), Sinan Ahmed. D. Dheyauldeen (University of Oslo, Norway), Sandra Diepenhorst (Vrije Universiteit, The Netherlands), Nagehan Emiralioglu (Hacettepe University, Turkey), Ela Erdem Eralp (Marmara University, Turkey), Pinar Ergenekon (Marmara University, Turkey), Nathalie Feyaerts (University Hospital Leuven, Belgium), Gavriel Georgiou (Nicosia General Hospital, Cyprus), Amy Glen (University of Southampton, UK), Christine van Gogh (Vrije Universiteit Amsterdam, the Netherlands), Yasemin Gokdemir (Marmara University, Turkey), Myrofora Goutaki (University of Bern, Switzerland), Onder Gunaydın (Hacettepe University, Turkey), Eric G Haarman (Vrije Universiteit Amsterdam, The Netherlands), Amanda Harris (University of Southampton, UK), Isolde Hayn (Charité-Universitätsmedizin Berlin, Germany), Simone Helms (University of Münster, Germany), Sara-Lynn Hool (University of Bern, Switzerland), Isabelle Honoré (RESPIRARE, France), Hasnaa Ismail Koch (University of Southampton, UK), Bülent Karadag (Marmara University, Turkey), Céline Kempeneers (University Hospital of Liège, Belgium), Synne Kennelly (University of Oslo, Norway), Elisabeth Kieninger (University of Bern, Switzerland), Sookyung Kim (AP-HP, France), Panayiotis Kouis (University of Cyprus, Cyprus), Yin Ting Lam (University of Bern, Switzerland), Philipp Latzin (University of Bern, Switzerland), Marie Legendre (RESPIRARE, France), Natalie Lorent (University Hospital Leuven, Belgium), Jane S Lucas (University of Southampton, UK), Bernard Maitre (RESPIRARE, France), Alison McEvoy (University of Southampton, UK), Rana Mitri-Frangieh (RESPIRARE, France), David Montani (RESPIRARE, France), Loretta Müller (University of Bern, Switzerland), Noelia Muñoz (La Fe University and Polytechnic Hospital, Spain), Heymut Omran (University of Münster, Germany), Ugur Ozcelik (Hacettepe University, Turkey), Beste Ozsezen (Hacettepe University, Turkey), Samantha Packham (University of Southampton, UK), Jean-François Papon (AP-HP, France), Clara Pauly (University Hospital of Liège, Belgium), Charlotte Pioch (Charité-Universitätsmedizin Berlin, Germany), Anne-Lise ML Poirrier (University Hospital of Liège, Belgium), Johanna Raidt (University of Münster, Germany), Ana Reula (La Fe University, Spain), Rico Rinkel (Vrije Universiteit Amsterdam, the Netherlands), Jobst Roehmel (Charité-Universitätsmedizin Berlin, Germany), Andre Schramm (University of Münster, Germany), Simone Tanner (Vrije Universiteit, the Netherlands), Guillaume Thouvenin (RESPIRARE, France), Woolf T Walker (University of Southampton, UK), Hannah Wilkins (University of Southampton, UK), Panayiotis Yiallouros (University of Cyprus, Cyprus), Ali Cemal Yumusakhuylu (Marmara University, Turkey), Niklas Ziegahn (Charité-Universitätsmedizin Berlin, Germany)

Table I. Characteristics o	.s of EPIC-PCD participants, overall and by age group (14–597)						
	Total N (%)	Age 0-6 y N (%)	Age 7-14 y N (%)	Age 15-30 y N (%)	Age 31-50 y N (%)	Age >50 y N (%)	<i>p</i> -value ^a
Number of participants	397 (100)	44 (100)	130 (100)	157 (100)	43 (100)	23 (100)	
Age, median (IQR)	15 (9–22)	4 (2–5)	10 (8–12)	16 (15–21)	38 (34–43)	57 (56–62)	
Male sex	211 (53)	23 (52)	72 (55)	81 (52)	24 (56)	11 (48)	0.937
Age of PCD diagnosis.	9 (4–17)	1 (0-2)	6 (1-8)	12 (8–17)	34 (29–36)	51 (43–55)	
median (IOR)	- (,	- ()	- ()	(• ,		(,	
Consanguinity							0.001
Yes	119 (30)	7 (16)	45 (35)	52 (33)	11 (26)	4 (17)	
No	136 (35)	13 (29)	40 (30)	61 (39)	19 (44)	3 (13)	
Not reported	142 (35)	24 (55)	45 (35)	44 (28)	13 (30)	16 (70)	
·	. ,	ζ, γ		ζ, γ	· · /		
Situs							<0.001
Situs inversus totalis	142 (36)	25 (57)	46 (35)	60 (38)	7 (16)	4 (17)	
Situs ambiguous	4 (1)	0 (0)	1(1)	3 (2)	0 (0)	0 (0)	
Situs solitus	245 (62)	18 (41)	82 (63)	94 (60)	32 (74)	19 (83)	
Not reported	6 (1)	1 (2)	1 (1)	0 (0)	4 (9)	0 (0)	
			. ,				
Cardiovascular malformat	ion present						0.088
Yes	34 (9)	7 (16)	11 (8)	14 (9)	2 (5)	0 (0)	
No	295 (74)	30 (68)	104 (80)	114 (73)	32 (74)	15 (65)	
Not reported	68 (17)	7 (16)	15 (12)	29 (18)	9 (21)	8 (35)	
		(-)	- ()	- (-)	- ()	- ()	
Neonatal rhinitis							0.189
Yes	125 (32)	15 (34)	47 (36)	49 (31)	12 (28)	2 (9)	
No	88 (22)	7 (16)	27 (21)	39 (25)	11 (26)	4 (17)	
Not reported	184 (46)	22 (50)	56 (43)	69 (44)	20 (46)	17 (74)	
	((= =)		()		()	
Neonatal cough							<0.001
Yes	129 (33)	15 (34)	44 (34)	59 (38)	10 (23)	1 (4)	
No	91 (23)	3 (7)	31 (24)	40 (25)	14 (33)	3 (13)	
Not reported	177 (44)	26 (59)	55 (42)	58 (37)	19 (44)	19 (83)	
·	. ,	ζ, γ		ζ, γ	· · /	· · ·	
Neonatal respiratory distr	ess						0.554
Yes	181 (46)	26 (59)	60 (46)	67 (43)	20 (46)	8 (35)	
No	143 (36)	11 (25)	49 (38)	60 (38)	15 (35)	8 (35)	
Not reported	73 (18)	7 (16)	21 (16)	30 (19)	8 (19)	7 (30)	
•	, , , , , , , , , , , , , , , , , , ,	()	()	()	ζ, γ	ζ,	
Active smoking							<0.001
Yes. daily	4 (1)	NA	NA	3 (2)	0 (0)	1 (4)	
Yes, rarely	5 (1)	NA	NA	3 (2)	1 (2)	1 (4)	
Fx-smoker	15 (4)	NA	NA	2 (1)	8 (19)	5 (22)	
Never smoker	188 (47)	NA	NA	141 (90)	32 (74)	15 (65)	
Not reported	185 (47)	NΔ	NA	8 (5)	2 (5)	1 (5)	
	100 (77)			0(0)	2 (3)	- (5)	
Smoking in household							0.002
Yes	70 (18)	6 (14)	24 (18)	34 (22)	4 (9)	2 (9)	
No	262 (66)	34 (77)	97 (75)	89 (56)	27 (63)	15 (65)	
Not reported	65 (16)	4 (9)	9 (7)	34 (22)	12 (28)	6 (26)	

Table 1: Characteristics of EPIC-PCD participants, overall and by age group (N=397)

EPIC-PCD: Ear-nose throat prospective international cohort of patients with primary ciliary dyskinesia. y: years. All characteristics are presented as N and column % with the exception of age as median and IQR: interquartile range. NA: not applicable; age-specific questionnaire version does not include question on active smoking in this agecategory. ^a: Chi-square test of independence.

	n voluo ^a
$\frac{1}{10} \frac{1}{10} \frac$	<i>p</i> -value
N (%)	
Number of participants $397(100)$ 44 (100) 130 (100) 157 (100) 43 (100) 23 (100)	
Nasal symptoms	0.189
Daily/often 204 (52) 19 (43) 68 (52) 72 (46) 29 (67) 16 (70)	
Sometimes/rarely 148 (37) 18 (41) 48 (37) 65 (41) 11 (26) 6 (26)	
Never45 (11)7 (16)14 (11)20 (13)3 (7)1 (4)	
Nasal symptoms 145 (37) 17 (39) 46 (35) 52 (33) 22 (51) 8 (35) persisting all the time ^b	0.786
Type of nasal symptoms ^c	
Rhinorrhoea 317 (90) 34 (92) 102 (88) 124 (91) 36 (90) 21 (95)	0.991
Blocked nose 240 (68) 17 (46) 83 (72) 96 (70) 32 (80) 12 (55)	0.459
Sneezing $76(16)$ $5(14)$ $19(16)$ $32(18)$ $9(10)$ $11(18)$	0.054
Anosmia/hyposmia $45(13)$ $0(0)$ $7(6)$ $14(10)$ $14(35)$ $10(45)$	<0.001
	0.001
Colour of nasal discharge in case of rhinorrhoea ^d	
Clear 65 (16) 9 (20) 19 (15) 27 (17) 6 (14) 4 (17)	0.940
White 60 (15) 6 (14) 23 (18) 21 (14) 6 (14) 4 (17)	0.922
Yellow 106 (27) 10 (23) 34 (26) 45 (29) 12 (28) 5 (22)	0.964
Green 75 (19) 9 (21) 25 (19) 26 (27) 9 (21) 6 (26)	0.907
Mixed with blood 11 (3) 0 (0) 1 (1) 5 (3) 3 (7) 2 (9)	0.077
Constinue	0.005
	0.005
Daily/ often 46 (12) 5 (11) 14 (11) 11 (7) 11 (26) 5 (22)	
Sometimes/rarely 144 (36) 20 (46) 51 (39) 48 (31) 15 (35) 10 (43)	
Never/ not reported 207 (52) 19 (43) 65 (50) 98 (62) 17 (39) 8 (35)	
Periods of snoring ^e	0.096
Almost every night 43 (23) 5 (20) 13 (20) 9 (15) 10 (38) 6 (40)	
Only during colds 48 (25) 9 (36) 19 (29) 11 (19) 7 (27) 2 (13)	
Sometimes also without 83 (44) 8 (32) 29 (45) 35 (59) 6 (23) 5 (34)	
colds	
Not reported 16 (8) 3 (12) 4 (6) 4 (7) 3 (12) 2 (13)	
Headache	<0.001
Daily/often $50(13) 0(0) 15(12) 20(13) 9(21) 6(26)$	
Sometimes/rarely $193 (19) = 9 (20) = 55 (12) = 20 (13) = 5 (21) = 0 (20)$	
Never / not reported $154 (30) = 35 (80) = 60 (46) = 46 (30) = 6 (14) = 7 (30)$	
$\frac{1}{100} \frac{1}{100} \frac{1}$	
Headache when bending 43 (11) 1 (2) 7 (5) 28 (18) 5 (12) 2 (9) down 2 (9) 2 (9)	0.014
Migraines	<0.001
Yes 27 (7) 0 (0) 3 (2) 10 (6) 7 (16) 7 (30)	
No 370 (93) 44 (100) 127 (98) 147 (94) 36 (84) 16 (70)	
EVICE 32 completed $140(25)$ $14(25)$ $20(25)$ $50(25)$ $50(25)$ $27(52)$	-0.004
Single 22 completed 140 (35) 14 (32) 29 (22) 56 (36) 27 (63) 14 (61) $27 (63) = 21 (62, 51) - 25 (45, 26) - 27 (45, 11) - 26 (46, 16) - 26 (25, 51) - 56 $	<0.001

Table 2: Upper respiratory symptoms of past 3 months reported by EPIC-PCD participants, overall and by age group (N=397)

EPIC-PCD: Ear-nose throat prospective international cohort of patients with primary ciliary dyskinesia. y: years. Symptoms are presented as N and column %. SNOT-22 presented as median and IQR: interguartile range. Sino Nasal Outcome Test-22 (SNOT-22) questionnaire on chronic rhinosinusitis related items scored 0–5 ("No problem" to "Problem as bad as it can be"), total score range 0–110, mild 0–20, moderate 21–50, severe ≥51. ^a: Chi-square test of independence. ^b: Among 352 people with chronic

nasal symptoms ^c: Among 352 people with chronic nasal symptoms, categories are not exclusive. ^d: Among 317 people with rhinorrhea, categories are not exclusive. ^e: Among 190 people with snoring. ^f: Kruskal-Wallis test.

	Total N (%)	Age 0-6 y	Age 7-14 y	Age 15-30 y	Age 31-50 y	Age >50 y	<i>p</i> -value ^a
		N (%)	N (%)	N (%)	N (%)	N (%)	•
ENT consultations on site	382 (100)	38 (100)	123 (100)	155 (100)	43 (100)	23 (100)	
Nose appearance							0.054
Normal	209 (55)	23 (60)	76 (62)	85 (55)	16 (37)	9 (39)	
Blocked	165 (43)	14 (37)	45 (37)	68 (44)	26 (61)	12 (52)	
Not recorded	8 (2)	1 (3)	2 (2)	2 (1)	1 (2)	2 (9)	
Nasal discharge present							0.767
Yes	286 (75)	28 (74)	94 (76)	112 (72)	34 (79)	18 (78)	
No	88 (23)	9 (24)	27 (22)	41 (27)	7 (16)	4 (18)	
Not recorded	8 (2)	1 (2)	2 (2)	2 (1)	2 (5)	1 (4)	
Type of nasal discharge ^b							0.866
Serous	87 (30)	10 (36)	30 (32)	32 (28)	9 (27)	6 (33)	
Sero-mucous	122 (43)	11 (39)	42 (45)	46 (41)	16 (47)	7 (39)	
Muco-purulent	66 (23)	6 (21)	19 (20)	29 (26)	8 (23)	4 (22)	
Mixed with blood	3 (1)	1 (4)	0 (0)	1 (1)	0 (0)	1 (6)	
Not recorded	8 (3)	0 (0)	3 (3)	4 (4)	1 (3)	0 (0)	
Nasal mucosa							0.018
Abnormal	169 (44)	13 (34)	53 (43)	68 (44)	21 (49)	14 (61)	
Normal	202 (53)	21 (55)	67 (55)	86 (55)	21 (49)	7 (30)	
Not recorded	11 (3)	4 (11)	3 (2)	1 (1)	1 (2)	2 (9)	
Nasal polyps							0.001
Yes	52 (14)	2 (5)	11 (9)	22 (14)	12 (28)	5 (22)	
No	301 (79)	28 (74)	100 (81)	127 (82)	29 (68)	17 (74)	
Not assessed	29 (7)	8 (21)	12 (10)	6 (4)	2 (5)	1 (4)	
Nasal polyps size ^{cd}							0.827
Fully blocking	9 (17)	1 (50)	3 (27)	6 (4)	2 (17)	0 (0)	
Partially blocking	39 (75)	1 (50)	7 (64)	17 (77)	9 (75)	5 (100)	
Not assessed	4 (8)	0 (0)	1 (9)	2 (9)	1 (8)	0 (0)	
Bilaterally ^{cd}							0.263
Fully blocking	17 (33)	1 (50)	3 (27)	6 (4)	4 (33)	3 (60)	
Partially blocking	2 (4)	0 (0)	2 (18)	0 (0)	0 (0)	0 (0)	
Not recorded	33 (64)	1 (50)	6 (55)	16 (72)	8 (67)	2 (40)	
Unilaterally ^{cd}							0.213
Fully blocking	39 (75)	2 (5)	7 (6)	15 (10)	11 (26)	4 (17)	-
Partially blocking	4 (8)	0 (0)	3 (2)	1 (1)	0 (0)	0 (0)	
Not recorded	9 (17)	0 (0)	1 (9)	6 (27)	1 (8)	1 (20)	

(NI_202)

EPIC-PCD: Ear-nose throat prospective international cohort of patients with primary ciliary dyskinesia. y: years. Symptoms are presented as N and column %.^a: Chi-square test of independence.^b: Among 286 people with nasal discharge.^c: Among 52 participants with nasal polyps.^d: Nasal polyps described as partially blocking or with Lildholdt score 1 or 2, fully blocking or with Lildholdt score 3.

Table 3 (continued): Sinonasal examination results of EPIC-PCD participants, overall and by age group (N=382)							
	Total N (%)	Age 0-6 y	Age 7-14 y	Age 15-30 y	Age 31-50 y	Age >50 y	<i>p</i> -value ^a
		N (%)	N (%)	N (%)	N (%)	N (%)	
ENT consultations on site	382 (100)	38 (100)	123 (100)	155 (100)	43 (100)	23 (100)	
Inferior nasal turbinates							0.003
Normal	221 (58)	21 (55)	67 (55)	100 (64)	21 (49)	12 (52)	
Hypertrophy	129 (34)	13 (34)	47 (38)	46 (30)	15 (35)	8 (35)	
Atrophy	3 (1)	0 (0)	0 (0)	0 (0)	3 (7)	0 (0)	
Not recorded	29 (7)	4 (11)	9 (7)	9 (6)	4 (9)	3 (13)	
Deviated nasal septum							<0.001
Yes	118 (31)	2 (5)	28 (23)	66 (42)	12 (28)	10 (43)	
Bulging forward	5 (1)	1 (3)	0 (0)	1 (1)	2 (5)	1 (5)	
No	235 (62)	28 (74)	88 (71)	81 (52)	28 (65)	10 (43)	
Not recorded	24 (6)	7 (18)	7 (6)	7 (5)	1 (2)	2 (9)	
Facial pain or sensitivity							<0.001
Yes	51 (13)	0 (0)	9 (7)	21 (14)	12 (28)	9 (39)	
No	302 (79)	27 (71)	107 (87)	127 (82)	28 (65)	13 (57)	
Not recorded	29 (8)	11 (29)	7 (6)	7 (4)	3 (7)	1 (4)	

EPIC-PCD: Ear-nose throat prospective international cohort of patients with primary ciliary dyskinesia. y: years. Symptoms are

presented as N and column %. a: Chi-square test of independence

Figure legends

Figure 1: Flowchart of study population of EPIC-PCD (ENT prospective international cohort of patients with primary ciliary dyskinesia).

Figure 2: Factors associated with sinonasal disease in EPIC-PCD participants (N=397).

Sinonasal disease defined by composite outcome score consisting of three variables: patient-reported headache while bending down as a proxy for sinusitis, and ENT examination findings of nasal polyps and facial pain. EPIC-PCD: Ear-nose throat prospective international cohort of patients with primary ciliary dyskinesia. Odds ratio (OR) indicated by squares and 95% confidence intervals by horizontal lines.

Figure S1: Proportion of patient-reported symptoms (snoring, headache, nasal symptoms) during the past three months among EPIC-PCD participants (N=397).

EPIC-PCD: Ear-nose throat prospective international cohort of patients with primary ciliary dyskinesia.

Figure S2: Sino-Nasal Outcome Test (SNOT)-22 score by frequency of patient-reported nasal symptoms during the past three months among EPIC-PCD participants (N=140).

SNOT-22 scores 0–5 ("No problem" to "Problem as bad as it can be"); total score range 0–110; mild 0–20, moderate 21–50; severe \geq 51. Score ranges indicated by horizontal lines. EPIC-PCD: Ear-nose throat prospective international cohort of patients with primary ciliary dyskinesia.

Figure S3: Association of ciliary ultrastructural defect with sinonasal disease in EPIC-PCD participants (N=197).

Sinonasal disease defined by composite outcome score consisting of three variables: patient-reported headache while bending down as a proxy for sinusitis, ENT examination findings of nasal polyps, and facial pain. EPIC-PCD: Ear-nose throat prospective international cohort of patients with primary ciliary dyskinesia. Odds ratio (OR) indicated by squares and 95% confidence intervals (CI) by horizontal lines.

References

1. Morgan LC, Birman CS. The impact of Primary Ciliary Dyskinesia on the upper respiratory tract. *Paediatr Respir Rev* 2016; 18: 33–38.

2. Goutaki M, Meier AB, Halbeisen FS, Lucas JS, Dell SD, Maurer E, Casaulta C, Jurca M, Spycher BD, Kuehni CE. Clinical manifestations in primary ciliary dyskinesia: systematic review and metaanalysis. *The European respiratory journal* 2016/08/06 ed. 2016; 48: 1081–1095.

3. Sommer JU, Schäfer K, Omran H, Olbrich H, Wallmeier J, Blum A, Hörmann K, Stuck BA. ENT manifestations in patients with primary ciliary dyskinesia: prevalence and significance of otorhinolaryngologic co-morbidities. *Eur Arch Otorhinolaryngol* 2011; 268: 383–388.

4. Hosie PH, Fitzgerald DA, Jaffe A, Birman CS, Rutland J, Morgan LC. Presentation of primary ciliary dyskinesia in children: 30 years' experience. *J Paediatr Child Health* 2015; 51: 722–726.

5. Behan L, Dimitrov BD, Kuehni CE, Hogg C, Carroll M, Evans HJ, Goutaki M, Harris A, Packham S, Walker WT, Lucas JS. PICADAR: a diagnostic predictive tool for primary ciliary dyskinesia. *The European respiratory journal* 2016/02/27 ed. 2016; 47: 1103–1112.

6. Goutaki M, Halbeisen FS, Barbato A, Crowley S, Harris A, Hirst RA, Karadag B, Martinu V, Morgan L, O'Callaghan C, Ozçelik U, Scigliano S, Ucros S, Yiallouros P, Schulzke SM, Kuehni CE. Late Diagnosis of Infants with PCD and Neonatal Respiratory Distress. *Journal of clinical medicine* 2020/09/10 ed. 2020; 9.

7. Stannard W, O'Callaghan C. Ciliary function and the role of cilia in clearance. *J Aerosol Med* 2006; 19: 110–115.

8. Bequignon E, Dupuy L, Zerah-Lancner F, Bassinet L, Honoré I, Legendre M, Devars du Mayne M, Escabasse V, Crestani B, Maître B, Escudier E, Coste A, Papon JF. Critical Evaluation of Sinonasal Disease in 64 Adults with Primary Ciliary Dyskinesia. *Journal of clinical medicine* 2019/05/10 ed. 2019; 8.

9. Goutaki M, Hüsler L, Lam YT, Koppe HM, Jung A, Lazor R, Müller L, Pedersen ESL, Kuehni CE. Respiratory symptoms of Swiss people with primary ciliary dyskinesia. *ERJ open research* 2022/04/14 ed. 2022; 8.

10. Mener DJ, Lin SY, Ishman SL, Boss EF. Treatment and outcomes of chronic rhinosinusitis in children with primary ciliary dyskinesia: where is the evidence? A qualitative systematic review. *Int Forum Allergy Rhinol* 2013; 3: 986–991.

11. Lucas JS, Burgess A, Mitchison HM, Moya E, Williamson M, Hogg C. Diagnosis and management of primary ciliary dyskinesia. *Archives of disease in childhood* 2014/04/29 ed. 2014; 99: 850–856.

12. Damseh N, Quercia N, Rumman N, Dell SD, Kim RH. Primary ciliary dyskinesia: mechanisms and management. *The application of clinical genetics* 2017/10/17 ed. 2017; 10: 67–74.

13. Lucas JS, Alanin MC, Collins S, Harris A, Johansen HK, Nielsen KG, Papon JF, Robinson P, Walker WT. Clinical care of children with primary ciliary dyskinesia. *Expert review of respiratory medicine* 2017/07/27 ed. 2017; 11: 779–790.

14. Kuehni CE GM Rubbo B, Lucas JS. Bronchiectasis (ERS Monograph) [Internet]. Sheffield: European Respiratory Society; 2018. Available from: http://books.ersjournals.com/content/9781849840989/9781849840989.

15. Rubbo B, Best S, Hirst RA, Shoemark A, Goggin P, Carr SB, Chetcuti P, Hogg C, Kenia P, Lucas JS, Moya E, Narayanan M, O'Callaghan C, Williamson M, Walker WT. Clinical features and management of children with primary ciliary dyskinesia in England. *Archives of disease in childhood* 2020/03/12 ed. 2020; 105: 724–729.

16. Goutaki M, Shoemark A. Diagnosis of Primary Ciliary Dyskinesia. *Clin Chest Med* 2022; 43: 127–140.

17. Alanin MC, Johansen HK, Aanaes K, Høiby N, Pressler T, Skov M, Nielsen KG, von Buchwald C. Simultaneous sinus and lung infections in patients with primary ciliary dyskinesia. *Acta Otolaryngol* 2015; 135: 58–63.

18. Licari A, Castagnoli R, Denicolò CF, Rossini L, Marseglia A, Marseglia GL. The Nose and the Lung: United Airway Disease? *Front Pediatr* 2017; 5: 44.

19. Alanin MC. Bacteriology and treatment of infections in the upper and lower airways in patients with primary ciliary dyskinesia: adressing the paranasal sinuses. *Danish medical journal* 2017/05/30 ed. 2017; 64.

20. Davis SD, Rosenfeld M, Lee H-S, Ferkol TW, Sagel SD, Dell SD, Milla C, Pittman JE, Shapiro AJ, Sullivan KM, Nykamp KR, Krischer JP, Zariwala MA, Knowles MR, Leigh MW. Primary Ciliary Dyskinesia: Longitudinal Study of Lung Disease by Ultrastructure Defect and Genotype. *Am J Respir Crit Care Med* 2019; 199: 190–198.

21. Kos R, Israëls J, van Gogh CDL, Altenburg J, Diepenhorst S, Paff T, Boon EMJ, Micha D, Pals G, Neerincx AH, Maitland-van der Zee AH, Haarman EG. Primary ciliary dyskinesia in Volendam: Diagnostic and phenotypic features in patients with a CCDC114 mutation. *American journal of medical genetics. Part C, Seminars in medical genetics* 2022/03/29 ed. 2022; 190: 89–101.

22. Halbeisen FS, Pedersen ESL, Goutaki M, Spycher BD, Amirav I, Boon M, Malena C-C, Crowley S, Emiralioglu N, Haarman EG, Karadag B, Koerner-Rettberg C, Latzin P, Loebinger MR, Lucas JS, Mazurek H, Morgan L, Marthin J, Pohunek P, Santamaria F, Schwerk N, Thouvenin G, Yiallouros P, Nielsen KG, Kuehni CE. Lung function from school age to adulthood in primary ciliary dyskinesia. *Eur Respir J* 2022; : 2101918.

23. Pappa AK, Sullivan KM, Lopez EM, Adams KN, Zanation AM, Ebert CS Jr, Thorp BD, Senior BA, Leigh MW, Knowles MR, Kimple AJ. Sinus Development and Pneumatization in a Primary Ciliary Dyskinesia Cohort. *American journal of rhinology & allergy* 2020/06/20 ed. 2021; 35: 72–76.

24. Fokkens WJ, Lund VJ, Hopkins C, Hellings PW, Kern R, Reitsma S, Toppila-Salmi S, Bernal-Sprekelsen M, Mullol J, Alobid I, Terezinha Anselmo-Lima W, Bachert C, Baroody F, von Buchwald C, Cervin A, Cohen N, Constantinidis J, De Gabory L, Desrosiers M, Diamant Z, Douglas RG, Gevaert PH, Hafner A, Harvey RJ, Joos GF, Kalogjera L, Knill A, Kocks JH, Landis BN, Limpens J, et al. European Position Paper on Rhinosinusitis and Nasal Polyps 2020. *Rhinology* 2020; 58: 1–464.

25. Campbell R. Managing upper respiratory tract complications of primary ciliary dyskinesia in children. *Curr Opin Allergy Clin Immunol* 2012; 12: 32–38.

26. Goutaki M, Lam YT, Alexandru M, Anagiotos A, Armengot M, Bequignon E, Boon M, Burgess A, Coste A, Emiralioglu N, Erdem E, Haarman EG, Harris A, Hool SL, Karadag B, Kim S, Latzin P, Lorent N, Ozcelik U, Reula A, Roehmel J, van Gogh C, Yiallouros P, Zappe SM, Papon JF. Study protocol: the ear-nose-throat (ENT) prospective international cohort of patients with primary ciliary dyskinesia (EPIC-PCD). *BMJ open* 2021/10/13 ed. 2021; 11: e051433.

von Elm E, Altman DG, Egger M, Pocock SJ, Gøtzsche PC, Vandenbroucke JP. The
Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement:
guidelines for reporting observational studies. *Journal of clinical epidemiology* 2008/03/04 ed. 2008;
61: 344–349.

28. Goutaki M, Papon JF, Boon M, Casaulta C, Eber E, Escudier E, Halbeisen FS, Harris A, Hogg C, Honore I, Jung A, Karadag B, Koerner-Rettberg C, Legendre M, Maitre B, Nielsen KG, Rubbo B, Rumman N, Schofield L, Shoemark A, Thouvenin G, Willkins H, Lucas JS, Kuehni CE. Standardised clinical data from patients with primary ciliary dyskinesia: FOLLOW-PCD. *ERJ open research* 2020/02/15 ed. 2020; 6.

29. Kennedy JL, Hubbard MA, Huyett P, Patrie JT, Borish L, Payne SC. Sino-nasal outcome test (SNOT-22): a predictor of postsurgical improvement in patients with chronic sinusitis. *Ann Allergy Asthma Immunol* 2013; 111: 246-251.e2.

30. Johansson L, Akerlund A, Holmberg K, Melén I, Stierna P, Bende M. Evaluation of methods for endoscopic staging of nasal polyposis. *Acta Otolaryngol* 2000; 120: 72–76.

31. Lucas JS, Barbato A, Collins SA, Goutaki M, Behan L, Caudri D, Dell S, Eber E, Escudier E, Hirst RA, Hogg C, Jorissen M, Latzin P, Legendre M, Leigh MW, Midulla F, Nielsen KG, Omran H, Papon JF, Pohunek P, Redfern B, Rigau D, Rindlisbacher B, Santamaria F, Shoemark A, Snijders D, Tonia T, Titieni A, Walker WT, Werner C, et al. European Respiratory Society guidelines for the diagnosis of primary ciliary dyskinesia. *The European respiratory journal* 2016/11/12 ed. 2017; 49.

32. Pedersen ESL, Goutaki M, Harris AL, Dixon L, Manion M, Rindlisbacher B, Patient Advisory Group CP, Lucas JS, Kuehni CE. SARS-CoV-2 infections in people with primary ciliary dyskinesia: neither frequent, nor particularly severe. *The European respiratory journal* 2021/04/10 ed. 2021; 58.

33. Pedersen ESL, Collaud ENR, Mozun R, Dexter K, Kruljac C, Silberschmidt H, Lucas JS, Goutaki M, Kuehni CE. Facemask Usage Among People With Primary Ciliary Dyskinesia During the COVID-19 Pandemic: A Participatory Project. *International journal of public health* 2022/01/04 ed. 2021; 66: 1604277.

34. Zawawi F, Shapiro AJ, Dell S, Wolter NE, Marchica CL, Knowles MR, Zariwala MA, Leigh MW, Smith M, Gajardo P, Daniel SJ. Otolaryngology Manifestations of Primary Ciliary Dyskinesia: A Multicenter Study. *Otolaryngology--head and neck surgery : official journal of American Academy of Otolaryngology-Head and Neck Surgery* 2021/06/23 ed. 2022; 166: 540–547.

35. Bequignon E, Dupuy L, Escabasse V, Zerah-Lancner F, Bassinet L, Honoré I, Legendre M, Devars du Mayne M, Crestani B, Escudier E, Coste A, Papon JF, Maître B. Follow-Up and Management of Chronic Rhinosinusitis in Adults with Primary Ciliary Dyskinesia: Review and Experience of Our Reference Centers. *Journal of clinical medicine* 2019/09/25 ed. 2019; 8.

36. Piatti G, Ambrosetti U, Aldè M, Girotto G, Concas MP, Torretta S. Chronic Rhinosinusitis: T2r38 Genotyping and Nasal Cytology in Primary Ciliary Dyskinesia. *The Laryngoscope* 2022/03/22 ed. 2022; .

37. Chiyonobu K, Xu Y, Feng G, Saso S, Ogawa S, Ikejiri M, Abo M, Kondo M, Gotoh S, Kubo H, Hosoki K, Nagao M, Fujisawa T, Nakatani K, Takeuchi K. Analysis of the clinical features of Japanese patients with primary ciliary dyskinesia. *Auris, nasus, larynx* 2021/08/30 ed. 2022; 49: 248–257.

38. Wolf JS, Papadimitriou JC, Morales RE, Califano JA, Kline NL, Bhatnagar K, Hebert AM, Taylor RJ. The association of active and passive tobacco smoke exposure with chronic rhinosinusitis symptom severity: A cross-sectional study. *Int Forum Allergy Rhinol* 2022; 12: 278–285.



