

A spatio-temporal mapping and bayesian modelling of risk factors of pneumonia symptoms in under-five children in Nigeria.

Atoloye, K. A. ^{*,1}, Lawal, T. V. ¹, Adebowale, A. S. ^{1,2} & Fagbamigbe, A.F. ¹

¹Department of Epidemiology and Medical Statistics, Faculty of Public Health, University of Ibadan, Ibadan, Nigeria

²Population Health and Research Entity, North-West University, Mafikeng, South Africa

*Corresponding author:

Email: kehinde.atoloye@gmail.com (AKA)

Abstract

Background

Pneumonia remains a public health challenge in most parts of the world, with Nigeria having the highest number of pneumonia-related deaths. Understanding the geographical distribution, trends and risk factors associated with pneumonia symptoms will aid an appropriate intervention of pneumonia and subsequently reduce its burden in Nigeria.

Method

This cross-sectional study used data from the 2008, 2013 and 2018 Nigeria Demographic Health Survey. In each of the survey round, a multi-stage cluster sampling technique was used to select the eligible respondents who are women of reproductive age. The outcome variables are the presence of key symptoms of pneumonia: fever, cough, and short rapid breaths. Optimized hotspot analysis was used to identify states with a significantly high prevalence of pneumonia symptoms, MCMC mixed-effect models were fitted to each symptom.

Results

The prevalence of cough was 12.1%, 10.1% and 16.9% in 2008, 2013 and 2018 respectively, 16.2%, 13.3%, and 25.7% for fever; and 41.7%, 42.5% and 6.5% for short rapid breaths respectively with variations across the states. The adjusted odds of having a cough among the children aged 6-11, 12-23, 24-35 and 36-47 months were 95% higher (adjusted odds ratio (aOR)=1.95, 95% Credible Interval (CrI): 1.77, 2.18), 92% higher (aOR=1.92, 95% CrI: 1.73, 2.12), 45% higher (aOR=1.45, 95% CrI: 1.31, 1.62) and 15% higher (aOR=1.15, 95% CrI: 1.03, 1.27) respectively, relative to ages 0-5 months. Similar patterns were noticed for fever and short rapid breaths. Mothers' education was significant for

cough and fever but not for short rapid breaths. Mothers' age was significant only for short rapid breaths at higher odds. Other significantly associated factors with symptoms include residence type, housing quality, wealth index and region.

Conclusion

Fever, cough, and short rapid Breaths are prevalent among under-five children in Nigeria. These symptoms are associated with different characteristics and varied across states in Nigeria. Therefore, it is pertinent that mothers improve on the available preventive and management strategies with the view to mitigating the consequences of pneumonia symptoms among under-five children in Nigeria.

Keywords: Pneumonia symptoms; Under-five children; Cough; Fever; Short Rapid Breadth; Nigeria

Introduction

Pneumonia is an acute respiratory infection in one or both lungs, characterized by cough, fever, shortness of breath, shallow breathing, low energy, etc. [1]. On an annual basis, the disease has accounted for over ten million hospital admissions and more than a million deaths among under-five children globally [2,3]. In 2017, pneumonia was identified as the 4th leading cause of mortality and may eventually become the 3rd by 2024 [4]. According to the World Health Organization (WHO), pneumonia accounts for about 14% of the disease burden globally [2].

In Africa, pneumonia is a potentially life-threatening ailment [2,5,6]. It is not just a major causal factor of mortality and morbidity, but it is as well associated with a non-negligible burden on healthcare infrastructures [7,8] and incomes of households [9]. Oftentimes, pneumonia involves several pathogens that are transmissible among individuals. Analyses have identified various pneumonia results to be

temporally seasonal [10,11]. Paynter et al. found that admissions based on pneumonia are clustered spatially on a high level aided by contact with infected individuals [12].

WHO estimated 868,000 deaths as a result of pneumonia in 2010 with 140,000 deaths on an annual basis among under-five children, the highest in Africa [13]. In Nigeria, pneumonia is one of the leading communicable diseases. Nigeria is one of the countries with the highest burden of pneumonia in childhood and accounts for 162,000 pneumonia-related deaths worldwide in 2018, constituting 20.2% of global deaths [14]. The prevalence of pneumonia in a study conducted in Nigeria in 2013 was 13.3% and 23.9% in another study in 2015 [15,16]. This high prevalence may attract poor health outcomes within the population if unchecked. Therefore, it is of utmost importance to abate the prevalence of pneumonia symptoms in Nigeria to ensure not just healthy living in children but for the future national sustainability as emphasized in the Sustainable Development Goal (SDG)-3, to “Ensure healthy lives and promote well-being for all and sundry and to bring to an end, avoidable deaths of under-five children with all countries aiming to abate neonatal deaths to as low as 12 per 1 000 live births and under-5 deaths to as low as 25 per 1 000 live births’ by 2030” [17].

Several factors have been identified as risk factors for childhood pneumonia, ranging from indoor air pollution due to biomass, outdoor air pollution as a result of vehicular emission and many other factors [18,19]. Previous studies have also documented that the air intake from polluting fuel is an influential risk factor for acute respiratory infection among under-five children in developing countries [20–27].

Other studies have shown that some socioeconomic characteristics like maternal education, region of residence, and household wealth status are important precursors to the risk factors for pneumonia symptoms among children [28–33]. Moreover, studies have also demonstrated that rural-urban residence [34], maternal age at the child’s birth and the child’s birth order (Mishra, 2003) and

water and sanitation facilities [35] are other risk factors for child pneumonia. However, most of these studies focused on the actual disease but not the individual possible symptoms. Addressing issues around the symptoms is more beneficial to the population's health than the disease itself.

It is also important to note that most of these earlier studies used only descriptive and traditional frequentist approaches. Concurrent assessment of the geographical distribution, trends and risk factors associated with pneumonia symptoms remains a grey area for research in Nigeria. Understanding the spatial-temporal distribution, trends and risk factors associated with pneumonia symptoms will aid the prevention of pneumonia and subsequently reduce its prevalence and attendant burden. Therefore, this study employed a Markov Chain Monte Carlo Random Intercept Logistic Regression to identify and explain the risk factors associated with pneumonia symptoms (cough, fever and short rapid breaths). This study makes an original contribution to existing knowledge on pneumonia by providing information on spatial-temporal distribution and drivers of the symptoms of pneumonia among under-five children in Nigeria. Our study outcomes will aid public health efforts toward reducing the incidence of pneumonia in children.

Methods

Study Area

This study was conducted in Nigeria - a West African country with an estimated population of about 215 million. Nigeria consists of six (6) geo-political zones (North-Central, North-East, North-West, South-East, South-South and South-West), which are further divided into 36 states and the Federal Capital Territory (FCT), where each state consists of a third-tier local government and wards. Temperature and humidity are fairly constant all-round year in the south, while the seasons vary considerably in the north. Nigeria's climate is relatively hot and is usually classified into two seasons, dry and wet. The dry season runs from November to March, whereas, the wet season begins in April and ends in October. These seasons come with the symptoms of pneumonia depending on individuals' susceptibility, adaptation, and body reactions to changes in environmental conditions.

Data Collection

The data used in this study were nationally representative and based on the Nigeria Demographic Health Survey (NDHS) conducted in 2008, 2013 and 2018. In the surveys, the unit of inquiry was women of reproductive age (15 to 49 years). They were asked questions regarding their reproductive health, births and children alongside other related topics. The sampling of the respondents during the survey was based on a 2-stage stratified cluster design (National Population Commission [Nigeria] and ICF International, 2014). Data were collected from all eligible women in selected households with the aid of a pretested questionnaire by trained interviewers. Data collected from women about their under-five children were specially processed to constitute the children's recode file which was utilized for analysis in this study. A detailed explanation of the field processes has been reported elsewhere [39–41]

Derivation of Variables

The outcome variables for this study are fever, cough, and short and rapid breaths as symptoms of pneumonia. Each outcome variable measures the presence of the symptoms as “yes” or “no” with value labels “1” or “0” respectively.

Based on existing literature, the explanatory variables were grouped into household, maternal and children characteristics. The maternal background characteristics are age, education, household wealth quintile, place of residence, and geopolitical zone. Children's characteristics include current age in months and sex. The household characteristics included cooking fuel (clean/unclean), drinking water source (improved or unimproved), housing material (improved or unimproved) and toilet facility (improved or unimproved). Clean fuel includes electricity, and liquefied natural gas/biogas and unclean fuel include wood, charcoal, kerosene, straw shrubs, animal dung and grass. The improved sources of drinking water include a protected well, borehole, bottled water and spring rainwater. The unimproved includes a spring tanker, unprotected well with drum, sachet water, surface water, and other sources. The housing material quality derivation was based on a composite score of the type of wall, floor and roof materials. If cement/carpet/rug/ceramic tiles/vinyl asphalt strips were used for the floor, the floor quality is coded 1 else it is coded 0. In the same vein, wall material quality is coded 1 if made of cement blocks/bricks else 0. If the roof material is made of calamine/cement roofing shingles/cement fibres/ceramic tiles/zinc, it is coded 1 else 0. Categories of the housing material were further aggregated and coded as poor (1), average (2) and good (3). For toilet facilities, it was coded as improved or unimproved depending on the type of toilet facility used.

Data Analyses

Descriptive statistics, Bayesian MCMC logistic regression and spatial analyses were used. The spatial variation of the prevalence of the symptoms of pneumonia was analysed using the Optimized Hotspot Analysis. It entails the creation of maps of statistically significant hot and cold spots using the Getis-Ord Gi* statistic employing incident points or weighted features (points or polygons). It evaluates the characteristics of the input feature class to produce optimal results. Optimized Hotspot analysis uses the Getis-Ord Gi* statistic using a fixed distance band in ArcGIS software [42].

The Bayesian MCMC logistic regression model is as specified below:

$$y_{i,j,k} = \mathbf{x}'_{i,j,k}\boldsymbol{\beta} + v_{i,j} + v_i + \varepsilon_{i,j,k}$$

For $k = 1 \dots n_{ij}$ level-1 units that are nested within $j = 1 \dots n_i$ level-2 units that are in turn nested within $i = 1 \dots n$ level-3 units.

where $\mathbf{x}'_{i,j,k}$ is the covariate vector, $\boldsymbol{\beta}$ are unknown regression parameters, v_i is the unknown random effect at level 3, $v_{i,j}$ is the unknown random effect at level-2, and $\varepsilon_{i,j,k}$ are the model residuals that follow a logistic distribution. The model is a random-intercept mixed-effect model. Level 1 is fixed and consists of fixed effect predictors while levels 2 and 3 are the random effect levels defined by the group-level variables. Level 2 group level is defined by the community groups/cluster group variable while the level 3 group is defined by the state group variable. No group-level predictor is involved as interest is more of the amount of variation contributed by the group-level variables. The absence of group-level predictors makes the model a random-intercept mixed model.

In fitting the MCMC model, parameters specified in the *brms* package in R included 4 chains, 4 cores, 4 threads, 50,000 iterations and a thinning rate of 50. A weakly informative prior, derived from the sample was also used.

Results

Distribution of Pneumonia Symptoms and the Demographic Characteristics of the Children and their Mothers.

Distribution of Cough

Table 1 shows the prevalence of cough between 2008 and 2013. The prevalence of cough is highest in 2018 (16.9%) and lowest in 2013 (10.1%). Across the survey years, the prevalence of cough was consistently highest among children aged 6 to 11 months and least among women with no formal education where the prevalence was 9.9%, 7.6%, and 14.7% in 2008, 2013, and 2018 survey year respectively.

Table 1. Prevalence of Cough by Selected Demographic Characteristics, per Year

	2008 (N=22139) n (%)	2013 (N=26076) n (%)	2018 (N=11375) n (%)
Total	2673 (12.1)	2630 (10.1)	1921 (16.9)
Characteristics			
Sex of Child	2.71	0.887	0.052
Male	1319 (11.7)	1350 (10.3)	968 (16.8)
Female	1354 (12.4)	1280 (9.9)	953 (17.0)
Child Age (In Months)	147.532*	268.016*	92.017*
0-5	257 (9.7)	205 (7.3)	164 (13.6)
6-11	399 (15.5)	429 (14.1)	271 (22.0)
12-23	684 (15.3)	763 (14.1)	496 (20.8)
24-35	553 (13.4)	513 (10.5)	360 (16.5)
36-47	424 (9.7)	407 (8.0)	362 (16.6)
48-59	356 (9.0)	313 (6.5)	268 (12.3)
Maternal Age	11.237	19.614*	17.875*
15-19	158 (13.4)	140 (11.3)	69 (15.3)
20-24	524 (12.3)	566 (11.3)	388 (19.2)
25-29	793 (12.5)	718 (9.8)	580 (17.9)
30-34	587 (12.3)	589 (10.2)	414 (15.6)
35-39	370 (11.2)	383 (9.3)	302 (15.6)
40-44	174 (11.0)	166 (8.5)	118 (15.2)
45-49	67 (9.7)	68 (9.4)	50 (17.0)
Maternal Education	99.797*	170.411*	29.459*
No Education	1063 (9.9)	935 (7.6)	646 (14.7)
Primary	702 (13.5)	622 (11.5)	375 (19.8)

Secondary	765 (15.0)	888 (13.0)	725 (17.9)
Higher	143 (12.6)	185 (12.2)	175 (16.8)
Source of Drinking Water	0.008	0.019	1.283
Improved	1372 (12.1)	1561 (10.1)	1153 (16.6)
Unimproved	1301 (12.1)	1069 (10.1)	768 (17.4)
Cooking Fuel Type	23.816*	1.991	2.481
Clean	466 (14.7)	451 (10.7)	361 (15.8)
Unclean	2207 (11.6)	2179 (10.0)	1560 (17.2)
Toilet Facility	0.406	0.441	2.21
Improved	1295 (12.2)	1268 (10.0)	1049 (17.4)
Unimproved	1378 (11.9)	1362 (10.2)	872 (16.3)
Housing Quality Material	30.726*	5.101	14.678
Poor	1084 (10.8)	1087 (9.9)	539 (19.2)
Average	374 (12.2)	392 (9.4)	326 (16.7)
Good	1215 (13.4)	1151 (10.5)	1056 (16.0)
Place of Residence	2.361	8.192*	0.701
Urban	783 (12.6)	921 (10.9)	733 (16.5)
Rural	1890 (11.9)	1709 (9.7)	1188 (17.1)
Wealth Index	36.247*	29.890*	13.817
Poorest	596 (10.9)	498 (8.4)	434 (19.2)
Poorer	560 (10.7)	614 (10.1)	362 (15.9)
Middle	544 (12.6)	605 (11.3)	419 (16.7)
Richer	537 (13.8)	489 (10.1)	408 (17.2)
Richest	436 (13.6)	424 (11.0)	298 (15.3)
Region	567.619*	745.567*	337.491
North Central	353 (8.4)	324 (8.2)	307 (15.3)
North East	817 (15.5)	914 (16.7)	544 (26.7)
North West	346 (6.3)	362 (4.4)	304 (10.9)
South East	408 (20.5)	364 (15.5)	298 (18.6)
South South	477 (19.8)	449 (14.5)	310 (24.7)
South West	272 (9.7)	217 (7.4)	158 (9.5)

***Chi-square values significant at 95% C.I.**

Distribution of Fever

Table 2 also shows that the prevalence of fever dipped between 2008 and 2013, from 16.2% to 13.3% and rose to 25.8% in 2018. Across the survey years, the data show that children in the age group 12-23 months mostly experienced fever more than their counterparts in the other age group while the least prevalence was found among those aged 0 to 5 months. The data further depict that in each survey year, fever's prevalence was persistently higher in children living in households where their source of drinking water was unimproved and those who used unclean fuel. Urban residents also had a lower prevalence of fever in under-five children compared to the rural residents and this pattern was observed across the survey rounds.

Table 2. Prevalence of Fever by Selected Demographic Characteristics of the Children, per Survey Year

	2008 (N=22164) n (%)	2013 (N=26116) n (%)	2018 (N=11374) n (%)
Total	3590 (16.2)	3483 (13.3)	2925 (25.8)
Characteristics			
Sex of Child	4.123*	2.792	0.111
Male	1880 (16.7)	1803 (13.7)	1473 (25.6)
Female	1710 (15.7)	1680 (13.0)	1452 (25.9)
Child Age (In Months)	236.131*	338.382*	149.025*
0-5	247 (9.3)	212 (7.5)	177 (14.7)
6-11	514 (19.9)	541 (17.8)	372 (30.2)
12-23	938 (20.9)	996 (18.4)	757 (31.8)
24-35	743 (18.0)	702 (14.3)	582 (26.6)
36-47	625 (14.3)	575 (11.3)	547 (25.0)
48-59	523 (13.3)	457 (9.5)	490 (22.4)
Maternal Age	10.013	8.509	9.537
15-19	221 (18.7)	180 (14.5)	127 (28.2)
20-24	701 (16.5)	699 (14.0)	561 (27.8)
25-29	981 (15.4)	985 (13.5)	809 (24.9)
30-34	759 (15.9)	739 (12.8)	652 (24.5)
35-39	547 (16.6)	513 (12.5)	504 (26.0)
40-44	262 (16.5)	261 (13.4)	201 (26.0)
45-49	119 (17.1)	106 (14.6)	71 (24.1)

Maternal Education	3.41	5.552	156.361*
No Education	1747 (16.3)	1659 (13.4)	1379 (31.4)
Primary	859 (16.5)	749 (13.9)	503 (26.6)
Secondary	822 (16.1)	899 (13.2)	864 (21.3)
Higher	162 (14.3)	176 (11.6)	179 (17.1)
Source of Drinking Water	19.767*	4.590*	10.145*
Improved	1717 (15.1)	2004 (13.0)	1717 (24.7)
Unimproved	1873 (17.3)	1479 (13.9)	1208 (27.4)
Cooking Fuel Type	7.640*	65.805*	151.739*
Clean	461 (14.5)	399 (9.5)	358 (15.7)
Unclean	3129 (16.5)	3084 (14.1)	2567 (28.2)
Toilet Facility	6.048*	0.778	55.942*
Improved	1651 (15.6)	1677 (13.1)	1378 (22.8)
Unimproved	1939 (16.8)	1806 (13.5)	1547 (29.0)
Housing Quality Material	16.895*	56.614*	197.277*
Poor	1737 (17.3)	1671 (15.2)	946 (33.7)
Average	477 (15.5)	512 (12.3)	596 (30.6)
Good	1376 (15.2)	1300 (11.9)	1383 (20.9)
Place of Residence	42.322*	4.232*	86.975*
Urban	847 (13.6)	1080 (12.7)	929 (20.9)
Rural	2743 (17.2)	2403 (13.6)	1996 (28.8)
Wealth Index	31.415*	64.975*	261.618*
Poorest	957 (17.4)	861 (14.5)	811 (35.8)
Poorer	886 (16.9)	883 (14.5)	668 (29.4)
Middle	721 (16.6)	773 (14.4)	634 (25.2)
Richer	603 (15.5)	580 (11.9)	503 (21.2)
Richest	423 (13.2)	386 (10.0)	309 (15.8)
Region	487.378*	740.839*	380.292*
North Central	396 (9.4)	300 (7.6)	412 (20.5)
North East	1079 (20.5)	1205 (22.0)	737 (36.1)
North West	847 (15.4)	848 (10.2)	885 (31.7)
South East	514 (25.8)	479 (20.4)	371 (23.1)
South South	498 (20.7)	442 (14.3)	329 (26.2)
South West	256 (9.2)	209 (7.1)	191 (11.4)

*Chi-square value significant at 95% C.I.

Distribution of Short Rapid Breaths

The prevalence of short rapid breath (SRB) among under-five children across the three survey rounds was presented in Table 3. The prevalence of SRB was highest in 2008 (41.7%) and least in 2018 (6.5%). There was a consistently high prevalence of SRB among under-five children living in households where unimproved sources of drinking water were used – the prevalence of SRB were 47.3%, 48.9%, and 7.4% in 2003, 2013, and 2018 respectively compared to households with improved sources of drinking water. SRB prevalence in under-five children was consistently higher in rural areas than in urban. In 2008 and 2013 data for instance, compared to under-five children living in the urban areas where the prevalence of SRB was 36.1% and 32.9%, about 44.0% and 47.6% were found in the rural areas respectively.

Table 3. Prevalence of Short Rapid Breaths by Selected Demographic Characteristics, per Year

	2008 (N=2620) n (%)	2013 (N=2605) n (%)	2018 (N=11376) n (%)
Total	1092 (41.7)	1106 (42.5)	741 (6.5)
Characteristics			
Sex of Child	0.042	0.06	0.858
Male	544 (41.9)	572 (42.7)	363 (6.3)
Female	548 (41.5)	534 (42.2)	378 (6.7)
Child Age (In Months)	7.212	12.832*	55.812*
0-5	107 (42.5)	86 (42.2)	67 (5.6)
6-11	165 (42.3)	190 (44.7)	107 (8.7)
12-23	299 (44.6)	334 (44.3)	206 (8.6)
24-35	202 (37.2)	233 (45.6)	138 (6.3)
36-47	177 (42.4)	147 (36.4)	140 (6.4)
48-59	142 (40.8)	116 (37.8)	83 (3.8)
Maternal Age	20.680*	19.719*	7.356
15-19	72 (47.4)	63 (45.3)	28 (6.2)
20-24	254 (49.1)	267 (47.7)	156 (7.7)
25-29	315 (40.8)	290 (41.0)	212 (6.5)
30-34	216 (37.4)	230 (39.2)	169 (6.4)
35-39	138 (38.0)	141 (37.2)	113 (5.8)
40-44	69 (40.1)	78 (47.3)	47 (6.1)

45-49	28 (42.4)	37 (54.4)	16 (5.4)
Maternal Education	129.945*	91.417*	22.122*
No Education	558 (54.0)	495 (53.4)	311 (7.1)
Primary	272 (39.4)	265 (43.3)	153 (8.1)
Secondary	235 (31.2)	295 (33.5)	231 (5.7)
Higher	27 (19.0)	51 (27.6)	46 (4.4)
Source of Drinking Water	32.737*	29.953*	10.361*
Improved	488 (36.3)	589 (38.1)	412 (5.9)
Unimproved	604 (47.3)	517 (48.9)	329 (7.4)
Cooking Fuel Type	77.870*	113.769*	28.153*
Clean	107 (23.3)	89 (19.8)	93 (4.1)
Unclean	985 (45.6)	1017 (47.2)	648 (7.1)
Toilet Facility	17.827*	15.629*	0.723
Improved	474 (37.5)	483 (38.5)	382 (6.3)
Unimproved	618 (45.6)	623 (46.1)	359 (6.7)
Housing Quality Material	123.180*	111.767*	56.336*
Poor	560 (52.6)	557 (51.8)	255 (9.1)
Average	173 (47.5)	196 (50.8)	149 (7.6)
Good	359 (30.1)	353 (30.9)	337 (5.1)
Place of Residence	13.815*	52.518*	13.447*
Urban	277 (36.1)	300 (32.9)	242 (5.5)
Rural	815 (44.0)	806 (47.6)	499 (7.2)
Wealth Index	141.450*	143.625*	57.612*
Poorest	305 (52.2)	263 (53.0)	218 (9.6)
Poorer	296 (54.0)	321 (53.1)	150 (6.6)
Middle	226 (42.5)	276 (46.1)	163 (6.5)
Richer	160 (30.5)	144 (29.6)	127 (5.4)
Richest	105 (24.4)	102 (24.3)	83 (4.2)
Region	228.194*	270.902*	421.444*
North Central	143 (41.4)	149 (46.6)	78 (3.9)
North East	469 (58.7)	553 (61.2)	323 (15.8)
North West	176 (52.9)	128 (35.9)	104 (3.7)
South East	112 (27.7)	132 (36.5)	102 (6.4)
South South	141 (30.0)	117 (26.2)	106 (8.4)
South West	51 (19.0)	27 (12.5)	28 (1.7)

***Chi-square values significant at 95% C.I.**

Optimized Hotspot Analysis of Prevalence of Pneumonia Symptoms

Figs 1-3 show the spatial and temporal distribution of states in Nigeria with a significantly high prevalence of pneumonia symptoms. For Cough, Fig 1 indicates a decrease in states with a significantly high prevalence of cough over time. In 2008, only Gombe has a significantly high prevalence at 90% CI while the other years has no states with significantly high prevalence. Fig 2 indicates an overall decrease in prevalence and states with a significantly high prevalence of fever over time. In 2008, Bayelsa, Rivers, Akwa Ibom, Edo, Anambra, Enugu, Ebonyi and Imo states had significantly high prevalence at 90% CI while in 2013, the south-south states had less significant prevalence which include Delta, Bayelsa and Rivers only at 90% CI. There was no state with a significantly high prevalence of fever in 2018. Fig 3 shows a decrease in the number of states with significantly high prevalence from 2008 to 2018. In 2008, Kaduna, Kano and Jigawa were the states with a significantly high prevalence of short rapid breath. In 2013 and 2018, only Yobe and Taraba has a significantly high prevalence of pneumonia symptoms respectively.

Fig 1. Significance of High Prevalence of Cough in under-five children in Nigeria, 2008-2018

Fig 2. Significance of High Prevalence of Fever in under-five children in Nigeria, 2008-2018

Fig 3. Significance of High Prevalence of Short, Rapid Breaths in under-five children in Nigeria, 2008-2018

Factors Associated with Cough, Fever and Short Rapid Breath

Table 4 presents the results of the adjusted MCMC models fitted for each of the symptoms. It is worthy of note that the factors included in these models were put through bivariate analysis to determine their independent association with the respective symptoms before they were seeded into the multiple models. The adjusted odds of cough reduced by 20% (adjusted odds ratio (aOR) =0.80, 95% Credible Interval (CrI): 0.76-0.85) in 2013 but rose by 52% (aOR =1.52, 95% CrI: 1.40, 1.63) in 2018 compared with the likelihood in 2008. Similarly, the adjusted odds of fever reduced by 24% (aOR =0.76, 95% CrI: 0.72, 0.80) in 2013 and nearly doubled (aOR =1.93, 95% CrI: 1.82, 2.05) in 2018. Whereas the adjusted odds of short rapid breaths were insignificant in 2013, but reduced by 91% (aOR =0.09, 95% CrI: 0.08, 0.11) in 2018 compared with 2008.

Table 4 Results of the Monte Carlo Markov Chain Models for each Symptom of Pneumonia.

		Cough	Fever	Short, Rapid Breaths
		Adjusted OR (95% CrI)	Adjusted OR (95% CrI)	Adjusted OR (95% CrI)
Variables				
Year	2008	Reference		
	2013	0.80(0.76-0.85)*	0.76(0.72-0.80)*	0.96(0.84-1.11)
	2018	1.52(1.40-1.63)*	1.93(1.82-2.05)*	0.09(0.08-0.11)*
Sex of Child	Male	Reference		
	Female	1.02(0.96-1.07)	0.94(0.90-0.99)*	1.01(0.91-1.11)
Child Age (months)	0-5	Reference		
	6-11	1.95(1.77-2.18)*	2.66(2.44-2.97)*	1.35(1.11-1.67)*
	12-23	1.92(1.73-2.12)*	2.86(2.61-3.16)*	1.34(1.12-1.63)*
	24-35	1.45(1.31-1.62)*	2.23(2.01-2.44)*	1.11(0.90-1.34)
	36-47	1.15(1.03-1.27)*	1.75(1.58-1.93)*	1.00(0.82-1.22)
	48-59	0.93(0.84-1.04)	1.51(1.36-1.67)*	0.79(0.64-0.98)*
Mother's Age (years)	15-19	Reference		
	20-24	1.11(0.98-1.27)	1.04(0.93-1.16)	1.42(1.12-1.80)*
	25-29	1.02(0.90-1.15)	1.01(0.90-1.13)	1.16(0.92-1.46)
	30-34	1.00(0.88-1.15)	1.02(0.91-1.15)	1.15(0.91-1.46)
	35-39	0.95(0.84-1.08)	1.03(0.92-1.16)	1.05(0.82-1.36)
	40-44	0.95(0.82-1.11)	1.07(0.93-1.22)	1.25(0.93-1.65)
	45-49	1.06(0.87-1.30)	1.08(0.90-1.28)	1.19(0.82-1.72)

Mother's Education

No Education	Reference		
Primary	1.39(1.28-1.52)*	1.13(1.05-1.22)*	1.13(0.98-1.31)
Secondary	1.35(1.23-1.48)*	1.08(1.00-1.17)	1.01(0.87-1.20)
Higher	1.32(1.15-1.51)*	1.06(0.94-1.21)	0.93(0.71-1.22)

Source of Drinking Water

Improved	Reference		
Unimproved	0.99(0.92-1.06)	0.99(0.94-1.05)	1.06(0.95-1.20)

Cooking Fuel Type

Clean	Reference		
Unclean	0.90(0.80-1.00)	0.95(0.86-1.06)	1.12(0.90-1.40)

Toilet Type

Improved	Reference		
Unimproved	0.98(0.91-1.04)	1.02(0.96-1.08)	0.99(0.88-1.13)

Housing Material Quality

Poor	Reference		
Average	0.99(0.90-1.08)	0.97(0.90-1.05)	0.98(0.82-1.16)
Good	0.89(0.80-0.99)*	1.01(0.92-1.11)	0.76(0.62-0.92)*

Type of Place of Residence

Urban	Reference		
Rural	1.03(0.95-1.12)	1.13(1.05-1.21)*	0.98(0.84-1.14)

Wealth Index

Poorest	Reference		
Poorer	1.06(0.97-1.16)	1.04(0.97-1.13)	1.12(0.94-1.31)
Middle	1.14(1.01-1.30)*	1.03(0.93-1.15)	1.14(0.91-1.42)
Richer	1.14(0.99-1.31)	0.93(0.82-1.06)	0.90(0.68-1.19)
Richest	1.09(0.91-1.31)	0.80(0.68-0.95)*	0.92(0.66-1.28)

Region

North Central	Reference		
North East	2.64(1.62-4.10)*	2.80(1.80-4.53)*	3.16(2.12-4.81)*
North West	0.76(0.49-1.22)	1.63(1.06-2.51)*	1.05(0.70-1.62)
South East	1.93(1.17-3.16)*	2.20(1.35-3.53)*	1.22(0.77-1.92)
South South	1.97(1.25-3.19)*	1.73(1.09-2.75)*	1.13(0.73-1.77)
South West	0.85(0.52-1.35)	0.83(0.51-1.31)	0.43(0.27-0.67)*

Random Effects (Intercepts)

Neighbourhood Level

Variance (CrI 95%)	0.23(0.43-0.52)*	0.21(0.42-0.50)*	0.37(0.53-0.69)*
% Variation Explained (CrI 95%)	0.06(0.03-0.10)*	0.04(0.02-0.07)*	0.03(-0.01-0.06)*

State Level

Variance (CrI 95%)	0.17(0.31-0.55)*	0.16(0.31-0.53)*	0.10(0.22-0.47)*
% Variation Explained (CrI 95%)	0.03(-0.09-0.14)*	0.03(-0.06-0.13)*	0.01(-0.06-0.08)*

OR: Odds Ratio, CrI: Credible Interval

Discussion

This study was designed to identify and explain the Spatio-temporal distribution and risk factors associated with pneumonia symptoms among under-five children in Nigeria using the Markov Chain Monte Carlo Random Intercept Logistic Regression. The study was conceptualized against the backdrop and gap observed in similar studies previously conducted in Nigeria. A temporal pattern was observed in the model. This corroborated the need for spatial analyses of the symptoms by survey years. The likelihood of cough and fever both reduced between 2008 and 2013 but the odds rose by 52% and 93% respectively in 2018. In sharp contrast, the adjusted odds of short rapid breaths were reduced by 91% in 2018 compared with 2013. The temporal pattern observed in the current study corroborates findings from similar studies conducted elsewhere [43,44].

The in-depth spatial hotspot analysis indicated a significant reduction in the prevalence of the symptoms of pneumonia from 2008 to 2018. This decrease may not be unconnected with the various interventional programme and efforts of the government of Nigeria in conjunction with the United Nations and other international partners. For instance, the introduction of the National Pneumonia Control Strategy and Implementation Plan (NPCSIP) by the Nigerian government in the year 2019 towards reducing the high number of childhood mortality due to pneumonia and the menace of pneumonia-related morbidities/deaths in under-five children [45] can explain the reason for our finding. The impact of the government's interventions may be responsible for the statistically significant reduction of these symptoms over the years. In our study, the gender of the child was found to be related to fever. Gender has been said to be a significant epidemiological factor for most diseases [46]. It is important to note that the males also had a higher prevalence of fever as a symptom of pneumonia for all the years under study. Similarly, Victoria et al. found gender to be significantly related to pneumonia but was expressed as the males having a higher odds of experiencing pneumonia, though

the odds of a male having pneumonia than a female in this study was lower [47]. The role of sex hormones in the immunity of individuals may have been responsible for this difference [46].

Children aged 6 to 47 months had higher odds of experiencing cough, fever, or short rapid breaths as symptoms of pneumonia relative to 0 to 5 months children. Previous studies also reported that the odds of contracting pneumonia was higher among children aged 6 to 48 months and lower among children aged 48 to 59 months [31]. This confirms the assertion that younger age corresponds to higher odds of contracting pneumonia in children [7]. Maternal age was significant only for short rapid breaths in mothers aged 20 to 24 where they had higher odds of bringing about short rapid breaths as a pneumonia symptom in their children. This may be due to inexperience in child care or their lack of knowledge or perception of pneumonia. Another study that examined how age affects mothers' perception and knowledge of pneumonia in children found that women aged 18-24 were grossly lacking in knowledge [48]. This poor knowledge may have accounted for mothers aged 20-24 ability to prevent the occurrence of pneumonia (short rapid breaths) in under-five children. Maternal education goes to a large extent in affecting their knowledge of pneumonia and its preventive measures. It is expected higher education among mothers would correspond to lower odds of the occurrence of pneumonia symptoms in their children. However, this may not necessarily be the case, as there may be a difference between knowledge and practice, particularly regarding health outcomes [49].

The sources of drinking water were not found to be significant for the three symptoms of pneumonia in this study contrary to earlier findings [50,51] despite a large-scale lack of quality water in Nigeria [52,53]. Unclean cooking fuel was found not to be significantly associated with cough, fever, and short rapid breaths as symptoms of pneumonia. This is against the expectation of the effect of pollution caused by unclean solid fuels. The findings were corroborated in a study of the household environment and symptoms of childhood acute respiratory tract infections in Nigeria [54].

The role that housing plays in the state of the health of the people cannot be overlooked and overemphasized [55]. Homes that are healthy with enough quality ventilation and insulation guarantees the health of the members of such households, free from pests and contaminant [56]. Findings from this study showed that the odds of cough and short rapid breaths as symptoms of pneumonia were lower among children residing in buildings with good housing material quality. Overcrowding, poor sleeping arrangements, and poor ventilation dominate most housing with poor quality in Nigeria, but findings on toilet facility contrasts previous literature that sanitation and hygiene is a major contribution to childhood pneumonia [57]. A possible basis for the disparity in the odds of the symptoms across the geopolitical regions can be attributed to variations in social and economic development in these regions [54,58,59].

Study Limitations and Strengths

The use of secondary data for this study limits the choice of variables we explored. The data might have suffered recall bias as this study was a cross-sectional study wherein respondents were made to recall past events without any means of verification by the data collectors. Notwithstanding, the data collection procedures and methods adopted by DHS minimized the effect of such bias. The DHS is well-known for its high quality as a result of the data collection approach used by the data originators. The data used is also nationally representative data covering nearly two decades. Also, the pneumonia symptoms available in the DHS dataset are not exhaustive and prevented the measurement of pneumonia from a diagnostic perspective which would have enabled us to have a variable measuring pneumonia as a disease. A major strength of our study is the assessment of factors associated with individual symptoms compared to other studies that studied their composite. The use of Bayesian models in place of frequentist models in understanding the associated risk factors helped to avoid the problems of the inconsistency of results associated with the frequentist approach.

Conclusion

This study provided evidence that variations exist in the prevalence of pneumonia symptoms among under-five children across the states in Nigeria. The northern states were more characterized by *short, rapid breaths* while the southern states were more characterized by *fever* but both regions have a fair share of *cough* among under-five children.

For the cough and fever symptoms, the increased prevalence between 2013 and 2018 indicates that the symptoms need more attention. Although care should be taken in interpreting the significance of a factor to at least one of the symptoms of pneumonia as significance to pneumonia, our findings

supported positive change in pneumonia prevention by providing evidence of risk factors influencing the symptoms of pneumonia. Child's age and region of residence were significant determinants of all pneumonia symptoms considered in the study. While place of residence was a significant predictor of fever, wealth and mother's education was significant for cough and fever.

Recommendations

Government and other relevant stakeholders should do more in the area of providing good housing conditions to its population as this will alleviate the prevalence of pneumonia symptoms in Nigeria. The interventions for pneumonia control towards the United Nation's Millennium Development Goal (MDG) 4 of reducing child mortality by two-thirds should be sustained and improved upon since these strategies have yielded good results. Effective case management and treatment at the community level should be ensured. The high prevalence of pneumonia symptoms in Nigeria as evidenced in this study points to the need for government to improve on the existing sensitization frameworks on home prevention of pneumonia symptoms in under-five children in Nigeria.

Declarations:

Ethics approval and consent to participate

The owners of the secondary data used in this analysis obtained the necessary ethical approval and consent prior to data collection. There was no need for further approvals on our part as open-source secondary data.

Competing interests

338 The authors declared no conflict of interest

339 **Funding**

340 Authors received no funding for this study

341 **Authors' contributions**

342 AKA, LTV and FAF conceptualized and designed the study. AKA and LTV analyzed and interpreted
343 the data while FAF and AAS were senior authors overseeing the analysis and interpretation of the
344 results. AKA and LTV drafted the manuscript FAF and AAS reviewed and edited the manuscript. All
345 authors have read and agreed to the published version of the manuscript.

346 **Acknowledgements**

347 The authors are grateful to ICF Macro, USA, for granting the authors the request to use the
348 Demographic and Health Survey data.

349

References

1. WHO. Pneumonia: the forgotten killer of children. 2006.
2. WHO. World Health Statistics. 2019.
3. Nair H, Simões EAF, Rudan I, Gessner BD, Azziz-Baumgartner E, Zhang JSF, et al. Global and regional burden of hospital admissions for severe acute lower respiratory infections in young children in 2010: A systematic analysis. *Lancet*. 2013;381: 1380–1390. doi:10.1016/S0140-6736(12)61901-1
4. Institute of Health Metrics and Evaluation. Global Burden of Disease Study 2017. *Lancet*. 2017;5: 1–27. Available: http://www.healthdata.org/sites/default/files/files/policy_report/2019/GBD_2017_Booklet.pdf
5. Franco J. Community-acquired Pneumonia. *Radiol Technol*. 2017;88: 621–636.
6. Di Pasquale MF, Sotgiu G, Gramegna A, Radovanovic D, Terraneo S, Reyes LF, et al. Prevalence and etiology of community-acquired pneumonia in immunocompromised patients. *Clin Infect Dis*. 2019;68: 1482–1493. doi:10.1093/cid/ciy723
7. Prina E, Ranzani OT, Torres A. Community-acquired pneumonia. *Lancet*. 2015;386: 1097–1108. doi:10.1016/S0140-6736(15)60733-4
8. Tan KK, Dang DA, Kim KH, Kartasasmita C, Kim HM, Zhang XH, et al. Burden of hospitalized childhood community-acquired pneumonia: A retrospective cross-sectional study in Vietnam, Malaysia, Indonesia and the Republic of Korea. *Hum Vaccines Immunother*. 2018;14: 95–105. doi:10.1080/21645515.2017.1375073
9. Sabin LL, Estrella B, Sempértegui F, Farquhar N, Mesic A, Halim N, et al. Household costs

associated with hospitalization of children with severe pneumonia in Quito, Ecuador. *Am J Trop Med Hyg.* 2020;102: 731–739. doi:10.4269/ajtmh.19-0721

10. Nimbalkar PM, Tripathi NK. Space-time epidemiology and effect of meteorological parameters on influenza-like illness in Phitsanulok, a Northern Province in Thailand. *Geospat Health.* 2016;11: 274–282. doi:10.4081/gh.2016.447

11. Farrar DS, Awasthi S, Fadel SA, Kumar R, Sinha A, Fu SH, et al. Seasonal variation and etiologic inferences of childhood pneumonia and diarrhea mortality in India. *Elife.* 2019;8: 1–23. doi:10.7554/eLife.46202

12. Paynter S, Ware RS, Weinstein P, Williams G, Sly PD. Childhood pneumonia: A neglected, climate-sensitive disease? *Lancet.* 2010;376: 1804–1805. doi:10.1016/S0140-6736(10)62141-1

13. Liu L, Johnson HL, Cousens S, Perin J, Scott S, Lawn JE, et al. Global, regional, and national causes of child mortality: An updated systematic analysis for 2010 with time trends since 2000. *Lancet.* 2012;379: 2151–2161. doi:10.1016/S0140-6736(12)60560-1

14. United Nations Children’s Fund (UNICEF). For every child, reimagine, UNICEF Annual Report. 2019. Available: https://www.unicef.org/sites/default/files/2020-06/UNICEF-annual-report-2019_2.pdf

15. Abdulkarim A, Ibraheem R, Adeboye M. Childhood pneumonia at the University of Ilorin Teaching Hospital, Ilorin Nigeria. *Niger J Paediatr.* 2013;40: 284-289–289.

16. Kuti B. Childhood Community - acquired Pneumonia at the Wesley Guild Hospital , Ilesa : Prevalence , Pattern , and Outcome Determinants. 2015; 98–104. doi:10.4103/1596-4078.182323

17. United Nations. The Sustainable Development Goals Report. 2017; 60.
18. WHO. World health statistics 2018: monitoring health for the SDGs, sustainable development goals. 2018. Available: <http://link.springer.com/10.1007/978-3-319-76887-8>
<http://link.springer.com/10.1007/978-3-319-93594-2>
<http://dx.doi.org/10.1016/B978-0-12-409517-5.00007-2>
<http://dx.doi.org/10.1016/j.jff.2015.06.018>
<http://dx.doi.org/10.1038/s41559-019-0877-3>
19. Smith KR, Pillarisetti A. Household Air Pollution from Solid Cookfuels and Its Effects on Health. Dis Control Priorities, Third Ed (Volume 7) Inj Prev Environ Heal. 2017; 133–152. doi:10.1596/978-1-4648-0522-6_ch7
20. Mishra V. Indoor air pollution from biomass combustion and acute respiratory illness in preschool age children in Zimbabwe. Int J Epidemiol. 2003;32: 847–853. doi:10.1093/ije/dyg240
21. Mishra V, Smith KR, Retherford RD. Effects of Cooking Smoke and Environmental Tobacco Smoke on Acute Respiratory Infections in Young Indian Children. Popul Environ. 2005;26: 375–396. Available: <http://www.jstor.org/stable/27503934>
22. Wichmann J, Voyi KV V. Impact of cooking and heating fuel use on acute respiratory health of preschool children in South Africa. South African J Epidemiol Infect. 2006;21: 48–54. doi:10.1080/10158782.2006.11441264
23. Kilabuko JH, Nakai S. Effects of cooking fuels on acute respiratory infections in children in Tanzania. Int J Environ Res Public Health. 2007;4: 283–288. doi:10.3390/ijerph200704040003

24. Sanbata H, Asfaw A, Kumie A. Association of biomass fuel use with acute respiratory infections among under- five children in a slum urban of Addis Ababa, Ethiopia. BMC Public Health. 2014;14: 1–8. doi:10.1186/1471-2458-14-1122
25. Buchner H, Rehfuess EA. Cooking and season as risk factors for acute lower respiratory infections in African children: A cross-sectional multi-country analysis. PLoS One. 2015;10: 1–21. doi:10.1371/journal.pone.0128933
26. Jean-Daniel LP. Effects of Biomass Fuel on Child Acute Respiratory Infections in Rural Areas of Cameroon and Gabon. Univers J Public Heal. 2016;4: 121–132. doi:10.13189/ujph.2016.040301
27. Capuno JJ, Tan CAR, Javier X. Cooking and coughing: Estimating the effects of clean fuel for cooking on the respiratory health of children in the Philippines. Glob Public Health. 2018;13: 20–34. doi:10.1080/17441692.2016.1202297
28. Azad KMAK. Risk Factors for Acute Respiratory Infections (ARI) Among Under-five Children in Bangladesh. J Sci Res. 2009;1: 72–81. doi:10.3329/jsr.v1i1.1055
29. Dahal GP, Johnson FA, Padmadas SS. Maternal smoking and acute respiratory infection symptoms among young children in nepal: Multilevel analysis. J Biosoc Sci. 2009;41: 747–761. doi:10.1017/S0021932009990113
30. Astale T, Chenault M. Help-seeking behavior for children with Acute respiratory infection in Ethiopia: Results from 2011 Ethiopia Demographic and Health Survey. PLoS One. 2015;10: 1–10. doi:10.1371/journal.pone.0142553
31. Tekle AG, Worku A, Berhane Y. Factors associated with acute respiratory infection in children under the age of 5 years: evidence from the 2011 Ethiopia Demographic and Health

Survey. *Pediatr Heal Med Ther*. 2015; 9. doi:10.2147/phmt.s77915

32. Adesanya OA, Chiao C. A multilevel analysis of lifestyle variations in symptoms of acute respiratory infection among young children under five in Nigeria. *BMC Public Health*. 2016;16: 880.
33. Pinzón-Rondón ÁM, Aguilera-Otalvaro P, Zárate-Ardila C, Hoyos-Martínez A. Acute respiratory infection in children from developing nations: a multi-level study. *Paediatr Int Child Health*. 2016;36: 84–90. doi:10.1179/2046905515Y.0000000021
34. Harerimana JM, Nyirazinyoye L, Thomson DR, Ntaganira J. Social, Economic and environmental risk factors for acute lower respiratory infections among children under fiveyears of age in Rwanda. *Arch Public Heal*. 2016;74: 1–7. doi:10.1186/s13690-016-0132-1
35. Jabessa S. Multilevel Analysis of Acute Respiratory Infection Symptoms among under Five Children in Ethiopia. *J Biom Biostat*. 2015;06. doi:10.4172/2155-6180.1000251
36. National Population Commission(NPC)[Nigeria] & ICF International. Nigeria Demographic and Health Survey, 2008. DHS Measure Macro, New York and Nigeria Population Commission, Abuja, Nigeria; 2009.
37. National Population Commission(NPC)[Nigeria] and ICF. 2019. Nigeria Demographic and Health Survey 2018. Abuja, Nigeria, And Rockville, Maryland, USA: NPC and ICF; 2019.
38. National Population Commission [Nigeria] and ICF International. Nigeria Demographic and Health Survey 2013. Abuja, Nigeria; 2014.
39. National Population Commission (NPC) [Nigeria] and ICF. Nigeria Demographic and Health Survey 2018. Abuja, Nigeria, and Rockville, Maryland, USA: NPC and ICF. 2019. Available: <https://www.dhsprogram.com/pubs/pdf/FR359/FR359.pdf>

40. National Population Commission (NPC) [Nigeria] and ICF. Nigeria Demographic and Health Survey 2013. Abuja, Nigeria, and Rockville, Maryland, USA: NPC and ICF International. 2014; 143–157. doi:10.4324/9780429042669-8
41. National Population Commission (NPC) [Nigeria] and ICF. Nigeria Demographic and Health Survey 2008. Abuja, Nigeria: National Population Commission and ICF Macro. 2009; [19]. Available: <http://measuredhs.com/publications/publication-sr171-summary-reports-key-findings.cfm>
42. Zhuang H, Zhang Y, Wang W, Ren Y, Liu F, Du J, et al. Optimized hot spot analysis for probability of species distribution under different spatial scales based on MaxEnt model: *Manglietia insignis* case. *Biodivers Sci*. 2018;26: 931–940. doi:10.17520/biods.2018059
43. Oloyede I, Ijezie E. Epidemiological and seasonal variations of pneumonia in children admitted in the University of Uyo Teaching Hospital, Uyo, Akwa Ibom State, Nigeria: A retrospective 5-year study (2013- 2017). *Niger J Paediatr*. 2020;47: 42–47.
44. McAllister DA, Liu L, Shi T, Chu Y, Reed C, Burrows J, et al. Global, regional, and national estimates of pneumonia morbidity and mortality in children younger than 5 years between 2000 and 2015: a systematic analysis. *Lancet Glob Heal*. 2019;7: e47–e57. doi:10.1016/S2214-109X(18)30408-X
45. Federal Ministry of Health. National Pneumonia Control Strategy and Implementation Plan. Abuja Nigeria; 2019. Available: https://stopppneumonia.org/wp-content/uploads/2020/02/National_Integrated_Pneumonia_Control_Strategy_Implementation_Plan.pdf
46. Falagas ME, Mourtzoukou EG, Vardakas KZ. Sex differences in the incidence and severity of

respiratory tract infections. *Respir Med.* 2007;101: 1845–1863.

doi:10.1016/j.rmed.2007.04.011

47. Victora CG, Fuchs SC, Flores JAC, Fonseca W, Kirkwood B. Risk Factors of Pneumonia Among Children in a Brazilian Metropolitan Area. 2021;93.

48. Pradhan S, Rao A, Pattanshetty S, Nilima A. Knowledge and perception regarding childhood pneumonia among mothers of under-five children in rural areas of Udupi Taluk, Karnataka: A cross-sectional study. *Indian J Heal Sci.* 2016;9: 35. doi:10.4103/2349-5006.183690

49. Simiyu DE, Wafula EM, Nduati RW. Mothers' knowledge, attitudes and practices regarding acute respiratory infections in children in Baringo District, Kenya. *East Afr Med J.* 2003;80: 303–307. doi:10.4314/eamj.v80i6.8706

50. Rakić A. Water Quality Control in the Water Supply System for the Purpose of Preventing Legionnaires' Disease. *Water Challenges an Urban World.* 2017. doi:10.5772/intechopen.74018

51. George CM, Brooks WA, Graziano JH, Nonyane BAS, Hossain L, Goswami D, et al. Arsenic exposure is associated with pediatric pneumonia in rural Bangladesh: a case control study. *Environ Heal.* 2015;14: 83. doi:10.1186/s12940-015-0069-9

52. Osuchukwu NC, Agba AMO, Emeh JU. Infectious Diseases and the Environment: Baseline Information From Stakeholders in Cross River State, Nigeria. *Int J Res - GRANTHAALAYAH.* 2017;5: 263–281. doi:10.29121/granthaalayah.v5.i1.2017.1889

53. Enebeli UU. The association between water, sanitation and hygiene practices and the occurrence of childhood diarrhoea in Abia state, Nigeria. *Int J Appl Res.* 2019;5: 104–109.

54. Akinyemi JO, Morakinyo OM. Household environment and symptoms of childhood acute

respiratory tract infections in Nigeria , 2003 – 2013 : a decade of progress and stagnation.

2018; 1–12. Available: doi: 10.1186/s12879-018-3207-5

55. Zanzuzdana A, Khan M, Kraemer A. Housing Satisfaction Related to Health and Importance of Services in Urban Slums: Evidence from Dhaka, Bangladesh. Soc Indic Res. 2013;112: 163–185. doi:10.1007/s11205-012-0045-5

56. World Health Organization. World Health Statistics 2010 Indicator compendium Interim version. World Health. 2010; 236.

57. Prüss-Ustün A, Wolf J, Bartram J, Clasen T, Cumming O, Freeman MC, et al. Burden of disease from inadequate water, sanitation and hygiene for selected adverse health outcomes: An updated analysis with a focus on low- and middle-income countries. Int J Hyg Environ Health. 2019;222: 765–777. doi:10.1016/j.ijheh.2019.05.004

58. Kandala N-B, Ji C, Stallard N, Stranges S, Cappuccio FP. Spatial analysis of risk factors for childhood morbidity in Nigeria. Am J Trop Med Hyg. 2007;77: 770–779.

59. Adedini SA, Odimegwu C, Imasiku ENS, Ononokpono DN. Ethnic differentials in under-five mortality in Nigeria. Ethn Health. 2015;20: 145–162. doi:10.1080/13557858.2014.890599

Supporting Information

S1 Table. Distribution of demographic characteristics for cough per year of survey




S2 Table. Distribution of demographic characteristics for fever per year of survey

S3 Table. Distribution of demographic characteristics for short rapid breaths per year of survey


S4 Table. Prevalence of cough by state and year of survey

S5 Table. Prevalence of fever by state and year of survey




525 **S6 Table. Prevalence of short rapid breaths symptoms by state and year of survey**




Prevalence of Cough






Figure




Prevalence of Fever



Figure



Prevalence of Short, Rapid Breaths



Figure