

The influence of acute dietary nitrate supplementation on skeletal muscle fatigue and recovery in older women

Running Title: Nitrate and muscle fatigue and recovery in older women

William S. Zougheb,¹ Richard L. Hoffman,¹ Brandon A. Yates,² Ranjani N. Moorthi,³
Kenneth Lim,³ and Andrew R. Coggan^{1,2}

1. Department of Kinesiology, School of Health & Human Sciences, Indiana University Purdue
University Indianapolis, Indianapolis, IN

2. Indiana Center for Musculoskeletal Health, Indiana University School of Medicine, Indianapolis
IN

3. Division of Nephrology and Hypertension, Department of Medicine, Indiana University School of
Medicine, Indianapolis IN

Corresponding Author: Andrew R. Coggan, PhD, Department of Kinesiology, Indiana University Purdue
University Indianapolis, IF 101C, 250 University Boulevard, Indianapolis, IN 46112; Phone: (317) 274-
0656; Fax: (317) 274-2041; e-mail: acoggan@iupui.edu

ABSTRACT

Older individuals fatigue more rapidly during, and recover more slowly from, dynamic exercise. Women are particularly vulnerable to these deleterious effects of aging, which increases their risk of falling. We have shown that dietary nitrate (NO_3^-), a source of nitric oxide (NO) via the $\text{NO}_3^- \rightarrow \text{nitrite } (\text{NO}_2^-) \rightarrow \text{NO}$ pathway, enhances muscle speed and power in older individuals in the non-fatigued state; however, it is unclear if it reduces fatigability and/or improves recoverability in this population. Using a double-blind, placebo-controlled, crossover design, we studied 18 older (age 70 ± 4 y) women who were administered an acute dose of beetroot juice (BRJ) containing either 15.6 ± 3.6 or <0.05 mmol of NO_3^- . Blood samples were drawn throughout each ~3 h visit for plasma NO_3^- and NO_2^- analysis. Peak torque was measured during, and periodically for 10 min after, 50 maximal knee extensions performed at 3.14 rad/s on an isokinetic dynamometer. Ingestion of NO_3^- -containing BRJ increased plasma NO_3^- and NO_2^- concentrations by 21 ± 8 and 4 ± 4 fold, respectively. However, there were no differences in muscle fatigue or recovery. Dietary NO_3^- increases plasma NO_3^- and NO_2^- concentrations but does not reduce fatigability during or enhance recoverability after high intensity exercise in older women.

29 **KEYWORDS**

30 nitric oxide, beetroot juice, isokinetic dynamometry, aging

31

INTRODUCTION

As humans age, there are detrimental changes in physical function, which are due in large part to changes in the properties of skeletal muscle. These include decreases in the maximal force, speed, and power of muscle contraction [Larsson et al. 2019], as well as reductions in mitochondrial respiratory capacity [Ferri et al. 2020] and in blood flow during exercise [Ridout et al. 2010; Casey et al. 2015]. The latter lead to a greater disturbance of cellular energetics during activity [Coggan et al. 1993; Lewsey et al. 2020], which along with the overall slowing of contractile properties result in increased fatigability during dynamic exercise [Christie et al. 2012], especially at higher movement velocities [Dalton et al. 2010; Callahan and Kent-Braun 2011]. The rate of recovery from fatiguing exercise, i.e., recoverability, is also diminished in older individuals [Schwendner et al. 1997; Yoon et al. 2012]. Since most activities of daily living are intermittent, not continuous, in nature, the latter may be of particular practical and clinical relevance. Women are especially susceptible to the deleterious effects of aging on physical function [Murtaugh and Hubert 2004; Freedman et al. 2016], with increased fatigability and reduced recoverability of muscle predisposing them to falls [Schwendner et al. 1997] and an increased risk hospitalization and even institutionalization.

Dietary nitrate (NO_3^-) supplementation may provide a viable means of offsetting some of these detrimental age-related changes in skeletal muscle function. Upon ingestion and concentration/secretion by the salivary glands, NO_3^- can be reduced by oral bacteria to form nitrite (NO_2^-), which once absorbed can in turn be reduced by, e.g., deoxyhemoglobin to generate nitric oxide (NO) [Lundberg and Weitzberg 2009]. Given the importance of NO in modulating muscle function, including contractility, mitochondrial respiration, blood flow, etc. [Stamler and Meissner 2001], numerous studies in recent years have therefore assessed the effects of dietary NO_3^- (usually in the form of beetroot juice (BRJ)) on exercise capacity. As reviewed elsewhere, such studies have demonstrated that acute or short-term dietary NO_3^- supplementation can improve the inherent contractile properties of muscle (Coggan et al. 2021; Esen et

al. 2022] as well as enhance performance during high intensity dynamic exercise [Silva et al. 2022], at least/especially during open-ended tests performed by non-athletes. Increasing NO bioavailability via NO₃⁻ intake could therefore potentially ameliorate at least some of the negative effects of aging described above. Indeed, we recently found that acute ingestion of BRJ containing 13.4 mmol of NO₃⁻ significantly increased maximal knee extensor speed and power in twelve 71 y old men and women [Coggan et al. 2020a]. However, there were no changes in fatigability during an “all-out”, 50 contraction (~1 min) fatigue test conducted at 3.14 rad/s on an isokinetic dynamometer. The latter finding could be due to the numerous factors likely contributing to muscle fatigue under such conditions (e.g., changes in high energy phosphate and/or H⁺ concentrations, altered Ca²⁺ handling kinetics, impaired excitation-contraction coupling, etc. [Kent-Braun et al. 2012]), only some of which might be influenced by increased NO bioavailability. However, they could also be due to the relatively small sample size, or to the fact that we studied older men as well as women when we have found that women appear to be more likely to benefit from NO₃⁻ supplementation [Coggan et al. 2018b].

Although dietary NO₃⁻ may not (or may) enhance muscle fatigue resistance in older persons, it might still improve the rate of recovery from fatiguing activities. As a potent vasodilator, NO could increase muscle blood flow post-exercise, when intramuscular pressure is no longer elevated by contractions. This might speed the “washout” of fatigue-related metabolites such as H⁺ from muscle [Kemp et al. 1997]. Furthermore, by delivering more O₂, higher post-exercise blood flow could increase the rate of mitochondrial ATP synthesis, thus resulting in more rapid restoration of high energy phosphate levels [Layec et al. 2013] and hence muscle function [Jansson et al. 1990]. In fact, two studies of healthy young men and women found that under hypoxic conditions dietary NO₃⁻ accelerated phosphocreatine (PCr) recovery kinetics following high intensity knee extensor exercise [Vanhatalo et al. 2011, 2014], indicative of increased O₂ delivery via enhanced blood flow and/or improved mitochondrial coupling. However, NO₃⁻ ingestion had no effect on PCr recovery in healthy older men and women studied under

normoxic conditions [Kelly et al. 2013]. Thus, whether dietary NO_3^- supplementation might augment recovery of muscle function following fatiguing exercise in older persons is unclear.

The purpose of the present study was to test the hypothesis that acute dietary NO_3^- supplementation would reduce fatigability and/or improve recoverability of muscle specifically in older women. If so, this could have broad implications for this population of individuals, who are particularly vulnerable to the deleterious effects of reductions in muscle function with aging.

MATERIALS AND METHODS

Participants

A total of 18 community-dwelling women with a mean age, height, weight, and body mass index of 70 ± 4 y, 1.65 ± 0.06 m, 67.1 ± 9.5 kg, and 24.6 ± 3.2 kg/m^2 , respectively, completed this registered clinical trial (NCT03595774). These women were studied after screening a total of 143 potential participants, of whom $n=98$ were deemed ineligible based on a phone screen and review of their medical records, $n=19$ declined to participate or were lost to follow-up, and $n=8$ failed to meet inclusion criteria based on an initial visit to the University Hospital Clinical Research Center (CRC). During this visit, potential participants underwent a health history, physical examination, resting ECG, and phlebotomy (CBC, CMP, fasting insulin, and lipids), and also practiced the isokinetic dynamometry testing protocol (see below).

Individuals were excluded if they were not between the ages of 65 and 79 y, were unable to provide informed consent, were currently smokers, pregnant, or lactating, were taking proton pump inhibitors, antacids, or xanthine oxidoreductase inhibitors, or had a history of major metabolic (thyroid disorders, type I or II diabetes), cardiovascular (e.g., moderate or severe valvular disease, myocardial/pericardial disease, stage II or greater hypertension, heart failure, myocardial infarction/ischemia), renal ($\text{eGFR} < 60$ mL/min/1.73 m^2 , or 61-90 mL/min/1.73 m^2 and albumin:creatinine ratio > 30), neuromuscular (e.g., cervical spondylotic radiculomyelopathy, lumbar spondylosis, amyotrophic lateral sclerosis, Guillain-Barre syndrome, acquired demyelinating polyneuropathies), or

liver (e.g., SGOT/SGPT >2x normal) diseases, were anemic (hematocrit < 30%), or had any other contraindications to vigorous exercise. Other exclusion criteria included whether the participant was on hormone replacement therapy or used phosphodiesterase inhibitors, since these can respectively diminish [Obach et al. 2004] or potentiate [Webb et al. 1999] the effects of dietary NO_3^- . All other participants were included. The Human Subjects Office at Indiana University approved this study and written informed consent was obtained from each individual.

Experimental Design and Protocol

Starting approximately 1 wk after completion of the initial screening visit, eligible participants were studied using a randomized, double-blind, placebo-controlled crossover design. Participants were asked to avoid foods high in NO_3^- (e.g., spinach, arugula, beets) for 24 h and alcohol, caffeine, or food for 12 h before testing. Otherwise, no dietary restrictions were imposed. Upon arrival at the CRC, an intravenous catheter was placed and a baseline blood sample taken for future analysis of plasma NO_3^- and NO_2^- concentrations as described below. The participant then ingested 140 mL of a concentrated beetroot juice (BRJ) supplement (Beet it Sport, James White Drinks, Ipswich, UK) containing (based on direct measurement) either 15.6 ± 3.6 or <0.05 mmol of NO_3^- . Extraction of NO_3^- from BRJ to create the latter placebo was performed by the manufacturer using a highly selective anion exchange resin. After 1 and 2 h of quiet rest, additional blood samples were collected, after which muscle function testing was performed as described below. Ten minutes later, a final blood sample was obtained, after which participants were fed lunch then released. They then returned to the CRC after a 1 wk washout period and repeated the experiment, this time ingesting the opposite BRJ supplement than before. Participants therefore made a total of three study visits, i.e., initial screening, placebo BRJ, NO_3^- -containing BRJ, with the latter two in random order.

Measurement of Muscle Function

Fatigability and recoverability of the knee extensor muscles were assessed using isokinetic dynamometry (Biodex System 4 Pro, Biodex Medical Systems, Shirley, NY). The femoral condyle of the participant's dominant leg was aligned with the dynamometer's axis of rotation while the lower leg, thigh, waist, and torso were securely restrained with straps to prevent any extraneous movement. The participant first performed three maximal knee extensions at each of five velocities (i.e., 0, 1.57, 3.14, 4.71, and 6.28 rad/s), with 120 s of rest between sets, to determine their peak torque-velocity relationship (data reported separately). After an additional 120 s of rest, they then performed 50 maximal knee extensions at a velocity of 3.14 rad/s. The velocity of knee flexion was set to 6.28 rad/s, allowing the participant to freely reposition their leg against minimal resistance between repetitions. Participants were instructed to go "all-out" during this fatigue test, i.e., to extend their leg as quickly and as forcefully as possible during each repetition and to not pace themselves in an attempt to maintain torque. Strong verbal encouragement was provided throughout each test. After completion of the fatigue test, participants performed sets of three additional maximal knee extensions at 3.14 rad/s after 30, 60, 150, 300, and 600 s (nominally) of recovery.

Data from the isokinetic dynamometer were processed by windowing, filtering, and smoothing as previously described [Coggan et al. 2015]. Fatigability was assessed based on the average torque, total work, work during the first and last 1/3rd of the fatigue test, and the ratio of the latter as reported by the manufacturer's software. The torque data were also exported and analyzed to determine the peak torque during each of the 50 contractions. Recoverability was assessed based on the degree of restoration of peak torque at the different time points following the fatigue test. Because it was not possible to measure peak torque at identical times in every participant, a monoexponential function was also fit to the peak torque-recovery duration data (including the final value from the fatigue test) for each individual to determine their time constant (τ) and half-life ($=\ln(2)/\tau$) of torque recovery:

Actual torque (t)

$= \text{Minimum torque}_{\text{predicted}}$

$+ (\text{Maximum torque}_{\text{predicted}} - \text{Minimum torque}_{\text{predicted}}) \times (1 - e^{(-\tau \times t)})$

where t = the exact time after the 50 contraction fatigue test and torque is expressed as a percentage of the maximum torque observed at the start of the fatigue test (which almost always occurred during the 2nd or 3rd of the 50 knee extensions).

Measurement of Plasma NO₃⁻ and NO₂⁻

A dedicated high-performance liquid chromatography system (ENO-30, Eicom USA, San Diego, CA) was used to measure plasma NO₃⁻ and NO₂⁻ concentrations as previously described in detail [Coggan et al. 2018a]. Briefly, 25 µL of thawed plasma was mixed 1:1 with methanol, centrifuged, and then a 10 µL aliquot of the protein-poor supernatant was injected into the HPLC. Plasma NO₃⁻ and NO₂⁻ concentrations were determined from calibration curves generated using NIST-traceable standard solutions.

Statistical Analysis

Data were analyzed using GraphPad Prism version 9.3.1 (GraphPad Software, La Jolla, CA). The normality of data distribution was tested using the D'Agostino-Pearson omnibus test. Plasma NO₃⁻ and NO₂⁻ and peak torque data were compared between trials using two-way (treatment x time or treatment x contraction number) repeated measures ANOVA. Post-hoc testing was performed using the Holm-Šidák multiple comparison procedure. Fatigue test summary statistics (e.g., average torque) and monoexponential recovery curve fit parameters were compared between trials using paired t-tests. Multiplicity-corrected P values less than 0.05 were considered significant. Deidentified data are available from the authors upon request.

RESULTS

Plasma NO_3^- and NO_2^- levels

There were no significant changes in plasma NO_3^- or NO_2^- levels following ingestion of the NO_3^- -depleted BRJ placebo (Table 1). In contrast, plasma NO_3^- and NO_2^- concentrations were significantly elevated 1 and 2 h post-ingestion of NO_3^- -containing BRJ, as well as 10 min after completion of the exercise testing (approximately 2.5 h post-ingestion).

Fatigability

There were no significant differences between placebo and NO_3^- trials in the time required to complete the 50 contraction fatigue test (i.e., 62 ± 6 vs. 64 ± 9 s; $P=0.153$) or in the total time spent extending (33 ± 2 vs. 34 ± 2 s; $P=0.078$) or flexing (29 ± 4 vs. 30 ± 7 s; $P=0.384$) the leg. There were also no significant differences between trials in the average torque, total work, work during the first $1/3^{\text{rd}}$ or last $1/3^{\text{rd}}$ of the test, or their ratio (Table 2). Finally, analysis of the raw data to determine the peak torque during each of the 50 contractions also revealed no significant treatment ($P=0.851$) or interaction ($P=0.963$) effects (Fig. 1).

Recoverability

Peak torque was measured after 35 ± 3 , 64 ± 5 , 159 ± 19 , 309 ± 20 , and 603 ± 4 s of recovery in the placebo trial and after 34 ± 3 , 65 ± 2 , 154 ± 3 , 302 ± 4 , and 603 ± 7 s of recovery in the NO_3^- trial (treatment effect, $P=0.161$; interaction effect, $P=0.239$). No significant differences due to NO_3^- ingestion were observed, regardless of whether torque was expressed relative to body mass (treatment effect, $P=0.326$; interaction effect, $P=0.281$) or as a percentage of the non-fatigued baseline (treatment effect, $P=0.275$; interaction effect, $P=0.500$) (Fig. 2). There were also no differences in the predicted minimum or maximum torque or time course of recovery of torque determined via exponential curve fitting (Table 3).

DISCUSSION

Aging results in more rapid fatigue during, and slower recovery from, dynamic exercise. Older women in particular are prone to these negative effects of aging, which increases their risk of falls. The purpose of

the present study was to determine whether acute dietary NO_3^- intake might ameliorate these age-related changes in muscle function. However, despite producing large changes in plasma NO_3^- and NO_2^- levels, and hence presumably in NO bioavailability, we found no effect of such supplementation on either the fatigability or recoverability of muscle in this population.

In the present study, the ingestion of BRJ containing 15.6 ± 3.6 mmol of NO_3^- increased the plasma concentration 21 ± 8 -fold, or to an individual peak of 783 ± 139 $\mu\text{mol/L}$ above baseline. Assuming essentially complete absorption, this corresponds to a volume of distribution of 20.1 ± 3.3 L, which agrees exactly with our recent novel pharmacokinetic modeling study of the combined $\text{NO}_3^-/\text{NO}_2^-$ system [Coggan et al. 2020b]. Plasma NO_2^- concentrations also rose, albeit to a lesser extent, i.e., 4 ± 4 -fold, due to the larger volume of distribution and almost 70-fold more rapid clearance of the latter and the fact that only 3% of ingested NO_3^- is converted to NO_2^- [Coggan et al. 2020b]. These changes in plasma NO_3^- and NO_2^- are consistent with numerous previous studies of dietary NO_3^- supplementation that have employed a similar dose [e.g., Jonvik et al. 2018]. We did not measure breath NO levels as we have in our previous studies of NO_3^- supplementation in older individuals [Coggan et al. 2020a; Gallardo et al., 2021], but based on such data the latter likely would have increased by at least 100%. It therefore seems reasonable to conclude that the ingestion of NO_3^- -containing BRJ was effective in increasing whole-body NO bioavailability.

Despite the above, acute NO_3^- supplementation has no impact on muscle fatigability during a 50 contraction (~ 60 s) isokinetic knee extensor exercise test. The present results corroborate and extend our prior study of a smaller group of older women and men combined [Coggan et al. 2020a]. Although direct comparisons are challenging due to potential differences between small muscle mass and whole-body exercise, they are also consistent with those of prior studies examining the effects of NO_3^- ingestion on, e.g., 30 s Wingate test performance in younger, healthier individuals [Rimer et al. 2016; Jonvik et al. 2018]. Indeed, a recent meta-analysis by Silva et al. [2022] concluded that dietary NO_3^- is effective at improving performance during dynamic exercise lasting 2-10 min, but not during shorter or longer efforts.

In contrast, Kadach et al. [2023] recently found that NO_3^- supplementation increased muscle torque throughout the first 90 s of a 300 s bout of repeated maximal isometric knee extensions (3 s contraction, 2 s rest). Notably, however, numerous other investigations have failed to demonstrate any effect of dietary NO_3^- on maximal isometric torque [Esen et al. 2022], making the study by Kadach et al. [2023] an exception. Regardless, based on the present as well as our previous results, it seems clear that although acute ingestion of NO_3^- enhances maximal muscle speed and hence power in older individuals [Coggan et al. 2020a; Gallardo et al. 2021], it does not alter fatigability during intense contractile activity in older persons.

A novel aspect of the present investigation was determination of the effects of dietary NO_3^- on recovery of muscle contractile function following fatiguing exercise. Specifically, we hypothesized that NO_3^- supplementation might enhance recovery of muscle torque by magnifying post-exercise hyperemia. Reductions in intramuscular pO_2 and/or pH during exercise should accelerate conversion of $\text{NO}_3^-/\text{NO}_2^-$ to NO [Lundberg and Weitzberg 2009], which in turn could lead to greater vasodilation and higher blood flow following exercise, once intramuscular pressure decreases. If so, this could result in more rapid restoration of muscle pH and/or high energy phosphate levels, and hence muscle function. Indeed, NO_3^- intake has been previously reported to increase PCr recovery kinetics in healthy young men and women studied in hypoxia [Vanhatalo et al. 2011, 2014], but not in younger [Vanhatalo et al. 2014] or older [Kelly et al. 2013] individuals studied in normoxia. Although we did not measure changes in muscle pH and/or high energy phosphate levels in the older women in the present study, acute NO_3^- ingestion had no effect on the recovery of muscle torque following fatiguing exercise. Thus, either any increase in post-exercise blood flow was insufficient to result in more rapid restoration of function, or increased flow simply did not improve recovery, reflecting the likely multifactorial nature of fatigue under the present conditions [Kent-Braun et al., 2012].

There are a number of strengths to the present study, including a relatively large sample size, careful screening of participants to avoid confounding the effects of aging with those due to chronic disease, and the rigorous evaluation of muscle fatigability/recoverability during/after high intensity dynamic exercise. There are, however, a number of limitations as well. Specifically, as indicated above we did not collect any mechanistic measurements (e.g., changes in blood flow, PCr kinetics, etc.) that could have provided further insight into the effects of acute NO_3^- intake on skeletal muscle physiology in this population, even if function was unchanged. It is also possible that we might have obtained different results had we utilized a different dose of NO_3^- , had studied the participants after multiple days of supplementation, and/or had studied men instead of women. Since the effects of aging on muscle fatigability are most evident at higher speeds of movement [], we might also have observed an effect of dietary NO_3^- on fatigability had we tested at a higher velocity of knee extension. Similarly, it is possible that NO_3^- supplementation might enhance recovery from fatigue induced by other types and/or durations/intensities of exercise.

CONCLUSION

In summary, we have assessed the effects of acute NO_3^- intake on maximal isokinetic knee extensor torque during and after high intensity dynamic exercise in older women. Despite resulting in significant elevations in plasma NO_3^- and NO_2^- levels, and hence presumably in NO levels, NO_3^- supplementation had no effect on muscle fatigability or recoverability under these conditions.

REFERENCES

- Callahan, D.M., J.A. Kent-Braun. 2011. Effect of old age on human skeletal muscle force-velocity and fatigue properties. *J. Appl. Physiol.* 111:1345-1352.
- Casey, D.P., S.M. Ranadive, M.J. Joyner. 2015. Aging is associated with altered vasodilator kinetics in dynamically contracting muscle: role of nitric oxide. *J. Appl. Physiol.* 119:232-241.
- Christie, A., E.M. Snook, J.A. Kent-Braun. 2012. Systematic review and meta-analysis of skeletal muscle fatigue in old age. *Med. Sci. Sports Exerc.* 43:568-577.
- Coggan, A.R., A.M. Abduljalil, S.C. Swanson, M.S. Earle, J.W. Farris, L.A. Mendenhall, P.-M. Robitaille. 1993. Muscle metabolism during exercise in young and older untrained and endurance-trained men. *J. Appl. Physiol.* 75:2125-2133.
- Coggan, A.R., M.N. Baranauskas, R.L. Hinrichs, Z. Liu, S.J. Carter. 2022. Effect of dietary nitrate on human muscle power: a systematic review and individual subject data meta-analysis. *J. Int. Soc. Sports Nutr.* 18:66.
- Coggan, A.R., S.R. Broadstreet, K. Mahmood, D. Mikhalkova, M. Madigan, I. Bole, J.L. Leibowitz, et al. 2018a. Dietary nitrate increases VO_2 peak and performance but does not alter ventilation or efficiency in patients with heart failure with reduced ejection fraction. *J. Card. Fail.* 24:65-73.

277 Coggan, A.R., S.R. Broadstreet, D. Mikhalkova, I. Bole, J.L. Leibowitz, A. Kadkhodayan, S. Park, et al. 2018b.
 278 Dietary nitrate-induced increases in human muscle power: high versus low responders. *Physiol. Rep.*
 279 6:e13575.

280 Coggan, A.R., R.L. Hoffman, D.A. Gray, R.N. Moorthi, D.P. Thomas, J.L. Leibowitz, D. Thies D, et al. 2020a.
 281 A single dose dietary of nitrate increases maximal muscle speed and power in healthy older men and
 282 women. *J. Gerontol. A Biol. Sci. Med. Sci.* 75:1154-1160.

283 Coggan, A.R., J.L. Leibowitz, A. Kadkhodayan, D.T. Thomas, S. Ramamurthy, C. Anderson Spearie, S. Waller,
 284 et al. 2015. Effect of acute dietary nitrate intake on knee extensor speed and power in healthy men and
 285 women. *Nitric Oxide* 48:16-21.

286 Coggan, A.R., S.B. Racette, D. Thies, L.R. Peterson, R.E. Stratford Jr. 2020b. Simultaneous pharmacokinetic
 287 analysis of nitrate and its reduced metabolite, nitrite, following ingestion of inorganic nitrate in a mixed
 288 patient population. *Pharm. Res.* 37,235.

289 Dalton, B.H., G.A. Power, A.A. Vandervoort, C.L. Rice. 2010. Power loss is greater in old men than young
 290 men during fast plantar flexion contractions. *J. Appl. Physiol.* 109:1441-1447.

291 Esen, O., N. Dobbin, M.J. Callaghan. 2022. The effect of dietary nitrate on the contractile properties of
 292 human skeletal muscle: A systematic review and meta-analysis. May 23;1-12. doi:
 293 10.1080/07315724.2022.2037475. Online ahead of print.

294 Ferri, E, E. Marzetti, R. Calvani, A. Picca, M. Cesari, B. Arosio. 2020. Role of age-related mitochondrial
295 dysfunction in sarcopenia. 21:5236.

296 Freedman, V.A., D.A. Wolf, B.C. Spillman BC. 2016. Disability-free life expectancy over 30 years: A growing
297 female disadvantage in the US population. Am. J. Public Health. 106:1079-1085.

298 Gallardo, E.J., D.A. Gray, R.L. Hoffman, B.A. Yates, R.N. Moorthi, A.R. Coggan. 2021. Dose-response effect
299 of dietary nitrate on muscle contractility and blood pressure in older subjects: a pilot study. J. Gerontol. A
300 Biol. Sci. Med. Sci. 76:591-598.

301 Jansson, E., G.A. Dudley, B. Norman, and P.A. Tesch. Relationship of recovery from intense exercise to
302 the oxidative potential of skeletal muscle. Acta Physiol. Scand. 139:147-152.

303 Jonvik, K.L., J. Nyakayiru, J.W. van Dijk, K. Masse, S.B. Ballak, J.M.G. Senden, L.J.C. van Loon, L.B. Verdijk.
304 2018. Repeated-sprint performance and plasma responses following beetroot juice supplementation do
305 not differ between recreational, competitive, and elite sprint athletes. Eur. J. Sport Sci. 7:1-10.

306 Kadach, S., J.W. Park, Z. Stoyanov, M.I. Black, A. Vanhatalo, M. Burnley, P.J. Walter, et al. 2023. ¹⁵N-labeled
307 dietary nitrate supplementation increases human skeletal muscle nitrate concentration and improves
308 muscle torque production. Acta Physiol. (Oxf.) Jan6:e13924. Online ahead of print.

309 Kelly, J., J. Fulford, A. Vanhatalo, J.R. Blackwell, O. French, S.J. Bailey, M. Gilchrist, et al. 2013. Effects of
310 short-term dietary nitrate supplementation on blood pressure, O₂ uptake kinetics, and muscle and
311 cognitive function in older adults. Am. J. Physiol. 304:R73-R83.

312 Kemp, G.J., C.H. Thompson, D.J. Taylor, G.K. Radda. 1997. Proton efflux in human skeletal muscle during
313 recovery from exercise. *Eur. J. Appl. Physiol.* 76:462-471.

314 Kent-Braun, J.A., R.H. Fitts, A. Christie. 2012. Skeletal muscle fatigue. *Compr. Physiol.* 2:997-1044.

315 Larsson, L., H. Degens, M. Li, L. Salviati, Y. il Lee, W. Thompson, J.L. Kirkland, M. Sandri. 2019. Sarcopenia:
316 Aging-related loss of muscle mass and function. *Physiol. Rev.* 99: 427–511.

317 Layec, G., L.J. Haseler, J.D. Trinity, C.R. Hart, X. Liu, Y. Le Fur, E.-K. Jeong, R.S. Richardson. 2013.
318 Mitochondrial function and increased convective O₂ transport: implications for the assessment of
319 mitochondrial respiration in vivo. *J. Appl. Physiol.* 115:803-811.

320 Lewsey, S.C., K. Weiss, M. Schär, Y. Zhang, P.A. Bottomley, T.J. Samuel, Q.-L. Xue, et al. 2020. Exercise
321 intolerance and rapid skeletal muscle energetic decline in human age-associated frailty. *JCI Insight* Oct
322 15;5(20):e141246.

323 Lundberg, J.O., E. Weitzberg. 2009. NO generation from inorganic nitrate and nitrite: Role in physiology,
324 nutrition, and therapeutics. *Arch. Pharm. Res.* 32:1119-1126.

325 Murtagh, K.N., H.B. Hubert. 2004. Gender differences in physical disability among an elderly cohort. *Am.*
326 *J. Public Health.* 94:1406-1411.

327 Ridout, S.J., B.A. Parker, S.L. Smithmyer, J.U. Gonzales, K.C. Beck, D.N. Proctor. 2010. Age and sex influence
328 the balance between maximal cardiac output and peripheral vascular reserve. *J. Appl. Physiol.* 108:483-
329 489.

330 Rimer, E. G., L. R. Peterson, A. R. Coggan, and J. C. Martin. 2016. Acute dietary nitrate supplementation
331 increases maximal cycling power in athletes. *Int. J. Sports Physiol. Perform.* 11:715–720.

332 Schwendner, K.I., A.E. Mikesky, W.S. Holt Jr, M. Peacock, D.B. Burr. 1997. Differences in muscle endurance
333 and recovery between fallers and nonfallers, and between young and older women. *J. Geront.* 52:M155-
334 M160.

335 Silva, K.V.C., B.D. Costa, A.C. Gomes, B. Saunders, J.F. Mota. 2022. Factors that moderate the effect of
336 nitrate ingestion on exercise performance in adults: A systematic review with meta-analyses and meta-
337 regressions. *Adv. Nutr.* 13:1866-1881.

338 Stamler, J.S., G. Meissner. 2001. Physiology of nitric oxide in skeletal muscle. *Physiol. Rev.* 81:209-237.

339 Vanhatalo, A., A.M. Jones, J.R. Blackwell, P.G. Winyard, J. Fulford. 2014. Dietary nitrate accelerates
340 postexercise muscle metabolic recovery and O₂ delivery in hypoxia. *J. Appl. Physiol.* 117:1460-1470.

341 Vanhatalo, A., J. Fulford, S.J. Bailey, J.R. Blackwell, P.G. Winyard, A.M. Jones. 2011. Dietary nitrate reduces
342 muscle metabolic perturbation and improves exercise tolerance in hypoxia. *J. Physiol.* 589:5517-5528.

343 Webb, D.J., S. Freestone, M.J. Allen, G.J. Muirhead. 1999. Sildenafil citrate and blood-pressure–lowering
 344 drugs: results of drug interaction studies with an organic nitrate and a calcium antagonist. *Am. J. Cardiol.*
 345 83:21C-28C.

346 Yoon, T., B. Schlinder-Delap, M.L. Keller, S.K. Hunter. 2012. Supraspinal fatigue impedes recovery from a
 347 low-intensity sustained contraction in old adults. *J. Appl. Physiol.* 112:849-858.

348

349 **FUNDING**

350 This work was supported by the Office of the Vice Provost for Research at Indiana University Purdue
351 University Indianapolis and by the National Institutes of Health (grant numbers R21 AG053606 to ARC,
352 K23 DK102824 to RNM, P30 AR072581 to Sharon Moe, and UL1 TR002529 to Anantha Shekhar). The
353 contents of this article are solely the responsibility of the authors and do not necessarily represent the
354 official view of the National Institutes of Health.

355 **DISCLOSURE STATEMENT**

356 The authors have nothing to disclose.

Table 1. Effects of NO₃⁻ supplementation on plasma NO₃⁻ and NO₂⁻ concentrations.

		Time of measurement			
		Pre	1 h post	2 h post	10 min post
		<u>ingestion</u>	<u>ingestion</u>	<u>ingestion</u>	<u>exercise</u>
Plasma [NO ₃ ⁻] (μmol/L)	Placebo	47.7	47.7	51.9	51.1
		±30.3	±23.5	±25.9	±24.4
	Nitrate	42.6	622.9	751.3	735.2
		±17.7	±227.6	±156.9	±158.2
P value		0.312	1.20 x 10 ⁻⁷	3.35 x 10 ⁻¹⁰	5.60 x 10 ⁻¹⁰
Plasma [NO ₂ ⁻] (μmol/L)	Placebo	0.343	0.363	0.387	0.411
		±0.230	±0.286	±0.358	±0.330
	Nitrate	0.376	0.786	0.968	1.209
		±0.286	±0.783	±0.803	±0.883
P value		0.404	2.28 x 10 ⁻²	4.50 x 10 ⁻³	2.45 x 10 ⁻³

Values are mean±S.D. for n=18.

Table 2. Fatigue test summary results.

<u>Trial</u>	Average Torque (Nm/kg)	Total work (J/kg)	Work first 1/3 rd (J/kg)	Work last 1/3 rd (J/kg)	Fatigue index (%)
Placebo	0.57 ±0.10	34.6 ±8.0	15.7 ±8.0	8.0 ±2.1	50.8 ±4.8
Nitrate	0.57 ±0.10	35.3 ±8.2	16.0 ±3.7	8.2 ±2.1	51.4 ±5.8
P value	0.969	0.260	0.204	0.229	0.593

Values are mean±S.D. for n=18.

Table 3. Curve fit parameters for recovery of isokinetic torque at 3.14 rad/s.

<u>Trial</u>	Minimum torque <u>(% of non-fatigued)</u>	Maximum torque <u>(% of non-fatigued)</u>	Time constant <u>(s⁻¹)</u>	Half-life <u>(s)</u>	S.E.E. <u>(%)</u>
Placebo	45.6	97.8	0.01515	72.4	4.3
	±7.9	±8.0	±0.00898	±82.0	±1.2
Nitrate	45.8	97.4	0.01457	58.1	4.7
	±6.2	±5.5	±0.00671	±26.7	±1.6
P value	0.882	0.856	0.785	0.506	0.336

Values are mean±S.D. for n=18.

FIGURE LEGENDS

Figure 1. Knee extensor peak torque during the fatigue test. Note that although 5 repetition averages are presented for clarity, statistical analysis was performed using the peak torque from each of the 50 repetitions. Values are mean±S.D. for n=18. No significant differences were observed.

Figure 2. Recovery of knee extensor peak torque following the fatigue test. Data are expressed either relative to body mass (*top panel*) or as a percentage of the maximum non-fatigued value (*bottom panel*). Values are mean±S.D. for n=18. No significant differences were observed.



