

1 **Hearing loss after exposure to vincristine and platinum-based**
2 **chemotherapy among childhood cancer survivors**

3 Sven Strebel, MSc Pharm.^{1,2,3}; Luzius Mader, PhD¹; Philippa Jörger, MSc HST^{1,3}; Nicolas
4 Waespe, MD, PhD^{1,2,4}; Seraina Uhlmann, BSc Pharm.¹; Nicolas von der Weid, MD⁵;
5 Marc Ansari, MD^{2,6}; Claudia E. Kuehni, MD, MSc^{1,4}✉

6 ¹ Institute of Social and Preventive Medicine, University of Bern, Bern, Switzerland

7 ² CANSEARCH research platform in pediatric oncology and hematology, Department of
8 Pediatrics, Gynecology and Obstetrics, University of Geneva, Geneva, Switzerland

9 ³ Graduate School for Health Sciences, University of Bern, Bern, Switzerland

10 ⁴ Division of Pediatric Hematology/Oncology, Department of Pediatrics, Inselspital, Bern
11 University Hospital, University of Bern, Bern, Switzerland

12 ⁵ Department of Pediatric Oncology and Hematology, University Children's Hospital Basel
13 (UKBB), University of Basel, Basel, Switzerland

14 ⁶ Department of Women, Child and Adolescent, Division of Pediatric Oncology and
15 Hematology, Geneva University Hospital, University of Geneva, Geneva, Switzerland

16 ✉Corresponding author

17 Correspondence to: Claudia E. Kuehni, Prof. MD; Institute of Social and Preventive
18 Medicine, University of Bern, Mittelstrasse 43, 3012 Bern, Switzerland; E-mail:
19 claudia.kuehni@ispm.unibe.ch; Telephone: +41 31 684 35 07

20 ABSTRACT

21 **Background:** Vincristine poses a suspected additional risk factor for hearing loss among
22 childhood cancer survivors (CCS) treated with platinum-based chemotherapy, yet evidence is
23 scarce since no study reports vincristine doses. We examined the association of vincristine
24 with hearing loss in a national cohort of CCS.

25 **Methods:** We included CCS registered in the Swiss Childhood Cancer Registry treated at
26 age ≤ 18 years with platinum-based chemotherapy between 1990–2014. All participants in our
27 retrospective cohort study had audiogram and treatment data from medical records. We
28 identified CCS exposed to vincristine and calculated the total cumulative dose. We defined
29 clinically relevant hearing loss as grade ≥ 2 using the International Society of Pediatric
30 Oncology Boston Ototoxicity Scale at latest follow-up.

31 **Results:** Our study population included 270 CCS (43% female; median age at cancer
32 diagnosis 6.8 years; interquartile range [IQR]: 2.1–11.7 years) with median age at audiogram
33 13.5 years (IQR: 9.3–17.0 years). Vincristine exposure was associated with an increased risk
34 of hearing loss in the multivariable logistic regression analysis (odds ratio [OR] 4.8; 95%
35 confidence interval [CI]: 1.8–12.9). We found no evidence of dose-response relationship (OR
36 1.0; 95% CI: 0.97–1.04) or effect modification from vincristine from other ototoxic
37 treatments, such as type of platinum agent, cranial radiotherapy, and hematopoietic stem cell
38 transplantation.

39 **Conclusion:** Vincristine is associated with a higher risk of hearing loss in CCS treated with
40 platinum-based chemotherapy. We suggest future studies investigate the underlying
41 mechanism and causality among CCS without exposure to other ototoxic cancer treatments.

42 **KEYWORDS:** vinca alkaloids, vincristine, hearing loss, ototoxicity, platinum compounds,
43 cranial radiotherapy, childhood cancer survivors

44 **1. Introduction**

45 Hearing loss is a side effect of platinum-based chemotherapy among children.[1] It is
46 usually irreversible and impairs neurocognitive functioning of childhood cancer survivors
47 (CCS).[2, 3] Several studies identified risk factors for platinum-induced hearing loss such as
48 age at diagnosis, type of platinum agent, total cumulative dose of platinum, concomitant
49 cranial radiotherapy (CRT), and hematopoietic stem cell transplantation (HSCT).[4-7]
50 However, such factors insufficiently explain interindividual variation of platinum-induced
51 hearing loss, possibly involving additional factors, such as vincristine.[4, 8, 9] Vincristine is
52 known as neurotoxic and reported to cause central neuropathy with paralysis of the auditory
53 nerve (cranial nerve VIII).[10, 11] In addition, vincristine-induced toxicity possibly affects
54 the medial olivocochlear bundle and—to a lesser extent—the outer hair cells.[12]

55 Only two studies examined the role of vincristine in platinum-induced hearing loss,
56 highlighting the need for further evidence.[4, 9] Both included vincristine as a co-variable in
57 their analysis but did not further investigate the effect of the total cumulative dose of
58 vincristine and whether there was effect modification between vincristine and other ototoxic
59 cancer treatments. Therefore, we analyzed data from a nationwide cohort of CCS with
60 treatment and audiogram data from medical records to 1) quantify the effect of vincristine on
61 platinum-induced hearing loss; 2) test for interactions between vincristine and other ototoxic
62 cancer treatments; and 3) determine a possible dose-response relationship of vincristine-
63 induced ototoxicity.

64 2. Methods

65 2.1 Study population

66 We analyzed data from CCS treated with platinum-based chemotherapy between 1990–
67 2014 in the nine specialized pediatric oncology clinics in Switzerland.[6] We included CCS 1)
68 registered in the national and population-based Swiss Childhood Cancer Registry
69 (SCCR),[13] 2) treated for cancer at age ≤ 18 years, 3) with no evidence of hearing loss
70 before start of cancer treatment in their medical records, and 4) available audiogram after
71 completing platinum-based chemotherapy.[6] Further details about identifying eligible CCS
72 and data collection are published elsewhere.[6, 14] The Ethics Committee of the Canton of
73 Bern approved the SCCR and the Swiss Childhood Cancer Survivor Study (KEK-BE:
74 166/2014; 2021-01462).[15]

75 2.2 Study procedure

76 We collected clinical information on cancer diagnosis and treatment from medical records
77 and the SCCR in 2015.[6, 14] We extracted clinical variables, including sex (female, male);
78 cancer diagnosis according to the International Classification of Childhood Cancer (3rd
79 edition)[16]; age at cancer diagnosis (< 5 , 5–9, and 10–18 years); period of cancer diagnosis
80 (1990–1995, 1996–2001, 2002–2007, and 2008–2014); type and total cumulative dose
81 (mg/m^2) of platinum agents (cisplatin, carboplatin); CRT (yes, no); and HSCT (yes, no).
82 During initial data collection in 2015, no treatment information about vincristine exposure
83 was collected. We went through all medical records and treatment protocols again in 2022 and
84 calculated vincristine exposure and total cumulative vincristine dose (mg/m^2) as treated or—
85 when unavailable—based on treatment protocol.

86 2.3 Measurement of hearing loss

87 We determined hearing loss at latest follow-up after completion of platinum-based
88 chemotherapy. We evaluated audiograms with the classification system of the International
89 Society of Pediatric Oncology (SIOP) Boston Ototoxicity Scale.[17] Based on the approach of
90 previous studies, we defined clinically relevant hearing loss (yes, no) as SIOP-Boston grade \geq
91 2.[4, 6] In cases of asymmetric hearing, we took the most affected ear as the reference for
92 grading. Further details on the evaluation of the audiograms were previously published.[6]

93 2.4 Statistical analysis

94 First, we compared the prevalence of clinically relevant hearing loss overall and stratified
95 by type of platinum-based chemotherapy between CCS with and without vincristine exposure.
96 We fitted univariable logistic regression models including sex, age at audiogram, age at
97 cancer diagnosis, period of cancer diagnosis, type of platinum-based chemotherapy, exposure
98 to vincristine, concomitant CRT, and HSCT as exposures and clinically relevant hearing loss
99 (yes, no) as outcome.[6] The multivariable regressions included a priori sex, age at most
100 recent audiogram, age at cancer diagnosis, and all other clinical characteristics previously
101 associated with hearing loss at $P < 0.05$ in the univariable models. To examine effect
102 modification of other ototoxic treatments, we included interaction terms between vincristine
103 exposure and type of platinum-based chemotherapy, CRT, and HSCT.[4, 18, 19] We also
104 included the total cumulative dose of cisplatin, carboplatin, and vincristine as a continuous
105 variable in a sub-analysis, which included only CCS who received vincristine, to determine a
106 possible dose-response relationship. We excluded survivors with missing values for the

107 cumulative dose of vincristine or platinum-based chemotherapy from the regression analysis.

108 We calculated global *P*-values using likelihood-ratio tests (LRT).

109 We performed a sensitivity analysis to control for additional ototoxic factors, such as brain
110 surgeries or cerebrospinal fluid shunts, associated with central nervous system (CNS) tumors
111 diagnosis.[1, 20] We included CNS tumor diagnosis (yes, no) as a co-variable in our
112 multivariable regression model.

113 We used Stata version 16.1 (StataCorp LP, Austin, TX, USA) for all analyses.

114 3. Results

115 3.1 Characteristics of study population

116 We included 270 CCS with a median age at cancer diagnosis of 6.8 years (interquartile
117 range [IQR]: 2.1–11.7 years) in our analysis (Table 1). A detailed flow-diagram of the study
118 population was published elsewhere.[6] Median time from cancer diagnosis to most recent
119 audiogram was 5 years (IQR 2.5–8.1 years). The most common cancer diagnoses were CNS
120 tumors (n = 104; 39%). Over half (n = 145; 54%) received vincristine with a median total
121 cumulative dose of 23 mg/m² (IQR 10.1–40.9 mg/m²) (Supplement Figure A.1). CCS
122 previously treated with vincristine were younger at cancer diagnosis, more likely survivors of
123 CNS tumors or neuroblastoma, and received more often carboplatin, CRT, and HSCT than
124 CCS without vincristine exposure.

125 3.2 Role of vincristine in platinum-induced hearing loss

126 Of the included CCS treated with vincristine, 49% (95% CI: 41-57) had hearing loss
127 compared with 22% (95% CI: 15-30) of CCS not treated with vincristine ($p < 0.001$) (Figure
128 1). Stratified by type of platinum-based chemotherapy, we saw the largest difference in
129 prevalence of hearing loss between CCS with and without vincristine exposure among those
130 treated with only cisplatin (68% vs. 21%).

131 Exposure to vincristine was associated with a higher risk for hearing loss in the univariable
132 (odds ratio [OR] 3.5, 95% CI: 2.0–6.0) and multivariable (OR 4.8, 95% CI: 1.8–12.9) logistic
133 regression analyses (Table 2). Using interaction terms, we found no evidence for effect
134 modification between exposure to vincristine and type of platinum-based chemotherapy,

135 CRT, and HSCT (all $P > 0.05$) (Table 3). In a sub-analysis including only CCS with vincristine
136 exposure, we found no evidence for a linear effect of total cumulative vincristine dose on the
137 risk of hearing loss (OR 1.0, 95% CI: 0.97–1.04) (Table 2) (Supplement Figure A.1).

138 In our sensitivity analysis, diagnosis with a CNS tumor (yes, no) was not associated with
139 an increased risk for hearing loss in the multivariable regression (OR 0.8, 95% CI: 0.2–2.7)
140 (Supplement Table A.1). The effect size of vincristine (yes, no) did not change compared with
141 the model without adjustment for CNS tumor diagnosis (OR 5.3, 95% CI: 1.7–17.0 vs. OR
142 4.8, 95% CI: 1.8–12.9).

143 **4. Discussion**

144 In our nationwide cohort study of CCS treated with platinum-based chemotherapy, we found
145 the prevalence of platinum-induced hearing loss higher among CCS with additional
146 vincristine than of CCS without vincristine exposure. Vincristine exposure increased the risk
147 of hearing loss fivefold when adjusting for clinical characteristics. However, we found no
148 evidence of a dose-response relationship of cumulative vincristine dose and the risk for
149 hearing loss, or evidence of effect modification by other ototoxic cancer treatments.

150 Our study is strengthened from analyzing the ototoxicity of vincristine in a large cohort of
151 CCS in terms of overall exposure and considering the total cumulative vincristine dose and
152 potential interactions with other ototoxic cancer treatments. Further strengths include the
153 well-described cohort with clinical data on hearing function, cancer diagnosis, and platinum-
154 based chemotherapy available. However, we cannot conclude whether vincristine is ototoxic
155 independently or only in combination with platinum agents because all included CCS were
156 treated with platinum-based chemotherapy. Furthermore, we had no data on brain surgery or

157 cerebrospinal fluid shunts among survivors of CNS tumors, which have been associated with
158 an increased risk of hearing loss in other studies.[1, 20, 21] Because 70% of CCS in our
159 cohort who received vincristine were diagnosed with CNS tumors and most patients with
160 CNS tumors were exposed to vincristine (97%), we cannot exclude that part of the association
161 of vincristine with hearing loss as possibly explained by other factors associated with the
162 diagnosis of CNS tumors or other concomitant treatments. However, in our sensitivity
163 analysis, CNS tumor diagnosis was not associated with an increased risk of hearing loss and
164 the effect of vincristine remained unchanged after controlling for CNS tumor diagnosis. We
165 also lack data on use of concomitant ototoxic drugs such as aminoglycoside antibiotics and on
166 prophylactic administration of sodium thiosulfate.[1, 22] However, sodium thiosulfate was
167 only recently been approved in the United States for clinical use; it is not yet approved in
168 Switzerland.[23] Another limitation is the retrospective study design and the use of non-
169 standardized routine data from different clinics.

170 Ototoxicity of vincristine was reported mainly in case studies of adult patients with cancer
171 and in two recent cohort studies involving CCS.[4, 9, 11, 24-27] Moke and colleagues
172 examined hearing loss in a large multi-center cohort of 1,481 children, adolescents, and young
173 adults treated with cisplatin.[4] They also collected data on vincristine exposure and reported
174 that vincristine (yes, no) increased the risk for cisplatin-induced hearing loss by a factor of
175 four (OR 3.6) at latest follow-up,[4] which is consistent with our estimate (OR 4.8). However,
176 a direct comparison must be made with caution because Moke and colleagues also included
177 adults in their analysis and the regression models differed in terms of selected co-variables. In
178 another cohort study of 368 pediatric patients with cancer, Meijer and colleagues
179 demonstrated vincristine exposure (hazard ratio 2.9) was an important determinant for the

180 cumulative incidence of cisplatin-induced hearing loss.[9] Neither of the two cohort studies
181 included data on the cumulative dose of vincristine.[4, 9]

182 We did not observe an effect from higher cumulative doses of vincristine on the risk of
183 hearing loss, yet we had no data on administration method, timing of administration with
184 other drugs (particularly platinum agents), pharmacokinetics, drug-drug interactions with
185 azole antifungals, and genetic factors possibly affecting total exposure and toxicity of
186 vincristine.[28-31] Two prospective studies with small sample sizes ($N \leq 23$) examined
187 ototoxicity of vincristine and included dose information.[12, 32] Lugassy and colleagues
188 enrolled adult patients with cancer ($N=23$) in their study and did not observe any deleterious
189 effects from vincristine on hearing among patients with moderate dose exposure (mean total
190 dose of 12 mg).[32] One patient on high-dose vincristine (total dose of 24 mg) developed
191 sensorineural hearing loss during the observation period.[32] Riga and colleagues enrolled
192 pediatric patients diagnosed with leukemia in their study ($N=15$) who had been treated
193 according to the Berlin-Frankfurt-Muenster-95 (BMF-95) protocol. In their study, exposure to
194 low and moderate doses of vincristine ($\leq 1.5 \text{ mg/m}^2$ per dose) did not result in any abnormal
195 changes ($\geq 15 \text{ dB}$) in mean hearing thresholds levels measured by pure tone audiometry at
196 latest follow-up.[12] But they found abnormalities in transient evoked otoacoustic emissions
197 (TEOAE) and a decrease in contralateral suppression after three cycles of vincristine
198 administration, indicating a measurable neurotoxic effect of vincristine on outer hair cells and
199 the medial olivocochlear bundle.[12] However, the study's short observation period of 22 days
200 does not allow prediction of the reversibility of observed effects or subsequent detection of
201 hearing loss.[12]

202 5. Conclusion

203 Our results support emerging evidence for an increased risk of platinum-induced hearing
204 loss with concomitant vincristine administration.[4, 9] Interindividual variability in hearing
205 loss after platinum-based chemotherapy is large and not yet fully understood.[8], which
206 emphasizes the importance of identifying additional risk factors for hearing loss among CCS,
207 such as the neurotoxic treatment vincristine. However, we suggest including TEOAE
208 measurements in further studies to better understand the mechanism and clinical course of
209 vincristine-induced neurotoxicity in the medial olivocochlear bundle and its influence on
210 platinum-induced hearing loss. In addition, we suggest investigating causality of vincristine
211 and hearing loss among CCS without platinum-based chemotherapy or CRT to determine
212 whether vincristine causes permanent hearing loss even without established ototoxic cancer
213 treatments and whether genetic factors possibly play a role.

214 **Clinical trial registration**

215 Not applicable.

216 **Funding source**

217 This study was supported by the Swiss Cancer League (HSR-4951-11-2019), the
218 CANSEARCH Foundation (<https://cansearch.ch/>), and the European Union's Seventh
219 Framework Program for research, technological development and demonstration under grant
220 agreement no. 602030. The EU Commission takes no responsibility for any use made of the
221 information set out. Further funding comes from Swiss Cancer Research (KLS/KFS-4825-01-
222 2019, KFS-5027-02-2020) and Kinderkrebshilfe Schweiz
223 (<https://www.kinderkrebshilfe.ch/de>).

224 **Declaration of competing interest**

225 The authors declare that they have no known competing financial interests or personal
226 relationships that could have appeared to influence the work reported in this paper.

227 **Acknowledgments**

228 We thank the study team of the Childhood Cancer Research Group, Institute of Social and
229 Preventive Medicine, University of Bern, and the Swiss Childhood Cancer Survivor Study,
230 the data managers of the Swiss Pediatric Oncology Group, the team of the Swiss Childhood
231 Cancer Registry, and our collaborators of the PanCareLIFE project. We further thank the
232 research group of the CANSEARCH research platform in pediatric oncology and hematology
233 of the University of Geneva for their inputs and support. We thank Kristin Bivens from the

234 editorial service of the Institute of Social and Preventive Medicine at the University of Bern
235 for her editorial suggestions.

236 **References**

- 237 [1] Landier W. Ototoxicity and cancer therapy. *Cancer*. 2016;122:1647-58.
- 238 [2] Bass JK, Liu W, Banerjee P, Brinkman TM, Mulrooney DA, Gajjar A, et al. Association
239 of Hearing Impairment With Neurocognition in Survivors of Childhood Cancer. *JAMA*
240 *oncology*. 2020;6:1363-71.
- 241 [3] Orgel E, O'Neil SH, Kayser K, Smith B, Softley TL, Sherman-Bien S, et al. Effect of
242 Sensorineural Hearing Loss on Neurocognitive Functioning in Pediatric Brain Tumor
243 Survivors. *Pediatric blood & cancer*. 2016;63:527-34.
- 244 [4] Moke DJ, Luo C, Millstein J, Knight KR, Rassekh SR, Brooks B, et al. Prevalence and
245 risk factors for cisplatin-induced hearing loss in children, adolescents, and young adults: a
246 multi-institutional North American cohort study. *The Lancet Child & adolescent health*.
247 2021;5:274-83.
- 248 [5] Clemens E, de Vries AC, Pluijm SF, Am Zehnhoff-Dinnesen A, Tissing WJ, Loonen JJ, et
249 al. Determinants of ototoxicity in 451 platinum-treated Dutch survivors of childhood cancer:
250 A DCOG late-effects study. *European journal of cancer (Oxford, England : 1990)*.
251 2016;69:77-85.
- 252 [6] Strebel S, Mader L, Sláma T, Waespe N, Weiss A, Parfitt R, et al. Severity of hearing loss
253 after platinum chemotherapy in childhood cancer survivors. *Pediatric blood & cancer*.
254 2022;69:e29755.
- 255 [7] Li Y, Womer RB, Silber JH. Predicting cisplatin ototoxicity in children: the influence of
256 age and the cumulative dose. *European journal of cancer (Oxford, England : 1990)*.
257 2004;40:2445-51.
- 258 [8] Langer T, am Zehnhoff-Dinnesen A, Radtke S, Meitert J, Zolk O. Understanding
259 platinum-induced ototoxicity. *Trends in pharmacological sciences*. 2013;34:458-69.
- 260 [9] Meijer AJM, Li KH, Brooks B, Clemens E, Ross CJ, Rassekh SR, et al. The cumulative
261 incidence of cisplatin-induced hearing loss in young children is higher and develops at an
262 early stage during therapy compared with older children based on 2052 audiological
263 assessments. *Cancer*. 2022;128:169-79.

- 264 [10] Madsen ML, Due H, Ejksjær N, Jensen P, Madsen J, Dybkær K. Aspects of vincristine-
265 induced neuropathy in hematologic malignancies: a systematic review. *Cancer chemotherapy*
266 *and pharmacology*. 2019;84:471-85.
- 267 [11] Mahajan SL, Ikeda Y, Myers TJ, Baldini MG. Acute acoustic nerve palsy associated with
268 vincristine therapy. *Cancer*. 1981;47:2404-6.
- 269 [12] Riga M, Psarommatis I, Korres S, Varvutsi M, Giotakis I, Apostolopoulos N, et al.
270 Neurotoxicity of vincristine on the medial olivocochlear bundle. *International journal of*
271 *pediatric otorhinolaryngology*. 2007;71:63-9.
- 272 [13] Michel G, von der Weid NX, Zwahlen M, Redmond S, Strippoli MP, Kuehni CE.
273 Incidence of childhood cancer in Switzerland: the Swiss Childhood Cancer Registry. *Pediatric*
274 *blood & cancer*. 2008;50:46-51.
- 275 [14] Weiss A, Kuonen R, Brockmeier H, Grotzer M, Candreia C, Maire R, et al. Audiological
276 monitoring in Swiss childhood cancer patients. *Pediatric blood & cancer*. 2018;65.
- 277 [15] Kuehni CE, Rueegg CS, Michel G, Rebholz CE, Strippoli MP, Niggli FK, et al. Cohort
278 profile: the Swiss childhood cancer survivor study. *International journal of epidemiology*.
279 2012;41:1553-64.
- 280 [16] Steliarova-Foucher E, Stiller C, Lacour B, Kaatsch P. International Classification of
281 Childhood Cancer, third edition. *Cancer*. 2005;103:1457-67.
- 282 [17] Brock PR, Knight KR, Freyer DR, Campbell KC, Steyger PS, Blakley BW, et al.
283 Platinum-induced ototoxicity in children: a consensus review on mechanisms, predisposition,
284 and protection, including a new International Society of Pediatric Oncology Boston
285 ototoxicity scale. *Journal of clinical oncology : official journal of the American Society of*
286 *Clinical Oncology*. 2012;30:2408-17.
- 287 [18] Gertson K, Hayashi SS, Trinkaus K, Wan F, Hayashi RJ. Prevalence of Ototoxicity
288 Following Hematopoietic Stem Cell Transplantation in Pediatric Patients. *Biology of blood*
289 *and marrow transplantation : journal of the American Society for Blood and Marrow*
290 *Transplantation*. 2020;26:107-13.
- 291 [19] Bass JK, Hua CH, Huang J, Onar-Thomas A, Ness KK, Jones S, et al. Hearing Loss in
292 Patients Who Received Cranial Radiation Therapy for Childhood Cancer. *Journal of clinical*
293 *oncology : official journal of the American Society of Clinical Oncology*. 2016;34:1248-55.
- 294 [20] Guillaume DJ, Knight K, Marquez C, Kraemer DF, Bardo DM, Neuwelt EA.
295 Cerebrospinal fluid shunting and hearing loss in patients treated for medulloblastoma. *Journal*
296 *of neurosurgery Pediatrics*. 2012;9:421-7.

- 297 [21] Weiss A, Sommer G, Kasteler R, Scheinemann K, Grotzer M, Kompis M, et al. Long-
298 term auditory complications after childhood cancer: A report from the Swiss Childhood
299 Cancer Survivor Study. *Pediatric blood & cancer*. 2017;64:364-73.
- 300 [22] Brock PR, Maibach R, Childs M, Rajput K, Roebuck D, Sullivan MJ, et al. Sodium
301 Thiosulfate for Protection from Cisplatin-Induced Hearing Loss. *The New England journal of*
302 *medicine*. 2018;378:2376-85.
- 303 [23] Brock P, Meijer A, Kogner P, Ansari M, Capra M, Geller J, et al. Sodium thiosulfate as
304 cisplatin otoprotectant in children: The challenge of when to use it. *Pediatric blood & cancer*.
305 2023:e30248.
- 306 [24] Lugassy G, Shapira A. Sensorineural hearing loss associated with vincristine treatment.
307 *Blut*. 1990;61:320-1.
- 308 [25] Aydogdu I, Ozturan O, Kuku I, Kaya E, Sevinc A, Yildiz R. Bilateral transient hearing
309 loss associated with vincristine therapy: case report. *Journal of chemotherapy (Florence,*
310 *Italy)*. 2000;12:530-2.
- 311 [26] Kalcioglu MT, Kuku I, Kaya E, Oncel S, Aydogdu I. Bilateral hearing loss during
312 vincristine therapy: a case report. *Journal of chemotherapy (Florence, Italy)*. 2003;15:290-2.
- 313 [27] Yousif H, Richardson SG, Saunders WA. Partially reversible nerve deafness due to
314 vincristine. *Postgraduate medical journal*. 1990;66:688-9.
- 315 [28] van de Velde ME, Kaspers GL, Abbink FCH, Wilhelm AJ, Ket JCF, van den Berg MH.
316 Vincristine-induced peripheral neuropathy in children with cancer: A systematic review.
317 *Critical reviews in oncology/hematology*. 2017;114:114-30.
- 318 [29] van de Velde ME, Uittenboogaard A, Yang W, Bonten E, Cheng C, Pei D, et al. Genetic
319 Polymorphisms Associated with Vincristine Pharmacokinetics and Vincristine-Induced
320 Peripheral Neuropathy in Pediatric Oncology Patients. *Cancers*. 2022;14.
- 321 [30] van de Velde ME, Kaspers GJL, Abbink FCH, Twisk JWR, van der Sluis IM, van den
322 Bos C, et al. Vincristine-Induced Peripheral Neuropathy in Pediatric Oncology: A
323 Randomized Controlled Trial Comparing Push Injections with One-Hour Infusions (The
324 VINCA Trial). *Cancers*. 2020;12.
- 325 [31] van de Velde ME, Panetta JC, Wilhelm AJ, van den Berg MH, van der Sluis IM, van den
326 Bos C, et al. Population Pharmacokinetics of Vincristine Related to Infusion Duration and
327 Peripheral Neuropathy in Pediatric Oncology Patients. *Cancers*. 2020;12.
- 328 [32] Lugassy G, Shapira A. A prospective cohort study of the effect of vincristine on audition.
329 *Anti-cancer drugs*. 1996;7:525-6.

330

TABLE 1 Demographic and clinical characteristics of study population

	Total cohort N=270		No Vincristine n=125		Vincristine n=145		P-value ^a	
	n	(%)	n	(%)	n	(%)		
Demographic characteristics								
<i>Sex</i>								
Male	154	(57)	66	(53)	88	(61)	0.192	
Female	116	(43)	59	(47)	57	(39)		
<i>Age at most recent audiogram</i>								
< 10 years	81	(30)	32	(26)	49	(34)	0.066	
10–15 years	108	(40)	47	(38)	61	(42)		
> 15 years	81	(30)	46	(37)	35	(24)		
Clinical characteristics								
<i>Age at cancer diagnosis</i>								
< 5 years	112	(41)	42	(34)	70	(48)	<0.001	
5–9 years	67	(25)	24	(19)	43	(30)		
10–18 years	91	(34)	59	(47)	32	(22)		
<i>Period of cancer diagnosis</i>								
1990–1995	48	(18)	24	(19)	24	(17)	0.454	
1996–2001	75	(28)	36	(29)	39	(27)		
2002–2007	82	(30)	32	(26)	50	(34)		
2008–2014	65	(24)	33	(26)	32	(22)		
<i>Diagnosis (ICCC-3)</i>								
III CNS and miscellaneous intracranial and intraspinal neoplasms	104	(39)	3	(2)	101	(70)	<0.001	
IV Neuroblastoma and other peripheral nervous cell tumors	39	(14)	13	(10)	26	(18)		
V Retinoblastoma	5	(2)	2	(2)	3	(2)		
VI Renal tumors	6	(2)	3	(2)	3	(2)		
VII Hepatic tumors	15	(6)	14	(11)	1	(1)		
VIII Malignant Bone tumors	62	(23)	62	(50)	0	(0)		
IX Soft tissue and other extraosseous sarcoma	12	(4)	1	(1)	11	(8)		
X Germ cell tumors, trophoblastic tumors, and neoplasms of gonads	27	(10)	27	(22)	0	(0)		
<i>Treatments^b</i>								
<i>Platinum agent</i>								
Cisplatin	151	(56)	98	(78)	53	(37)	<0.001	
Carboplatin	62	(23)	17	(14)	45	(31)		
Cisplatin and Carboplatin	57	(21)	10	(8)	47	(32)		
<i>Cumulative cisplatin dose categories</i>								
No cisplatin	62	(23)	17	(14)	45	(31)	<0.001	
≤ 300 mg/m ²	51	(19)	11	(9)	40	(28)		
301–450 mg/m ²	71	(26)	38	(30)	33	(23)		
> 450 mg/m ²	86	(32)	59	(47)	27	(19)		
<i>Cumulative carboplatin dose categories</i>								
No carboplatin	151	(56)	98	(78)	53	(37)	<0.001	
< 1500 mg/m ²	29	(11)	3	(2)	26	(18)		
1500–3000 mg/m ²	53	(20)	14	(11)	39	(27)		
> 3000 mg/m ²	32	(12)	6	(5)	26	(18)		
Missing	5	(2)	4	(3)	1	(1)		
<i>Cranial radiation</i>								
No	172	(64)	115	(92)	57	(39)	<0.001	
Yes	98	(36)	10	(8)	88	(61)		
<i>HSCT</i>								
No	244	(90)	120	(96)	124	(86)	0.004	
Yes	26	(10)	5	(4)	21	(14)		

Abbreviations: CNS, central nervous system; ICCC-3, International Classification of Childhood Cancer–Third Edition; HSCT, hematopoietic stem cell transplantation.

^aP-values calculated from chi-square statistics comparing childhood cancer survivors with and without vincristine exposure.

^bEach subject could have had more than one treatment modality.

TABLE 2 Results from univariable and multivariable^a logistic regression analyses

	SIOP-Boston grade ≤ 1 n=172		SIOP-Boston grade ≥ 2 n=98		Univariable n=270		Multivariable ^b n=270		Multivariable ^c n=139	
	n	(%)	n	(%)	OR ^d (95%-CI)	P-value ^e	OR ^d (95%-CI)	P-value ^e	OR ^d (95%-CI)	P-value ^e
<i>Sex</i>						0.778		0.726		0.791
Female	75	(44)	41	(42)	<i>Reference</i>		<i>Reference</i>		<i>Reference</i>	
Male	97	(56)	57	(58)	1.1 (0.7–1.8)		1.1 (0.6–2.2)		1.1 (0.4–2.9)	
<i>Age at audiogram</i>						0.006		<0.001		0.034
< 10 years	63	(37)	18	(18)	<i>Reference</i>		<i>Reference</i>		<i>Reference</i>	
10–15 years	62	(36)	46	(47)	2.6 (1.4–5.0)		7.8 (2.7–22.7)		7.6 (1.4–40.0)	
> 15 years	47	(27)	34	(35)	2.5 (1.3–5.0)		8.0 (2.3–27.2)		4.5 (0.8–25.5)	
<i>Age at cancer diagnosis</i>						0.001		<0.001		0.014
< 5 years	61	(35)	51	(52)	3.0 (1.6–5.5)		9.0 (3.0–27.2)		10.1 (1.8–57.2)	
5–9 years	40	(23)	27	(28)	2.4 (1.2–4.8)		2.4 (0.9–6.1)		1.4 (0.4–5.0)	
10–18 years	71	(41)	20	(20)	<i>Reference</i>		<i>Reference</i>		<i>Reference</i>	
<i>Period of cancer diagnosis</i>						<0.001		0.011		0.037
1990–1995	18	(10)	30	(31)	7.4 (3.1–17.3)		5.4 (1.8–16.1)		7.9 (1.6–39.3)	
1996–2001	46	(27)	29	(30)	2.8 (1.3–6.1)		1.7 (0.6–4.7)		1.3 (0.3–5.4)	
2002–2007	55	(32)	27	(28)	2.2 (1.0–4.7)		1.4 (0.6–3.7)		2.5 (0.7–8.7)	
2008–2014	53	(31)	12	(12)	<i>Reference</i>		<i>Reference</i>		<i>Reference</i>	
<i>Treatments</i>										
<i>Platinum-based chemotherapy</i>						<0.001		<0.001		
Carboplatin only	54	(31)	8	(8)	<i>Reference</i>		<i>Reference</i>		n.a.	
Cisplatin only	94	(55)	57	(58)	4.1 (1.8–9.2)		15.2 (5.0–45.8)		n.a.	
Both platinum agents	24	(14)	33	(34)	9.3 (3.7–23.1)		8.8 (2.8–27.4)		n.a.	
<i>Cumulative cisplatin dose</i>										0.008
Per 100 mg/m ² increase	n.a.	n.a.	n.a.	n.a.	n.a.		n.a.		1.6 (1.1–2.3)	
<i>Cumulative carboplatin dose</i>										0.023
Per 500 mg/m ² increase	n.a.	n.a.	n.a.	n.a.	n.a.		n.a.		0.8 (0.7–1.0)	
<i>Vincristine</i>						<0.001		0.002		
No	98	(57)	27	(28)	<i>Reference</i>		<i>Reference</i>		n.a.	
Yes	74	(43)	71	(72)	3.5 (2.0–6.0)		4.8 (1.8–12.9)		n.a.	
<i>Cumulative vincristine dose</i>										0.871
Per 1 mg/m ² increase	n.a.	n.a.	n.a.	n.a.	n.a.		n.a.		1.0 (0.97–1.04)	
<i>Cranial radiation</i>						0.001		0.286		0.790
No	122	(71)	50	(51)	<i>Reference</i>		<i>Reference</i>		<i>Reference</i>	
Yes	50	(29)	48	(49)	2.3 (1.4–3.9)		1.7 (0.6–4.4)		1.2 (0.3–4.7)	
<i>HSCT</i>						<0.001		0.013		0.006
No	166	(97)	78	(80)	<i>Reference</i>		<i>Reference</i>		<i>Reference</i>	
Yes	6	(3)	20	(20)	7.1 (2.7–18.4)		5.2 (1.4–19.4)		12.1 (1.7–83.6)	

Abbreviations: CI, confidence interval; HSCT, hematopoietic stem cell transplantation; n.a., not applicable; OR, odds ratio; SIOP, International Society of Pediatric Oncology.

^aFactors associated with hearing loss at $P < 0.05$ were included in the multivariable analyses. Sex, age at audiogram, and age at cancer diagnosis were included in the multivariable model independent of the strength of the association in the respective univariable model.

^bMain analysis including vincristine (yes, no) and exposure to platinum-based chemotherapy (cisplatin, carboplatin, both platinum agents) as categorical variables (N=270).

^cSub-analysis including only childhood cancer survivors (CCS) exposed to vincristine with total cumulative doses of chemotherapy as continuous variables (N=139). CCS with missing dose information (n=6) were excluded from analysis.

^dOdds ratio from univariable and multivariable logistic regression models: OR < 1 indicates that SIOP-Boston grade ≤ 1 is more likely. OR > 1 indicates that SIOP-Boston grade ≥ 2 is more likely.

^eGlobal P -value calculated from likelihood-ratio test.

TABLE 3 Interactions between vincristine and other ototoxic cancer treatments

	Multivariable ^a	
	OR ^b (95%-CI)	<i>P</i> -value ^c
Model 1		
<i>Test for interaction between platinum-based chemotherapy and vincristine</i>		
Carboplatin	<i>Reference</i>	0.274
Cisplatin × vincristine exposure	2.8 (0.2–35.2)	
Both platinum agents × vincristine exposure	0.5 (0.03–11.0)	
Model 2		
<i>Test for interaction between CRT and vincristine</i>		
No CRT	<i>Reference</i>	0.200
CRT × vincristine exposure	0.3 (0.04–2.0)	
Model 3		
<i>Test for interaction between HSCT and vincristine</i>		
No HSCT	<i>Reference</i>	0.354
HSCT × vincristine exposure	4.4 (0.18–105.8)	

Abbreviations: CI, confidence interval; CRT, cranial radiotherapy; HSCT, hematopoietic stem cell transplantation; OR, odds ratio.

^aEach interaction between vincristine and the variables platinum-based chemotherapy, CRT, or HSCT was analyzed in a separate multivariable model (Models 1–3) (N=270). Each model was adjusted for sex, age at audiogram, age at diagnosis, period of diagnosis, other ototoxic cancer treatments (i.e. platinum-based chemotherapy, CRT, or HSCT).

^bOdds ratio from univariable and multivariable logistic regression models: OR < 1 indicates that SIOP-Boston grade ≤ 1 is more likely. OR > 1 indicates that SIOP-Boston grade ≥ 2 is more likely.

^cGlobal *P*-value calculated from likelihood-ratio test.




FIGURE 1 Comparison of prevalence of clinically relevant hearing loss between childhood cancer survivors with and without vincristine exposure (N=270). Prevalence is shown for the overall cohort and stratified by type of platinum-based chemotherapy.