1 Title: Disordered balance of T cell subsets in arterial tertiary lymphoid organs in

2 immunoglobulin G4-related vascular disease

3

4 **Author information:**

- 5 Satomi Kasashima, MD, PhD^{1,2,3}; Atsuhiro Kawashima, MD, PhD^{2,3}; Nozomu Kurose, MD,
- 6 PhD^{2,3}; Satoru Ozaki, PhD¹; Fuminori Kasashima, MD, PhD⁴; Yasushi Matsumoto, MD,
- 7 PhD⁴; Hirofumi Takemura, MD, PhD⁵; Hiroko Ikeda, MD, PhD⁶; Ken-ichi Harada, MD,
- 8 PhD^7
- ⁹ ¹Department of Clinical Laboratory Science, Graduate School of Health Science,
- ¹⁰ ⁷Department of Human Pathology, Graduate School of Medicine, Kanazawa University,
- 11 Kanazawa, Japan; ⁵Department of Cardiovascular Surgery, ⁶Department of Pathology,
- 12 Kanazawa University Hospital, Kanazawa, Japan; ²Department of Pathology, ³Department of
- 13 Clinical Laboratory and ⁴Department of Cardiovascular Surgery, National Hospital
- 14 Organization, Kanazawa Medical Center, Kanazawa, Japan
- 15
- 16 Short title: Disordered balance of T cell subsets in IgG4-VD
- 17
- 18 Corresponding author: Satomi Kasashima, M.D., Ph.D.
- 19 Department of Clinical Laboratory Science, Kanazawa University, 5-11-80 Kodatsuno
- 20 Kanazawa 920-0942, Japan Tel: 076-265-2602 Fax: 076-264-4369
- 21 E-mail: sato-kasa@staff.kanazawa-u.ac.jp
- 22 ORCID number 0000-0002-9158-9898
- 23
- 24 **The total word count of the manuscript :** 4173words
- 25

26 Abstract

27	Background: Arterial tertiary lymphoid organs (ATLOs) are ectopic lymphoid structures
28	that control local arterial immune responses and are characterized by germinal centers (GCs).
29	T follicular helper cells (Tfh), resident in GCs, regulate immunoglobulin production and GC
30	development. They consist of Tfh1, Tfh2, and Tfh17 subsets. T follicular regulatory (Tfr)
31	cells possess suppressive functions as Tregs and migrate into GCs as Tfh cells.
32	Immunoglobulin G4 (IgG4)-related diseases are systemic inflammatory fibrous lesions with
33	elevated serum IgG4 and infiltration of IgG4-positive plasmacytes. These manifest in
34	vascular lesions as frequently formed aneurysms with prominent adventitia thickening
35	(IgG4-related abdominal aortic aneurysm; IgG4-AAA). IgG4-AAAs contain several ATLOs.
36	However, the association of IgG4-AAA pathogenesis and ATLOs or ATLOs and immune
37	cell subsets is poorly understood.
38	Methods: We performed whole-slide immunohistochemical image analysis in surgical
39	specimens of IgG4-AAA (n = 21), non-IgG4-related inflammatory AAA (non-IgG4-IAAA, n
40	= 17), atherosclerotic AAA (aAAA, $n = 10$), and Takayasu aortitis (TKA, $n = 5$). We
41	examined the shape of ATLOs, the T cell subsets within and outside ATLOs, and the
42	histological activity of IgG4-relataed disease.
43	Results: IgG4-AAA was characterized by numerous, large, irregular-shaped ATLOs, and
44	higher numbers of Tfr and Tfh2 cells than Tfh1 cells were present compared to patients with
45	other vascular diseases. The morphological abnormalities (in number, area, and form) of
46	ATLOs in IgG4-AAA and the increased number of Tfr cells are closely related to the activity
47	of IgG4-related disease, including the number of IgG4-positive plasmacytes and the degree of
48	adventitial thickness. All T cell subsets (Treg, Th1, Th2, and Th17) were more enriched
49	within ATLOs than outside ATLOs. In particular, an increase in Tfr cells in IgG4-AAA was

- 50 associated with ATLO formation. Increased Tfh17 cells were found in TKA, and aAAA and
- 51 non-IgG4-IAAA were characterized by increased Tfh1 cells.
- 52 **Conclusions:** In the classification of vascular lesions, considering the imbalance in T cell
- 53 subsets, IgG4-AAA should be positioned as adventitial vasculitis with predominant Tfr and
- 54 Tfh2 cells, accompanied by the abnormal appearance of ATLOs.
- 55
- 56 **Keywords:** Immunoglobulin G4-Related disease; IgG4-related abdominal aortic aneurysm;
- 57 Arterial tertiary lymphoid tissues; Germinal center; Follicular helper T cells; Follicular
- 58 regulatory T cells
- 59

60 Nonstandard abbreviations and acronyms

- 61 immunoglobulin G4-related abdominal aortic aneurysm (IgG4-AAA)
- 62 non-immunoglobulin G4-related inflammatory abdominal aortic aneurysm (non-IgG4-IAAA)
- 63 arterial/aortic tertiary lymphoid organs (ATLOs)
- 64 follicular helper T cell (Tfh)
- 65 type 1 helper T cell (Th1)
- 66 type 2 helper T cell (Th2)
- 67 type 17 helper T cell (Th17)
- 68 type 1 follicular helper T cell (Tfh1)
- 69 type 2 follicular helper T cell (Tfh2)
- 70 type 17 follicular helper T cell (Tfh17)
- 71 interfollicular type 1 helper T cell (ifTh1)
- 72 interfollicular type 2 helper T cell (ifTh2)
- 73 interfollicular type 17 helper T cell (ifTh17)
- 74 regulatory T cells (Tregs)

75	follicular	regulatory	T cells ((Tfr)

- 76 interfollicular regulatory T cells (ifTreg)
- 77
- 78 Clinical Perspective:
- 79 What is New?
- •This prospective study uses a new histopathological method for whole slide analysis. We
- 81 characterized IgG4-related vascular disease according to imbalances in T cell subsets
- 82 related to morphological changes of arterial/aortic tertiary lymphoid organs.
- •Compared to Takayasu arteritis and atherosclerotic changes, imbalances in T cell
- 84 subsets—according to the predominance of follicular helper type 2 T cells and follicular
- 85 regulatory T cells in IgG4-related vascular disease—suggested that IgG4-related vascular

86 disease should be defined as adventitial vasculitis.

87

88 What are the clinical implications?

- •This relatively straightforward method for diagnosing vascular diseases based on
- 90 imbalances in specific T cell subsets and arterial/aortic tertiary lymphoid organ changes can
- 91 inform the pathological diagnosis of vasculitis, such as IgG4-related vascular disease and
- 92 Takayasu arteritis.

•The hallmark of an imbalance in T cell subsets can enable serological diagnosis of vasculitis

94 and use of circulating T cells to monitor disease activity.

95

96 Introduction

97 Elevated serum IgG4 levels and chronic fibro-inflammation in affected systemic organs,

- 98 including the salivary glands, pancreas, and retroperitoneum, characterize immunoglobulin
- 99 G4-related disease (IgG4-RD).¹⁻³ Vascular lesions such as inflammatory abdominal aortic

100	aneurysms	(IAAA), a type	of aneurysm	characterized l	by marked	adventitial fibrous
-----	-----------	----------------	-------------	-----------------	-----------	---------------------

- 101 thickening that accounts for 5%–10% of all resected abdominal aortic aneurysms, commonly
- 102 occur in IgG4-RD.^{4,5} About 50% of IAAA cases are represented by IgG4-related abdominal
- 103 aortic aneurysms (IgG4-AAA).^{4,5}
- 104 Tertiary lymphoid organs (TLOs) are ectopic lymphoid structures with morphologic features
- similar to secondary lymphoid organs that allow homing of naïve cells in the T cell area, the
- 106 interface between the T- and B-cell zones, and the germinal center (GC) areas that present
- 107 follicular dendritic cells (FDCs).^{6.7} TLOs are major control sites enabling immune
- 108 surveillance and innate immune responses. ⁶⁻⁸
- 109 Arterial/aortic tertiary lymphoid organs (ATLOs) are TLOs in the media and/or the adventitia
- 110 of the aorta and artery. Resident macrophages, T cells, B cells, mast cells, and dendritic cells

111 in the adventitia have different distributions inside and outside ATLOs.^{9,10} ATLOs have been

- 112 described in autoimmunity, microbial infection, vasculitis, and atherosclerosis.¹¹⁻¹³ ATLOs
- 113 may serve essential functions in IgG4-AAA because the number of ATLOs of IgG4-AAA is
- 114 characteristically increased compared to atherosclerotic AAA (aAAA).^{4,5,14}
- 115 Regarding TLOs/ATLOs, follicular helper T (Tfh) cells, subsets of Th cells expressing
- 116 CXCR5, BCL-6, and IL-21, are primarily located in GCs.¹⁵ Tfh cells control the formation of
- 117 GCs and drive the maturation of long-lived, high-affinity B cells into memory B cells and
- 118 plasmacytes to produce IgG.⁷ Corresponding to Th subsets (type1 Th, Th1; type 2 Th, Th2;
- and type 17 Th, Th17), Tfh cells also consist of at least three subsets (type 1 Tfh, Tfh1; type
- 120 2 Tfh, Tfh2; and type 17 Tfh, Tfh17). ^{15,16} Follicular regulatory T cells (Tfr) are a subset of
- 121 forkhead box P3 (Foxp3)+ regulatory T cells (Tregs) that adopt features of Tfh to enable their
- 122 migration into GC while maintaining their suppressive function as Tregs for immune
- 123 cells.^{17,18} The ratio of Tfh and Tfr cells and the balance of Th1, Th2, and Th17 cells have
- 124 attracted attention in the pathogenesis and progression of several immune diseases.^{19,20}

125	In IgG4-RD, the inflammatory and fibrotic processes are controlled by cytokines secreted
126	mainly by Th2 cells and Tregs. ^{21,22} Additionally, a disrupted balance between Th1 and Th2
127	cells and increased Th2 and Th17 cell counts may contribute to the pathogenesis of
128	IgG4-RD. ²³⁻²⁷ Furthermore, the essential role of Tfh cells in IgG4-RD has attracted research
129	attention. 26,28,29 Akiyama et al. reported that activated peripheral circulating Tfh2 cell counts
130	are high in IgG4-RD. ²³ This change was positively correlated with the IgG4-RD responder
131	index, the number of affected organs along with the disease state, and circulating Tfh1 cells,
132	whereas Tfh17 cells did not affect serum IgG4 levels. ²³ Recent reports on the disordered
133	balance of Tfh cells and Th cells outside the GC showed that it might be closely related to the
134	morphological irregularity of TLOs of IgG4-RD in the salivary gland. ^{30,31}
135	In vascular disease, the association of T cell subsets in ATLOs with the pathogenesis and
136	progression of disease has not been precisely examined. This retrospective study
137	histologically examined Tfh subsets, including Tfh1, Tfh2, and Tfh17 cells, and Tregs using
138	whole-slide analysis in IgG4-AAA compared to aAAA, TKA, non-IgG4-IAAA, and normal
139	aorta.
140	
141	Methods
142	Case selection
143	We selected 38 cases of IAAA diagnosed by histologic dense adventitia with severe
144	sclerosing inflammation and fibrosis (>2 mm thick) among patients undergoing surgery for
145	aneurysm at Kanazawa University Hospital and Kanazawa Medical Center (Kanazawa,
146	Japan) between January 2006 and December 2016. We included 21 cases of IgG4-AAAs
147	based on the comprehensive histologic diagnostic criteria for IgG4-related diseases as
148	follows ² : (1) dense lymphoplasmacytic inflammatory infiltrates with increased numbers of
149	IgG4-positive plasma cells (>50/high-power fields [HPFs; 10 eyepiece, 40 object lenses] and

150	IgG4/IgG ratio of >50%) (Figure 1A–C), (2) storiform pattern of fibrosis, and (3) obliterative
151	vasculitis. The 17 remaining IAAAs were classified into non-IgG4-related IAAAs
152	(non-IgG4-IAAA). No significant differentiation in the radiological findings was observed
153	between IgG4-AAAs and non-IgG4-IAAAs. Ten aAAA cases were selected sequentially
154	between January 2006 and December 2006. The patients with Takayasu arteritis (TKA) were
155	diagnosed according to the American College of Rheumatology 1990 criteria. ³² Five patients
156	with TKA could be examined histopathologically and were included in this study (Figure
157	1D-F). aAAA was defined as aneurysms with severe atherosclerosis in the intima and media,
158	thin adventitia of <1 mm, and minimal inflammatory cell infiltration (Figure 1G–1I). Ten
159	normal control aortas (no or slight atherosclerotic abdominal aorta without dilatation) were
160	obtained from consecutive autopsies performed at Kanazawa Medical Center between
161	January and August 2012.
162	In the non-IgG4-IAAA, TKA, aAAA, and control aorta groups, IgG4-positive plasma cells
163	were scarce, and serum IgG4 was within the normal range. Patients who received steroids
164	within 6 months of surgery or biopsy were excluded from all groups. No patient had a
165	ruptured aneurysm or a medical history of another vasculitis (such as giant cell aortitis and
166	antineutrophil cytoplasmic antibody-related vasculitis). The clinicopathological features of
167	patients with IgG4-AAA, non-IgG4-IAAA, TKA, aAAA, and control aorta are summarized
168	in Table 1. Some of these cases were included in our previous reports. ¹⁴
169	
170	Pathological measurements
171	Immunohistochemistry

172 Immunohistochemistry was performed with a fully automated immunostainer (Roche

173 Diagnostics, Tokyo, Japan; VENTANA BenchMark ULTRA) using antibodies against IgG4

174 (Nichirei Bioscience, Tokyo, Japan; clone HP6025, ×2), CD21 (Leica Biosystems, Wetzlar,

- 175 Germany; clone 2G9, ×100) (Figures 2A, 2D), Foxp3 (Abcam; clone236A/E7, ×100)
- 176 (Figures 2B and 2C), Tbet (Abcam, clone EPR9301, ×200) (Figure 2E), GATA3 (Abcam,
- 177 clone EPR16651, ×400) (Figure 2F), RORγ (Abcam, clone EPR20006, ×200) (Figure 2G),
- and BCL-6 (Abcam, Cambridge, UK; clone EPR11410-43, ×1500). CD21 is a marker of
- 179 FDCs, whereas BCL-6 is a marker of lymphocyte presence in GCs.^{8,33}
- 180
- 181 Image data acquisition
- 182 All histopathological glass slides were scanned and converted into virtual slides using a
- 183 whole-slide scanner at ×40 magnification with a resolution of 0.25 µm/pixels (Nano Zoomer,
- 184 Hamamatsu Photonics, Hamamatsu, Japan).
- 185
- 186 Software
- 187 Whole-slide image analysis was performed using QuPath's interface to the OpenSlide library
- 188 (https://qupath.github.io). ^{34,35}
- 189

190 ATLOs analysis

- 191 Aggregations of FDCs indicated by CD21 immunopositivity in the media and adventitia were
- 192 defined as ATLOs (Figures 1C, 1F, and 1I). Using CD21- and BCL-6-immunostained slides,
- 193 all ATLO regions on each slide were annotated by an experienced pathologist (K.S.) (Figure
- 194 2D). A batch script was subsequently applied to automatically identify the area and
- 195 circumference of each ATLO and the number of ATLOs in the whole-slide images. The
- 196 circularity of each ATLO was defined as $4\pi S/L^2$, wherein S and L are the area and
- 197 circumference, respectively. The circularity of a circle is 1, and decreased circularity of the
- 198 shape refers to the shape's irregularity. The area of the ATLOs was calculated as the ratio of
- 199 the ATLO area to the whole aortic wall area.

200

201 Analysis of T cell subsets

Foxp3, Tbet, GATA3, and ROR γ expression was analyzed using QuPath's 'simple tissue detection' and 'fast cell counts' commands. The number of positive cells and the area were used to calculate the average number of positive cells/mm².

205

206 Definition of T cell subsets

207 In contrast to Tfr and Tfh cells, T cells outside ATLOs in the adventitia were defined as

208 interfollicular T (ifT) cells. ATLO annotations using CD21-immunostained slides were

209 reconstructed to Foxp3-, Tbet-, GATA3-, and RORγ-stained slides, which were continuous

210 section slides. QuPath commands were used to calculate the numbers of Foxp3-, GATA3-,

211 Tbet-, and RORγ-immunopositive cells within the ATLO areas in the whole-slide images.

212 Foxp3-immunopositive cells distributed within ATLOs were defined as Tfr cells, and those

213 outside ATLOs were considered ifTreg cells. Tbet-immunopositive cells distributed within

214 ATLOs were defined as Tfh1 cells, and those outside ATLOs were considered ifTh1 cells.

215 GATA3-immunopositive cells distributed within ATLOs were defined as Tfh2 cells, and

216 those outside ATLOs were considered if Th2 cells. RORy-immunopositive cells distributed

217 within ATLOs were defined as Tfh17 cells, and those outside ATLOs were considered

218 ifTh17 cells. Then, Tfr cells were confirmed by double immunostaining for Foxp3 and

219 BCL-6; Tfh1 cells were confirmed by double immunostaining for Tbet and BCL-6; Tfh2

220 cells were confirmed by double immunostaining for GATA3 and BCL-6; and Tfh17 cells

221 were confirmed by double immunostaining for RORy and BCL-6. After this, the ratios of

222 Tfr/if Treg, Tfh1/ifTh1, Tfh2/ifTh2, and Tfh17/ifTh17 cells were calculated.

223

224 Statistical analysis

225 Kruskal–Wallis analysis of variance was used to compare the groups. Multivariate logistic

226 regression analysis was performed to determine factors that had a significant effect in the

227 univariate analysis and calculate the adjusted odds ratio (OR) and 95% confidence interval

228 (CI). Spearman's correlation coefficients were used to test the associations between

229 continuous variables. All analyses were performed using SPSS v.28 (IBM, Armonk, NY,

230 USA). Statistical significance was indicated by p < 0.05.

231

232 Ethical approval and consent to participate

233 The study was approved by the Human Investigation Review Committee of Kanazawa

234 Medical Center (No. 30-63) and Kanazawa University Hospital (No. 973-1). It conformed to

the principles outlined in the Declaration of Helsinki. The patients provided informed consent

before the study.

237

238 Results

239 ATLOs

ATLOs were rarely observed in the control aorta (10%) but were sometimes seen in aAAA

241 (40%) and TAK cases (60%). However, all IgG4-AAA samples possessed several ATLOs.

242 ATLOs seen in IgG4-AAA comprised a thicker network of FDCs and were large, distorted,

243 numerous, and evenly distributed in the adventitia (Figure 1C). In ATLOs of the IgG4-AAA

244 group, immunoblasts were sparse, the polarization of recognizable light and dark zones was

unclear, and they often collapsed and fused. In contrast, ATLOs seen in the other groups

246 were small and nearly round (Figures 1F and 1I).

247 The median area and median area rate of the ATLOs were significantly higher in the

IgG4-AAA group (2.3 mm² and 1.29%, respectively) than the non-IgG4-IAAA (1.1 mm² and

0.30%, respectively), TKA (0.88 mm² and 0.22\%, respectively), aAAA (0.31 mm² and 0.53\%,

250 respectively), and normal aorta (0.12 mm^2 and 0.03%, respectively) (Figure 3). The median 251 circularity of the ATLOs in the IgG4-AAA group (0.64) was significantly lower than that in 252 the non-IgG4-IAAA (0.71, p = 0.005), TKA (0.95, p = 0.001), aAAA (0.93, p < 0.001), and 253 normal aorta (0.95, p < 0.001) groups. The median number of ATLOs was higher in the 254 IgG4-AAA group (3.7/mm²). However, there was no significant difference between each 255 subgroup. The median number, area, and area rate of ATLOs in the aAAA group were nearly 256 intermediate between those of the normal aorta and non-IgG4-IAAA groups. 257 258 Number of T cells within ATLOs (Tfr, Tfh1, Tfh2, and Tfh17 subsets) in the whole 259 aortic wall 260 The median number of Tfr cells was significantly higher in the IgG4-AAA group $(448.5/\text{mm}^2)$ than in the non-IgG4-IAAA (211.8/mm², p = 0.007), aAAA (1020/mm², p =261 262 0.002), and TKA (11.9/mm², p < 0.001) groups (Figure 4A). The median number of Tfr cells 263 in the TKA group was significantly lower than that in the IgG4-AAA (p = 0.001) and aAAA 264 (p = 0.047) groups. 265 The median number of Tfh2 cells was also significantly higher in the IgG4-AAA group $(182.4/\text{mm}^2)$ than in the non-IgG4-IAAA (115.9/mm², p = 0.031), aAAA (62.0mm², p =266 0.008), and TKA (9.33/mm², p = 0.01) groups. The median number of Tfh2 cells in the TKA 267 268 group was significantly lower than that in the IgG4-AAA (p = 0.004) and aAAA (p = 0.047) 269 groups. 270 The median number of Tfh1 cells was significantly higher in the non-IgG4-IAAA group 271 $(114.3/\text{mm}^2)$, TKA group $(1.11/\text{mm}^2, p = 0.007)$, IgG4-AAA (62.9/mm², p = 0.056), and aAAA (92.4/mm², p = 0.773) groups. The median number of Tfh1 cells in the TKA group 272 273 was significantly lower than that in the IgG4-AAA (p = 0.001) and aAAA (p = 0.047) 274 groups.

- 275 The median number of Tfh17 cells was significantly higher in the TAK group (12.3/mm²)
- 276 (Figure 1E, 1H, 1K) than in the IgG4-AAA ($1.6/\text{mm}^2$, p = 0.016), non-IgG4-IAAA
- 277 ($0.58/\text{mm}^2$, p = 0.007), and aAAA ($2.23/\text{mm}^2$, p = 0.047) groups. The median number of
- 278 Tfh17 cells in the IgG4-AAA group was higher than that in the non-IgG4-IAAA group (p =
- 279 0.042).
- 280

Number of T cells outside ATLOs (ifTreg, ifTh1, ifTh2, and ifTh17 subsets) in the

- whole aortic wall
- 283 The median numbers of ifTreg, ifTh1, and ifTh2 cells in the whole aortic wall were higher in
- the IgG4-AAA group (10.0, 5.6, and 35.1/mm², respectively) than in the other groups,
- whereas the median numbers of ifTh17 cells in all groups were low (less than 1/mm²) with no
- 286 significant differences
- 287

288 The ratio of T cells within to outside ATLOs (Tf/if T)

- 289 In all groups, the numbers of intrafollicular T cell subsets were greater than the
- 290 corresponding number of extrafollicular T cell subsets; the density of each T cell subset
- 291 increased within ATLOs (Figure 4B, 4C). However, the concentration of T cell subsets
- 292 within ATLOs differed in each group of vascular lesions, whereas the ratios of
- intrafollicular/interfollicular T cells of all T cell subsets were equally very low in the controlaorta.
- 295 The median Tfr/ifTreg ratio was significantly and characteristically higher in the IgG4-AAA
- group (44.9) than in the other groups. The median Tfh1/ifTh1 ratio was similarly high in the
- aAAA (37.4) and non-IgG4-IAAA (36.9) and was significantly higher than that in the
- IgG-AAA group (11.2; p = 0.042, p = 0.034, respectively). The median Tfh2/ifTh2 ratio was
- slightly higher in the IgG4-AAA group. However, there were no significant differences in the

- 300 other groups. The median Tf17/ifTh17 ratio was significantly higher in the TKA group (22.0)
- 301 than in the other groups. The median Tf17/ifTh17 ratios in the remaining four groups were
- 302 low (controls, 0; aAAA, 2.8; non-IgG4-IAAA, 0.1; IgG4-AAA, 1.6).
- 303 In the control aorta, the median ratios of intrafollicular/interfollicular T cells were low for all
- 304 T cell subsets (Tfr/ifTreg ratio, 5.6; Tfh1/ifTh1, ratio 3.8; Tfh2/ifTh2, ratio 5.7; Tfh17/ifTh17
- ratio, 0). In the TKA group, the median ratios of Tfr/ifTreg, Tfh1/ifTh1, and Tfh2/ifTh2 (2.7,
- 306 0.2, and 0.7, respectively) cells were the lowest among all groups. The median ratio of
- 307 intrafollicular/interfollicular T cell subsets was similar between the aAAA and
- 308 non-IgG4-IAAA groups, and there were no significant differences (aAAA vs.
- 309 non-IgG4-IAAA, Tfr/ifTreg ratio, 20.8 vs. 27.2; Tfh1/ifTh1 ratio, 37.4 vs. 36.9; Tfh2/ifTh2
- 310 ratio, 10.2 vs. 7.0; Tfh17/ifTh17 ratio, 2.8 vs. 0.1).
- 311

312 Correlations

- 313 Correlations between IgG4-RD pathological parameters and ATLO features
- 314 Among all cases, the number of IgG4+ cells significantly correlated with the average area (R
- 315 = 0.541, p = 0.001), area ratio (R = 0.491, p = 0.003), ATLO irregularity (R = 0.517, p = 0.003)
- 316 0.014), the number of Tfr cells (R = 0.446, p = 0.008), and the Tfr/ifTfr ratio (R = 0.334, p =
- 317 0.002) (Figure 5A). The IgG4/IgG ratio correlated with the average area (R = 0.504, p =
- 318 0.009), area ratio (R = 0.614, p = 0.001), ATLO irregularity (R = 0.562, p = 0.034), number
- 319 of Tfr cells (R = 0.452, p = 0.020), Tfr/ifTfr ratio (R = 0.443, p = 0.021), and Tfh2/ifTfh2
- ratio (R = 0.443, p = 0.002), is reversely correlated with the number of Tfh1 cells (R = -0.394,
- 321 p = 0.047). Adventitial thickness also significantly correlated with the average area (R =
- 322 0.361, p = 0.026), ATLO irregularity (R = 0.517, p = 0.011), number of Tfr cells (R = 0.461,
- 323 p = 0.004), number of Tfh2 cells (R = 0.461, p = 0.042), and Tfr/ifTfr ratio (R = 0.389, p = 0.004)
- 324 0.021).

325	Among IgG4-AAA cases	, the number of I	gG4+ cells	significantly	v correlated	l with the area	l
		,	50. 	Digititite without	,		-

- ratio (R = 0.270, p = 0.048), ATLO irregularity (R = 0.338, p = 0.030), and the number of
- 327 Tfh2 cells (R = 0.304, p = 0.034), is reversely correlated with the number of Tfh1 cells (R =
- -0.458, p = 0.037) and Tfh1/ifTfh1 ratio (R = -0.343, p = 0.003). The IgG4/IgG ratio
- 329 correlated with ATLO irregularity (R = 0.346, p = 0.045) is reversely correlated with the
- number of Tfh1 cells (R = -0.458, p = 0.040). Adventitial thickness also significantly
- 331 correlated with the average area (R = 0.277, p = 0.048) and ATLO irregularity (R = 0.291, p
- 332 = 0.044), is reversely correlated with the number of Tfh1 cells (R = -0.526, p = 0.036) and the
- 333 Tfh1/ifTfh1 ratio (R = -0.328, p = 0.032).
- 334 The ATLO irregularity in IgG4-AAA cases was significantly associated with the average
- 335 area (R = 0.442, p = 0.021), area ratio (R = 0.432, p = 0.023), and the number of ATLOs (R =
- 0.331, p = 0.034), whereas ATLO irregularity in all cases was not associated with these
- 337 factors.
- 338

339 Correlations between ATLO features and T cell subsets

- 340 Among all cases, the average area and area ratio of ATLOs were significantly correlated with
- 341 the number of Tfr cells (R = 0.343, p = 0.030; R = 0.304, p = 0.046, respectively) and the
- 342 Tfr/ifTreg ratio (R = 0.372, p = 0.028; R = 0.302, p = 0.048, respectively) (Figure 5B). ATLO
- 343 irregularity was significantly correlated with the number of Tfr cells (R = 0.638, p < 0.001),
- 344 Tfr/ifTreg ratio (R = 0.472, p = 0.003), and the number of Tfh2 cells (R = 0.580, p < 0.001).
- 345 Among IgG4-AAA cases, the average area (R = 0.349, p = 0.043), areal ratio (R = 0.386, p =
- 346 0.044), and ATLO irregularity (R = 0.473, p = 0.022) significantly correlated with the
- number of Tfr cells (R = 0.495, p = 0.043). The area ratio was significantly related to the
- 348 number of Tfh2 cells (R = 0.499, p = 0.027).

349

Compared with aAAA, IgG4-AAA was associated with ATLO irregularity, with an adjusted

OR of 16.61 (95% CI; 12.34–2.6132, p = 0.001), the area ratio of ATLOs, with an adjusted

350 Multivariable logistic regression

351

352

353	OR of 6.43 (95% CI; 0.345–0.781, $p = 0.11$); the number of Tfr cells, with an adjusted OR of
354	2.55 (95% CI; 0.894–0.984, $p = 0.01$); and the Tfr/ifTreg ratio, with an adjusted OR of 2.01
355	(95% CI; 0.789–0.967, <i>p</i> = 0.023).
356	Compared with the other IgG4-AAA cases, IgG4-AAA was significantly associated with
357	ATLO irregularity, with an adjusted OR of 18.95 (95% CI; 231.8–1.28, $p = 0.006$) and the
358	number of Tfr cells, with an adjusted OR of 5.21 (95% CI; 1.99–3.55, $p = 0.008$).
359	
360	Discussion
361	Recently, several reports have indicated that a disrupted balance of T cell subsets contributes
362	to the systemic and vascular manifestations of vascular disease. Th1 and Th17 cytokine
363	responses correlate with TAK activity. ³⁶ An increase in serum IL-6 characterizes TAK, and
364	IL-6 is a key cytokine involved in the differentiation of Th17 cells and regulates the balance
365	between Th17 and Treg cells. Gao et al. reported that abnormal T cell maturation into
366	Th2-like Tregs and decreased circulating mature Tregs could contribute to TKA
367	pathogenesis. ³⁷ Our findings demonstrate that the number of Tfh17 cells was higher in TKA
368	than in other vascular groups. The increase in Th17 cells in TAK might be due to the higher
369	number of Tf17 cells.
370	Flow cytometry-based peripheral blood analysis revealed that uncontrolled inflammation
371	involving Tfh and Tfr cells underlies IgG4-RD etiology. ³⁸ A disrupted balance between Th1
372	and Th2 cells and increased Th2 and Th17 cell counts may contribute to IgG4-RD
373	pathogenesis. ^{15,23-28} Grados et al. reported that IgG4-RD is characterized by a shift in
374	circulating T cells toward Th2/Tfh2 and Th17/Tfh17 polarization. ³⁹ However, previous

15

375 studies of circulating T cells via flow cytometry have not evaluated intrafollicular T cells

- 376 because Tfr and Tfh cells present in GCs have different phenotypes from circulating Tfr and
- 377 Tfh cells, which are primarily defined by CXCR5 positivity.²⁰
- 378 Using whole-slide analysis, this is the first report to describe the T cell subsets in IgG4-AAA,
- 379 particularly regarding ATLOs. In IgG4-AAA, the number of Tfr, ifTreg, Tfh2, and ifTh2
- 380 cells were significantly high; therefore, all Treg and Th2 cells were also more numerous than
- 381 other groups. Moreover, the Tfr/ifTreg ratio was considerably higher than that of the other
- 382 groups. The Tfr cell count and the Tfr/ifTreg ratio were closely related to IgG4-AAA activity
- 383 (adventitia thickness, IgG4+ cell count, and IgG4/IgG ratio). Thus, the increase in Treg cell
- 384 concentrations within ATLOs was closely associated with the pathogenesis and progression
- 385 of IgG4-AAA and is a novel pathological hallmark of IgG4-AAA. Regarding T cell subsets,
- the Tfh2 cell number increase and Tfh1 cell number decrease might assist in the pathological
- 387 diagnosis of IgG4-AAA. Reliable pathological hallmarks of IgG4-AAA are needed given that
- 388 diagnosis of IgG4-RD using serum IgG4 levels or tissue IgG4+plasmacytes is challenging.
- 389 The number of circulating Tfr cells or the Tfr/Treg ratio, not just Treg cells, should be
- 390 assessed to diagnose or estimate the activity of IgG4-AAA using flow cytometry.
- 391 Considering that the interaction of Tfr and Tfh cells regulates the production of IgG within
- 392 the GC, controlling Tfr cells creates opportunities to develop novel therapeutic approaches in
- immune disease^{40,41}. Our findings indicate that Tfr therapy might be a promising therapeutic
- 394 method for IgG4-AAA in the future.
- 395 ATLOs are rarely present in the normal aorta, and their presence suggests that there is a
- 396 vascular lesion. Previous reports indicated that ATLOs may organize pro- and atherogenic
- 397 immune responses, and disruption in the balance of immune cell subsets may trigger
- 398 atherosclerosis. ^{42,43} In this study, the ATLOs in aAAA were sparsely distributed, small, and
- 399 nearly round, similar to TLOs in chronic sialadenitis. These are thought to be a response to

400	local chronic stimulation. ⁴³ Clemet et al. noticed that active TKA with increased aortic
401	18F-fuluoro-deoxyglucose uptake contained several ATLOs, whereas a clear association
402	between pathogenesis and ATLOs in TKA was not observed. ¹³ In contrast, ATLOs in
403	IgG4-AAA were significantly larger, more numerous, and more irregular-shaped, with a
404	thick network of FDCs, compared to other vascular groups. Larger ATLO size in IgG4-AAA
405	was associated with a more irregular shape and a higher number of ATLOs. Additionally,
406	they were fused, or divided and collapsed, not simply enlarged. Moreover, the area, number,
407	and ATLO irregularity in IgG4-AAA significantly correlated with the activity of IgG4-RD,
408	such as the number of IgG4+ plasmacytes and serum IgG4 levels. Therefore, we believe that
409	the morphological deviation of ATLOs in IgG4-AAA might be associated with the active
410	phase of IgG4-AAA, ¹⁴ which could be useful for the pathological diagnosis of IgG4-AAA.
411	Recently, inflammation evaluated by 18F-PET/CT has been noted in IgG4-related vascular
412	diseases. ⁴⁴ Based on TKA, ³⁶ the activity of IgG4-AAA evaluated by FDG uptake observed by
413	PET/CT might reflect the amount or activity of ATLOs in IgG4-AAA.
414	Tfr cells are more likely to regulate the function and output of TLOs/ATLOs at an established
415	GC. In contrast, if Treg cells control initial GC formation; hence, the balance between Tfr and
416	ifTreg cells could contribute to GC development and decay by regulating the production of
417	several cytokines, such as IL-10 and IL-21.38,45 In IgG4-SS, Tfr/ifTreg imbalance is
418	associated with the invagination of the surrounding T-B-cell border into TLOs, causing
419	TLOs to have irregular shapes. ^{30,31} Similarly, in IgG4-AAA, both the number of Tfr cells and
420	the Tfr/if Treg ratio are closely related to the average area, area ratio, and ATLO irregularity.
421	Thus, the Tfr/ifTreg imbalance reflects characteristic ATLOs dysplasia in IgG4-AAA.
422	The morphological dysplasia of TLOs in IgG4-SS is greater (larger and more irregular
423	ATLOs with more prominent immunoblasts) than the ATLOs in IgG4-AAA. ^{30,31} Compared
424	to IgG4-AAA, IgG4-SS cases might be in a more active phase; they often have high serum

425	IgG4 levels and are prone to multiple organ lesions of IgG4-RD. In IgG4-SS, the Tfh2/ifTh2
426	and Tfh17/ifTh17 ratios are high, and the Tfh17/ifTh17 ratio is significantly related to the
427	number of IgG4 cells in the area, the area size, and TLO irregularity. ³¹ In contrast,
428	IgG4-AAA shows a higher degree of fibrosis and is more closely related to inflammatory
429	features, including elevated serum CRP and IL-6 levels related to local IL-6 production. ¹⁴
430	This report illustrates that in IgG4-AAA, the number of Tfh17 cells and the Tfh17/ifTh17
431	ratio were low, and the Th17 cell component was not related to IgG4-AAA activity and
432	ATLO morphology. Therefore, the differences in the morphological abnormalities of
433	ATLOs/TLOs in IgG4-AAA and IgG4-SS may be due to differences in the activity of
434	IgG4-RD and pathogenesis and progression factors.
435	The number of each T cell subset in aAAA was uncharacteristic. However, the ratios of
436	intrafollicular/interfollicular T cell subsets in aAAA were strikingly different from those of
437	IgG4-AAA, showing a moderate increase in the Tfr/ifTreg ratio, a high Tfh1/ifTh1 ratio, and
438	low ratios of Tfh2/ifTh2 and Tfh17/ifTh17. Interestingly, the ratios of
439	intrafollicular/interfollicular T cell subsets in non-IgG4-IAAA were similar to those of aAAA,
440	while the number of each T cell subset in non-IgG4-IAAA was higher than those of aAAA.
441	Thus, aAAA and non-IgG4-IAAA were similar, and non-IgG4-IAAA might be an
442	accelerated inflammatory state of aAAA.
443	In conclusion, the increases in T cell subsets inside ATLOs were characteristic for each type
444	of vascular lesion. Tfr and Tfh2 cells were predominant over Tfh1 cells in IgG4-AAA. The
445	increase in Tfr and the decrease in Tfh1 cells represent novel hallmarks of IgG4-AAA that
446	are associated with morphologically abnormal ATLO formation and disease pathogenesis. An
447	increase in Tfh17 cells was an attribute of TKA, and aAAA and non-IgG4-IAAA were

448 characterized by an increase in Tfh1 cells. In the classification of vascular lesions, based on

18

449	the imbalance in T cell subsets, IgG4-AAA should be positioned as vasculitis in predominant
450	adventitia accompanied by morphological abnormalities of ATLOs.
451	
452	Acknowledgements: The authors would like to sincerely thank the Master's students
453	Madoka Igarashi, Miyu Okuda, Ms. Hiromi Kojima, and Ms. Youko Umehara for their
454	technical assistance.
455	
456	Declarations
457	Funding: This work was supported by Grant-in AUC for Scientific Research (C) (JSPS
458	KAKENHI) Grant Number 20K9142.
459	
460	Conflict of Interest: The authors declare that they have no conflicts of interest.
461	
462	Author Contributions: SK performed data selection, experiments, data analysis, and
463	manuscript writing. NK, FK, YM, and HI participated in the selection of cases. AK and KH
464	supervised the work and were involved in setting up the experimental design. All authors
465	reviewed the manuscript and agreed with its contents.
466	
467	References:
468	1. Stone JH, Zen Y, Deshpande V. IgG4-related disease. N Engl J Med. 2012.366: 539-551.
469	https://doi: 10.1056/NEJMra1104650.
470	2. Deshpande V, Zen Y, Chan JK, Yi EE, Sato Y, Yoshino T, et al. Consensus statement on
471	the pathology of IgG4-related disease. Consensus statement on the pathology of IgG4-related
472	disease. Mod Pathol. 2012;25:1181-1192.

- 473 3. Wallace ZS, Naden RP, Chari S, Choi H, Della-Torre E, Jean-Francois Dicaire J, et al.
- 474 American College of Rheumatology/European League Against Rheumatism IgG4-Related
- 475 Disease Classification Criteria Working Group. The 2019 American College of
- 476 Rheumatology/European League Against Rheumatism Classification Criteria for
- 477 IgG4-Related Disease. Arthritis Rheumatol. 2020.72(1): 7-19. https://doi: 10.1002/art.41120.
- 478 4. Kasashima S, Zen Y, Kawashima A, Endo M, Matsumoto Y, Kasashima F. Inflammatory
- 479 abdominal aortic aneurysms: Close relationship to IgG4-related periaortitis. Am J Surg
- 480 Pathol. 2008.32:197-204.
- 481 5. Kasashima S, Zen Y, Kawashima A, Endo M, Matsumoto Y, Kasashima F. A new
- 482 clinicopathological entity of IgG4-related inflammatory abdominal aortic aneurysm. *J Vasc*
- 483 Surg. 2009.49:1264-1271.
- 484 6. Neyt K, Perros F, GeurtsvanKessel CH, Hammad H, Lambrecht BN. Tertiary lymphoid
- 485 organs in infection and autoimmunity. *Trends Immunol.* 2012.33:297-305.
- 486 7. Victora GD, Nussenzweig MC. Germinal Centers. Annu Rev Immunol. 2012.30:429–457.
- 487 doi: 10.1146/annurev-immunol-020711-075032
- 488 8. Gars E, Butzmann A, Ohgami R, Balakrishna JP, O'Malley DP. The life and death of the
- 489 germinal center. Ann Diagn Pathol. 2020.44:151421. https://doi:
- 490 10.1016/j.anndiagpath.2019.151421.
- 491 9. Bobryshev YV, Lord RS, Pärsson H. Immunophenotypic analysis of the aortic aneurysm
- 492 wall suggests that vascular dendritic cells are involved in immune responses. *Cardiovasc*
- 493 Surg. 1998.6:240-249.
- 494 10. Swedenborg J, Mäyränpää MI, Kovanen PT. Mast cells: important players in the
- 495 orchestrated pathogenesis of abdominal aortic aneurysms. Arterioscler Thromb Vasc Biol.
- 496 2011.31:734-740. doi: 10.1161/ATVBAHA.110.213157.

- 497 11. Wollenberg I, Agua-Doce A, Hernandez A, Almeida C, Oliveira VG, Faro J, Graca L.
- 498 Regulation of the germinal center reaction by Foxp3+ follicular regulatory T cells. *J Immunol*.
- 499 2011.187: 4553-4560. https://doi: 10.4049/jimmunol.1101328.
- 500 12. Yin C, Mohanta SK, Srikakulapu P, Weber C, Habenicht AJ. Artery tertiary lymphoid
- 501 organs: powerhouses of atherosclerosis immunity. Front Immunol. 2016.7:387.
- 502 13. Clement M, Galy A, Bruneval P, Morvan M, Hyafil F, Benali K, et al. Tertiary lymphoid
- 503 organs in Takayasu arteritis. Front Immunol. 2016.7:158. doi: 10.3389/fimmu.2016.00158.
- 504 eCollection 2016.
- 505 14. Kasashima S, Kawashima A, Zen Y, Ozaki S, Kasashima F, Endo M, et al. Upregulated
- 506 Interleukins (IL-6, IL-10, and IL-13) in Immunoglobulin G4-Related Aortic Aneurysm
- 507 Patients. J Vasc Surg. 2018. 67(4):1258-1262. https://doi.org/10.1016/j.jvs.2016.12.140
- 508 15. Vinuesa CG, Linterman MA, Yu D, MacLennan IC. Follicular Helper T Cells. Ann Rev
- 509 *Immunol.* 2016.20(34): 335-368. https://doi:10.1146/annurev-immunol-041015-055605.
- 510 16. Shulman Z, Gitlin AD, Targ S, Jankovic M, Pasqual G, Nussenzweig MC, Victora GD. T
- 511 Follicular Helper Cell Dynamics in Germinal Centers. *Science*. 2013. 9; 341(6146): 673–677.
- 512 doi: 10.1126/science.1241680
- 513 17. Sage PT, Sharpe AH. T follicular regulatory cells in the regulation of B cell responses.
- 514 Trends Immunol. 2015. 36: 410-418. https://doi: 10.1016/j.it.2015.05.005.
- 515 18. Laidlaw BJ, Lu Y, Amezquita RA, Weinstein JS, Vander Heiden JA, Gupta NT,
- 516 Kleinstein SH, Kaech SM, Craft J. Interleukin-10 from CD4(+) follicular regulatory T cells
- 517 promotes the germinal center response. *Sci Immunol.* 2017. 2(16): eaan4767. https://doi:
- 518 10.1126/sciimmunol.aan4767.
- 519 19. Xu B, Wang S, Zhou M, Huang Y, Fu R, Guo C, Chen J, Zhao J, Gaskin F, Fu SM, Yang
- 520 N. The ratio of circulating follicular T helper cell to follicular T regulatory cell is correlated

- 521 with disease activity in systemic lupus erythematosus. *Clin Immunol.* 2017.183: 46-53.
- 522 https://doi: 10.1016/j.clim.2017.07.004.
- 523 20. Deng J, Wei Y, Fonseca VR, Graca L, Yu D T follicular helper cells and T follicular
- 524 regulatory cells in rheumatic diseases. *Nat Rev Rheumatol.* 2019.15(8): 475-490. https://doi:
- 525 10.1038/s41584-019-0254-2.
- 526 21. Mattoo H, Della-Torre E, Mahajan VS, Stone JH, Pillai S. Circulating Th2 memory cells
- 527 in IgG4-related disease are restricted to a defined subset of subjects with atopy. *Allergy*.
- 528 2014.69(3):399-402. doi: 10.1111/all.12342.
- 529 22. Heeringa JJ, Karim AF, van Laar JAM, Verdijk RM, Paridaens D, van Hagen PM, van
- 530 Zelm MC. J Expansion of blood IgG4+ B, TH2, and regulatory T cells in patients with
- 531 IgG4-related disease. *Allergy Clin Immunol.* 2018.141(5):1831-1843.e10. doi:
- 532 10.1016/j.jaci.2017.07.024
- 533 23. Akiyama M, Yasuoka H, Yamaoka K, Suzuki K, Kaneko Y, Kondo H, et al. Enhanced
- 534 IgG4 Production by Follicular Helper 2 T Cells and the Involvement of Follicular Helper 1 T
- 535 Cells in the Pathogenesis of IgG4- Related Disease. Arthritis Res Ther .2016.18:167. doi:
- 536 10.1186/s13075-016- 1064-4
- 537 24. Akiyama M, Suzuki K, Yasuoka H, Kaneko Y, Yamaoka K, Takeuchi T. Follicular
- helper T cells in the pathogenesis of IgG4-related disease. *Rheumatology (Oxford).* 2018.57:
- 539 236-245. https://doi: 10.1093/rheumatology/kex171.
- 540 25. Chen Y, Lin W, Yang H, Wang M, Zhang P, Feng R, et al. Aberrant expansion and
- 541 function of follicular helper T cell subsets in IgG4-related disease. Arthritis Rheumatol.
- 542 2018.70:1853–1865. doi: 10.1002/art.40556
- 543 26. Kubo S, Nakayamada S, Zhao J, Yoshikawa M, Miyazaki Y, Nawata A, et al. Correlation
- of T follicular helper cells and plasmablasts with the development of organ involvement in

545 patients with IgG4- related disease. *Rheumatology*. 2018. 57:514–524.

- 546 doi:10.1093/rheumatology/kex455
- 547 27. Cargill T, Makuch M, Sadler R, Lighaam LC, Peters R, van Ham M, et al. Activated
- 548 T-Follicular Helper 2 Cells are associated with disease activity in IgG4-related sclerosing
- 549 cholangitis and pancreatitis. *Clin Transl Gastroenterol*. 2019.10(4): e00020. doi:
- 550 10.14309/ctg.000000000000020
- 551 28. Liu J, Yin W, Westerberg LS, Lee P, Gong Q, Chen Y, et al. Immune dysregulation in
- 552 IgG4-related disease. Front Immunol. 2021.01. https://doi.org/10.3389/fimmu.2021.738540
- 553 29. Kamekura R, Takano K, Yamamoto M, Kawata K, Shigehara K, Jitsukawa Sl. Cutting
- edge: a critical role of regional T follicular helper cells in the pathogenesis of IgG4-related
- 555 disease. J Immunol. 2017.199: 2624-2629. https://doi: 10.4049/jimmunol.1601507.
- 556 30. Kasashima S, Kawashima A, Kurose N, Ozaki S, Ikeda H, Harada K. Regional
- 557 disturbance of the distribution of T regulatory cells and T helper cells associated with
- 558 irregular-shaped germinal centers in immunoglobulin G4-related sialadenitis. *Virchows Arch.*
- 559 2021.479(6):1221-1232. doi: 10.1007/s00428-021-03187-2.
- 560 31. Kasashima S, Kawashima A, Kurose N, Ozaki S, Ikeda H, Harada K. The disturbance of
- the distribution of T helper cell subsets in the mantle area surrounding germinal centers in
- 562 immunoglobulin G4-related sclerosing sialadenitis. Virchows Arch. 2022.481(5):767-777.
- 563 doi: 10.1007/s00428-022-03384-7.
- 564 32. Grayson PC, Ponte C, Suppiah R, Robson JC, Gribbons KB, Judge A, Craven A, Khalid
- 565 S, Hutchings A, Danda D, Luqmani RA, Watts RA, Merkel PA; DCVAS Study Group. 2022
- 566 American College of Rheumatology/EULAR Classification Criteria for Takayasu Arteritis.
- 567 Arthritis Rheumatol. 2022.74(12):1872-1880. doi: 10.1002/art.42324.
- 568 33. Chung Y, Tanaka S, Chu F, Nurieva RI, Martinez GJ, Rawal S, Wang Y, Lim H,
- 569 Reynolds JM, Zhou X, Fan H, Liu Z, Neelapu SS, Dong C. Follicular regulatory T cells

- 570 expressing Foxp3 and Bcl-6 suppress germinal center reactions. *Nat Med.* 2021.17(8):
- 571 983-988. https://doi: 10.1038/nm.2426.
- 572 34. Schneider CA, Rasband WS, Eliceiri KW. NIH Image to Image J: 25 years of image
- 573 analysis. *Nat Methods*. 2012. 9: 671-675. https://doi: 10.1038/nmeth.2089.
- 574 35. Satyanarayanan M, Goode A, Gilbert B, Harkes J, Jukic D. OpenSlide: A vendor-neutral
- 575 software foundation for digital pathology. J Pathol Inform. 2013.4:27. https://doi:
- 576 10.4103/2153-3539.119005.
- 577 36. Saadoun D, Garrido M, Comarmond C, Desbois AC, Domont F, Savey L. Th1 and Th17
- 578 cytokines driveinflammation in Takayasu arteritis. Arthritis
- 579 Rheumatol.2015.67(5):1353-1360. doi: 10.1002/art.39037
- 580 37. Gao N, Cui W, Zhao LM, Li TT, Zhang JH, Pan LL. Contribution of Th2-like Treg cells
- to the pathogenesis of Takayasu's arteritis. *Clin Exp Rheumatol.* 2020.38 Suppl 124(2):48-54.
- 582 38. Ito F, Kamekura R, Yamamoto M, Takano K, Takaki H, Yabe H, Ikegami I, Shigehara K,
- 583 Himi T, Takahashi H, Ichimiya S. IL-10+ T follicular regulatory cells are associated with the
- pathogenesis of IgG4-related disease. *Immunol Lett.* 2019. 207:56-63. doi:
- 585 10.1016/j.imlet.2019.01.008.
- 586 39. Grados A, Ebbo M, Piperoglou C, Groh M, Regent A, Samson M. T Cell Polarization
- 587 toward TH2/TFH2 and TH17/TFH17 in Patients with IgG4-Related Disease. Front Immunol.
- 588 2017.8: 235. doi: 10.3389/fimmu.2017.00235
- 589 40. Dudreuilh C, Basu S, Scottà C, Dorling A, Lombardi G. Potential Application of
- 590 T-Follicular Regulatory Cell Therapy in Transplantation. Front Immunol. 2021.2;11:612848.
- 591 doi: 10.3389/fimmu.2020.612848.
- 41. Yao Y, Chen CL, Yu D, Liu Z. Roles of follicular helper and regulatory T cells in allergic
- 593 diseases and allergen immunotherapy. *Allergy*. 2021.76(2):456-470. doi: 10.1111/all.14639.

- 42. Mohanta SK, Yin C, Peng L, Srikakulapu P, Bontha V, Hu D, Falk Weih, Weber C,
- 595 Gerdes N, Habenicht AJR. Artery tertiary lymphoid organs contribute to innate and adaptive
- immune responses in advanced mouse atherosclerosis. *Circ Res.* 2014.114:1772-1787.
- 43. Weih F, Gräbner R, Hu D, Beer M, Habenicht AJ. Control of dichotomic innate and
- 598 adaptive immune responses by artery tertiary lymphoid organs in atherosclerosis. Front
- 599 Physiol. 2012.3:226.
- 600 44. Oyama-Manabe N, Yabusaki S, Manabe O, Kato F, Kanno-Okada H, Kudo K.
- 601 IgG4-related Cardiovascular Disease from the Aorta to the Coronary Arteries: Multidetector
- 602 CT and PET/CT. *Radiographics*. 2018.38(7):1934-1948. doi: 10.1148/rg.2018180049.
- 45. Fahlquist Hagert C, Degn SE. T follicular regulatory cells: Guardians of the germinal
- 604 center? *Scand J Immunol.* 2020.92: e12942. https://doi: 10.1111/sji.12942.

605

606 Figures legends

607	Figure 1.	IgG4-related	inflammatory	abdominal	l aortic aneu	ırysm (A	.–C), Ta	ikayasu arteritis	5
-----	-----------	--------------	--------------	-----------	---------------	----------	----------	-------------------	---

- 608 (D–F), and atherosclerotic abdominal aortic aneurysm (G–I) histopathological features.
- 609 Characteristic massive fibro-inflammation with lymphoplasmacytic aggregation and
- 610 thickening adventitia in IgG4-related inflammatory abdominal aortic aneurysm (A, B),
- 611 degenerative elastic fibers of the media in Takayasu arteritis (D, E), and fibrous intimal
- 612 thickening and thin media seen in atherosclerotic abdominal aortic aneurysm (G, H). CD21
- 613 immunohistochemistory shows follicular dendritic cells distributed across the germinal
- 614 centers as arterial/aortic tertiary lymphoid organs; large and irregular-shaped germinal

615 centers seen in IgG4-related inflammatory abdominal aortic aneurysm (C); few, small, and

oval- to round-shaped germinal centers seen in Takayasu arteritis (F) and atherosclerotic

617 abdominal aortic aneurysm (I). (A, D, and G: Hematoxylin-Eosin staining; B, E, and H:

618 Elastica van Gieson stain; C, F, and I: CD21 immunohistochemistory; A-C: original

619 magnification $\times 12.5$; D–I: original magnification $\times 20$; A–I: scale bar = 500 µm).

620

621 Figure 2. (A–D) QuPath's methods were used to perform regional annotation using CD21

622 immunohistochemical-stained slides. The red annotation line is the germinal center's

623 boundary line and is determined by using CD21 immunohistochemistory (A) and transferred

624 to Foxp3 immunohistochemistory (B). (C) Using QuPath's cell detection, with the inside red

625 line as the germinal center, Foxp3 immunopositive cells were selected as red-marked cells to

- 626 calculate the follicular regulatory T-cell number. (D) The Foxp3 immunohistochemistory
- 627 whole-slide image analysis of IgG4-related inflammatory abdominal aortic aneurysm shows
- 628 red annotation lines, which are the germinal centers' boundary lines. Follicular regulatory T
- 629 cells are calculated in each germinal center, and interfollicular regulatory T cells are

630 calculated outside of germinal centers. T-bet (E), GATA3 (F), and ROR-γ (G) indicate type 1

- helper T-cells, type 2 helper T cells, and type 17 helper T-cells, respectively. QuPath's
- analyses were performed in the same manner as that for regulatory T cells using Foxp3
- 633 immunohistochemistory (A–C and E–G: original magnification \times 400, scale bar = 100 μ m
- and D: original magnification $\times 10$, scale bar = 1000 μ m).
- 635

636 Figure 3. The morphological features of arterial/aortic tertiary lymphoid organs in patients

637 with IgG4-related inflammatory abdominal aortic aneurysm (IgG4-AAA, n = 21), non

638 IgG4-related inflammatory abdominal aortic aneurysm (Non-IgG4-AAA, n = 17), Takayasu

639 arteritis (TKA, n =5), atherosclerotic abdominal aortic aneurysm (aAAA, n = 10), and control

640 aorta (n = 5). Statistically significant differences between groups are indicated at * and **p < 100

641 0.01.

642

- 643 Figure 4. (A) The numbers of T cells subsets in patients with IgG4-related inflammatory
- 644 abdominal aortic aneurysm (IgG4-AAA, n = 21), non IgG4-related inflammatory abdominal
- 645 aortic aneurysm (Non-IgG4-AAA, n = 17), Takayasu arteritis (TKA, n = 5), atherosclerotic

abdominal aortic aneurysm (aAAA, n = 10), and control aorta (n = 5). (B) The ratio of each T

647 cell subset and (C) the ratio of T cell subsets within ATLOS and outside ATLOs. Tfr,

648 follicular regulatory T cell; Tfh1, type 1 follicular helper T cell; Tfh2, type 2 follicular helper

- T cell; fTh17, type 17 follicular helper T cell; ifTreg, interfollicular regulatory T cell; ifTh1,
- type 1 interfollicular helper T cell; ifTh2, type 2 interfollicular helper T cell; ifTh17, type 17;
- 651 ATLOs, arterial tertiary lymphoid organs. Statistically significant differences between groups
- 652 are indicated if p < 0.05 and p < 0.01.

653

- Figure 5. (A) Summary of correlations between IgG4-related disease activity factors and T
- 655 cell subsets or morphological features of arterial/aortic tertiary lymphoid organs. (B)
- 656 Summary of correlations between T cell subsets and features of arterial/aortic tertiary
- 657 lymphoid organs. In each cell, the higher the positive correlation coefficient, the deeper the
- red color, and the stronger the negative correlation, the deeper the blue. The numbers are
- 659 correlation coefficients; the bold, dark colored, underlined numbers represent p < 0.05, and
- 660 the lightly colored numbers are not significant.

	aAAA	non-IgG4-IAAA	IgG4-AAA	Takayasu disease
	(10 cases)	(17 cases)	(21 cases)	(5 cases)
Age (years)	68.3 (56–72)	70.2 (56–81)	70.5 (70)63–79	32.4 (34)22–54
Gender (male/female)	5'/5	8'/3	20'/1	0/5
Aneurysmal diameter (mm)	52.1 (34–75)	49.2 (30–100)	53.5 (23-87)	30 (20–45)
Low grade fever	0	5 (29.4%)	10 (47.6%)	2 (40%)
Abdominal or lumber pain	1 (10%)	5 (29.4%)	15 (71.4%)	0
C-reactive protein (mg/dL)	0.41 (0.1–1.1)	0.69 (0.21-8.04)	1.44 (0.01–5.22)	0.78 (0.1–1.51)
White blood cell (× $10^3/\mu$ L)	3.4 (3.2–6.7)	7.9 (3.4–19.5)	7.1 (4.7–9.5)	8.9 (6.8–20.2)
Serum IgG4 (mg/dL)	34.5 (12–59)	24.3 (15.0–58.9)	91.5 (33.9–559.0)	23.2 (0-46.9)
Serum IgG4/IgG (%)	2.3 (2.1–2.9)	1.9 (1.1–6.4)	8.1 (2.4–29.7)	2.8 (2.1–3.1)
Adventitial thickness (mm)	0.4 (0.1–1.3)	3.3 (2.2–6.0)	4.7 (0.6–14.0)	1.3 (0.3–1.8)
Histological IgG4 (/HPF)	6.3 (6.1–43.5)	17.1 (0-41.5)	83.3 (42.3–175.7)	12.1 (8.1–22.4)
Histological IgG4/IgG (%)	7.9 (0–38.7)	26.3 (0-40.9)	80.2 (58.3–98.7)	11.4 (3.5–18.4)

Table 1. Clinicopathological features of aAAA, non-IgG4-IAAA, IgG4-AAA, and Takayasu disease.

aAAA, atherosclerotic abdominal aortic aneurysm; non-IgG4-IAAA, non-IgG4-IAAA, non-IgG4-related inflammatory abdominal aortic aneurysm; IgG4-AAA, IgG4-related inflammatory abdominal aortic aneurysm; HPF, high power field

data are present as median (range). Three cases of Takayasu disease could measure aneurysmal diameter, serum IgG4, and serum IgG4/IgG.





























D







A	ATLOs features							T cell subtype						
		all cases	number	average area	area ratio	irregularity	Tfr	Tfh1	Tfh2	Tf17	Tfr/lfTreg	Tfh1/lfTh1	Tfh2/lfTh2	Tfh17/lfTh17
	Q sa	adventitial thickenss	0.220	0.361	0.108	0.410	0.461	0.131	0.332	-0.114	0.389	-0.113	0.443	0.104
	tur	number of gG4+ cells	0.123	0.541	<u>0.491</u>	<u>0.517</u>	0.446		0.277	0.105	0.334	0.111	0.332	0.115
	feg	IgG4+ cell/IgG+ cell ratioo	0.179	0.504	<u>0.414</u>	0.462	0.452	<u>-0.394</u>	0.333	0.287	0.433	-0.223	0.443	0.133

	IgG4-AAA cases	number	average area	area ratio	irregularity	Tfr	Tfh1	Tfh2	Tf17	Tfr/lfTreg	Tfh1/lfTh1	Tfh2/IfTh2	Tfh17/lfTh17
C sa	adventitial thickenss	0.405	0.277	-0.244	0.291	0.108	<u>-0.526</u>	0.222	-0.123	0.345	<u>-0.328</u>	0.123	0.122
IgG4-F featur	number of gG4+ cells	0.164	0.214	0.270	0.376	0.212	<u>-0.458</u>	0.306	0.078	0.322	<u>-0.343</u>	0.263	0.332
	IgG4+ cell/IgG+ cell ratioo	-0.144	0.391	0.182	0.340	0.359	<u>-0.374</u>	0.088	0.137	0.332	-0.332	0.322	0.277

	all cases	Tfr	Tfh1	Tfh2	Tf17	Tfr/IfTreg	Tfh1/lfTh1	Tfh2/lfTh2	Tfh17/lfTh17
	number	0.034	-0.244	-0.111	0.095	0.078	-0.243	0.056	-0.057
-Os	average area	0.343	-0.190	0.186	0.065	0.372	-0.190	0.124	-0.038
ATI	area ratio	<u>0.304</u>	-0.003	0.123	-0.057	<u>0.302</u>	-0.003	0.178	0.040
-	irregularity	0.638	0.171	<u>0.580</u>	-0.038	0.472	-0.137	0.456	0.120
	IgG4-AAA cases	Tfr	Tfh1	Tfh2	Tf17	Tfr/IfTreg	Tfh1/lfTh1	Tfh2/ifTh2	Tfh17/lfTh17
	number	0.335	-0.316	0.334	0.007	0.332	-0.343	0.223	0.023
-Os	average area	<u>0.349</u>	-0.208	<u>0.382</u>	0.107	0.345	-0.146	0.332	0.122
ATI	area ratio	0.386	-0.218	<u>0.499</u>	0.065	0.332	-0.230	0.389	-0.057
-	irregularity	0.473	0.132	0.144	0.241	0.459	-0.137	0.223	-0.038

Correlation coefficient

