

1 Efficacy of multilevel exercise programmes with different intensities on development
2 and metabolic homeostasis in overweight and obese children: study protocol for a
3 randomized controlled trial

4

5 Yimin Zhao^{1, #}, Chao Liu^{2, #}, Jiaying Wang², Yangong Liu², Haiyi Yu², Tianqi Chang², Chenyu
6 Fan², Wenxiu Wang¹, Yueying Li¹, Hu Wang², Yuzhou Xue², Wei Zhao^{2, 4}, Chuan Ren², Ligang
7 Cui³, Xinheng Feng², He Huang⁸, Dongwei Fan⁵, Xiang Yao⁶, Tongyan Han⁷, Tao Huang¹, and
8 Ming Xu^{2, *}

9

10 ¹ Department of Epidemiology and Biostatistics, School of Public Health, Peking University,
11 Beijing, China;

12 ²Department of Cardiology, Institute of Vascular Medicine, NHC Key Laboratory of
13 Cardiovascular Molecular Biology and Regulatory Peptides, Key Laboratory of Molecular
14 Cardiovascular Science, Ministry of Education, Beijing Key Laboratory of Cardiovascular
15 Receptors Research, Peking University Third Hospital, Beijing, China;

16 ³ Department of Ultrasound, Peking University Third Hospital, Beijing, China;

17 ⁴ Physical Examination Center, Peking University Third Hospital, Beijing, China;

18 ⁵ Department of Orthopedics, Peking University Third Hospital, Beijing, China;

19 ⁶ School of Psychological and Cognitive Sciences, Peking University, Beijing, China;

20 ⁷ Department of Pediatrics, Peking University Third Hospital, Beijing, China.

21 ⁸ Shanghai Key Laboratory of Metabolic Remodeling and Health, Institute of Metabolism and
22 Integrative Biology, Shanghai, China

23

24 #Yimin Zhao and Chao Liu contributed equally to this work.

25

26 **Corresponding authors:**

27 Ming Xu, MD, PhD

28 Department of Cardiology, Institute of Vascular Medicine, NHC Key Laboratory of

29 Cardiovascular Molecular Biology and Regulatory Peptides, Key Laboratory of Molecular

30 Cardiovascular Science, Ministry of Education, Beijing Key Laboratory of Cardiovascular

31 Receptors Research, Peking University Third Hospital, Beijing, China

32 Email: xuminghi@bjmu.edu.cn

33 **Abstract**

34 **Background:** The prevalence of childhood obesity is rapidly increasing worldwide. Exercise
35 has been suggested as the primary approach to manage body weight, but an effective
36 exercise program tailored for school-age children is still lacking.

37 **Objective:** This trial aims to evaluate the efficacy of a novel multilevel (children, families, and
38 schools)-coordinated exercise programmes with different intensities on adiposity and
39 development among overweight and obese school-age children. It also aims to elucidate the
40 complex interplays between the exercise interventions, multi-omic biomarkers, development,
41 and metabolic homeostasis.

42 **Methods:** This is a multilevel -coordinated two-stage randomized double-blind controlled trial,
43 comprising 366 overweight and obese children aged between 8 and 12 years from two
44 elementary schools in Beijing, China. The novel multilevel exercise programs involve families
45 and schools monitoring children's exercise and diet via a specially designed smartphone
46 application. The children will be randomly assigned to receive three school-based exercise
47 programmes (exercise intervention stage), namely low-intensity exercise, moderate-intensity
48 exercise, and intermittent vigorous exercise, for 12 weeks, followed by a 9-month
49 post-intervention follow-up (follow-up stage). The primary outcomes of this trial are the
50 changes in adiposity, development, and metabolic homeostasis during the two stages. The
51 secondary outcomes include but are not restricted to genomics and changes in blood
52 metabolomics, gut microbiota, and exercise performance.

53 **Discussion:** This trial has been approved by the ethics committee of Peking University Third
54 Hospital (ethnic approval number 2021-283-06). The findings of this trial will help establish

55 specialized multilevel coordinated school-based exercise programmes for overweight and
56 obese school-age children in China and quantitatively and qualitatively evaluate the efficacy
57 of the novel interventions.

58 **Trial registration:** This trial was prospectively registered at ClinicalTrials.gov with the
59 identifier of NCT04984005 on July 21, 2021.

60

61 Introduction

62 The prevalence of childhood obesity is rapidly increasing worldwide. According to recent data,
63 the number of obese children and adolescents has increased from 11 million in 1975 to 124
64 million in 2016.^{1,2} This issue is becoming particularly prevalent in developing countries such
65 as China, where the prevalence of childhood obesity and related non-communicable chronic
66 diseases is rising rapidly.³ It is projected that by 2030, 6.64 million children under the age of
67 seven, and 18.91 million children over the age of seven in China, will be obese.^{4,5} In Beijing,
68 the prevalence of overweight and obesity was found to be 17.6% and 29.1% in boys and 17.6%
69 and 18.0% in girls aged between 6 to 18 years, respectively.⁵

70
71 Childhood obesity has been linked to an increased risk of type 2 diabetes, heart disease, and
72 chronic kidney disease in later life.⁶⁻¹⁰ While exercise intervention remains the primary
73 approach for managing body weight, it may not achieve sustained desirable body weight
74 reduction on its own for obese children who also have to manage academic stress.¹¹⁻¹⁴ Indeed,
75 a recent multicomponent lifestyle intervention successfully reduced adiposity and obesity
76 prevalence among children.⁴ However, whether the involvement of schools and families could
77 improve the effectiveness of exercise intervention among school-age obese children is still
78 less studied.¹⁵ Notably, the parents and teachers may hold the opinions that strict exercise
79 intervention may impede growth and development and academic performance of the children,
80 thus reducing the effectiveness of exercise. In addition, it remains unknown whether a more
81 intense exercise programme would bring about additional health benefits in terms of
82 development and metabolic homeostasis beyond body weight during puberty for obese

83 children.^{1 14}

84

85 In this study, we aim to compare the effects of multilevel-coordinated exercise programmes
86 with different intensities on adiposity, development, and metabolic homeostasis, as well as
87 related mechanisms. Specifically, we aim to determine which exercise programme is more
88 effective in reducing body weight and improving development and metabolic homeostasis
89 among overweight and obese children, as well as identifying the factors that mediate
90 exercise-induced changes in these areas. Additionally, we aim to establish a regionally
91 representative school-family-hospital evaluation model for exercise intervention.

92

93 **Method and Analysis**

94 **Trial design**

95 This is a multilevel coordinated (children, families, and schools) two-stage 12-month
96 three-arm randomized double-blind controlled trial.

97

98 **Methods: Participants, interventions, and outcomes**

99 This protocol is reported following the guidelines of the Standard Protocol Items:

100 Recommendations for Interventional Trials (SPIRIT) 2013 Statement.¹⁶

101

102 **Study setting**

103 This trial will be conducted in two government-funded elementary schools in Beijing, the
104 capital city of China. The steering committee of this study included paediatricians, physicians,

epidemiologists, and statisticians from Peking University Third Hospital and School of Public Health at Peking University. All study procedures will be performed in accordance with the Declaration of Helsinki.

Eligibility criteria

Eligible participants must meet the following inclusion criteria:

- 1) overweight or obese (see **Table 1** for age- and sex-specific BMI cutoff points);
- 2) 8 to 12 years old;
- 3) did not participate in any weight loss programmes, including medication, exercise, and lifestyle modification, within the last three months before baseline assessment;
- 4) provide written informed consent by both themselves and their statutory guardians.

Participants will be excluded if they have any of the following conditions:

- 1) have any severe chronic disease, including asthma, cardiovascular disease, and cancer.
- 2) have drug-induced secondary obesity;
- 3) have Cushing syndrome, Prader-Willi syndrome, or other medical conditions that may lead to obesity;
- 4) have intellectual, mental, or behavioural disorders;
- 5) feel difficulty communicating with physical education (PE) teachers, paediatricians, and physicians;
- 6) have any other conditions that may prevent exercising safely, which will be assessed by paediatricians and physicians.

127

128 **Who will take informed consent?**

129 A group of research staff, including paediatricians and physicians, will collect written informed
130 consent from both children and their statutory guardians before evaluating the eligibility of
131 potential participants. During the consent process, potentially eligible participants and their
132 parents will be notified that this trial will require the collection of fasting blood, urine, and
133 faeces samples.

134

135 **Interventions**

136 **Explanation for the choice of comparators**

137 Eligible participants will be randomly assigned into three groups receiving low-intensity
138 exercise, moderate-intensity exercise, and intermittent vigorous exercise, respectively. The
139 low-intensity exercise is chosen as the comparator because exercise intervention, even at a
140 low intensity, has been proven effective in improving energy expenditure and cardiometabolic
141 health among children and adolescents. The novel intervention programmes emphasize the
142 involvement of family and school to assist children's exercise and food intake using a
143 specifically designed smartphone application. The health educators, the pediatrician in the
144 school hospital, will be trained by the project staff before the intervention. During the
145 intervention, the health educator will deliver two health education sessions, including exercise
146 and healthy eating, to children every month. Family members and school PE teachers at each
147 site will be separately invited to attend the health education sessions two times per month,
148 where the project staff shared the information on children's exercise performance and

strengthened the family and school involvement to assist children's exercise.

Intervention description

The levels of intensity will be determined according to the percentages of heart rate reserve (HRR) during exercise, which is calculated by deducting the resting heart rate (HR_{resting}) from the maximum heart rate (HR_{max}).^{17 18} The HR_{resting} and the HR_{max} will be evaluated on a COSMED Quark PFT pulmonary function testing system in Peking University Third Hospital at baseline.

All participants will engage in a school-based exercise programme, instructed by PE teachers, at the frequency of 30 minutes/time $\times$ two times/day $\times$ three days/week for 12 weeks. Each 30-minute exercise will comprise a 7-minute warm-up session, a 16-minute exercise session, and a 7-minute cool-down session. In the warm-up and cool-down sessions, all groups of participants will do walking and stretching exercises, including arm circles, overhead arm reaches, and hip rotations, to reach a target heart rate of around $0.3 \times HRR + HR_{\text{resting}}$. During the 16-minute exercise session:

- 1) for the low-intensity exercise group, the participants will be instructed to take 800-meter quick walking and high knees to reach a target heart rate between $0.3 \times HRR + HR_{\text{resting}}$ and $0.4 \times HRR + HR_{\text{resting}}$;
- 2) for the moderate-intensity exercise group, the participants will be instructed to take 30-meter S-shaped quick walking, 30-meter S-shaped running, and 800-meter alternating between running and walking to reach a target heart rate between $0.4 \times HRR + HR_{\text{resting}}$

and $0.69 \times \text{HRR} + \text{HR}_{\text{resting}}$;

3) for the intermittent vigorous exercise group, the participants will be instructed to take various 30-second jumping and quick running activities to reach a target heart rate between $0.7 \times \text{HRR} + \text{HR}_{\text{resting}}$ and $0.85 \times \text{HRR} + \text{HR}_{\text{resting}}$, divided by a 60-second relaxation above $0.6 \times \text{HRR} + \text{HR}_{\text{resting}}$.

In the 2-week run-in period prior to intervention period, all participants will receive the same low-intensity exercise intervention as described above.

Criteria for discontinuing or modifying allocated interventions

We allow participants to discontinue the exercise intervention at any time for any reason. The research staff will document the reasons for dropping out if provided by the participants. We will not recruit new participants to fill the vacancy after the implementation of the exercise intervention.

Strategies to improve adherence to interventions

Participants will be instructed to wear a calibrated smartwatch (Huawei Band 6) during the run-in and intervention stages, which is provided by Huawei Technologies for free. During each 30-minute exercise session, the smartwatch will return 30 reads of real-time heart rate at a 1-minute interval. Adherence to the exercise programme is assessed by counting the real-time heart rate reads that are within the target range for each participant. We assume that all 16 reads collected during the 16-minute exercise session are located within the target

heart rate zone in each group. In addition, we will reward participants who satisfactorily complete the 12-week exercise intervention with a gift, including a pen and pencil box.

Relevant concomitant care permitted or prohibited during the trial

During the 2-week run-in period and 12-week exercise intervention period, all participants will be instructed to maintain their usual physical activity and diet using the specifically designed smartphone application. We will monitor their diet compositions via a three-day dietary recall, and the nutrient intake will be analyzed according to China Food Composition Tables.¹⁹ We will assess their physical activity, excluding exercise intervention, via the Huawei band 6. These two questionnaires will be performed by trained research staff face-to-face. In addition, participants and their parents will receive notifications via a mobile phone application, which will notify the participants to maintain their usual diet and physical activity.

Provisions for post-trial care

We will implement a 9-month post-intervention follow-up to observe the dynamic changes in adiposity among the participants.

Outcomes

Primary outcomes

The primary outcomes of this trial are the changes in adiposity and biomarkers of development and homeostasis, including anthropometric measurements, sexual hormones, body compositions, cardiopulmonary functions, and biochemical measurements.

215

216 **Secondary outcomes**

217 Secondary outcomes include changes in:

- 218 1. Exercise performance;
- 219 2. Hepatic hemodynamic index;
- 220 3. Bone mineral density;
- 221 4. Inflammatory biomarkers;
- 222 5. Thyroid function;
- 223 6. Oxidative stress biomarkers;
- 224 7. Oral and gut microbiota;
- 225 8. Blood and faecal metabolomics;
- 226 9. Flow-mediated dilation
- 227 10. Attention performance;
- 228 11. Sleep quality;
- 229 12. Diets and nutrients;
- 230 13. Whole genome sequencing;
- 231 14. Viromics;
- 232 15. Proteomics;
- 233 16. Volatile organic compounds in breath;

234

235 **Participant timeline**

236 **Figure 1** shows the schedule of enrollment, allocation, interventions, and assessments.

237

238 **Sample size**

239 We estimated the sample size using the following equation:

$$N_d = \frac{1641.6 \times \lambda}{(\sin^{-1}\sqrt{p_{max}} - \sin^{-1}\sqrt{p_{min}})^2}$$

240 , where N_d is the minimum sample size, P_{max} and P_{min} are the highest and lowest cure rates,

241 and λ depends on α , β , and degree of freedom. In this trial, α is 0.05, β is 0.1, and the degree

242 of freedom is 2. Thereby, λ is 12.65. We assumed that 65%, 50%, and 40% of overweight and

243 obese participants that receive low-intensity exercise, moderate-intensity exercise, and

244 intermittent vigorous exercise would successfully lower their body weight, achieving a healthy

245 BMI, respectively. A minimum sample size of 122 is required for each group accounting for a

246 10% drop-out rate. A total of 366 children will be included in this trial.

247

248 **Recruitment**

249 From January to March 2022, we included two elementary schools in Beijing by inviting the

250 school principals to participate in this trial. The two schools agreed with the study protocol. We

251 will recruit eligible participants, aged between 8 and 12 years, from seven school campuses.

252 All students and their parents (or their statutory guardians) will receive an advertisement for

253 this trial from the principal and their PE teachers. Potential eligible participants who provide

254 written informed consent by themselves and statutory guardians will be asked to complete a

255 nurse-administrated questionnaire concerning their health-related information and undergo a

256 physical examination according to standard operating procedure.

257

258 **Assignment of interventions: allocation**

259 **Sequence generation**

260 We will perform sex- and BMI-stratified randomization via computer-generated random
261 numbers.

262

263 **Concealment mechanism**

264 The participants will be randomized using a computer-generated random sequence, which will
265 be sealed in an envelope until the end of this trial.

266

267 **Implementation**

268 An independent statistician from the School of Public Health, Peking University, who will not
269 participate in recruitment, allocation, baseline assessment, and intervention, will conduct the
270 randomization.

271

272 **Assignment of interventions: Blinding**

273 **Who will be blinded**

274 All participants, their parents, the PE teachers, the paediatricians, the physicians, and the
275 statisticians will be blinded to the allocation.

276

277 **Procedure for unblinding if needed**

278 Not applicable to this trial.

279

Data collection and management

Plans for assessment and collection of outcomes

All data will be collected and processed according to standard operating procedures. The demographic information of participants will be collected at baseline. Trained nurses will conduct the physical examination under the supervision of paediatricians. The biochemical measurements will be conducted at the clinical laboratory of Peking University Third Hospital. The oral and gut microbiota compositions will be analyzed using the QIIME2 pipeline (<https://qiime2.org>). The metabolomics analysis will be conducted on a liquid chromatography-electrospray ionization tandem mass spectrometry system (QTRAP 6500+ system).^{20,21} Diet will be evaluated via a 3-day diet recall at baseline and at the end of intervention period. Meanwhile, the participants will be instructed to take photo of meals to facilitate nutrient intake analysis. Nutrients intake will be analyzed according to the China Food Composition Tables by two independent investigators.¹⁹

Plans to promote participant retention and complete follow-up

No other strategy is planned to promote participant retention and complete follow-up.

Data management

A double entry and validation approach is planned for data management. The personal information and medical records of each participant will be kept in an individual folder and stored in a secure storage area at the School of Public Health, Peking University. We will return the results of the physical examination and laboratory biochemical measurement to

participants at the end of the exercise intervention/follow-up.

Confidentiality

The personal information and medical records of participants will be treated as strictly confidential. The research team will ensure the privacy and confidentiality of the personal information and medical records of participants and store all data in an encrypted database with all identifiers removed.

Plans for collection, laboratory evaluation and storage of biological specimens for genetic or molecular analysis in this trial/future use

At baseline and the end of exercise intervention, 6 mL overnight fasting venous blood will be collected by trained nurses from Peking University Third Hospital under the supervision of a paediatrician. The participants will also provide urine and faeces samples. All biospecimens will be immediately delivered to the clinical laboratory of Peking University Third Hospital and stored at -80°C until analysis.

Statistical methods

Statistical methods for primary and secondary outcomes

Statistical analysis will be performed using Stata/MP version 17.0 by statisticians from Peking University Health Science Center, which will not participate in participant enrollment, allocation, and exercise intervention. The intention-to-treatment analysis will be used, which includes all participants who complete the baseline assessment and is subject to

randomization and exercise intervention. Between-group differences in continuous and categorical variables at baseline will be analyzed using one-way analysis of variance and chi-square test, respectively. Within-group changes from baseline to the end of the intervention will be analyzed using paired t test for each group. Between-group differences of changes in the continuous variables will be analyzed using one-way ANOVA, one-way analysis of covariance, or Mann-Whitney rank sum test when appropriate. Correlations of changes in continuous variables will be analyzed using Pearson's or Spearman's correlation analysis when appropriate. Statistical significance is set at $\alpha = 0.05$.

Interim analyses

No interim analysis will be performed due to the short duration of this trial.

Methods for additional analyses (e.g. subgroup analyses)

In the sensitivity analysis, per protocol analysis is planned for participants that satisfactorily complete both the baseline assessment and the two-stage intervention.

Methods in analysis to handle protocol non-adherence and any statistical methods to handle missing data

Missing covariates will be estimated using the multiple imputation method.

Plans to give access to the full protocol, participant level-data and statistical code

Access to the full study protocol, participant-level data, and statical codes will be granted for

authorized research staff and will not be publicly available.

Oversight and monitoring

Composition of the coordinating centre and trial steering committee

The principal investigator (Prof. Ming Xu in the author list) has full responsibility for the design and conduct of the trial. The steering committee is responsible for agreeing on the final protocol and reviewing the progress of the trial and potential changes to the protocol.

Composition of the data monitoring committee, its role and reporting structure

The data monitoring committee is composed of pedestrians and statisticians and is responsible for assessing adherence to exercise intervention of each participant, monitoring and reporting any adverse events during exercise intervention.

Adverse event reporting and harms

The research team is responsible for reporting any adverse events. During exercise intervention days, a field team comprised of a paediatrician (or a physician) and a research staff from the site management organization will be sent to each school to deal with potential adverse events. The PE teachers and other elementary school teachers will assist in addressing potential anxiety and mood issues during exercise intervention. All severe adverse events will be reported to the ethics committee of Peking University Third Hospital, which will decide whether to continue or terminate this trial.

Frequency and plans for auditing trial conduct

The principal and lead investigators will meet biweekly to review the progress of the trial and audit the trial conduct.

Plans for communicating important protocol amendments to relevant parties (e.g. trial participants, ethical committees)

Any amendments to the study protocol must receive approval from the Ethics Committee of Peking University Third Hospital. All participants, their parents, PE teachers, and research staff will be promptly informed of any changes made to the study protocol. In case of a potential COVID-19 outbreak in the school, the intervention will be suspended and resumed in compliance with the school and government policies.

Dissemination plans

The results of this trial will be presented at domestic and international conferences and published in peer-reviewed academic journals. Progress and final reports will be submitted to the funding agency and regulatory bodies.

Conclusions

The present trial will provide a novel multilevel-coordinated exercise intervention targeting obese and overweight children. This study aims to yield quantitative and qualitative results which will shed light on the effectiveness of multi-level exercises with various intensities in promoting cardiometabolic fitness and reducing adiposity in overweight and obese

school-aged children. Moreover, this study will utilize a multi-omic approach and systematic epidemiological design to investigate the possible molecular mechanisms of the exercise intervention in modulating adiposity at multiple levels. The outcomes of this trial will offer clinical evidence for establishing bodyweight management guidelines for Chinese schools.

Trial status

This trial has been registered at ClinicalTrials.gov with the identifier NCT04984005.

Acknowledgements

Our special thanks go to the children, parents and physical education teachers who are participating in this study. We also acknowledge the ethics committee of Peking University Third Hospital. We thank Huawei Technologies Co., Ltd. for donating the smartwatch and the technical support of the Huawei Research platform to this study. We also thank Beijing BOE Health Technology Co., Ltd. for providing the equipment for the physical examination.

Author contributions

YZ, CL, TH, and MX contributed to conception and design of the study. JW, YL, and HY organized the database. YZ, CL, TC, CF, WW, YL, HW, YX, WZ, CR LC, XF, DF, HH, XY, and TH performed the study. YZ and CL wrote the first draft of the manuscript. All authors contributed to manuscript revision, read, and approved the submitted version.

Funding

This trial is funded by National Key Research and Development Program (grant numbers: 2020YFA0803803 and 2020YFA0803800) and CAMS Innovation Fund for Medical Sciences (grant number: 2021-I2M-5-003).

Availability of data and materials

Study data will be available from the principal investigator upon reasonable request and will not be made publicly available.

Ethics approval and consent to participate

This study has been approved by the ethics committee of Peking University Third Hospital (ethic approval number 2021-283-06).

Consent for publication

Not applicable.

Competing interests

The authors declare no competing interests.

References

1. Jebeile H, Kelly AS, O'Malley G, et al. Obesity in children and adolescents: epidemiology, causes, assessment, and management. *The lancet Diabetes & endocrinology* 2022;10(5):351-65. doi: 10.1016/s2213-8587(22)00047-x [published Online First: 2022/03/07]
2. Di Cesare M, Sorić M, Bovet P, et al. The epidemiological burden of obesity in childhood: a worldwide epidemic requiring urgent action. *BMC Med* 2019;17(1):212. doi: 10.1186/s12916-019-1449-8 [published Online First: 2019/11/26]

3. Gupta N, Goel K, Shah P, et al. Childhood obesity in developing countries: epidemiology, determinants, and prevention. *Endocr Rev* 2012;33(1):48-70. doi: 10.1210/er.2010-0028 [published Online First: 2012/01/14]
4. Liu Z, Gao P, Gao AY, et al. Effectiveness of a Multifaceted Intervention for Prevention of Obesity in Primary School Children in China: A Cluster Randomized Clinical Trial. *JAMA Pediatr* 2022;176(1):e214375. doi: 10.1001/jamapediatrics.2021.4375 [published Online First: 2021/11/09]
5. Pan XF, Wang L, Pan A. Epidemiology and determinants of obesity in China. *The lancet Diabetes & endocrinology* 2021;9(6):373-92. doi: 10.1016/s2213-8587(21)00045-0 [published Online First: 2021/05/23]
6. Flynn JT, Urbina EM, Brady TM, et al. Ambulatory Blood Pressure Monitoring in Children and Adolescents: 2022 Update: A Scientific Statement From the American Heart Association. *Hypertension (Dallas, Tex : 1979)* 2022;79(7):e114-e24. doi: 10.1161/hyp.0000000000000215 [published Online First: 2022/05/24]
7. Myette RL, Feber J, Blinder H, et al. Blood pressure variability in children with obesity and sleep-disordered breathing following positive airway pressure treatment. *Pediatr Res* 2022;92(3):810-15. doi: 10.1038/s41390-021-01841-8 [published Online First: 2021/11/18]
8. Chrysaidou K, Kotsis V, Chainoglou A, et al. Impact of ambulatory SBP and overweight on executive function performance in children and adolescents. *Journal of hypertension* 2020;38(6):1123-30. doi: 10.1097/hjh.0000000000002371 [published Online First: 2020/05/07]
9. Twig G, Tirosch A, Leiba A, et al. BMI at Age 17 Years and Diabetes Mortality in Midlife: A Nationwide Cohort of 2.3 Million Adolescents. *Diabetes care* 2016;39(11):1996-2003. doi: 10.2337/dc16-1203 [published Online First: 2016/10/14]
10. Twig G, Yaniv G, Levine H, et al. Body-Mass Index in 2.3 Million Adolescents and Cardiovascular Death in Adulthood. *N Engl J Med* 2016;374(25):2430-40. doi: 10.1056/NEJMoa1503840 [published Online First: 2016/04/14]
11. Martin A, Booth JN, Laird Y, et al. Physical activity, diet and other behavioural interventions for improving cognition and school achievement in children and adolescents with obesity or overweight. *Cochrane Database Syst Rev* 2018;3(3):Cd009728. doi: 10.1002/14651858.CD009728.pub4 [published Online First: 2018/03/03]
12. Cadenas-Sanchez C, Idoate F, Cabeza R, et al. Effect of a Multicomponent Intervention on Hepatic Steatosis Is Partially Mediated by the Reduction of Intermuscular Abdominal Adipose Tissue in Children With Overweight or Obesity: The EFIGRO Project. *Diabetes care* 2022;45(9):1953-60. doi: 10.2337/dc21-2440 [published Online First: 2022/09/01]
13. Wang Y, Wang K, Du M, et al. Maternal consumption of ultra-processed foods and subsequent risk of offspring overweight or obesity: results from three prospective cohort studies. *Bmj* 2022;379:e071767. doi: 10.1136/bmj-2022-071767 [published Online First: 2022/10/06]
14. Hargreaves D, Mates E, Menon P, et al. Strategies and interventions for healthy adolescent growth, nutrition, and development. *Lancet (London, England)* 2022;399(10320):198-210. doi: 10.1016/s0140-6736(21)01593-2 [published Online First: 2021/12/03]
15. Mead E, Brown T, Rees K, et al. Diet, physical activity and behavioural interventions for the treatment of overweight or obese children from the age of 6 to 11 years. *Cochrane Database Syst Rev* 2017;6(6):Cd012651. doi: 10.1002/14651858.Cd012651 [published Online First: 2017/06/24]

16. Chan AW, Tetzlaff JM, Gøtzsche PC, et al. SPIRIT 2013 explanation and elaboration: guidance for protocols of clinical trials. *Bmj* 2013;346:e7586. doi: 10.1136/bmj.e7586 [published Online First: 2013/01/11]
17. Pelliccia A, Sharma S, Gati S, et al. 2020 ESC Guidelines on sports cardiology and exercise in patients with cardiovascular disease. *European heart journal* 2021;42(1):17-96. doi: 10.1093/eurheartj/ehaa605 [published Online First: 2020/08/30]
18. Blair SN, Gibbons LW. Guidelines for exercise testing and prescription. *Journal of Cardiopulmonary Rehabilitation and Prevention* 1986;6(8):315-16.
19. Yang Y. China Food Composition Table 2002.
20. Zhang F, Wan Y, Zuo T, et al. Prolonged Impairment of Short-Chain Fatty Acid and L-Isoleucine Biosynthesis in Gut Microbiome in Patients With COVID-19. *Gastroenterology* 2022;162(2):548-61.e4. doi: 10.1053/j.gastro.2021.10.013 [published Online First: 2021/10/24]
21. Yuan M, Breitkopf SB, Yang X, et al. A positive/negative ion-switching, targeted mass spectrometry-based metabolomics platform for bodily fluids, cells, and fresh and fixed tissue. *Nat Protoc* 2012;7(5):872-81. doi: 10.1038/nprot.2012.024 [published Online First: 2012/04/14]
22. China BNHCotPsRo. WS/T 586—2018, Screening for overweight and obesity among school-age children and adolescents, 2018.

500 **Table 1.** Age- and sex-specific BMI cutoff points of childhood overweight and obesity.²²

	Boys		Girls	
Age, years	Overweight	Obesity	Overweight	Obesity
8.0 - < 8.5	17.8 - < 19.7	≥ 19.7	17.6 - < 19.4	≥ 19.4
8.5 - < 9.0	18.1 - < 20.3	≥ 20.3	18.1 - < 19.9	≥ 19.9
9.0 - < 9.5	18.5 - < 20.8	≥ 20.8	18.5 - < 20.4	≥ 20.4
9.5 - < 10.0	18.9 - < 21.4	≥ 21.4	19.0 - < 21.0	≥ 21.0
10.0 - < 10.5	19.2 - < 21.9	≥ 21.9	19.5 - < 21.5	≥ 21.5
10.5 - < 11.0	19.6 - < 22.5	≥ 22.5	20.0 - < 22.1	≥ 22.1
11.5 - 12.0	19.9 - < 23.0	≥ 23.0	20.5 - < 22.7	≥ 22.7

501

502

503 **Figure 1** Schedule of enrollment, allocation, interventions, and assessments.

	Enrolment	Allocation	Run-in		Intervention			Follow-up		
Timepoint, week	-	-	1	2	4	8	12	24	36	48
Enrolment										
Informed consent	x									
Eligibility screen	x									
Baseline assessment	x									
Allocation		x								
Intervention										
Light-intensity exercise			x	x	x	x	x			
Moderate-intensity exercise			x	x	x	x	x			
Intermittent vigorous exercise			x	x	x	x	x			
Assessment										
Adiposity		x			x	x	x	x	x	x
Sexual hormones		x			x	x	x	x	x	x
Body compositions		x			x	x	x	x	x	x
Cardiopulmonary function		x			x	x	x	x	x	x
Biochemical measurement		x			x	x	x	x	x	x
Exercise performance		x					x			x
Hepatic hemodynamic index		x					x			x
Bone mineral density		x					x			x
Inflammatory biomarkers		x					x			x
Thyroid function		x					x			x
Oxidative stress biomarkers		x					x			x
Oral and gut microbiota		x					x			x
Blood and faecal metabolomics		x					x			x
Attention performance		x					x			x
Sleep quality		x					x			x
Diets and nutrients		x					x			x
Whole genome sequencing		x					x			x
Viromics		x					x			x
Proteomics		x					x			x
Volatile organic compounds in breath		x					x			x
Anthropometric measurement of the parents		x					x	x		
Whole genome sequencing of the parents;								x		
Disease history of the parents								x		

504