

1 **RESEARCH ARTICLE**

2 **Global seasonal activities of respiratory syncytial virus before the COVID-19 pandemic:**
3 **a systematic review**

4 Songwei Shan^{1,2}, Weixin Zhang¹, Huizhi Gao¹, Pei-Yu Huang¹, Zhanwei Du^{1,2}, Yuan Bai^{1,2},
5 Yiu-Chung Lau^{1,2}, Dongxuan Chen^{1,2}, Eric HY Lau^{1,2}, Joshua Nealon¹, Peng Wu^{1,2}

6

7 ¹World Health Organization Collaborating Centre for Infectious Disease Epidemiology and
8 Control, School of Public Health, Li Ka Shing Faculty of Medicine, The University of Hong
9 Kong, Hong Kong SAR, China.

10 ²Laboratory of Data Discovery for Health, Hong Kong Science and Technology Park, Hong
11 Kong SAR, China.

12

13

14 Corresponding author: Peng Wu, School of Public Health, The University of Hong Kong,
15 email: pengwu@hku.hk, Tel: 3917 6746

16

17 Running head: Global RSV seasonality

18

19 Word count (Abstract): 248

20 Word count (main text): 3,356

21 Keywords: activity, respiratory syncytial virus, seasonality, global

22 ABSTRACT

23 **Background:** Varied seasonal patterns of respiratory syncytial virus (RSV) have been
24 reported worldwide. We aimed to review the patterns of RSV activity globally before the
25 COVID-19 pandemic and to explore factors potentially associated with RSV seasonality.

26 **Methods:** We conducted a systematic review on articles identified in PubMed reporting RSV
27 seasonality based on data collected before 1 January 2020. Information on the timing of the
28 start, peak, and end of an RSV season, study location, study period, and details in study
29 methods were extracted. RSV seasonal patterns were examined by geographic location,
30 calendar month, analytic method and meteorological factors including temperature and
31 absolute humidity. Correlation and regression analyses were conducted to explore the
32 relationship between RSV seasonality and study methods and characteristics of study
33 locations.

34 **Results:** RSV seasons were reported in 209 articles published in 1973-2023 for 317 locations
35 in 77 countries. Variations were identified in types of data, data collection and analytical
36 methods across the studies. Regular RSV seasons were similarly reported in countries in
37 temperate regions, with highly variable seasons identified in subtropical and tropical
38 countries. Durations of RSV seasons were relatively longer in subtropical and tropical
39 regions than from temperate regions. Longer durations of RSV seasons were associated with
40 a higher daily average mean temperature and daily average mean absolute humidity.

41 **Conclusions:** The global seasonal patterns of RSV provided important information for
42 optimizing interventions against RSV infection. Heterogeneity in study methods highlighted
43 the importance of developing and applying standardized approaches in RSV surveillance and
44 data reporting.

45 INTRODUCTION

46 Respiratory syncytial virus (RSV) is a leading cause of hospitalization for acute lower
 47 respiratory tract infection (LRTI) in young children less than 5 years old ¹⁻³, as well as an
 48 increasingly recognized important pathogen in older adults (aged ≥ 65) ^{4,5}. Globally, RSV
 49 was estimated to cause nearly 25 million LRTI episodes and more than 70,000 deaths every
 50 year ⁶. In addition to Palivizumab, a monoclonal antibody recommended for high-risk infants
 51 ^{7,8} providing a short-term protection against RSV-associated severe infections ^{9,10}, a new
 52 monoclonal antibody nirsevimab with a prolonged protection ^{11,12} has been approved by the
 53 EU and the UK ¹³. Recently the first RSV vaccine (Arexvy, RSVPreF3 from GSK) approved
 54 for the use in older adults and several other vaccine candidates in their final stage of testing
 55 demonstrated promising efficacy in preventing RSV-associated LRTI and severe infections
 56 ^{14,15}.

58 Activities of RSV varied temporally and geographically. In temperate regions, the highest
 59 viral activities have been observed during winter months ^{16,17} with similar intensities across
 60 different years. Two-year cycles of RSV activity have been reported for some countries in
 61 Northern Europe, with a major epidemic wave in winter followed by a minor one in spring
 62 next year ¹⁸. In subtropical and tropical regions, seasonal patterns of RSV were more diverse
 63 and less clear than those reported for temperate countries ¹⁹. Knowledge about the seasonality
 64 of RSV, can guide the formulation of prevention strategies and the planning of public health
 65 resources. In particular, accurate estimates of the RSV season start and its duration can help
 66 to determine the optimal timing of initial doses and optimal dosing interval for prophylactic
 67 monoclonal antibodies and vaccination schedules.

68

We conducted a systematic review on RSV seasonality to characterize the global patterns of RSV activity by summarizing the reported start, peak and end of RSV seasons worldwide and describe approaches applied in analyzing the seasonality. We aimed to provide a better understanding of the seasonal patterns of RSV circulation to inform policy and decisions in prevention, diagnosis, and treatment of RSV infections.

METHODS

Literature search strategy and selection criteria

This systematic review was conducted following the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines²⁰. We focused on studies in which seasonal patterns of RSV were characterized by providing the time of the start, peak and/or end of a season based on laboratory-confirmed RSV infection data. We searched published articles from PubMed on January 17, 2023 using the combination of following search items (1 and 2) in “Title/Abstract” (Figure 1):

1. “RSV” or “Respiratory Syncytial Virus”
2. “seasonality” or “season*”

Two authors (S.S. and W.Z.) performed the literature search (#1 and #2) and screened retrieved articles independently. Full-text screening was conducted after title/abstract screening. Two authors (S.S. and W.Z.) independently identified articles for inclusion. Any disagreement in article selection and data retrieval was resolved by a third author (P.W.).

Eligible articles were studies reporting the seasonality of RSV based on monthly or weekly laboratory-confirmed RSV infections collected for a specified place, i.e., a city, a province or a country, for three years or longer. We excluded articles if: 1) the full text was not available

or not in English, 2) the analysis was based on less than three years' data, 3) the average number of reported RSV infections per year in a study was less than 50, 4) the time of the start, peak and/or end of RSV seasons was not specified in a calendar month or week. If an investigation on RSV seasonality covered the time period beyond 2019, the study period before the COVID-19 pandemic ≥ 3 years would be included in our analysis, and the RSV data collected during the pandemic would be excluded. Articles without original data analysis on RSV activities, such as reviews, commentaries, letters, were excluded, but we reviewed the references of these articles for additional studies to include for screening.

Data extraction and analysis

Two authors (S.S. and W.Z.) independently extracted data into a standardized data extraction form. Data fields included the timing (start, peak, and end) of an RSV season, study location (city, region or country), study period, information on study characteristics (such as the source of cases, case definitions, laboratory testing methods and analytic methods applied to determine RSV seasons). We classified the sources of study cases into outpatient, inpatient, outpatient and inpatient, case definitions into influenza-like-illness (ILI) or acute respiratory infection (ARI), severe acute respiratory infection (SARI) and acute LRTI and grouped laboratory testing methods for RSV confirmation into virus detection, antibody detection, antigen detection, and nucleic acid detection.

One publication might contain several different investigations on seasonal patterns of RSV, either based on different data sources, or based on the same dataset but with different analytic methods. If more than one estimate on timing of RSV season was reported in a single publication, using different data sources or methods, all were extracted and included in our analyses. It was common that one investigation on RSV seasonality was conducted based on

more than one type of case source, case definition and/or testing method, and all the methodological details were extracted and documented for analyses. Analytic methods used by different studies to determine the RSV seasons were classified as qualitative methods and quantitative methods based on whether the analysis on RSV seasonality was carried out using a statistical or mathematical approach, and the quantitative methods were further divided into three groups, threshold-based, coverage-based and model-based (*Supplementary Materials*).

In addition to the study-specific information, we also documented the latitude, daily average mean temperature and daily average mean relative humidity for all study locations reported in each investigation using Google Geocoding API ²¹ and R package GSOR ²²(*Supplementary materials*). The extracted daily average mean temperature and relative humidity data were used to derive the daily average mean absolute humidity for each location. We also extracted the geoclimatic information of each study site, and classified the investigations of RSV seasonality into three groups based on the latitude of the study sites, temperate zone (latitude > 35° or latitude < -35°), subtropical zone (23.5° < latitude ≤ 35° or -35° ≤ latitude < -23.5°) and tropical zone (-23.5° ≤ latitude ≤ 23.5°).

We described the reported RSV seasons as a function of geographical location, meteorological factors and characteristics of study design, and explored potential associations using Pearson correlation analyses and Kendall's rank correlation between the duration of the season and the latitude, daily average mean temperature and daily average mean absolute humidity of study sites as well as the analytic method and climate zone. Finally, linear regression models were applied to explore the independent associations between the duration of RSV seasons and the abovementioned factors as covariates. The statistical analyses were

conducted using R software Version 4.0.3 (R Foundation for Statistical Computing, Vienna, Austria).

RESULTS

We identified 2,510 citations from the PubMed through the keyword search and additional 19 citations from other sources. In total, 209 publications were included after screening based on the inclusion and exclusion criteria (Figure 1) and 552 investigations of RSV seasonality were identified and included in analyses.

The identified studies were published between 1973 and 2023, with the majority (147/209, 70%) from 2010 onwards (Supplementary Figure 1). RSV seasonality was reported for 317 distinct study sites in 77 countries from Asia, Europe, North America, South America, Oceania and Africa (Figure 2). Most investigations (342/552, 62%) were conducted with a study period of 3-6 years (Supplementary Figure 1), with the longest one from Malaysia using surveillance data over 27 years²³. Among the identified publications, more than half (124/209, 59%) were from temperate regions (latitude > 35° or latitude < -35°), followed by 31% (65/209) and 22% (47/209) in subtropical (23.5° < latitude ≤ 35° or -35° ≤ latitude < -23.5°) and tropical (-23.5° ≤ latitude ≤ 23.5°) regions, respectively (Table 1, Supplementary Table 1).

The information on the case source, case definition, and laboratory testing methods was missing in some investigations (case source: 260/552, 47%; case definition: 310/552, 56%; testing method: 127/552, 23%). Among those with information of case source and case definition provided, the majority of investigations (152/292, 52%) identified RSV cases from inpatients, and ARI or ILI (150/242, 62%) was the most frequently used syndromic

definitions to screen patients for RSV testing, and antigen (325/552, 59%) and nucleic acid detections (323/552, 59%) were the most commonly used laboratory testing methods in more recent publications (Table 1, Supplementary Table 1, Supplementary Figures 2-3). The majority of investigations (373/552, 68%) applied quantitative methods to determine the time of the start, peak and/or end of seasons, and among these studies, over half of the investigations used the threshold-based method (205/373, 55%) (Supplementary Tables 2-3).

The majority of the investigations were conducted using data collected from locations in the Northern Hemisphere (NH) and indicated one RSV season every year (Figure 2). A small number of investigations (11/552, 2%) reported two RSV peaks identified in one year (Supplementary Table 4) while 18 (3%) investigations indicated unique RSV seasonal patterns repeated every 2 years (Supplementary Table 5). Among study sites reporting one RSV season annually, locations in temperate countries were fairly consistent in the time of the start, peak and end of RSV seasons in either the NH or Southern Hemisphere (SH) (Figure 3). In temperate countries, RSV activities generally started from late autumn or early winter, i.e., October-December in NH, May-June in SH, peaked in winter (December-February in NH, and June to July in SH), and ended in late winter or spring (February-May in NH, and July-September in SH). (Figure 3, Supplementary Figure 4). Larger variations were observed in the start, peak and end month of RSV seasons in subtropical and tropical countries at a lower latitude, with high RSV activities observed in July/August-November in tropical locations (Supplementary Figures 5-6). The durations of RSV seasons were relatively shorter in study sites in the temperate region (median: 3.97 months, interquartile range (IQR): 3.00-5.00) compared with the estimates from subtropical (4.00, 3.52-6.14) and tropical countries (5.00, 4.00-6.00) (Figure 2, Supplementary Figure 6).

The Pearson correlation analysis indicated an overall negative association between the duration of the RSV season (in months) and the latitude both in the NH ($p < 0.001$) and the SH ($p = 0.002$) (Supplementary Table 7), and the statistical significance in the association was only indicated with the estimates derived from quantitative methods ($p < 0.001$). Positive associations were indicated in the correlation analyses between the daily average mean temperature (correlation coefficient: 0.33; 95% CI: 0.22, 0.44) and absolute humidity (0.34; 0.22, 0.45) of the study locations and the durations of RSV season estimated by the quantitative methods ($p < 0.001$) but not for those estimated by the qualitative methods (Supplementary Table 7). Durations of RSV seasons appeared to be longer in tropical /subtropical regions than the estimates from temperate regions, and the estimated RSV seasons were longer from the threshold-based methods, but shorter from the coverage-based method (Table 2, Supplementary Table 8).

In the main analysis with the linear regression model, we identified that climatic zone appeared to be independently associated with the estimated duration of RSV seasons, and analytic method used in estimating RSV seasons, annual daily average mean absolute humidity at the study location could also explain some variations (Adjusted $R^2 = 0.33$ for estimates from quantitative approaches, 0.25 for all estimates) in the estimated durations of RSV seasons (Table 2, Supplementary Table 8). Considering the close intercorrelations between climatic zone, latitude and temperature, we included the climatic zone (Table 2, Supplementary Table 8), latitude (Supplementary Tables 9-10) and daily average mean temperature (Supplementary Tables 11-12) of study locations into different linear regression models separately. The duration of RSV season would decrease by 0.041 (95% CI: 0.017, 0.064) months with every increase of 1 degree in latitude, and increase by 0.058 (95% CI: 0.0077, 0.11) months with every increase of 1 degree in daily average mean temperature.

Durations of RSV seasons estimated for the subtropical and tropical locations were on average 1.06 (95% CI: 0.45, 1.66) and 1.65 (95% CI: 0.77, 2.54) months longer than for the study sites in the temperate region. Compared to the coverage-based methods, the threshold-based methods tended to produce longer estimates of the duration of RSV season by 1.43 (95% CI: 0.96, 1.90) months. Overall, the duration of RSV season prolonged by 0.013 (95% CI: -0.0048, 0.030) months with every 1 g/m³ increase in absolute humidity at the study location (Table 2).

DISCUSSION

RSV seasonal patterns have been studied widely in particular since the last decade, with most of the data reported from temperate countries in the Northern Hemisphere. Our review indicated that largely similar seasonal patterns of RSV were observed in locations from the same climatic region either in the NH or the SH. In temperate regions, RSV activities primarily showed regular annual patterns starting in winter seasons, and the estimated durations of RSV season were relatively shorter than the reported from tropical and subtropical regions where prolonged RSV circulations were often more than half a year with greatly varied times of season start and/or peak.

Human infections of RSV are likely driven by two major antigenic groups, A and B²⁴ between which cross-immunity has been suggested^{25,26}. However, RSV typing is often not included in most of the established surveillance systems²⁷. The increasing susceptibility in infants due to the loss of protection from waning maternal antibodies leads to a higher risk of infection particularly in young children with high-risk conditions^{1,28,29}, potentially driving the transmission of RSV in the population³⁰ although the virus can infect people at any age including older adults^{31,32}. RSV shows comparable environmental survivability and

transmissibility to influenza viruses with a similar estimate of the basic reproduction number (R_0) and household second attack rate³³. Increasing RSV activities were commonly reported in cold seasons starting from late autumn through to early spring in temperate regions, and in rainy seasons in tropical/subtropical areas, and the associations between RSV activities and environmental factors including temperature, relative/absolute humidity, rainfall were inconsistently reported from studies conducted in tropical areas³⁴⁻³⁶.

RSV seasonality characterised in different studies appeared to be associated with the approaches used in data collection and analysis¹⁹. Previous research demonstrated that the timing of RSV seasons determined with laboratory surveillance data might be different from the results based on hospitalized patient data although the two estimates were highly correlated³⁷. The improvement of laboratory testing methods could also affect the estimates of RSV seasons^{38,39}. Our review showed that estimates of the timing of RSV seasons were correlated with the analytic approaches used in the studies, with shorter durations of RSV seasons estimated from “coverage-based method” (defined as the shortest time period when the majority of confirmed (severe) RSV infections occurred) than the estimates from the threshold-based methods which although were highly dependent on the often arbitrarily selected threshold of varied measures for RSV activities, such as proportion of virus detections or patients presenting ILI/ARI symptoms⁴⁰. Therefore, prudent comparison and interpretation of RSV seasons estimated from different studies or with different approaches is necessary.

RSV surveillance is often established based on the existing routine surveillance on influenza, more commonly seen in developed economies including the US⁴¹, Canada⁴² and some European countries⁴³. The WHO initiated a pilot RSV surveillance scheme by leveraging the

current Global Influenza Surveillance and Response System covering over 100 countries to explore possible surveillance strategies including case definitions, laboratory methods and target population, etc.⁴⁴. Current RSV surveillance largely applied similar syndromic case definitions as those used for influenza surveillance, i.e., sampling patients showing influenza-like-illness and testing for the virus, while the validity of using such case definitions for RSV surveillance has been questioned⁴⁵. In some countries/areas RSV surveillance was not conducted throughout a year but only during restricted period(s) following the influenza surveillance protocol given the regular annual influenza seasons in those locations^{17,46}. However, this might lead to an underreporting of RSV infections during the non-surveillance periods⁴⁷.

During the COVID-19 pandemic, circulation of respiratory viruses including RSV changed considerably worldwide largely due to implementation of non-pharmaceutical interventions and changes in human behaviours in response to the pandemic^{48,49}. RSV remained at a low level in many countries in 2020 when public health and social measure were widely adopted globally to contain the spread of SARS-CoV-2, but caused surges of infections with a large number of severe cases admitted into hospital in some countries when some interventions were relaxed from 2021 onwards^{50,51}, including the US during 2021-2023⁵², some geographically separated locations in Australia in 2020-21⁵³ and the UK in 2021-22⁵⁴. The inter-seasonal viral activities were perhaps related to the increased susceptibility in the population due to the lack of exposure to RSV during the previous year in response to COVID-19⁵⁵.

Monoclonal antibodies approved for prophylactic therapy provide immediate protection of high-risk individuals particularly young children against RSV-associated severe infections^{7,13}.

Five monthly injections of palivizumab and a single dose of nirsevimab can substantially reduce the risk of hospitalizations due to RSV in individuals receiving the therapy throughout RSV seasons especially in temperate regions where RSV seasons would normally not be longer than 5 months. The duration of protection provided by RSV vaccines has not been fully determined although promising efficacy for short-term prevention from infections was demonstrated in clinical trials⁵⁶⁻⁵⁸. The effectiveness of the abovementioned pharmaceutical interventions for high-risk population is determined by the duration of protection of the monoclonal antibodies and the vaccines, the timing of administration and RSV activities in specific locations. Ideally these preventive measures should be administered before an RSV season starts, and the protection can maintain throughout the season, likely during November - April, and May - October in temperate regions of the NH and the SH, respectively. However great challenges in implementing these efficacious interventions would be seen in locations where there were irregular seasons particularly with prolonged viral circulations⁵⁹⁻⁶¹. Based on the reported seasonal patterns in locations from northern temperate regions in this review, the Studies incorporating epidemiologic analysis on RSV seasonality into the effectiveness and cost-effectiveness of these RSV preventive strategies are urgently needed⁶².

There are several limitations in this study. First, data of RSV seasonality that were published and included in the review were skewed to high-income countries with well-established RSV surveillance systems. These countries are largely in temperate regions where RSV circulation might be different from the underrepresented middle- and low-income countries due to varied population structure, contact patterns and geoclimatic features⁶³. Second, RSV antigenic information could not be incorporated into the current analysis of seasonality due to the lack of data. Expanding RSV surveillance by collecting antigenic and genetic data is urgently needed to improve our understanding of the transmission dynamics of RSV locally and

globally⁶⁴. Third, except for the analytic method, we could not examine potential associations between RSV seasonality and other study characteristics such as case sources, case definitions, and laboratory methods due to the lack of information and non-standardized laboratory protocols used for RSV surveillance, or explore the association between RSV seasonality and population structure particularly birth rate given the lack of data although a high disease burden of RSV identified in young children⁶⁵⁻⁶⁷ implied a possible driving force from the young population on RSV activities. Lastly, the reported data of seasonality in this review did not allow us to investigate the RSV-associated disease burden occurred during the characterised RSV seasons which is a critical metric for evaluation of the health impact related to RSV circulation, particularly when there were large uncertainties in the start and the end of RSV seasons in a specific location and the infection risk is less likely to be homogeneous in a population over the season.

CONCLUSIONS

This study provided a global overview of RSV seasonality before the COVID-19 pandemic. The seasonal patterns illustrated in this review may inform optimal use of preventive and therapeutic interventions against RSV infection particularly in temperate regions of the NH and SH. Considerable variations in methods used by different studies highlighted the importance of developing and applying standardized approaches in RSV surveillance and data reporting. Further research is needed to understand the underlying mechanisms driving RSV seasonal patterns locally and globally.

340

341 **ACKNOWLEDGEMENTS**

342 The authors thank Prof. Julian W Tang, and Dr. Eeva Broberg for kindly sharing the data of
343 RSV seasonality from their publications. The authors also thank Stella SW Lai for technical
344 assistance.

345

346 **FUNDING**

347 This project was supported by the Health and Medical Research Fund (project no. 19200861)
348 from Food Bureau of Hong Kong SAR Government, and the AIR@InnoHK administered by
349 the Innovation and Technology Commission of Hong Kong SAR Government. The funding
350 bodies had no role in study design, data collection and analysis, preparation of the manuscript,
351 or the decision to publish.

352

353 **AUTHOR CONTRIBUTIONS**

354 All authors meet the International Committee of Medical Journal Editors criteria for
355 authorship. SS, PW, BY, DZ conceived and designed the study; SS, WZ, HG, PH collected
356 and extracted data; SS, PW conducted the data analysis and wrote the first draft of the
357 manuscript. All authors contributed to the interpretation of the results and revision of the
358 manuscript, and have approved the final version of the manuscript.

359

360 **DECLARATION OF INTERESTS**

361 JN was previously employed by and owns shares in Sanofi. The authors report no other
362 potential conflicts of interest.

363

REFERENCES

1. Li Y, Wang X, Blau DM, et al. Global, regional, and national disease burden estimates of acute lower respiratory infections due to respiratory syncytial virus in children younger than 5 years in 2019: a systematic analysis. *Lancet* 2022; **399**(10340): 2047-64.
2. Nair H, Nokes DJ, Gessner BD, et al. Global burden of acute lower respiratory infections due to respiratory syncytial virus in young children: a systematic review and meta-analysis. *Lancet* 2010; **375**(9725): 1545-55.
3. Shi T, McLean K, Campbell H, Nair H. Aetiological role of common respiratory viruses in acute lower respiratory infections in children under five years: A systematic review and meta-analysis. *J Glob Health* 2015; **5**(1): 122-31.
4. Falsey AR, McElhaney JE, Beran J, et al. Respiratory syncytial virus and other respiratory viral infections in older adults with moderate to severe influenza-like illness. *J Infect Dis* 2014; **209**(12): 1873-81.
5. Falsey AR, Walsh EE. Respiratory syncytial virus infection in elderly adults. *Drugs Aging* 2005; **22**(7): 577-87.
6. Troeger C, Forouzanfar M, Rao PC, et al. Estimates of the global, regional, and national morbidity, mortality, and aetiologies of lower respiratory tract infections in 195 countries: a systematic analysis for the Global Burden of Disease Study 2015. *Lancet Infect Dis* 2017; **17**(11): 1133-61.
7. Brady MT, Byington CL, Davies HD, et al. Updated Guidance for Palivizumab Prophylaxis Among Infants and Young Children at Increased Risk of Hospitalization for Respiratory Syncytial Virus Infection. *Pediatrics* 2014; **134**(2): E620-E38.
8. Reeves RM, van Wijhe M, Lehtonen T, et al. A Systematic Review of European Clinical Practice Guidelines for Respiratory Syncytial Virus Prophylaxis. *J Infect Dis* 2022; **226**(Suppl 1): S110-S6.
9. Hammitt LL, Dagan R, Yuan Y, et al. Nirsevimab for Prevention of RSV in Healthy Late-Preterm and Term Infants. *N Engl J Med* 2022; **386**(9): 837-46.

10. Simoes EAF, Madhi SA, Muller WJ, et al. Efficacy of nirsevimab against respiratory syncytial virus lower respiratory tract infections in preterm and term infants, and pharmacokinetic extrapolation to infants with congenital heart disease and chronic lung disease: a pooled analysis of randomised controlled trials. *Lancet Child Adolesc Health* 2023; **7**(3): 180-9.
11. Garegnani L, Styrmisdottir L, Roson Rodriguez P, Escobar Liquitay CM, Esteban I, Franco JV. Palivizumab for preventing severe respiratory syncytial virus (RSV) infection in children. *Cochrane Database Syst Rev* 2021; **11**(11): CD013757.
12. Null D, Bimle C, Weisman L, et al. Palivizumab, a humanized respiratory syncytial virus monoclonal antibody, reduces hospitalization from respiratory syncytial virus infection in high-risk infants. *Pediatrics* 1998; **102**(3): 531-7.
13. Keam SJ. Nirsevimab: First Approval. *Drugs* 2023; **83**(2): 181-7.
14. Mazur NI, Terstappen J, Baral R, et al. Respiratory syncytial virus prevention within reach: the vaccine and monoclonal antibody landscape. *Lancet Infect Dis* 2023; **23**(1): E2-E21.
15. Valero MV. Green light from US regulatory agency for GSK's jab caps off a decades-long quest. 2023. <https://www.nature.com/articles/d41586-023-01529-5> (accessed May 6 2023).
16. Li Y, Reeves RM, Wang X, et al. Global patterns in monthly activity of influenza virus, respiratory syncytial virus, parainfluenza virus, and metapneumovirus: a systematic analysis. *Lancet Glob Health* 2019; **7**(8): e1031-e45.
17. Obando-Pacheco P, Justicia-Grande AJ, Rivero-Calle I, et al. Respiratory Syncytial Virus Seasonality: A Global Overview. *J Infect Dis* 2018; **217**(9): 1356-64.
18. Broberg EK, Waris M, Johansen K, Snacken R, Penttinen P, European Influenza Surveillance N. Seasonality and geographical spread of respiratory syncytial virus epidemics in 15 European countries, 2010 to 2016. *Euro Surveill* 2018; **23**(5).
19. Staadegaard L, Caini S, Wangchuk S, et al. Defining the seasonality of respiratory syncytial virus around the world: National and subnational surveillance data from 12 countries. *Influenza Other Respir Viruses* 2021; **15**(6): 732-41.
20. Moher D, Shamseer L, Clarke M, et al. Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015 statement. *Syst Rev* 2015; **4**(1): 1.

21. Geocoding API. <https://developers.google.com/maps/documentation/geocoding>.
22. GSODR: Global Surface Summary of the Day ('GSOD') Weather Data Client. <https://cran.r-project.org/web/packages/GSODR/index.html>.
23. Khor CS, Sam IC, Hooi PS, Quek KF, Chan YF. Epidemiology and seasonality of respiratory viral infections in hospitalized children in Kuala Lumpur, Malaysia: a retrospective study of 27 years. *Bmc Pediatr* 2012; **12**.
24. Sullender WM. Respiratory syncytial virus genetic and antigenic diversity. *Clin Microbiol Rev* 2000; **13**(1): 1-15, table of contents.
25. Bhattacharyya S, Gesteland PH, Korgenski K, Bjornstad ON, Adler FR. Cross-immunity between strains explains the dynamical pattern of paramyxoviruses. *P Natl Acad Sci USA* 2015; **112**(43): 13396-400.
26. White LJ, Waris M, Cane PA, Nokes DJ, Medley GF. The transmission dynamics of groups A and B human respiratory syncytial virus (hRSV) in England & Wales and Finland: seasonality and cross-protection. *Epidemiol Infect* 2005; **133**(2): 279-89.
27. Chadha M, Hirve S, Bancej C, et al. Human respiratory syncytial virus and influenza seasonality patterns-Early findings from the WHO global respiratory syncytial virus surveillance. *Influenza Other Resp* 2020; **14**(6): 638-46.
28. Hall CB, Weinberg GA, Iwane MK, et al. The burden of respiratory syncytial virus infection in young children. *N Engl J Med* 2009; **360**(6): 588-98.
29. Meissner HC. Selected populations at increased risk from respiratory syncytial virus infection. *Pediatr Infect Dis J* 2003; **22**(2): S40-S5.
30. Hogan AB, Glass K, Moore HC, Anderssen RS. Exploring the dynamics of respiratory syncytial virus (RSV) transmission in children. *Theor Popul Biol* 2016; **110**: 78-85.
31. Savic M, Penders Y, Shi T, Branche A, Pircon JY. Respiratory syncytial virus disease burden in adults aged 60 years and older in high-income countries: A systematic literature review and meta-analysis. *Influenza Other Resp* 2023; **17**(1).

32. Shi T, Denouel A, Tietjen AK, et al. Global Disease Burden Estimates of Respiratory Syncytial Virus-Associated Acute Respiratory Infection in Older Adults in 2015: A Systematic Review and Meta-Analysis. *J Infect Dis* 2020; **222**: S577-S83.
33. Leung NHL. Transmissibility and transmission of respiratory viruses. *Nat Rev Microbiol* 2021; **19**(8): 528-45.
34. Moriyama M, Hugentobler WJ, Iwasaki A. Seasonality of Respiratory Viral Infections. *Annu Rev Virol* 2020; **7**: 83-101.
35. Paynter S. Humidity and respiratory virus transmission in tropical and temperate settings. *Epidemiol Infect* 2015; **143**(6): 1110-8.
36. Tang JLW, Loh TP. Correlations between climate factors and incidence-a contributor to RSV seasonality. *Rev Med Virol* 2014; **24**(1): 15-34.
37. Hampp C, Asal N, Lipowski E, et al. Validity of Laboratory-based Surveillance for Detection of Respiratory Syncytial Virus Seasons. *Am J Epidemiol* 2013; **177**(8): 841-51.
38. Ambrose CS, Steed LL, Brandon M, Frye K, Olajide IR, Thomson G. National and regional modeling of distinct RSV seasonality thresholds for antigen and PCR testing in the United States. *J Clin Virol* 2019; **120**: 68-77.
39. Midgley CM, Haynes AK, Baumgardner JL, et al. Determining the Seasonality of Respiratory Syncytial Virus in the United States: The Impact of Increased Molecular Testing. *J Infect Dis* 2017; **216**(3): 345-55.
40. Rose EB, Wheatley A, Langley G, Gerber S, Haynes A. Respiratory Syncytial Virus Seasonality - United States, 2014-2017. *Mmwr-Morbid Mortal W* 2018; **67**(2): 71-6.
41. National Respiratory and Enteric Virus Surveillance System. <https://www.cdc.gov/surveillance/nrevss/index.html>.
42. Flu (influenza): FluWatch surveillance - Canada.ca. <https://www.canada.ca/en/public-health/services/diseases/flu-influenza/influenza-surveillance.html>.
43. Surveillance Atlas of Infectious Diseases. <https://atlas.ecdc.europa.eu/public/index.aspx>.

44. Broor S, Campbell H, Hirve S, et al. Leveraging the Global Influenza Surveillance and Response System for global respiratory syncytial virus surveillance-opportunities and challenges. *Influenza Other Resp* 2020; **14**(6): 622-9.
45. Hirve S, Crawford N, Palekar R, Zhang WQ, Grp WRS. Clinical characteristics, predictors, and performance of case definition-Interim results from the WHO global respiratory syncytial virus surveillance pilot. *Influenza Other Resp* 2020; **14**(6): 647-57.
46. Teirlinck AC, Broberg EK, Berg AS, et al. Recommendations for respiratory syncytial virus surveillance at the national level. *Eur Respir J* 2021; **58**(3).
47. Mollers M, Barnadas C, Broberg EK, et al. Current practices for respiratory syncytial virus surveillance across the EU/EEA Member States, 2017. *Euro Surveill* 2019; **24**(40).
48. Baker RE, Park SW, Yang W, Vecchi GA, Metcalf CJE, Grenfell BT. The impact of COVID-19 nonpharmaceutical interventions on the future dynamics of endemic infections. *Proc Natl Acad Sci U S A* 2020; **117**(48): 30547-53.
49. Oh DY, Buda S, Biere B, et al. Trends in respiratory virus circulation following COVID-19-targeted nonpharmaceutical interventions in Germany, January - September 2020: Analysis of national surveillance data. *Lancet Reg Health Eur* 2021; **6**: 100112.
50. Bozzola E, Barni S, Villani A. Respiratory Syncytial Virus Pediatric Hospitalization in the COVID-19 Era. *Int J Environ Res Public Health* 2022; **19**(23).
51. Fairbank R. RSV wave hammers hospitals - but vaccines and treatments are coming. *Nature* 2022.
52. Lindsey Wang PBD, Nathan A Berger, David C Kaelber, Nora D Volkow, Rong Xu. Disruption in seasonality, patient characteristics and disparities of respiratory syncytial virus infection among young children in the US during and before the COVID-19 pandemic: 2010-2022. medRxiv; 2022.
53. Eden JS, Sikazwe C, Xie RP, et al. Off-season RSV epidemics in Australia after easing of COVID-19 restrictions. *Nat Commun* 2022; **13**(1).
54. Bardsley M, Morbey RA, Hughes HE, et al. Epidemiology of respiratory syncytial virus in children younger than 5 years in England during the COVID-19 pandemic, measured by laboratory,

clinical, and syndromic surveillance: a retrospective observational study. *Lancet Infect Dis* 2023; 23(1): 56-66.

55. Garg I, Shekhar R, Sheikh A, Pal S. Impact of COVID-19 on the Changing Patterns of Respiratory Syncytial Virus Infections. *Infect Dis Rep* 2022; 14(4): 558-68.

56. Falsey AR, Williams K, Gymnopoulos E, et al. Efficacy and Safety of an Ad26.RSV.preF-RSV preF Protein Vaccine in Older Adults. *N Engl J Med* 2023; 388(7): 609-20.

57. Kampmann B, Madhi SA, Munjal I, et al. Bivalent Prefusion F Vaccine in Pregnancy to Prevent RSV Illness in Infants. *N Engl J Med* 2023; 388(16): 1451-64.

58. Walsh EE, Perez Marc G, Zareba AM, et al. Efficacy and Safety of a Bivalent RSV Prefusion F Vaccine in Older Adults. *N Engl J Med* 2023; 388(16): 1465-77.

59. Bauman J, Eggleston M, Oquist N, Malinoski F. Respiratory syncytial virus: Seasonal data for regions of Florida and implications for palivizumab. *South Med J* 2007; 100(7): 669-76.

60. Hsu CH, Lin CY, Chi H, et al. Prolonged Seasonality of Respiratory Syncytial Virus Infection among Preterm Infants in a Subtropical Climate. *Plos One* 2014; 9(10).

61. Janet S, Broad J, Snape MD. Respiratory syncytial virus seasonality and its implications on prevention strategies. *Hum Vacc Immunother* 2018; 14(1): 234-44.

62. Getaneh AM, Li X, Mao Z, et al. Cost-effectiveness of monoclonal antibody and maternal immunization against respiratory syncytial virus (RSV) in infants: Evaluation for six European countries. *Vaccine* 2023; 41(9): 1623-31.

63. Tang JW, Lam TT, Zaraket H, et al. Global epidemiology of non-influenza RNA respiratory viruses: data gaps and a growing need for surveillance. *Lancet Infect Dis* 2017; 17(10): E320-E6.

64. Staadegaard L, Meijer A, Rodrigues AP, et al. Temporal Variations in Respiratory Syncytial Virus Epidemics, by Virus Subtype, 4 Countries. *Emerg Infect Dis* 2021; 27(5): 1537-40.

65. Glatman-Freedman A, Kaufman Z, Applbaum Y, et al. Respiratory Syncytial Virus hospitalization burden: a nation-wide population-based analysis, 2000-2017. *J Infect* 2020; 81(2): 297-303.

526 66. Johannesen CK, van Wijhe M, Tong S, et al. Age-Specific Estimates of Respiratory Syncytial
527 Virus-Associated Hospitalizations in 6 European Countries: A Time Series Analysis. *J Infect Dis*
528 2022; **226**(Suppl 1): S29-S37.

529 67. Tong S, Amand C, Kieffer A, Kyaw MH. Incidence of respiratory syncytial virus related
530 health care utilization in the United States. *J Glob Health* 2020; **10**(2): 020422.

531

532 **Table 1. Characteristics of publications and investigations included in the systematic**
533 **review.**

Characteristics	No. of Publications (N=209) ^a n (%)	No. of Investigations (N = 552) n (%)
Climatic zone		
Temperate (>35°)	123 (58.9)	335 (60.7)
Subtropical (23.5° - 35°)	65 (31.1)	134 (24.3)
Tropical (equator - 23.5°)	49 (23.4)	83 (15.0)
Continent		
Asia	64 (30.6)	140 (25.4)
Europe	68 (32.5)	155 (28.1)
North America	38 (18.2)	164 (29.7)
South America	14 (6.7)	49 (8.9)
Africa	12 (5.7)	24 (4.3)
Oceania	5 (2.4)	19 (3.4)
Others ^b	8 (3.8)	1 (0.2)
Study period^c		
Before Palivizumab approved	88 (42.1)	147 (26.6)
After Palivizumab approved	75 (35.9)	320 (58.0)
Others ^d	46 (22.0)	85 (15.4)
Length of study period		
3-6 years	127 (60.8)	342 (62.0)
7-10 years	53 (25.4)	123 (22.3)
> 10 years	37 (17.7)	87 (15.8)
Type of RSV data		
Weekly data	57 (27.3)	294 (53.3)
Monthly data	155 (74.2)	258 (46.7)
Analytic method		
Qualitative	146 (69.9)	179 (32.4)
Quantitative	66 (31.6)	373 (67.6)

Threshold-based	54 (25·9)	205 (37·1)
Coverage-based	8 (3·8)	142 (25·7)
Model-based	6 (2·9)	26 (4·7)
RSV case source		
Inpatient	94 (45·0)	152 (27·5)
Outpatient	14 (6·7)	25 (4·5)
Both	53 (25·4)	115 (20·8)
Unknown	48 (23·0)	260 (47·1)
RSV case definition ^e		
ARI or ILI	82 (39·2)	150 (27·2)
SARI	23 (11·0)	45 (8·2)
ALRI	48 (23·0)	59 (10·7)
Clinical judgement	9 (4·3)	30 (5·4)
Unknown	70 (33·5)	310 (56·2)
Testing method ^f		
Virus detection	45 (21·5)	168 (30·4)
Antigen detection	107 (51·2)	303 (54·9)
Antibody detection	8 (3·8)	42 (7·6)
Nucleic acid detection	110 (52·6)	323 (58·5)
Unknown	33 (15·8)	127 (23·0)

Note: ARI, acute respiratory infection; ILI, influenza-like illness; SARI, severe acute respiratory infection; ALRI, acute lower respiratory infection

^aAn investigation was identified as an analysis with a specified method on the reported RSV data for the seasonality of RSV activities. Analyses on the same data with different analytic methods were recognized as different investigations. A publication might include several investigations describing the seasonality of RSV activities by analysing different sets of data, or applying different analytic methods, etc. Therefore, the sum of investigations might be greater than the total number of publications.

^bRefers to the publications/investigations with study sites involving more than one continent.

^cFor different locations, the time for palivizumab to be licensed for prophylactic use are different. For the countries where palivizumab has not been licensed, such as China, all the publications/investigations of this country would be recognized as “before palivizumab approved”.

^dRefers to the publications/investigations in which the study period contains the periods both before and after the year when Palivizumab was approved in the study locations.

^eA publication/investigation on RSV seasonality might be based on multiple sources of cases, e.g., ARI, ILI and SARI perhaps were used together or in different combinations to identify patients for RSV testing. Some investigations/studies might be counted multiple times, and the sum of the proportions of publications/investigations from different case definition categories may exceed 100%.

^fA publication/investigation on RSV seasonality might adopt multiple testing methods for RSV detection. Some investigations/studies might be counted multiple times, and the sum of the proportions of publications/investigations from different testing method categories may exceed 100%.

Table 2. Associations identified in the linear regression analysis between durations of RSV seasons estimated from quantitative approaches and characteristics of the studies.


Variable	Change in duration of RSV season (months)	95% confidence interval	p-value
Climatic zone			
Temperate region (<-35° or >35°)	0	Referent	
Subtropical region (-35° - -23.5° or 23.5° - 35°)	1.06	0.45, 1.66	<0.001
Tropical region (-23.5° – 23.5°)	1.65	0.77, 2.54	<0.001
Analysis method			
Coverage-based method	0	Referent	
Threshold-based method	1.43	0.96, 1.90	<0.001
Model-based method	1.98	-0.99, 4.95	0.19
Meteorological factor			
Daily average mean absolute humidity	0.013	-0.0049, 0.030	0.15


FIGURE LEGENDS

Figure 1. Flowchart of study identification and selection for the systematic review.

Figure 2. Geographic distribution of study locations and the reported RSV seasonal patterns from all included investigations using either quantitative or qualitative approaches.

Figure 3. The timing of one-peak RSV seasons reported in investigations using either quantitative or qualitative approaches against the latitude of the study site located in the Northern Hemisphere (NH, centred at the north pole) in panels A, C and E, and Southern Hemisphere (SH, centred at the south pole) in panels B, D and F. The start (A), peak (C) and end (E) of RSV seasons in NH. The start (B), peak (D) and end (F) of RSV seasons in SH. The centre of the radius circles represents the north pole or the south pole (latitude = 90°) and the external border represents the equator (latitude = 0°). The circles were divided by the dashed radiuses into 12 sections to represent 12 months of a year. Two radius circles between the north/south pole and the external border indicating the latitudes at 23.5° and 35° are used to define the temperate region (35°-90°), subtropical region (23.5°-35°) and tropical region (equator-23.5°).






Northern Hemisphere


A.

Month of season start




Southern Hemisphere

B.




C.

Month of season peak




D.



E.

Month of season end



F.

