

TRENDS, IN-HOSPITAL OUTCOMES, AND INDEPENDENT PREDICTORS OF
ACUTE KIDNEY INJURY IN PATIENTS ADMITTED FOR MYOCARDIAL
INFARCTION WITH PERCUTANEOUS CORONARY INTERVENTION: An insight
from the National Inpatient Sample Database

Akanimo Antia, MD.¹, Daniel Ubokudom, MD.², Olanrewaju Adabale, MD.³, Ovie
Okorare, MD.⁴, Emmanuel Daniel, MD.⁵, Endurance Evbayekha, MD.⁶, Chinwendu Angel
Onuegbu, MD.⁷, Kenneth Ong, MD.⁸

Department of Medicine, Lincoln Medical Center, Bronx,

New York, United States of America¹

Department of Medicine, Thomas Hospital, Fairhope,

Alabama, United States of America²

Department of Medicine, East Carolina University Health Medical Center, Greenville

North Carolina, United States of America³

Department of Medicine, Vassar Brothers Medical Center

New York, United States of America⁴

Department of Medicine, Trinity Health Ann Arbor, Ypsilanti,

Michigan, United States of America⁵

Department of Medicine, St. Luke's Hospital, St. Louis,

Missouri, United States of America ⁶

Department of Medicine, Montefiore Medical Center,

New York, United States of America ⁷

Chief of Cardiology, Department of Cardiovascular Medicine

Lincoln Medical Center ⁸

Corresponding author:

Akanimo Antia, MD

Lincoln Medical Center

234 E 149th Street

Bronx, New York

Tel: 646- 355 7508

Email: akanimousenantia@gmail.com

Conflicts of Interest: All authors declare no conflict of interests

Key Words: Acute kidney Injury, Percutaneous Coronary Intervention, Myocardial Infarction, Temporal Trends, Mortality.

Abstract Word Count: 351

Manuscript Word Count (excluding references): 4129

Number of References: 26

Number of Figures: 1

Number of Tables: 3

ABSTRACT:

Background: Acute kidney injury (AKI) is an important risk factor associated with adverse outcomes in cardiovascular illnesses, more importantly, myocardial infarction (MI). This study describes the Trends, in-hospital outcomes, and independent predictors of Acute Kidney Injury (AKI) in patients admitted for Myocardial infarction with Percutaneous Coronary Intervention (PCI).

Methods: This retrospective study used patient records from the 2016-2020 National In-patient Database (NIS). We identified patients who were admitted for the management of an MI who had a PCI procedure and an AKI and evaluated their associated socio-demographic and comorbid factors using International Classification of Diseases- 10 (ICD-10) codes. The chi-square test was used to compare baseline characteristics between our populations with and without AKI and outcomes and multivariate logistic regression to identify independent predictors of AKI.

Results: There were 1,551,630 patients admitted for an MI and PCI, with 15% having an AKI. We observed that our population with AKI were older on admission and were more likely to be whites than blacks. A higher percentage were males. Our subpopulation was likely to have heart failure, atrial fibrillation, coronary artery disease, obesity, CKD and Charlson comorbidity index ≥ 3 . A diagnosis of AKI was associated with higher in-hospital mortality rates [adjusted odds ratio (aOR): 2.84, CI: 2.7–3.02, $p < 0.001$], longer mean length of stay (LOS) and higher hospital costs. We noted an

increasing trend in the percentage of patients who had an AKI, from about 13.5% in 2016 to 16.5% in 2020.

Conclusion: Acute Kidney Injury is strongly associated with worse hospital outcomes in patients admitted for MI and PCI, with higher mortality rates, a longer mean length of stay, and a higher hospitalization cost. A more concise look at preventive measures is recommended to minimize these outcomes.

INTRODUCTION:

Acute Kidney Injury (AKI) following percutaneous coronary intervention (PCI) is a common complication occurring in > 15% of STEMI patients who undergo primary PCI^{1,2}. The use of more contrast agents during the procedure has been linked to a higher incidence of AKI after primary PCI versus elective PCI.^{3,4} Emergent primary percutaneous coronary intervention (primary PCI) remains the gold-standard ST-segment elevation myocardial infarction (STEMI) treatment. Recent studies have shown improved outcomes and prognosis for patients with STEMI, with one-year mortality at 11%⁵.

AKI can occur in patients with both pre-existing impaired renal function and in patients with normal baseline renal function⁶. In general, acute myocardial infarction (AMI) is one of the critical conditions that can trigger AKI. The most common etiologies stem from hemodynamic instability causing renal hypoperfusion, radiocontrast toxicity, and athero-embolism. About 5% of patients undergoing PCI experience a transient increase

in the plasma creatinine value of more than 1.0 mg/dL because of contrast use. The risk is greatest in patients with a history of diabetes mellitus and moderate to severe impaired renal function.

In the past decades, the morbidity of AKI has increased from 3/1000 to 5/1000 in the United States ⁷. More so, acute myocardial infarction (AMI) complicated by cardiogenic shock may increase the incidence of AKI by more than 50% ⁸. Among patients with AMI, those with AKI had a 20- to 40-fold higher mortality rate in comparison to those without AKI. Patients with AKI also have prolonged hospital stays, increased hospital charges, and more long-term complications, including recurrent AMI, heart failure, chronic kidney disease progression, and long-term mortality ^{8,9}.

The aim of our retrospective study is to describe the trends, in-hospital outcomes, and independent predictors of acute kidney injury in patients admitted for myocardial infarction with percutaneous coronary intervention.

METHODS:

This study is reported following the strengthening of the reporting of observational studies in Epidemiology (STROBE) reporting guidelines.

Study Design

Data Source: In this retrospective study, we analyzed hospitalizations between January 1st, 2016, and December 31st, 2020, from the National Inpatient Sample (NIS) in the

United States (US). The NIS was created and is maintained by the Agency for Healthcare Research and Quality and is the largest publicly available all-payer in-patient database in the United States of America (US). It was designed as a stratified probability sample representing all non-federal acute care hospitals nationwide. Hospitals are stratified according to ownership/control, bed size, teaching status, urban/rural location, and geographic region.

A multistage 20% probability sample of all hospitals within each stratum is then collected. All discharges from these hospitals are recorded and then weighted to ensure they are nationally representative. Data from 47 statewide data organizations (46 States plus the District of Columbia) encompassing more than 97% of the US population is included in the NIS 2016 – 2020 sampling frame. As many as 30 discharge diagnoses for each hospitalization were recorded using the International Classification of Diseases, Tenth Revision, Clinical Modification (ICD-10) in NIS 2016, and 40 discharge diagnoses and 25 procedures were coded in the NIS 2020 database. In the NIS, diagnoses are divided into principal and secondary. A principal diagnosis was the main ICD-10 code for hospitalization. Secondary diagnoses were any ICD-10 code other than the principal diagnosis.

Since all patient data in NIS are de-identified and publicly available, we waived the institutional review board approval.

Inclusion Criteria and study variables: The study population consisted of all in-patient hospitalizations with a primary diagnosis or secondary diagnosis of Acute Kidney Injury (AKI) during admission for myocardial infarction and a Coronary percutaneous

intervention recorded between 2016 and 2020 following the exclusion of patients below 18 years. The diagnosis was identified using ICD 10 codes recommended by the American college of and the American Association of Cardiology Respectively.

Demographic Study variables, including Age, race, Median household income, Primary insurance, and co-morbidities (computed from the Charlson comorbidities index), were identified as variables already present in the data set. Other variables were identified using The American ICD10-CM (diagnosis) medical billing codes identified from the review of other nationwide studies on cardiovascular diseases (**Table 1**). The ICD-10 codes used to identify our principal diagnoses, secondary diagnoses, and comorbidities are listed in the supplementary Table 1

Outcomes: We compared the sociodemographic differences in the population with and without AKI as highlighted in **Table 1**. We further analyzed the trend in the incidence of AKI and outcomes inclusive of (cardiogenic shock, cardiac arrest, and in-hospital mortality) across the years, from January 1st, 2016, to December 31st, 2020.

Data Analysis: Our data was analyzed using STATA, version 17 (Stata Corp, Texas, USA). We performed all analyses using weighted samples for national estimates following the Healthcare Cost and Utilization Project (HCUP) regulations for using NIS databases. Based on descriptive analysis of our data, continuous variables were presented with mean, and standard deviation and the differences were tested using a T-test. Categorical variables were presented with numbers (percentages) and compared with the chi-square test. Trends in incidence, mortality, and complications of AKI in our population were accounted for using conditional marginal effects of the variable "YEARS." Outcomes were

analyzed to obtain adjusted Odds ratios (AOR) using multiple logistic regression and linear regression models to account for potential confounders. The P-values considered significant in the multivariate analysis were two-sided, with < 0.05 as the threshold for statistical significance.

RESULTS:

There were 1,551,630 patients admitted for an MI and PCI, with about 232,180 (15%) having an AKI compared to 1,319,450 (85%) without AKI.

Our patients with AKI were notably older with a mean age of 68.9 years as compared to those without AKI with a male predominance in both groups.

The white race (75.8%) was more represented than blacks (9.5%) but when isolating each race, AKI was more common in the blacks (12.2% vs 8.97%) than in whites (71.3% vs. 76.5%) ($p < 0.0001$).

Our subpopulation was likely to have heart failure (53.8% vs. 21%, $p < 0.0001$), atrial fibrillation (24.6% vs. 12%, $p > 0.0001$), coronary artery disease (2.5% vs. 1.44%, $p < 0.0001$), COPD (18% vs. 12.5%, $p > 0.0001$), history of CABG (9.6% vs. 7.14%, $p < 0.0001$), obesity (21.7% vs. 19.9%, $p < 0.0001$), CKD (49.2% vs. 11.73%, $p < 0.0001$), Charlson comorbidity index ≥ 3 (74.3% vs. 35.6%, $p < 0.0001$) and anemia (34.4% vs. 10.9%, $p < 0.0001$). They were also more likely to be obese.

Our population without a history of AKI had higher prevalence of Dyslipidemia (71.6% vs 67.3%), Nicotine use (27.6% vs 26.8%), Dialysis dependence (2.5% vs 1.3%) and Diabetes (17.9% vs 12%)

The independent Predictors of AKI in the study population include chronic kidney disease (aOR: 5.3, CI: 5.1 – 5.5, $p < 0.001$), cardiogenic shock (aOR: 4.9, CI: 4.7 – 5.0, $p < 0.001$), cardiac arrest (aOR: 3.1, CI: 2.9 – 3.3, $p < 0.001$), Charlson comorbidity index of ≥ 3 (aOR: 2.3, CI 2.2 – 2.4, $p < 0.001$) and Heart Failure (aOR: 2.1, CI 2.0 – 2.1, $p < 0.001$)

A diagnosis of AKI was associated with higher in-hospital mortality rates [adjusted odds ratio (aOR): 2.84, CI: 2.7–3.02, $p < 0.001$], longer mean Length of Stay (7.1 days vs. 3.0 days, $p < 0.001$) and higher hospital costs (\$183,785.8 vs. \$101,291.2, $p < 0.0001$).

We noted an increasing trend in the percentage of patients who had an AKI, from about 13.5% in 2016 to 16.5% in 2020.

DISCUSSION:

Acute kidney Injury is a serious and fatal complication of Acute Myocardial Infarction. Deterioration of renal status in patients hospitalized for Acute Myocardial infarction could be influenced by many factors, which include patient hemodynamic status, cardiac function, use of contrast agents and drugs, hypertension or Diabetes Mellitus, underlying CKD, and the performance of CABG ^{1,9-12}.

AKI was defined, according to the European Society of Urogenital Radiology as an increase of 25% in serum creatinine from baseline creatinine within 72 hours from

contrast media administration ². In the KDIGO criteria, an increase in creatinine was characterized into stages. Stage 1 is an increase of creatinine by 1.5-1.9 from baseline, and stage 3 is a threefold increase from baseline. This was translated into a severity score with presumptions that a 25% increase based on the European Society of Radiology translates to stage 1 KDIGO ¹³.

Risk factors for AKI have been reported in Patients undergoing PCI, such as old age, Diabetic Nephropathy, time from onset of symptoms to Primary PCI, pre-existing renal disease, congestive heart failure, contrast volume, and length of the procedure ¹⁴⁻¹⁸. In a recent study with 1656 STEMI patients, congestive heart failure and reduced left ejection was identified as individual predictors of AKI ¹⁹. This is probably due to decreased cardiac output in patients with myocardial infarction, especially in large infarct size and those with anterior myocardial infarction complicated by cardiogenic shock. Patients with acute coronary syndrome (ACS) tend to also have an increased risk of plaque disruption, particularly cholesterol crystals, subsequent systemic embolism and if the renal arterioles are involved, the risk of AKI increases ²⁰.

In addition to the above, our study observed cardiac arrest and a Charlson comorbidity index ≥ 3 to be strong predictors of AKI in patients that have a PCI procedure done (**table 2**). Contrast volume demonstrated a strong correlation with AKI. In earlier studies, higher risks of AKI were reported in patients receiving 153 -378mls in comparison to those with a lower contrast volume use (111-114 mls) ^{4,16,17}. Contrast nephropathy has been associated with decreased renal blood flow, causing both intrarenal AKI and prerenal AKI.

Medications like furosemide have also been observed as a risk factor for AKI. The use of these loop diuretics has been associated with decreased left ventricular ejection fraction, agitation of the renin-angiotensin system, leading to increased peripheral vascular resistance, renal hypoperfusion, and AKI. Furosemide is commonly used in MI patients presenting with cardiogenic shock and acute pulmonary Edema. Data regarding the use of ACEI and ARBS following early and acute stages of Acute Myocardial Infarction remains controversial. ACEI and ARBs have had great mortality benefits by preventing cardiac remodeling and reducing urine protein. Some studies have shown that the use of ACEI and ARB in patients with insufficient renal perfusion might lead to a decline in glomerular filtration rate, thereby inducing AKI ^{21,22}. Data from a study showed reduced use of anti-RAS agents by doctors due to hyperkalemia and elevated creatinine levels ²³. A clinical study of 6867 cases showed that the application of anti-RAS therapy in Patients with CKD with ACS could improve 90-day mortality rates ²⁴. Diabetes Mellitus per se is not an independent risk factor for developing AKI in patients with MI when examined independently. However, when combined with renal insufficiency in patients with MI, the risk increases²⁵. Recent studies have shown admission hyperglycemia to be an independent risk factor for AKI following primary PCI ^{20 26}.

Time plays a critical role in acute myocardial Infarction. Early PCI has been shown to limit infarct size, reduce complications of heart failure and decrease risk of AKI¹⁸.

Our research provided additional evidence that a diagnosis of acute kidney injury is linked to a mortality risk increase of over 50% and longer hospital stays. Furthermore, we noted a substantial increase in the overall hospital expenses incurred by this group

of patients (**Table 3**). Efforts have been made to negate the risk of AKI in patients undergoing percutaneous coronary intervention. Some of these measures include reducing contrast volume, Hydrating patients with normal saline infusion and pretreatment with acetylcysteine. Studies by Muller et al. in a large randomized prospective study, showed potential benefits of isotonic saline in reducing AKI especially in unstabilized cardiac patients undergoing PCI ²⁶.

Despite the above preventative measures, our study showed an increasing trend in the rates for AKI in patients receiving PCI procedures (**Figure 1**). This increasing trend could be attributed to the overall rising prevalence of cardiovascular comorbidities in our patient populations.

In lieu of these trends, we further emphasize, as previously stated, that infusions should be started early and continued long enough to ensure down-trending creatinine levels. Control of Blood glucose in patients presenting with hyperglycemia is necessary to reduce the formation of free radicals which could worsen renal function and myocardial injury. The use of medications such as ACEI and ARBs though beneficial should be individualized keeping in view baseline creatinine. An increase in creatinine of about 25-30% from baseline following use of ACEI and ARBs is acceptable in clinical practice. Intra-aortic balloon pump placement in patients with worsening cardiogenic shock not amenable to diuretics may help improve circulation, reducing the risk of AKI.

Although our study has notable strengths, it also has limitations. Firstly, due to our lack of access to individual-level data, we were unable to assess the consistency of the criteria used to diagnose acute kidney injury, which could potentially result in an over or

underestimation of cases. Additionally, as our database only includes in-hospital events, we could not investigate the long-term effects of AKI hospitalization. It is possible that the size of the infarct and patients with lower ejection fractions will have increased risks of AKI but this could not be assessed during this study.

On the other hand, the power of our study lies in its large sample size, which was obtained from a population database, enhancing the credibility of our results.

CONCLUSION:

The burden of acute kidney injury following PCI for acute myocardial infarction is significant, with up to 15% of patients developing AKI following the procedure. It is associated with higher in-hospital mortality, longer mean lengths of stay and higher and higher hospitalization costs. Our retrospective study has shown that independent predictors for AKI following PCI include CKD, cardiogenic shock, cardiac arrest, heart failure, Charlson comorbidity index ≥ 3 , closer attention and monitoring needs to be paid to patients with these risk factors in order to prevent AKI. We also found an increasing trend in the percentage of those that developed AKI from 2016 to 2020 underscoring the need for increased preventative measures.

Abbreviations:

ACEI – Angiotensin Converting Enzyme inhibitor

ACS - Acute Coronary Syndrome

AKI – Acute Kidney Injury

AMI – Acute Myocardial Injury

aOR – adjusted Odds Ratio

ARB – Angiotensin Receptor blockers

CABG – Coronary Artery Bypass Graft

CCI – Charlson Comorbidity Index

CHF – Congestive Heart Failure

CI – Confidence Interval

CKD – Chronic Kidney Disease

COPD – Chronic Obstructive Pulmonary Disease

HCUP - Healthcare Cost and Utilization Project

KIDGO – Kidney Disease Improving Global Outcomes

PCI – Percutaneous Coronary Intervention

STEMI – ST-Elevation Myocardial Infarction

Conflict of interest:

All authors declare that they have no competing interests (financial and non-financial).

Ethics declarations

The study was not submitted for research ethics approval as the activities described were conducted as part of the Nationwide Inpatient Sample Database (NIS), which is part of the family of databases and software tools developed for the Healthcare Cost and Utilization Project (HCUP) and uses de-identified data collected from hospitalized patients. Consent was not obtained, given the use of a de-identified database. All the experiments in our study were under the guidelines and agreement regulations of the Agency Healthcare Research and Quality (AHRQ)

Authors' Contributions

Akanimo Antia contributed to the conception and design of the research; Chinwendu Angel Onuegbu contributed to the design of the research; Olanrewaju Adabale contributed to the acquisition and analysis of the data; Ovie Okorare contributed to the interpretation of the data; Daniel Ubokudom, Emmanuel Daniel and Endurance Evbayekha helped draft the manuscript. Kenneth Ong made the final reviews. All authors critically revised the manuscript, agreed to be fully accountable for ensuring the integrity and accuracy of the work, and read and approved the final manuscript.

Consent for Publication

Not applicable. All data using the National Readmission Database Sample is de-identified.

Funding:

None

Data and Materials Availability

The datasets generated and analyzed during the current study are available in the Healthcare Cost and Utilization Project National Data Registry. This Data Use Agreement (“Agreement”) governs the disclosure and use of data in the HCUP Nationwide Databases from the Healthcare Cost and Utilization Project (HCUP), which the Agency maintains for Healthcare Research and Quality (AHRQ). Accordingly, HCUP Databases may only be released in “limited data set” form, as the Privacy Rule defines that term, 45 C.F.R. § 164.514(e). In addition, AHRQ classifies HCUP data as protected health information under the HIPAA Privacy Rule, 45 C.F.R. § 160.103. The datasets generated and analyzed during the current study are not publicly available except for the corresponding author who purchased the data and signed the HCUP Data Use agreement training. Researchers should readily be able to publicly purchase the same databases we did to conduct research.

REFERENCES:

1. Tsai TT, Patel UD, Chang TI, Kennedy KF, Masoudi FA, MathenyME, et al. Contemporary incidence, predictors, and outcomes of acute kidney injury in patients undergoing percutaneous coronary interventions: insights from the NCDR Cath-PCI registry. *JACC Cardiovasc Interv* 2014; 7:1–9. doi: 10.1016/j.jcin.2013.06.016.
2. Morcos S, Thomsen H, Webb J. Contrast-media-induced nephrotoxicity: a consensus report. *Eur Radiol* [Internet]. Springer-Verlag; 1999 [cited 2015 Jan 5]; 9(8):1602–13. Available from: <https://doi.org/10.1007/s003300050894> PMID: 10525875
3. Silvain J, Collet J-P, Montalescot G. Contrast-induced nephropathy: the sin of primary percutaneous coronary intervention? *Eur Heart J* [Internet]. 2014 Jun 14 [cited 2014 Jul 12]; 35(23):1504–6. Available from <http://www.ncbi.nlm.nih.gov/pubmed/24722806> <https://doi.org/10.1093/eurheartj/ehu126> PMID:24722806.
4. Marenzi G, Lauri G, Assanelli E, Campodonico J, De Metrio M, Marana I, et al. Contrast-induced nephropathy in patients undergoing primary angioplasty for acute myocardial infarction. *J Am Coll Cardiol* [Internet]. 2004 Nov 2 [cited 2014 Sep 4]; 44(9):1780–5. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/15519007> <https://doi.org/10.1016/j.jacc.2004.07.043> PMID: 15519007
5. Steg PG, James SK, Atar D, Badano LP, Blomstrom-Lundqvist C, Borger M a, et al. ESC Guidelines for the management of acute myocardial infarction in patients

presenting with ST-segment elevation. Eur Heart J [Internet]. 2012 Oct [cited 2014 Jul 9]; 33(20):2569–619. Available from:

<http://www.ncbi.nlm.nih.gov/pubmed/22922416>

<https://doi.org/10.1093/eurheartj/ehs215> PMID: 22922416

6. Rihal CS. Incidence and Prognostic Importance of Acute Renal Failure After Percutaneous Coronary Intervention. Circulation [Internet]. 2002 Apr 22 [cited 2014 Aug 27];105(19):2259–64. Available from:

<http://circ.ahajournals.org/cgi/doi/10.1161/01.CIR.0000016043.87291.33> PMID: 12010907

7. Tepel M, Aspelin P, Lameire N. Contrast-induced nephropathy: a clinical and evidence-based approach. Circulation [Internet]. 2006 Apr 11 [cited 2014 Sep 5]; 113(14):1799–806. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/16606801> <https://doi.org/10.1161/CIRCULATIONAHA.105.595090> PMID: 16606801

8. Hwang SH, Jeong MH, Ahmed K, Kim MC, Cho KH, Lee MG, et al. Different clinical outcomes of acute kidney injury according to acute kidney injury network criteria in patients between ST-elevation and non-ST elevation myocardial infarction. Int J Cardiol 2011; 150:99– 101. Doi: 10.1016/j.ijcard.2011.03.039.

9. Shacham Y, Leshem-Rubinow E, Steinvil A, Assa EB, Keren G, Roth A, et al. Renal impairment according to acute kidney injury network criteria among ST elevation myocardial infarction patients undergoing primary percutaneous

intervention: a retrospective observational study. Clin Res Cardiol 2014; 103:525

532. doi: 10.1007/s00392-014-0680-8

10. Koreny M, Karth GD, Geppert A, Neunteufl T, Priglinger U, Heinz G, et al.

Prognosis of patients who develop acute renal failure during the first 24 hours of cardiogenic shock after myocardial infarction. AmJMed 2002; 112:115–119. doi:

10.1016/S0002-9343(01)01070-1.

11. Levin A, Warnock DG, Mehta RL, Kellum JA, Shah SV, Molitoris BA, et al.

Improving outcomes from acute kidney injury: report of an initiative. Am J Kidney Dis 2007; 50:14. doi: 10.1053/j.ajkd.2007.05.008.

12. Fliser D, Laville M, Van Biesen W. A European Best Practice (ERBP) position

statement on the Kidney Disease Improving Global Outcomes (KDIGO) Clinical

Practice Guidelines on Acute Kidney Injury: Part1: definitions, conservative

management and contrast-induced nephropathy. Nephrol Dial Transplant [Internet].

2012 Oct 8; 0:1–10.

13. Tepel M, Aspelin P, Lameire N. Contrast-induced nephropathy: a clinical and

evidence-based approach. Circulation [Internet]. 2006 Apr 11 [cited 2014 Sep 5];

113(14):1799–806. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/16606801>

<https://doi.org/10.1161/CIRCULATIONAHA.105.595090> PMID: 16606801

14. Barrett B. Contrast nephrotoxicity. J Am Soc Nephrol. 1994; 5:125–37. PMID:

7993992

15. Narula A, Mehran R, Weisz G, Dangas GD, Yu J, Ge'ne'reux P, et al. Contrast-induced acute kidney injury after primary percutaneous coronary intervention: results from the HORIZONS-AMI substudy. *Eur Heart J* [Internet]. 2014 Jun 14 [cited 2014 Jul 14]; 35(23):1533–40. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/24603308>
<https://doi.org/10.1093/eurheartj/ehu063> PMID: 24603308
16. Celik O, Ozturk D, Akin F, Ayca B, Yalaki AA, Erturk M, et al. Association Between Contrast Media Volume-Glomerular Filtration Rate Ratio and Contrast-Induced Acute Kidney Injury After Primary Percutaneous Coronary Intervention. *Angiology*. 2014 Jul 8.
17. Schoolwerth AC, Sica DA, Ballermann BJ, Wilcox CS. Renal Considerations in Angiotensin Converting Enzyme Inhibitor Therapy. *Circulation*. 2001; 104:1985–91. <https://doi.org/10.1161/hc4101.096153> PMID: 11602506
18. Shacham Y, Leshem-Rubinow E, Gal-Oz A. Acute Cardio-Renal Syndrome as a Cause for Renal Deterioration Among Myocardial Infarction Patients Treated with Primary Percutaneous Intervention. *Can J Cardiol*. 2015; 31(10):1240–44. <https://doi.org/10.1016/j.cjca.2015.03.031> PMID: 26163472
19. Mahesh B, Yim B, Robson D, Pillai R, Ratnatunga C, Pigott D. Does furosemide prevent renal dysfunction in high-risk cardiac surgical patients? Results of a double-blinded prospective randomized trial. *Eur J Cardiothorac Surg* 2008; 33:370–376. doi: 10.1016/j.ejcts.2007.12.030.

20. Ya-feng Zhang, Da-dong Liu, Yue Zhou, Ji-zhuang Lou; Acute Kidney Injury in Patients with Acute Coronary Syndrome after Percutaneous Coronary Intervention: Pathophysiologies, Risk Factors, and Preventive Measures. *Cardiology* 1 December 2021; 146 (6): 678–689. <https://doi.org/10.1159/000517991>

21. Mansfield KE, Nitsch D, Smeeth L, Bhaskaran K, Tomlinson LA. Prescription of renin-angiotensin system blockers and risk of acute kidney injury: a population-based cohort study. *BMJ Open* 2016;6: e012690. doi: 10.1136/bmjopen-2016-012690. Wetmore JB, Tang F, Sharma A, Jones PG, Spertus JA.

22. Wetmore JB, Tang F, Sharma A, Jones PG, Spertus JA. The association of chronic kidney disease with the use of renin-angiotensin system inhibitors after acute myocardial infarction. *Am Heart J* 2015; 170:735–743. doi: 10.1016/j.ahj.2015.07.019.

23. Reddan DN, Szczech L, Bhapkar MV, Moliterno DJ, Califf RM, Ohman EM, et al. Renal function, concomitant medication use and outcomes following acute coronary syndromes. *Nephrol Dial Transplant* 2005; 20:2105–2112. doi: 10.1093/ndt/gfh981.

24. Rudnick MLR, Goldfarb S, Wexler L. Nephrotoxicity of ionic and nonionic contrast media in 1196 patients: A randomized trial. *Kidney Int.* 1995; 47:254–61. <https://doi.org/10.1038/ki.1995.32> PMID:7731155

25. Shacham Y, Gal-Oz A, Leshem-Rubinov E. Admission Glucose Levels and the Risk of Acute Kidney Injury in Nondiabetic ST Segment Elevation Myocardial Infarction Patients Undergoing Primary Percutaneous Coronary Intervention. *Cardiorenal Med.* 2015; 5:191–8. <https://doi.org/10.1159/000430472> PMID: 26195971

26. Mueller C. Prevention of contrast media-associated nephropathy. *Arch Intern Med.* 2002; 162:329–36. <https://doi.org/10.1001/archinte.162.3.329> PMID: 11822926

Table 1. Socio-Demographic and Co-morbid Differences in baseline characteristics of patients admitted for MI and PCI with and without AKI

Variables	Myocardial Infarction and Percutaneous Intervention N= 1, 551, 630 (%)		P-values
	AKI N= 232,180 (15%)	No AKI N= 1, 319,450 (85%)	
PATIENT CHARACTERISTICS			

Age, years, mean $\pm$ SE	68.9 $\pm$ 0.1	63.5 $\pm$ 0.03	<0.0001
SEX			<0.001
male	67.5	68.3)	
Female	32.5	31.7	
RACE			<0.0001
White	71.3	76.54	
Black	12.21	8.97	
Hispanic	9.4	8.1	
Asian	3.1	2.6	
Native American	0.6	0.7	
Other	0.3	0.3	
CHARLSON COMORBIDITY INDEX SCORE			<0.0001

1	9.6	34.2	
2	16.1	30.2	
≥3	74.3	35.6	
COMORBIDITIES			
Congestive Heart Failure	53.8	21.1	<0.0001
Atrial Fibrillation	24.61	12.04	<0.0001
CKD	49.2	11.7	<0.0001
Anemia	34.4	10.8	<0.0001
COPD	18.5	12.9	<0.0001
Obesity	21.7	19.9	<0.0001
History of CABG	9.0	7.14	<0.0001
Carotid Artery Disease	2.5	1.4	<0.0001

Dyslipidemia	67.3	71.6	<0.0001
Smoking	26.8	27.6	0.0018
Dialysis Dependent	1.3	2.5	<0.0001
Hypertension	23.4	53.6	<0.0001
Diabetes	12.0	17.9	<0.0001

*Table 2: Independent predictors of AKI in patients admitted for MI and a PCI
(multivariate analysis)*

Variables	Independent predictors of AKI	
	[aOR; CI]	p- values
CKD	5.296; 5.1 – 5.49	<0.001
Cardiogenic shock	4.85; 4.68 – 5.0	<0.001
Cardiac arrest	3.09; 2.93 – 3.27	<0.001

CCI ≤ 3	2.25; 2.15 – 2.35	<0.001
CHF	2.06; 2.0 – 2.1	<0.001

Table 3: In-hospital outcomes of patients with AKI on admission for MI and PCI

OUTCOMES	Myocardial Infarction and Percutaneous Intervention (%), [aOR, CI]	
	AKI	NO AKI
Mortality	11.96 (2.8; 2.67 – 3.01)	1.72 (ref)
Length of Stay	7.1 (2.24; 2.17 – 2.32)	3.0 (ref)
Total Hospital charge (\$)	183,785 (61,959; 59451.6 – 64467.9)	101291 (ref)




Figure 1: Trends of AKI in patients admitted for MI and a PCI from 2016 - 2020.

Supplemental Tables 1: International Classification of Diseases, Tenth Edition, Clinical Modification (ICD-10-CM) Codes Used To Identify Comorbidities, and Hospital Outcomes

Diagnosis codes (primary and secondary)	ICD-10 codes
Myocardial Infarction	I21
Percutaneous Coronary Intervention	0270346, 0272346, 027034Z, 027234Z, 0270356, 0272356, 027035Z, 027235Z, 0270366, 0272366, 027036Z, 027236Z, 027036Z, 0270376, 0272376, 027037Z, 027237Z, 02703D6, 02723D6, 02703DZ, 02723DZ, 02723D6, 0273466, 027146Z, 027346Z, 0271476, 0273476, 027147Z, 027347Z, 02714D6, 02734D6, 02734DZ, 02714E6, 02734E6, 02714EZ, 02734EZ, 02714F6, 02734F6, 02714FZ, 02734FZ, 02714G6, 02734G6, 02714GZ, 02734GZ, 02714Z6, 02734Z6, 02734ZZ, "02703Z6, 02723Z6, 02703ZZ, 02723ZZ, 0270446, 0272446, 027044Z, 02723ZZ, 02723ZZ, 0270446, 0272446, 027044Z, 027244Z, 027244Z, 0270456, 0272456, 027045Z, 027245Z, 0270466, 0272466, 027046Z, 027246Z, 0270476, 0272476, 027047Z, 027247Z, 02704D6, 02724D6, 02704DZ
Acute Kidney Injury	N17
Comorbidities	
Carotid artery disease	I652
Dyslipidemia	E78
Anemia	D50, D51, D52, D53, D55, D56, D57, D58, D59, D60, D61, D62, D63, D64
Atrial Fibrillation	I48

Chronic Kidney Disease	N18
Obesity	E660, E6601, E6609, E661, E662, E668, E669
History of CABG	Z951
Smoking	Z87891, F17200
Hypertension	I10
Diabetes	E119, E089, E109, E109, E108, E139, E1010
Dialysis Dependent	Z992
COPD	J41, J42, J43, J44
Heart Failure	I50
Other Variables	
Cardiogenic Shock	R570
Cardiac Arrest	I462, I469, I468
Abbreviations: MI: Myocardial infarction, CABG: Coronary artery bypass graft, COPD: Chronic Obstructive Pulmonary Disease	

