

Discontinuation of long-acting reversible contraceptive methods and associated factors among reproductive-age women in Shashemane town, Oromia, Ethiopia.

Fikru Letose^{1*}, Alemtsehay Tusa², Degemu Sahlu³ and Yohannis Miherite³

Affiliation

¹Amref Health Africa in Ethiopia, Gambella Field Office, Ethiopia.

²Melk Oda G/Hospital, Shashemane town, Oromia, Ethiopia.

³Department of Public Health, College of Health Science, Salale University, Fiche, Ethiopia.

Email addresses: FL: fikruletose6@gmail.com, AT: alemtsahytusa@gmail.com, DS: kergadegemu@gmail.com, YM: yohannes1979@gmail.com

*Corresponding author: Fikru Letose; Gambella Peoples' Regional National State Health Bureau, Gambella, Ethiopia; PO Box: 109; fikruletose6@gmail.com; Mobile Phone; +251921416858.

Research Type: Research Article

Abstract

Background: The early termination of long-acting reversible contraceptives raises issues for the healthcare system and has the potential to affect public health. Long-acting reversible contraception is now more widely available and used, although a sizable percentage of women still stop using it. Therefore, this study aims to assess factors associated with the discontinuation of the long-acting reversible contraceptive method among female users of health facilities in Shashemane Town.

Methods: A facility-based cross-sectional study was done in Shashemane town among 410 study participants. The study participants were selected by using a systematic random sampling method. Data were collected by using structured interviewer-administered questionnaires and entered into epi-data version 4.6.0.2 and exported to SPSS version 25 for analysis. Bivariate and multivariable logistic regressions were used to examine the association between independent variables and discontinuation of the long-acting reversible contraceptives. The results were presented using the Odds Ratio at 95% CI. $P < 0.05$ were used to dictate statistical significance.

Result: The overall prevalence of women who removed the long-acting reversible contraceptive method before the due date was 57.2%. Factors such as having an occupation as a housewife, desire to become pregnant, unwarned side effects, effectiveness, and dissatisfaction with the service provided were positively associated with discontinuation of the contraception.

Conclusion and Recommendation: The prevalence of the discontinuation of reversible long-acting contraceptives was high. Pre-insertion effective counseling about the benefits, follow-up care and management for side effects, and client reassurance are recommended.

Keywords: Discontinuation, Reversible long-acting, Contraceptive, Reproductive age women.

Introduction

An unintended pregnancy is defined as a pregnancy that occurred when no children or no more children were desired (Unwanted) or the pregnancy occurred earlier than desired (Mistimed)[1]. It is associated with an increased risk of problems for the mother and baby. According to the World Health Organization (WHO), 74 million women in low- and middle-income countries (LMIC) become pregnant unintentionally each year. This results in 47,000 maternal fatalities and

25 million unsafe abortions each year. This leads to 25 million unsafe abortions and 47,000 maternal deaths every year[2].

One of the most affordable health promotions to lower mother and child mortality by about 30% and 10%, respectively, is the use of modern contraceptive techniques. [3]. Therefore, by ensuring that every pregnancy is intended and every birth is safe, the vast majority of maternal and infant deaths can be avoided.[4]. Long-acting reversible contraceptives (LARCs) are a type of modern birth control method that prevents unintended conception for a long time without forcing the client to exert additional effort for at least three years of continuous use.[5]. Intra-Uterine Copper Devices (IUCDs), the levonorgestrel (LNG)-releasing intrauterine system, and LNG-releasing and etonogestrel (ENG)-releasing subdermal implants are categorized as LARC methods [6-8]. Long-acting reversible contraceptive discontinuation is the removal from using and ceasing it or switching to other methods for any reason before completion of the duration(1).

Despite the improvement in the availability and utilization of long-acting reversible contraception, early discontinuation is becoming a major problem[9]. In low-income countries (LICs), a systematic review report revealed that discontinuation of LARCs within one year of insertion was 20%[10]. In a study conducted on 21 LICs, LARCs comprised only 10% or less, of which 9% of implant users and 15% of IUCD users discontinued the method within the first year of use[11]. Besides, 13.1%, 26.3%, and 36.7% of IUD users discontinued at the first 12, 24, and 36 months, respectively[12]. The Ethiopian Demographic Health Survey (EDHS) revealed that the overall discontinuation of contraceptives at 12 months was 35%, of which 13% were IUD users and 11% were implant users[13]. Other studies conducted in Ethiopia showed that the prevalence of discontinuation of LARCs use was 20%([14], 36.94%([15], 50%[16], and 66.3%[17].

Contraceptive discontinuation may be associated with external factors, individual and/or partner factors [18-20]. The quality of family planning services and obstetric factors like past abortion history, number of living children, desire for more children, preferred family size, sex preference, irregular vaginal bleeding, lower abdominal pain, and abnormal vaginal discharge were linked to LARCs discontinuation, according to a small number of studies[14, 21-26]. Socio-demographic variables such as age, marital status, educational status, and religious concern were found to influence LARC discontinuation. Regarding method-related variables, studies revealed that side effects[11, 14], counseling, satisfaction with the service given, and information about family planning were factors associated with the discontinuation of LARCs[17, 21-23].

In Ethiopia, providing LARCs is a free but highly effective method of preventing unintended pregnancies [27]. Discontinuation while still in need is particularly problematic for women which results in 15% to 20% times at risk of unwanted pregnancy for LARC users three months after discontinuation[11]. Long-acting reversible contraceptives when discontinued to increased fertility rate, unintended pregnancy, and associated complications [11, 28]. The high LARCs discontinuation coupled with low uptake results in a big challenge to achieve the targeted contraceptive prevalence in Ethiopia[29].

Depending on cultural differences and time elapse, the causes for the termination of LARCs may be extremely contextual. Additionally, reducing the discontinuance of LARCs is a great way to prevent or at least minimize unintended pregnancy[30]. In countries like Ethiopia with high fertility and unmet need for family planning[13], it is important to understand how long women continue to use LARCs and what factors are associated with its discontinuation helps to improve the reproductive health of women. However, limited information regarding methods

discontinuation from using LARCs and associated factors in the study area. Therefore, the main aim of this study was to determine the discontinuation of LARCs and associated factors among reproductive-age women in Shashemane town, Oromia, Ethiopia.

Method and materials

Study area and period

The study was conducted in Shashemene town among women 15-49 years of age. Shashemene town is the capital city of the West Arise zone in the Oromia Region which is 250 Km far from Addis Ababa. The total population of the town is estimated to be 286,447 of which 49.5% are females and 34.7% are reproductive age group women. Currently, the town has two public hospitals, five health centers, two Nongovernmental organizations (NGOs) clinics, 1 private hospital, more than 50 private clinics, and pharmacies are providing health services. From these health facilities, all hospitals, health centers, private hospital, NGO clinics, and nine private clinics provide LARCs services. The study was conducted from May 16 to June 16, 2022.

Study design

The institution-based cross-sectional study design was applied.

Populations

Source population

All reproductive-age women who were using LARCs in the health facilities of Shashemene Town.

Study population

All randomly selected reproductive-age women who were using LARCs and come to selected health facilities of Shashemene Town with contraceptive-related issues during the study period.

Inclusion criteria and exclusion criteria

Inclusion criteria All reproductive-age women who were using LARCs and come to the selected health facilities for any issue concerning the method before completion of duration; like removal, side effects, and follow-up during the actual data collection period. Those women who lived in the study area at least for six months were included in the study.

Exclusion criteria: All women who have used LARCs outside of Shashemane town and come to the study area for the removal service. Those women who come to change Implanon to Jadelles, Jadelles to Implanon, and implants to IUCD or vice versa were excluded from the study.

Sample size determination

The required sample size was calculated by using the population proportion formula equals =

$$\frac{(z_{\alpha/2})^2 \times P(1 - P)}{d^2}$$

Where n is the minimum sample size, $p=39\%$ prevalence of Implanon discontinuation in Ambo town[31], $z=1.96$ (95% CI), and $d=0.05$ margin of error. Applying the formula by substituting these values into the equation gives 373. Ten percent was added for the non-response rate and the final Sample size become 410.

Sampling Technique

All public health facilities and non-governmental specialty clinics in Shashemene town which provides LARC services were included in the study. Namely, they are two public Hospitals (Referral Hospital and Melk oda G/Hospital), five public health centers (Abosto, Awasho, Bulchana, Dida Boke, and Arada,) and two non-governmental specialty clinics (Family Guidance Association and Marie stops International-Shashemene). Private health facilities were not included in this study because they were not providing full LARC services (insertion and

removal). To proportionally allocate the calculated sample size, average client flow for 6 months (December 2021 to April 2022) at each health institution was considered before data collection. The information obtained from the six-month enrollment record of the family planning registration book indicated that a total of 4016 women booked for LARCs at public and non-governmental specialty clinics. The average 6-month client flow identified from the family planning registration book at Referral Hospital, Melk oda General Hospital, Abosto HC, Awasho HC, Bulchana HC, Dida Boke HC, Arada HC, Family Guidance Association and Marie stops International-Shashemene were 452, 419, 403, 410, 354, 349, 322, 618, and 689 respectively. The calculated sample size was proportionally allocated to each facility. Then, the study participants were identified by systematic random sampling method. Data collectors approached and recruited LARC users who had come to the selected clinics for contraceptive-related reasons.

Therefore, every other woman was included in the sample until the total sample size for this study was obtained. The procedure was continued throughout the data collection period until the required sample size was achieved (**Figure 1**).

Data Collection Tool and procedure

The data were collected using structured questionnaires through the face-to-face interview which is developed after different literature related to the topic under study are reviewed. It was prepared in English and translated to Afan Oromo and Amharic for data collection. The questionnaire contains four parts: Socio-demographic characteristics, Contraceptive and counseling-related characteristics, Health system characteristics, and Obstetric related characteristics. Two days of training were given to data collectors and supervisor about the study

objective, procedure, and research ethical rules. Informed consent was taken from selected individual women and getting permission for the data collection takes 15-20 minutes.

Study variables

The dependent variable was the discontinuation long acting reversible contraceptive method and Independent Variables were Socio-demographic characteristics which are age, marital status, religion, occupation, education, and family income; Health system related characteristics, Health extension workers' availability and Time taken to reach to health facilities; Obstetrics related characteristics including Parity, Living children, History of abortion, Desire of pregnancy and Time to get pregnancy; Contraceptive and counseling related characteristics Information about contraceptive like History of family planning before used, Type of contraceptive used before LARCs, Responsible body for choosing LARCs, Counseling and Satisfaction with service given.

Data Quality assurances

To assure the quality of data, the questionnaire was pre-tested during May 9-11/2022 at 01 kebele (Hage) health center which was found outside of the actual study area. Five percent of the calculated sample of women were enrolled to check the validity and reliability of the questionnaire. After pre-testing the questionnaire, revisions and amendments were made accordingly. The training was given to the data collectors on how to interview and check the questionnaire for completeness. The supervisors and principal investigator checked and revised the completeness of the questionnaire and offer necessary feedback to data collectors.

Data management and analysis

The collected data was entered using Epi-Data version 4.6 and exported to SPSS 23 software for cleaning, recording, categorizing, and analyzing. A bivariate analysis was done to see the

association between independent and outcome variables. Those variables with a P-value of 0.2 during the bivariate analysis were included in the multiple logistic regression analysis to assess the relative effect of confounding variables. Since the outcome variable is categorical, the adjusted odds ratio was calculated using a multiple logistic regression model. After multivariate analysis had been done, the adjusted odds ratio (OR) was used to measure the strength of the association between the dependent variable and the independent variable, while the 95% CI and P-value were used to assess whether the association was significant.

Ethical consideration

An ethical assurance letter was received from Salale University (SLU) College of Health Sciences, Department of Public Health. It was presented to the Shashemene town health office to grant official permission to undertake research activities in the selected health facility. Written informed consent was obtained from each participant after the investigator had explained the nature, purpose, and procedure of the study. The signature of each participant was used on the willingness confirmation. The entire study's participants were informed that data was kept private and confidential and used only for research purposes. The participants were also assured that they have the right to refuse or withdraw if they are not comfortable at any time.

Results

Socio-demographic Characteristics

A total of 402 women responded to the questionnaire forming, a response rate of 98%. The mean age of the respondents was 26 years ($\pm$ 4.6). About 241(60.0%) were Oromo by Ethnicity, and 150(37.3%) were Muslims by religion. Most of the respondents 268(91.5%) were married. Two hundred fifty-three (62.9%) respondents attended secondary education. Regarding occupation,

the majority of the respondents 230(57.2%) were housewives. More of the respondents have a monthly income of 2000-3000 ETB per month (**Table-1**)

Health system-related characteristics

The majority of women 342(85.1%) have access to health extension workers in their catchment and more than half of the women 252(62.7%) live 30-60 minutes walking distance from the health facilities.

Obstetric-Related Characteristics

Among the women who ever used LARCs, only three (1%) have no history of live birth. Besides, 9 (2.2%) have no child, 122(30.3%) have a history of abortion, and 177(44%) desire to become pregnant (**Table- 2**).

Contraceptive and counseling-related characteristics

From the total number of women interviewed, about three-fourths (74.4%) used other contraceptive methods before LARCs. For instance, of these women, about 150(50.3%) had been using injectables. Regarding method selection, nearly two-thirds (64.7%) of women decided to use LARCs by themselves. Among the type of LARCs, 220(54.7%) women used Implanon. Among the total study participants, those women counseled about the benefit, side effects, and effectiveness of LARCs were 199 (49.5%), 203 (50.5%), and 227(56.5%) respectively. One hundred ninety-five (48.5%) of respondents mentioned having been satisfied with the service given before the insertion (**Table-3**).

The magnitude of long-acting reversible contraceptive discontinuation

The study revealed that among the study participants who ever used LARCs, 18.4%, 57.2%, and 81% of women discontinued the service at 12, 24, and 36 months, respectively. Besides, 38.8%

and 23.8% of women discontinued it at 12-24 and 24-36 months, respectively. Among the study participants, only 76(19%) of women used LARCs for at least 36 months duration. Regarding the types of LARCs discontinued at 12 months, the proportion of women who discontinued Implanon, Jadelle, and IUCD was 51.4%, 31.1%, and 17.6% respectively (**Figures 2, 3, and 4**).

Reason for discontinuation of long-acting reversible contraceptives

More than one-third (37.4%) of women discontinued LARCs within three years of utilization, mainly due to emerging side effects, followed by 84 (25.7%) who desired to have a child (**Figure -5**). The most common complaints among women who developed side effects were menstrual disruption 58(47.7%), weight gain 24(19.8%), insertion site arm pain 17(14.0%), unusual headache 16(12.8%), infection 6(4.7%) and method inconvenience1(1.2%) (**Figure -5**).

Factors associated with discontinuation of reversible long-acting contraceptives

From the bivariate logistic regression analysis, factors such as age, educational status, occupational status, desire to become pregnant, counseling about the benefit of LARCs, counseling on the effectiveness of LARCs, and satisfaction with the service given have a p -value ≤ 0.25 and are considered for multivariable logistic regression.

Factors from bivariate logistic regression analysis with a p -value ≤ 0.25 were entered into multivariable logistic regression analysis. With multivariable logistic regression analysis, considering a p -value ≤ 0.05 , the independent predictors of LARCs discontinuation were maternal occupation, desire to be pregnant shortly, lack of counseling about the side effects of LARCs, no counseling on the effectiveness of LARCs, and not being satisfied with LARCs service provision (**Table 4**).

Among socio-demographic variables, the occupational status of women was found to be associated with the discontinuation of LARCs. Women having the occupation of housewife were 10% times less likely to discontinue LARCs than women having the occupation of merchant (AOR=0.09; CI: 0.04, 0.17). The odds of discontinuing LARCs among women who desire to be pregnant were 3.7 times higher than counterparts (AOR=3.7; CI: 1.90, 7.14). Regarding counseling status of women about the side effects of LARCs, those who were not obtained counseling would be 2.6 times more likely to discontinue LARCs than their counterparts (AOR=2.6; CI: 1.10, 6.01). Those Women who were not counseled about the effectiveness of LARCs were 2.8 times more likely to discontinue LARCs than their counterparts (AOR=2.8; CI: 1.43, 5.39). Women who were not satisfied with the service given were 2.8 times more likely to discontinue LARCs than their counterparts (AOR=2.8; CI: 1.15, 6.95) (**Table 4**).

Discussions

Nevertheless, 30% of maternal and 10% of infant deaths reduced by LARCs which is one of the most practical and cost-effective method by ensuring that every pregnancy is desired and every birth is safe[3, 4], the overall prevalence of LARCs were found to be 18.4%. Current study finding was consistent with the studies conducted at Hawassa City from Southern Ethiopia of 17.7% [29], Bahir Dar City from Northwest Ethiopia of 17.2% [32] and slightly less than urban public health facilities of Ethiopia[32].

The current study's percentage of LARCS discontinuation at 36 months was lower than the 65% found in a study conducted in Debre Tabor, Northwest Ethiopia. This variation could be a result of the subjects or the design of the study. Participants from Debre Tabor, in contrast to the current study, included both urban and rural populations. Urban inhabitants might be more aware of the consequences of contraception. Additionally, efforts may have been taken to enhance

counseling, particularly for the Women in the current study who had issues with weight gain and menstruation disruption. The current study result was also lower than studies conducted in Ambo 38.2%(26), Dale District 23.4%(27), and Arsi zone 25%(19). This discrepancy might be due to differences in the study design, sample size, duration of the study, lack of pre-insertion counseling, satisfaction with the given service, and sociocultural differences of respondents across the study areas.

Among the types of LARCs discontinued in this study, the prevalence of Implanon (41.8%) and IUCD (15.2%) were found to be less than secondary data analyzed from EDHS of 61% and 45% by 36 months respectively (12). This might be due to the discrepancy in the population structure of the national pattern and variation in the duration of utilization before discontinuation.

The study revealed that the main reason for discontinuation of LARCs was emerging side effects (37.4%), desire to have a child (25.7%), method shift (21.7%), and health concern (13.5%). Menstrual disruption (47.7%) was the main side effect faced among discontinuers. Menstrual irregularity may not cause serious health problems but interferes with daily activity, especially sexual activity with their husband. Discontinuation due to side effects was a result of inadequate counseling and information on possible side effects before and following the insertion of LARCs. Giving adequate counseling and information on the side effects and effectiveness of LARCs likely improves service satisfaction and ensure the continuation of LARCs [13, 16, 29].

Among the independent predictors, the occupational status of women was found to be associated with discontinuation LARCs. Women in the occupation of housewives were 91% more likely to discontinue LARCs than merchants. This may be because women having the occupation of

merchant might be more likely to be educated and have access to information about modern family planning from different sources.

The current study also revealed that the odds of discontinuing LARC among women who desired to be pregnant shortly were 3.6 times higher than their counterparts. This is consistent with studies conducted in Kenya[33], Bahir Dar City [34], Debre Markos town[35], and Hawassa City[29]. This might be because the culture of the community influences married women to bring a child to ensure the ability of the women to bear a child. In addition to this, 62.4% of women in the present study were aged 18–29 years. Since the majority of study participants in this study were young, they might intend to have more children and discontinue the contraceptives.

The present study showed that women who were not counseled about the side effects of the LARCs were 2.6 times more likely to discontinue LARCs compared to their counterparts which are consistent with the study done in Ethiopia, systematic review[13], Hawassa[29], and Kersa district[36]. Besides, the current study also revealed that counseling women about the effectiveness of LARCs were negatively associated with LARCs discontinuation. The most cited reason for this effect was effective counseling at insertion time on possible side effects of the method will help the mother to accept the possible side effects and minimize removal of the method early. In addition to this, the use of appropriate counseling type and timing might also help the women to cope with minor side effects and strengthen the continuation of the method.

According to the study, women who were unsatisfied were 2.8 times more likely to stop using long-acting reversible contraceptives (LARCs) than those who were satisfied. This might be the result of incorrect pre-insertion counselling and a lack of seclusion, which led to frustration and quitting. This finding is consistent with a study conducted in Makelle City, Dale district,

Kambata zone in southern Ethiopia, and Vietnam(18,21,27,36). This might be due to women's lack of interest in the method chosen. Furthermore, the confidentiality of the service, elucidation of the service supplier, communication skills, and other service provisions during the insertion of LARCs may contribute to being discontinuity of long-acting reversible contraceptives. Therefore, providing quality reproductive health services to clients can lead to better satisfaction and use of the services.

Limitation

The study's cross-sectional design prevented it from determining the cause-and-effect link. In addition, the possibility of recall bias could not be ruled out, and to study determinant factors, a cross-sectional design study has limitations.

Conclusion

In this study, the overall discontinuation of LARC among women was high (57.2%). The independent predictors of LARC discontinuation among women were maternal occupation as a housewife, the desire for pregnancy, not counseling on side effects, not counseling on the effectiveness of LARCs, and not being satisfied with the service given.

Recommendation

Address low community awareness through community-level awareness creation programs and health education to ensure long-acting reversible contraceptive use. Provide pre-insertion counseling on side effects, effectiveness, and benefits. Early management and reassurance can decrease discontinuation and enhance retention. A further comprehensive study of this area that includes health professionals and is community-based is recommended.

Consent for publication

All authors read the manuscript and have provided their consent to publish

Availability of data

The data sets used and/or analyzed during the current study are available from the corresponding author upon reasonable request.

Competing Interest

All authors declare that they have no Competing Interest

Funding

The funding source for per diem of data collectors and supervisors of this research was Salale University. The funding body has no role in the design of the study and collection, analysis, and interpretation of data and in writing this manuscript.

References

1. <https://www.cdc.gov/reproductivehealth/contraception/unintendedpregnancy/index.htm>. Accessed Date. 2023 March 21].
2. <https://www.who.int/news/item/25-10-2019-high-rates-of-unintended-pregnancies-linked-to-gaps-in-family-planning-services-new-who-study>. Accessed date [cited 2023 March 21].
3. Cleland J, et al., *Contraception and health*. The Lancet, 2012. **380**(9837): p. 56.
4. USAID, *Holistic Approach Enhances Family Planning Programs. Respond's Experience with the SEED Programming Model*. 2014. **27**: p. 1-8.
5. Kolman KB, Hadley SK, and Jordahliafrato MA., *Long-acting reversible contraception: who, what, when, and how*. J Fam Pract 2015. **64**(479): p. 84.
6. Peterson HB and Curtis KM., *Clinical practice. long-acting methods of contraception* N Engl J Med, 2005. **353**: p. 75.
7. *Long-Acting Reversible Contraception: Implants and Intrauterine Devices*. November 2017 2021 [cited 2023 March 26].
8. Wolters Kluwer Health Inc. *ACOG Committee opinion no. 450: increasing use of contraceptive implants and intrauterine devices to reduce unintended pregnancy*. Obstet Gynecol 2018 [cited 2023 March 26].
9. Blane A, Curtis S, and Croft T. *Does contraceptive discontinuation matter? Quality of care and fertility consequences*, in *MEASURE Evaluation Technical Report*1999, Carolina Population Center: University of North Carolina at Chapel Hill.
10. Staveteig, et al., *Uptake and Discontinuation of LongActing Reversible Contraceptives (LARCs) in Low-Income Countries*. DHS Analytical Studies; Rockville, Maryland, USA. ICF International, 2015(54).

11. Ali, et al., *Long-term contraceptive protection, discontinuation, and switching behavior: intrauterine device (IUD) use dynamics in 14 developing countries*. London: World Health Organization and Marie Stopes International. 2011.
12. Central Statistical Agency., *Ethiopia demographic and health survey 2016, 2017 CSA and ICF*: Addis Ababa, Ethiopia.
13. Gebeyehu NA, et al., *Discontinuation of long-acting reversible contraceptive use and its determinants among women in Ethiopia: Systematic review and meta-analysis*. Front. Public Health, 2022.
14. Ezegwui HU, et al., *The discontinuation rate and reasons for discontinuation of implanon at the family planning clinic of University of Nigeria teaching hospital (UNTH) Enugu, Nigeria* Niger J Med, 2011. **20**: p. 448–50.
15. Haregwa Asnake Weldekidan, et al., *Discontinuation rate of long-acting reversible contraceptives and associated factors among reproductive-age women in Butajira town, Central Ethiopia*. Women's Health 2022. **18**: p. 1–11.
16. Tadele Girum Adal., *Early Discontinuation of Long Acting Reversible Contraceptives among Married and in Union Women: A Systematic Review and Meta-analysis*. Ann Med Health Sci Res, 2017. **7**: p. 113-118.
17. Grunloh DS, et al., *Characteristics associated with discontinuation of long-acting reversible contraception within the first 6 months of use*. Obstet Gynecol 2013. **122**: p. 1214–21.
18. Cohen R, Sheeder J, and Teal SB., *Cohen R, Sheeder J, Teal SB. Predictors of Discontinuation of Long-Acting Reversible Contraception Before 30 Months of Use by Adolescents and Young Women*. J Adolesc Health 2019. **65**: p. 302-295.
19. Tesfaye K., *Early discontinuation of implant and its associated factors among women who ever used implant in 2017/2018 in Hawassa Health Centers, Southern Ethiopia, 2019*. J Med Case Rep Rev, 2019. **2**(7): p. 320-330.
20. Asari VG., *Determinants of contraceptive use in Kerala: the case of son/daughter preference*. J Fam Welf, 1994. **40**: p. 19-25.
21. Bhat PN and Halli SS., *Factors influencing continuation of IUD use in South India: evidence from a multivariate analysis*. J Biosoc Sci. , 1998. **30**: p. 297-319.
22. Nageso A and Gebretsadik A., *Discontinuation rate of Implanon and its associated factors among women who ever used Implanon in Dale District, Southern Ethiopia*. BMC Womens Health, 2018. **18**(1): p. 189.
23. Park MH, N. TH, and Ngo TD., *Dynamics of IUD use in Vietnam: implications for family planning services at primary health care level*. Int J Womens Health 2011. **3**: p. 429–434.
24. Youssef RM., *Youssef RM. Contraception use and probability of continuation: a community-based survey of women in southern Jordan*. East Mediterr Health J, 2005. **11**: p. 545-558.
25. Federal Democratic Republic Ethiopia., *National Guideline for Family Planning Services in Ethiopia* 2018: Addis Ababa, Ethiopia.
26. Jain AK and Winfrey W., *Contribution of contraceptive discontinuation to unintended births in 36 developing countries*. Stud Fam Plann 2017; **48**: p. 269–78.
27. Fekadu GA, et al., *Factors associated with early long-acting reversible contraceptives discontinuation in Ethiopia: evidence from the 2016 Ethiopian demographic and health survey*. Arch Public Health, 2020. **78**(1): p. 1-10.

28. Bahamondes L, et al., *Long-acting reversible contraceptive (LARCs) methods*. Best Pract Res Clin Obstet Gynaecol, 2020. **66**: p. 28-40.
29. Belay Amare Abebe, Nega Assefa, and Bezatu Mengistie., *Discontinuation of Reversible Long-Acting Contraceptive and Associated Factors among Female Users in Health Facilities of Hawassa City, Southern Ethiopia: Cross-Sectional Study*. Open Access Journal of Contraception, 2020. **11** p. 113–123.
30. <https://www.who.int/news/item/25-10-2019-high-rates-of-unintended-pregnancies-linked-to-gaps-in-family-planning-services-new-who-study>. Accessed date. [cited 2023 March 24].
31. Mamo K, Siyoum M, and et al., *Premature Implanon Discontinuation and Associated Factors Among Implanon User Women in Ambo town, Central Ethiopia*. J Heal Med Nurs 2019. **58**: p. 39–46.
32. Mihretie GS, A.S., Abebaw Y, et al., *Factors associated with discontinuation among long-acting reversible contraceptive users: a multisite prospective cohort study in urban public health facilities in Ethiopia*. BMJ Open, 2022. **12**.
33. Kungu W, Agwanda A, and Khasakhala A., *Prevalence of and factors associated with contraceptive discontinuation in Kenya*. Afr J Prm Health Care Fam Med, 2022. **14**(1): p.
34. Melesse YD, et al., *Discontinuation of Long-Acting Reversible Contraceptive Methods and Associated Factors among Women in Health Facilities of Bahir Dar City, Northwest Ethiopia: A Cross-Sectional Study*. Reprod Syst Sex Disord, 2022. **11**(319).
35. Melese Siyoum, et al., *Implanon Discontinuation Rate and Associated Factors among Women who Ever Used Implanon in the Last Three Years in Debre Markos Town, Northwest Ethiopia, 2016, Cross-Sectional Study*. ARC Journal of Public Health and Community Medicine 2017. **2**(1): p. 8-16.
36. Gali Nega, Muluemebet Abera, and Afework Tadele., *Discontinuation rate and associated factors among contraceptive implant users in Kersa district, southwestern Ethiopia*. BMC Archives of Public Health, 2021. **79**(75).

Table 1: Socio-demographic characteristics of long-acting reversible contraceptive users in health facilities of Shashemene town, Southern Ethiopia, 2022(n=402).

Variable	Frequency	Percentage
Age, years		
<20	25	6.2
20-29	226	56.2
30-39	117	29.1
40+	34	8.5
marital status		
Married	368	91.5
Single	22	5.5
Divorced/separated	7	1.7
Widowed	5	1.2
Educational status (n=402)		
no formal education	22	5.5
Primary education	253	62.9
Secondary education	91	22.6
Higher education/ university/College	36	9.0
Occupational (n=402)		
Housewife	230	57.2
Government employed	43	10.7
Private employed	44	10.9
Merchant	85	21.1
Family income		
<2000	29	7.2
2000-3000	220	54.7
3000+	153	38.1

Table 2: Obstetric history of women who ever used LARC 2021/2022, in Shashemene town, Oromia, South Ethiopia, 2022(n=402).

Variable	Frequency	Percentage
Parity		
0	3	0.7
1-2	137	34.1
3-4	203	50.5
5+	59	14.7

Living children (n=198)		
0	9	2.2
1-2	139	34.6
3-4	198	49.3
5+	56	13.9
History of abortion		
Yes	122	30.3
No	280	69.7
Desire of pregnancy		
Yes	177	44.0
No	225	56.0

medRxiv preprint doi: <https://doi.org/10.1101/2023.07.25.23293161>; this version posted July 27, 2023. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY 4.0 International license.

Table 3: Contraceptive and counseling-related characteristics of women who ever used LARCs in 2021/2022 in Shashemene town, Oromia, Southern Ethiopia, 2022 (n=402).

Variable	Frequency	Percentage
Contraceptive use before LARCs		
Yes	299	74.4
No	103	25.6
Type of contraceptive used before LARCs		
Injection	150	50.2
OCP	93	31.1
OCP nad Injection	45	15.2
Others	10	3.5
Method selection		
My self	260	64.7
My husband	12	3.0
Health professional	83	20.6
Others (Health Extension workers, Neighbor...)	47	11.7
Type of LARC used		
Implanon	220	54.7
Jaddles	110	27.4
IUD	72	17.9
Counseled about benefit of LARC		
Yes	199	49.5
No	203	50.5
Counseled about side effect		
Yes	203	50.5
	199	49.5

No		
Counseled about effectiveness		
Yes	227	56.5
No	175	43.5
Information participants got during counselling (Satisfied)		
Yes	216	53.7
No	186	46.3


Table 4: Bivariate and Multivariate analysis of factors associated with discontinuation of long-acting reversible contraceptives in Shashemene town southern Ethiopia, 2022 (n=402).

Variables	Discontinuation of LARCs		COR (95%CI)	P-value	AOR (95%CI)	P-value
	Yes%	No%				
Age						
<20	13(52.0%)	12(48.0%)	1			
20-29	162(71.7%)	64(28.3%)	0.43(0.18- 0.98)	0.047	0.59(0.17 - 1.98)	0.394
30-39	41(35.0%)	76(65.0%)	2.00(0.84 -4.80)	0.117	2.34(0.68 - 8.02)	0.175
40+	14(41.2%)	20(58.8%)	1.54(0.54 -4.38)	0.411	5.53(0.26 -24.16)	0.073
Educational status						
no formal education	10(45.5%)	12(54.5%)	1		1	
Primary education	191(75.5%)	62(24.5%)	0.27(0.11- 0.65)	0.004	0.47(0.14 - 1.56)	0.220
Secondary education	24(26.4%)	67(73.6%)	2.33(0.89 -6.08)	0.085	4.82(0.24 - 18.67)	0.093
Higher education	5(13.9%)	31(86.1%)	5.17(1.46-18.27)	0.011	8.24(0.41 - 47.96)	0.099
Occupational status						
Housewife	184(80.0%)	46(20.0%)	0.09(0.05 -0.17)	0.000	0.09(0.04 - 0.17) *	0.000
Government employee	4(9.3%)	39(90.7%)	3.83(1.23-11.90)	0.020	0.69(0.15 - 3.09)	0.630
Private employee	18(40.9%)	26(59.1%)	0.56(0.26 - 1.22)	0.147	0.45(0.16 - 1.27)	0.135
Merchant	24(28.2%)	61(71.8%)	1		1	
Desire of pregnancy						
Yes	104(46.2%)	121(53.8%)	2.87(1.89 - 4.36)	0.000	3.69(1.90-7.14) *	0.000
No	126(71.2%)	51(28.8%)	1		1	
Counseled about benefit of LARC						
Yes	74(37.2%)	125(62.8%)	5.60(3.63 -8.66)	0.000	0.79(0.29 - 2.12)	0.643
No	156(76.8%)	47(23.2%)	1		1	
Counseled side effect of LARC						
Yes	152(76.4%)	47(23.6%)	1		1	
No	78(38.4%)	125(61.6%)	5.18(3.36 -7.98)	0.000	2.57(1.10 -6.01) *	0.029
Counseled effectiveness						

of LARC						
Yes	122(69.7%)	53(30.3%)	1		1	
No	108(47.6%)	119(52.4%)	2.53(1.67 - 3.83)	0.000	2.77(1.43 - 5.39) *	0.003
Satisfied with the service given						
Yes	162(78.3%)	127(65.1%)			1	
No	68(34.9%)	45(21.7%)	6.72(4.31-10.46)	0.000	2.83(1.15 -6.95) *	0.023

*p<0.05, OR- Odds Ratio, AOR- Adjusted Odds Ratio

medRxiv preprint doi: <https://doi.org/10.1101/2023.07.25.23293161>; this version posted July 27, 2023. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a [CC-BY 4.0 International license](#).



medRxiv preprint doi: <https://doi.org/10.1101/2023.07.25.23293161>; this version posted July 27, 2023. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY 4.0 International license.

Figure 1: Six months' enrollment in Health institutions of Shashemene town administration in two public hospitals, five HCs and two NGO clinics from family planning registration book, for sample selection (December - June/2022).




Figure 2: proportion of LARC discontinuation rate among LARC users in Shashemene town, Southern Ethiopia, 2022.




Figure 3. LARC discontinuation <2 years by method




Figure 4. Duration of LARCs utilization before discontinuation

Reason for discontinuation of LARCs





Figure 4- Reason for discontinuation among women who ever used LARC in 2021/2022 in Shashemene town, Southern Ethiopia (n=230).



medRxiv preprint doi: <https://doi.org/10.1101/2023.07.25.23293161>; this version posted July 27, 2023. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY 4.0 International license.

Figure 5. side effects reported by women