

1 **The plasma proteome of plant-based diets: analyses of 1463 proteins in 50,000 people.**

2 Tammy Y.N. Tong¹, Karl Smith-Byrne¹, Keren Papier¹, Joshua R. Atkins¹, Mahboubeh
3 Parsaeian¹, Timothy J. Key¹ and Ruth C. Travis¹

4 *1. Cancer Epidemiology Unit, Nuffield Department of Population Health, University of*
5 *Oxford, Oxford, UK*

6

7 Corresponding author: Tammy Y.N. Tong

8 Postal address: Cancer Epidemiology Unit, University of Oxford, Richard Doll Building, Old

9 Road Campus, Oxford, OX3 7LF

10 Tel: +44 (0) 1865 289653

11 Email: tammy.tong@ndph.ox.ac.uk

12 **Abstract**

13 *Background and objectives:* Circulating proteins are integral to many biological processes.

14 We aimed to assess differences in the plasma proteome between people of different dietary
15 groups defined by degree of animal food consumption.

16 *Methods:* The UK Biobank recruited middle-aged adults (mostly 40 to 69 years) throughout
17 the UK between 2006-2010. Relative concentrations of 1463 plasma proteins were quantified
18 using the Olink Proximity Extension Assay on blood samples from 49,326 participants, who
19 were also asked to report their ethnicity and consumption of red and processed meat, poultry,
20 fish, dairy and eggs. We defined six diet groups among the white British participants (23,116
21 regular meat eaters, 23,323 low meat eaters, 484 poultry eaters, 1074 fish eaters, 722
22 vegetarians, and 54 vegans), and two diet groups among the British Indians (390 meat eaters
23 and 163 vegetarians). We used multivariable-adjusted linear regressions to assess differences
24 in protein concentrations between diet groups, with correction for multiple testing.

25 *Results:* We observed significant differences in many plasma proteins by diet group (683
26 proteins in white British participants, 1 in British Indians), in particular many proteins that
27 are majority expressed in the digestive system. Of the biggest differences, compared with
28 regular meat eaters, the non-meat eaters had significantly higher FGF21 (e.g. +0.40 SD in
29 vegetarians), GUCA2A (+0.33), FOLR1 (+0.32), IGFBP2 (+0.31) and DSG2 (+0.30); all
30 groups except the vegans had lower HAVCR1 (-0.38 in vegetarians). The observed
31 differences were generally similar in direction in both ethnicities.

32 *Discussion:* In this first comprehensive assessment of plasma proteins by diet group, we
33 identified many differences in proteins between groups defined by animal food consumption;
34 this variation in protein levels suggests differences in various biological activities, including

35 gastrointestinal tract and kidney function, which may relate to differences in future disease

36 risk.

37 *Keywords:* vegetarians, vegans, proteomics

38 **Introduction**

39 Proteins are essential for many bodily functions including supporting cell and tissue growth
 40 and structural integrity, and for enabling a range of enzymatic, biochemical, signalling and
 41 transport functions across different systems in the human body. Proteomics describes the
 42 large-scale study of multiple proteins and their interconnected pathways, the investigation of
 43 which allows a more comprehensive understanding of disease mechanisms [1]. Protein
 44 expression can be altered by environmental factors [2], including diet, which is an important
 45 exogenous source of dietary protein, and which provides the nine essential amino acids that
 46 cannot be synthesised endogenously but are nonetheless necessary for protein synthesis in the
 47 human body. Previous studies have shown that differences in dietary habits can substantially
 48 impact dietary intakes of protein and both dietary intakes and circulating concentrations of
 49 amino acids [3]. However, there is scant evidence on how diet may affect the proteome,
 50 partly due to the limited availability of proteomics data in large scale studies, and no prior
 51 studies have investigated how vegetarian and vegan diets may influence the proteome.
 52 Nonetheless, protein levels have an established role in the aetiology of multiple diseases [4],
 53 and the examination of dietary influences on the proteome may offer unique insights into
 54 understanding how vegetarian and vegan diets may affect future disease risk.

55 The aim of this study is to provide a detailed description of circulating protein concentrations
 56 in people with varying degrees of animal food consumption, using data from the UK
 57 Biobank.

58

59 **Methods**

60 *Study Population*

61 The UK Biobank is a prospective cohort study of around 500,000 middle-aged people
 62 (recruitment target 40-69 years), recruited from across the United Kingdom between 2006
 63 and 2010. The scientific rationale and design of the UK Biobank have been described in
 64 detail previously [5]. In brief, participants were identified from National Health Service
 65 registers, and were invited to join the study if they lived within travelling distance (~25 km)
 66 of one of the 22 assessment centres in England, Wales and Scotland. People who consented
 67 to participate in the study attended a baseline visit at the assessment centre where they
 68 completed a touchscreen questionnaire which asked about their lifestyle (including diet,
 69 alcohol consumption, smoking status, physical activity), socio-demographic characteristics
 70 and general health and medical history. All participants were also given a verbal interview,
 71 and had their physical measurements and blood samples taken by trained staff. Permission for
 72 access to patient records for recruitment was approved by the Patient Information Advisory
 73 Group (now the National Information Governance Board for Health and Social Care) in
 74 England and Wales and the Community Health Index Advisory Group in Scotland, and all
 75 participants gave informed consent to participate using a signature capture device at the
 76 baseline visit.

77

78 *Ethnicity classification*

79 On the touchscreen questionnaire, participants were asked to self-identify their ethnicity as
 80 ‘White’, ‘Mixed’, ‘Asian or Asian British’, ‘Black or Black British’, ‘Chinese’, ‘Other ethnic
 81 group’, ‘Do not know’, or ‘Prefer not to answer’, with further sub-categories under each
 82 option. Participants were included for our analyses if they self-identified as ‘White’, or as
 83 ‘Asian or Asian British’ and subsequently as ‘Indian’, hereafter referred to as ‘white British’
 84 and ‘British Indian’. The white British population was included as it made up the majority of

the participants in UK Biobank, while the British Indian population was included because of the large proportion of vegetarians in this population group (25% compared to less than 2% in the overall population). The number of vegetarians in the other ethnic groups was too small to allow valid comparisons by diet groups, and therefore people of other ethnic groups were excluded from these analyses.

90

Diet group classification

Participants were classified into diet groups based on self-reported dietary data from the touchscreen questionnaire completed at recruitment. Participants were asked to report their frequency of consumption of processed meat (including processed poultry), unprocessed red meat (beef, lamb or mutton, pork), unprocessed poultry (such as chicken or turkey), oily fish, and other types of fish, with the possible responses ranging from “never” to “once or more daily”. Participants were also asked whether they never consumed dairy and eggs or foods containing eggs. Based on their responses to these questions, the white British participants were classified into six diet groups: regular meat eaters (red and processed meat consumption >3 times per week), low meat eaters (red and processed meat consumption ≤3 times per week), poultry eaters (participants who ate unprocessed poultry but no red and processed meat), fish eaters (participants who ate fish, but not red and processed meat, or poultry), vegetarians (participants who did not eat any meat or fish), and vegans (participants who did not eat any meat, fish, dairy products or eggs). The British Indian participants were classified into two diet groups: meat eaters (ate any combination of red or processed meat or poultry) and vegetarians; the numbers of fish eaters and vegans were low in the British Indians, and thus were not included in these analyses.

108

109 *Assay for plasma proteins*

110 Non-fasting blood samples were collected from all UK Biobank participants by trained
 111 personnel (either a phlebotomist or a nurse) except in a small proportion (0.3%) of
 112 participants who declined, were deemed unable to, or where the attempt was abandoned for
 113 technical or health reasons. Proteomic profiling was conducted using the Olink Proximity
 114 Extension Assay on blood plasma samples in 54,306 participants selected as part of the UK
 115 Biobank Pharma Proteomics Project (UKB-PPP). The samples were shipped on dry ice to
 116 Olink Analyses Service, Uppsala, Sweden for analysis; details of the selection procedures for
 117 inclusion in UKB-PPP and the technical details of the proteomics assays have been described
 118 elsewhere [6]. In brief, the Olink Explore 1536 platform used in this study is an antibody-
 119 based assay which measures the relative abundance of 1,472 protein analytes, including 1,463
 120 unique proteins, distributed across four 384-plex panels: inflammation, oncology,
 121 cardiometabolic, and neurology. The Olink Explore platform is based on proximity extension
 122 assays that are highly sensitive and reproducible with low cross-reactivity. Relative
 123 concentrations of the 1,463 unique proteins were readout by next-generation sequencing.
 124 Measurements are expressed as normalized protein expression (NPX) values which are log-
 125 base-2 transformed. Protein values below the limit of detection (LOD) were replaced with the
 126 LOD divided by the square root of 2 [7]. Protein concentrations were subsequently inverse
 127 rank normal transformed. All results for differences in protein concentrations by diet groups
 128 may be interpreted as SD differences.

129

130 *Inclusion and exclusion criteria*

131 Of the 54,306 participants selected for proteomic profiling, 52,705 participants remained after
 132 the quality control procedures. Participants were further excluded if they were not of white

British or British Indian ethnicity (n=2964), could not be classified into one of the pre-specified diet groups (n=414), or had missing information for fasting time (n=1). After the exclusions, 49,326 participants (48,773 white British and 553 British Indian) were included in the analyses. A participant flow chart of the inclusion criteria is shown in **Supplementary figure 1**.

Statistical analyses

Baseline characteristics of UK Biobank participants included in these analyses were tabulated by six diet groups in the white British population and by two diet groups in the British Indian population, as mean (SD) for continuous variables and number (%) for categorical variables. We used multivariable-adjusted linear regressions to estimate differences in protein concentrations by diet group, separately by ethnicity, using regular meat eaters as a reference group in the white British participants, and meat eaters as a reference group in the British Indians. The model was adjusted for age at recruitment (5 year categories), sex, region (London, North-West England, North-East England, Yorkshire, West Midlands, East Midlands, South-East England, South-West England, Wales, Scotland), fasting status (0–1, 2, 3, 4, 5, 6–7, ≥8 hours), body mass index (BMI; <20, 20.0–22.4, 22.5–24.9, 25.0–27.4, 27.5–29.9, 30.0–32.4, 32.5–34.9, ≥35 kg/m², unknown), alcohol consumption (<1, 1–7, 8–15, ≥16 g/d, unknown), smoking status (never, previous, current <15 cigarettes/day, current ≥15 cigarettes/d, unknown) and physical activity (<10, 10–49.9, ≥50 excess metabolic equivalent of task, hr/wk, unknown). Wald tests were used to assess overall heterogeneity between diet group in each ethnicity, and for pairwise comparisons of each of the other diet groups compared to the reference group (regular meat eaters in white British participants/meat eaters in British Indians). Heterogeneity between vegetarians and vegans in the white British

population was assessed based on post-estimation linear combinations of parameters for all proteins.

To account for multiple testing while considering the high correlations between the circulating proteins, we conducted a principal component analysis of the circulating proteins in the complete proteins dataset before exclusions for these analyses, and determined that the first 639 principal components explained 95% of the total variation in the exposure data. Consequently, the effective number of independent tests [8] was determined to be 639, and the statistical significance level was set to be $p\text{-value } 0.05/639 = 0.000078$. For the top proteins identified (the top 10 significant proteins in each pairwise comparison against the reference group, based on ranking of p-values), we conducted additional sensitivity analyses to evaluate the extent to which the associations may be influenced by key covariates including BMI, smoking and alcohol consumption, by presenting models with and without adjustment of these variables. Additionally, we also restricted the analyses to people who self-reported to be in good and excellent health. All analyses were performed using Stata version 17.0 (StataCorp, TX, USA). All figures were generated using R version 4.2.1, the forest plots using “Jasper makes plots” package version 2-266 [9].

Integrating publicly available information on gene expression

To further understand the biological context for the proteins of interest, we extracted single cell mRNA expression from the Human Protein Atlas [10] to describe expression for any proteins that were significantly different in one or more comparisons. Normalized expression levels were extracted for genes in 30 different human tissues and 82 cell-types. Gene expression specificity at the cell or tissue type level was calculated as the ratio of each gene cell-type or tissue expression to the total expression of each gene across all cell or tissue

types. We subsequently identified genes for proteins that were majority expressed (more than 50% of total expression) in at least one cell or tissue type, and mapped these to their likely candidate cell and tissue of origin where possible.

Results

Baseline characteristics

The baseline characteristics of the study population are shown by ethnicity and diet group in Table 1. Compared with white British regular meat eaters, the non-meat eaters were on average slightly younger, more likely to be women, and to reside in London. They were less likely to be overweight, reported lower alcohol consumption, lower current smoking, and more physical activity. Compared with British Indian meat eaters, the vegetarians were more likely to be women and reported lower alcohol consumption, lower current smoking and slightly lower physical activity, but BMI was not noticeably different between the two groups. Fasting time was not meaningfully different by diet group in both ethnicities.

Differences in plasma proteins by diet group

The plasma proteins that were significantly different in white British vegetarians and vegans compared with regular meat eaters are shown as volcano plots in **Figure 1**. Significant differences in proteins in white British low meat eaters, poultry eaters and fish eaters compared with regular meat eaters, and in British Indian vegetarians compared with meat eaters are shown as volcano plots in **Supplementary figures 2-6**. Overall, 683 plasma proteins were significantly different by diet group (based on $p\text{-heterogeneity} < 0.000078$) in the white British participants, including 535 plasma proteins that were significantly different

in one or more pairwise comparisons between regular meat eaters and at least one of the other diet groups. This includes 296 proteins (23 higher, 273 lower) in low meat eaters compared with regular meat eaters, 59 (9 higher, 50 lower) in poultry eaters, 157 (104 higher, 53 lower) in fish eaters, 219 (194 higher, 25 lower) in vegetarians and 15 (12 higher, 3 lower) in vegans; and 2 proteins were significantly different between vegetarians and vegans (AHSP and FOLR2 were both higher in vegans). The proteins identified are involved in a range of different biological pathways. Of the 535 plasma proteins that were significantly different in one or more pairwise comparisons, 97 are majority expressed (protein mRNA >50% of total body mRNA expression) in one tissue type, while the other proteins have limited tissue specificity. In particular, these included 39 proteins that are majority expressed in the digestive system (including 16 in the liver, 8 in the pancreas, 8 in the small intestine), 30 in the nervous system (27 in the brain), and 12 in the respiratory system (7 in the lung) (**Supplementary figure 7**).

In the British Indian population, only 1 protein (SUMF2, which is higher in vegetarians) was significantly different when comparing vegetarians and meat eaters, likely owing to the much smaller number of British Indians in the study. However, the differences in many proteins were directly consistent between the two ethnicities, as illustrated by the forest plots of top proteins by diet group and ethnicity, with the six diet groups in the white British participants and the two diet groups in British Indians (**Figures 2-3, Supplementary figures 8-10**). The complete results of all proteins are shown in **Supplementary tables 1**.

Generally, in the multivariable-adjusted model, many proteins showed a gradient effect in magnitude of differences across diet groups by degree of animal food exclusion, from regular meat eaters to vegans. Figure 2 shows the top 10 proteins in white British vegetarians (based on the ranking of pairwise p-values in vegetarians compared with regular meat eaters), and Figure 3 shows the top 10 proteins in white British vegans. Of the top proteins in vegetarians

(Figure 2), compared with regular meat eaters, vegetarians and vegans both had significantly higher FGF21 (+0.40 SD in vegetarians; +0.74 SD in vegans), GUCA2A (+0.33; +0.79), FOLR1 (+0.32; +0.61) and IGFBP2 (+0.31; +0.55) after correction for multiple testing, whereas vegetarians but not vegans had significantly higher ART3 (+0.34), LAYN (+0.32), PCSK9 (+0.32), and lower CDHR2 (-0.26), HAVCR1 (-0.38) and CNDP1 (-0.39). In addition to the aforementioned proteins, vegans also had substantially higher FOLR2 (+0.69), GAS6 (+0.68), DSG2 (+0.68), THY1 (+0.62), MANSC1 (+0.58), but lower EGFR (-0.55) than regular meat eaters (Figure 3); the differences in these proteins were directionally consistent but less extreme in the vegetarians.

The top 10 proteins in white British low meat eaters, poultry eaters and fish eaters compared with regular meat eaters are shown in Supplementary figures 8, 9 and 10 respectively. Of these top proteins, none was uniquely different in one diet group only when compared with regular meat eaters. Similar to the observations in vegetarians, all three groups had substantially higher DSG2 and lower HAVCR1 compared with regular meat eaters; in both low meat eaters and poultry eaters HAVCR1 is the protein that exhibited the largest magnitude in difference across all proteins, while in fish eaters FGF23 showed the biggest difference. In addition, low meat eaters and poultry eaters both had lower OSM and CD99L2, the latter of which was higher in vegetarians. Low meat eaters, fish eaters and vegetarians all had lower ACP5 than regular meat eaters; this protein was lower but not significantly different in poultry eaters. In sensitivity analyses of the top proteins with different levels of covariate adjustment and limited to people of good and excellent health, results were similar (Supplementary tables 2).

Discussion

In this large British cohort, we observed differences in the levels of many proteins by diet group. This is the first comprehensive study on plant-based diets and plasma proteomics, and many of the substantial differences reported here are novel. In particular, we saw substantial differences in some proteins potentially associated with disease states such as HAVCR1 and DSG2 (associated with kidney disease [11] and inflammatory bowel disease, IBD [12], respectively), as well as proteins associated with nutritional status including IGFBP2 [13, 14], and possibly FGF21 [15] and the folate receptors FOLR1 and FOLR2 [16]. The magnitudes of differences in many proteins showed a gradient effect across the diet groups, and were directionally consistent between the white British and British Indian populations.

In terms of tissue mRNA expression, most of the proteins identified are not majority expressed in one specific tissue type, which suggests that they may be involved in a range of biochemical pathways and processes. Of the 97 proteins that were significantly different by diet group and also majority expressed in one tissue type, many were identified to be those expressed in the digestive system, which may signify dietary impact on digestive functions. We also see a number of proteins with high expression in the brain and lung, which may benefit from further research. Overall, the substantial differences in the protein concentrations observed likely reflect physiological differences between the different diet groups, as well as differences in underlying disease pathology and future disease risk.

No previous studies were found that examined plasma proteomics in vegetarians and vegans. A few previous studies have reported on proteomic profiles by other dietary patterns, including the Dietary Approaches to Stop Hypertension (DASH) diet, the Healthy Eating Index and the Mediterranean diet, as well as other data derived dietary patterns [17–20]. Whilst vegetarian and vegan diets share some common features with these other healthy dietary patterns, such as higher intakes of fruit, vegetables and whole grains, the diets are also inherently different in many other aspects. For example, while both the DASH diet and

Mediterranean diet recommend reducing red and processed meat consumption, they generally do not pose any restrictions on consumption of fish and fish products, or dairy products (particularly low-fat versions). In contrast, vegetarian and vegan diets are defined by the complete exclusion of animal food groups. Therefore, it is not surprising that while studies on the DASH diet also reported higher FOLR2 [17] and DSG2 [18] in people with higher adherence, we generally do not find similarities in the results.

FGF21

There are only a limited number of studies examining vegetarian and vegan diets with individual proteins. In support of our findings, higher concentrations of FGF21 in vegetarians and vegans have been shown in a small study of 36 each of omnivores, vegetarians and vegans [21]. FGF21 has been hypothesised to be high in vegans as a downstream effect of low intake of methionine [15], an essential amino acid that is known to be limiting in vegan diets [3]. Studies of FGF21 administration in rodents have found favourable effects on adiposity, lipid profiles, and non-alcohol fatty liver disease, but possible adverse effects on bone homeostasis [22]; randomised trials of FGF21 analogues also support the reduction in triglycerides [23] and improvement of fibrosis in patients with non-alcoholic steatohepatitis [24]. Consistent with these observations, we have previously reported lower BMI and body fat [25], more favourable lipid profiles [26] and lower heart disease risk [27], but also lower bone mineral density [25] and higher fracture risk [28] in vegetarians and vegans when compared with meat eaters; another study has reported lower odds of non-alcohol fatty liver disease in vegetarians than non-vegetarians [29].

GUCA2A

GUCA2A was substantially higher in all diet groups compared to regular meat eaters, with the exception of low meat eaters. GUCA2A activates the Guanylate Cyclase C receptor, which has a role in the maintenance of gut physiology including the increase of water movement into the intestinal lumen [30]. Previous studies have found that fasting plasma levels of GUCA2A are significantly lower in patients who have been diagnosed with Crohn's disease compared with healthy controls [31], and that loss of GUCA2A expression may be an important determinant of colorectal cancer [32]. Consistent with the observations in this study, a previous EPIC-Oxford study reported a higher frequency of bowel movements in vegetarians and vegans [33].

FOLR1 and FOLR2

We found both FOLR1 and FOLR2 to be higher in non-meat eaters, with vegans having particularly high levels. As folate receptors, FOLR1 and FOLR2 both bind to and import folic acid into cells, and are usually downregulated with repletion of folate, however, this process may also be mediated by homocysteine in a positive direction [16]. While vegetarians and vegans typically have high folate, they may also have high homocysteine due to low vitamin B12 intakes [34, 35], though no other studies have reported on folate receptor expression in vegetarians and vegans for comparison.

IGFBP2

Similar to our findings, a previous study in EPIC-Oxford has found that compared with meat-eaters, both vegetarians and vegans had higher concentrations of IGFBP2, a binding protein for IGF-1 [13]. The mechanism for higher IGFBP2 in vegetarians and vegans is not well established but has been suggested to be related to lower dietary intakes of essential amino

acids [13], protein (especially from dairy) [14], or lower total energy intake [36], which are all characteristic of vegetarian and vegan diets [3, 25, 37].

DSG2

Low and non-meat eaters, particularly vegans, had higher DSG2 in the current study. DSG2 belongs to the family of desmosomal cadherins, which includes both desmogleins (DSGs) and desmocollins (DSCs), with DSG2 and DSC2 being the major isoforms in humans, and the only isoforms expressed in the intestinal epithelium [12]. Consistently, the current study also showed that DSC2 was higher in non-meat eaters particularly vegans. Though the two proteins are structurally and functionally similar, laboratory evidence in mouse models suggest that DSG2 may have a more prominent role in the maintenance of the integrity of the intestinal epithelial barrier than DSC2 [38]. In humans, several small studies on IBD patients have found that lower intestinal protein levels of DSG2 in IBD patients, suggesting a role of the protein in IBD pathogenesis [12].

HAVCR1

HAVCR1 has been recognised as a biomarker of kidney injury and possibly an early diagnostic marker of kidney disease [11]. In mice, high-fat diets or the saturated fatty acid palmitate have been shown to upregulate expression of HAVCR1 in the proximal tubules of the kidney [39–41]. In humans, vegetarians and vegans tend to have lower intake of saturated fat, which is consistent with the observation of lower HAVCR1 in the non-meat eaters (though not significantly so in vegans) in the current study. Additionally, high protein diets, especially diets high in protein from animal sources, have been suggested to have adverse effects on kidney health [42]. A previous study suggested a lower risk of chronic kidney

disease in vegetarians after accounting for baseline risk of diabetes and hypertension [43].

Furthermore, as HAVCR1 is expressed on the surface of immune cells, it is also believed to

have a role regulating immune responses via activation and proliferation of immune cells

[11].

PCSK9

The observation of higher PCSK9 in vegetarians is of interest. PCSK9 is a well-established

drug target for treating hypercholesterolaemia, whereby PCSK9 inhibitors lowers low-

density-lipoprotein cholesterol (LDL-C) concentrations by preventing the binding of PCSK9

to LDL receptors, which subsequently enables the LDL receptors to bind and remove LDL-C

from the bloodstream instead [44, 45]. As a result, higher PCSK9 is often accompanied by

high LDL-C, whereas the vegetarians in the current [26] and other studies [46] typically have

significantly lower LDL-C, likely partly due to vegetarians having lower saturated fat intake

[47]. It may therefore be plausible that the relatively higher PCSK9 expression in people with

low LDL-C may be related to maintaining the minimum level of LDL-C necessary to support

cellular function and structural integrity, but further studies are needed to confirm this.

Other proteins

The functions of many other proteins identified are not well established. CNDP1 is a

dipeptidase, and one previous study reported higher CNDP1 in people on high protein diets

[48], which would be consistent with our observation of lower CNDP1 in non-meat eaters

(not statistically significant in vegans) who generally have lower protein intake. Several of

the proteins identified such as ART3, GAS6, THY1 and EGFR are involved in a range of

signalling pathways and progression of tumour tissues [49–52], while other proteins

including LAYN and CDHR2 are believed to have roles in cell adhesion [53, 54]. AHSP, which was significantly higher in vegans than vegetarians, has a role in stabilising haemoglobin and its loss of function has been linked to inherited anaemia [55], but its relevance for nutritional anaemia is unclear. Overall, the reasons that these proteins appear to be influenced by diet and their implications require further research. As the first study on diet group and a large panel of circulating proteins, the current study focused on the biggest differences by diet groups, and many other differences were observed that could not be described in detail, but still deserve further investigation. The study demonstrates the utility of investigating protein differences by diet groups; future research should measure these and other proteins using multiple technologies in relation to dietary factors, to replicate and expand on our findings.

Strengths and limitations

The key strength of this study was that it showed for first time differences in circulating proteins between people of different habitual diet groups in a large population, and many of the findings were completely novel. We have defined six diet groups in the white British population and two diet groups in the British Indian population, which allowed a detailed comparison by both dietary habits and ethnicity. Of the limitations, as with all observational studies, some level of self-selection bias may be present, which limits the generalisability of the findings. While we have adjusted for several important confounders, residual confounding by other dietary and non-dietary factors may still be present, though sensitivity analyses showed that results were consistent across all adjustment models. We are also unable to infer causality due to the cross-sectional nature of the studies.

398 *Conclusions*

399 In this large population-based cohort in the United Kingdom, many differences in circulating
400 protein concentrations were observed between different diet groups. These proteins are
401 involved in a range of different biological pathways and processes, particularly related to
402 gastrointestinal tract and kidney function. These differences likely reflect physiological
403 differences between the different diet groups, and the implications of these differences for
404 future disease risk require further investigation.

405 **References**

- 406 1. Cui M, Cheng C, Zhang L (2022) High-throughput proteomics: a methodological
407 mini-review. *Lab Invest* 102:1170–1181. <https://doi.org/10.1038/s41374-022-00830-7>
- 408 2. Al-Amrani S, Al-Jabri Z, Al-Zaabi A, et al (2021) Proteomics: Concepts and
409 applications in human medicine. *World J Biol Chem* 12:57–69.
410 <https://doi.org/10.4331/wjbc.v12.i5.57>
- 411 3. Schmidt JA, Rinaldi S, Scalbert A, et al (2016) Plasma concentrations and intakes of
412 amino acids in male meat-eaters, fish-eaters, vegetarians and vegans: a cross-sectional
413 analysis in the EPIC-Oxford cohort. *Eur J Clin Nutr* 70:306–312.
414 <https://doi.org/10.1038/ejcn.2015.144>
- 415 4. Pietzner M, Wheeler E, Carrasco-Zanini J, et al (2021) Mapping the proteo-genomic
416 convergence of human diseases. *Science* (80-) 374:.
417 <https://doi.org/10.1126/science.abj1541>
- 418 5. Fry A, Littlejohns TJ, Sudlow C, et al (2017) Comparison of sociodemographic and
419 health-related characteristics of UK Biobank participants with those of the general
420 population. *Am J Epidemiol* 186:1026–1034. <https://doi.org/10.1093/aje/kwx246>

- 421 6. Sun B, Chiou J, Traylor M, et al (2022) Genetic regulation of the human plasma
422 proteome in 54,306 UK Biobank participants. bioRxiv.
423 <https://doi.org/10.1101/2022.06.17.496443>
- 424 7. Albanes D, Alcalá K, Alcalá N, et al (2023) The blood proteome of imminent lung
425 cancer diagnosis. Nat Commun 14:3042. <https://doi.org/10.1038/s41467-023-37979-8>
- 426 8. Gao X, Starmer J, Martin ER (2008) A multiple testing correction method for genetic
427 association studies using correlated single nucleotide polymorphisms. Genet
428 Epidemiol 32:361–369. <https://doi.org/10.1002/gepi.20310>
- 429 9. Arnold M (2022) Jasper: Jasper makes plots. 2020
- 430 10. Thul PJ, Lindskog C (2018) The human protein atlas: A spatial map of the human
431 proteome. Protein Sci 27:233–244. <https://doi.org/10.1002/pro.3307>
- 432 11. Liu S, Tang W, Cao J, et al (2022) A Comprehensive Analysis of HAVCR1 as a
433 Prognostic and Diagnostic Marker for Pan-Cancer. Front Genet 13:1–11.
434 <https://doi.org/10.3389/fgene.2022.904114>
- 435 12. Schlegel N, Boerner K, Waschke J (2021) Targeting desmosomal adhesion and
436 signalling for intestinal barrier stabilization in inflammatory bowel diseases—Lessons
437 from experimental models and patients. Acta Physiol 231:1–15.
438 <https://doi.org/10.1111/apha.13492>
- 439 13. Allen NE, Appleby PN, Davey GK, et al (2002) The associations of diet with serum
440 insulin-like growth factor I and its main binding proteins in 292 women meat-eaters,
441 vegetarians, and vegans. Cancer Epidemiol Biomarkers Prev 11:1441–1448
- 442 14. Crowe FL, Key TJ, Allen NE, et al (2009) The association between diet and serum
443 concentrations of IGF-I, IGFBP-1, IGFBP-2, and IGFBP-3 in the European

- 444 Prospective Investigation into Cancer and Nutrition. *Cancer Epidemiol Biomarkers*
445 *Prev* 18:1333–1340. <https://doi.org/10.1158/1055-9965.EPI-08-0781>
- 446 15. McCarty MF (2017) The moderate essential amino acid restriction entailed by low-
447 protein vegan diets may promote vascular health by stimulating FGF21 secretion.
448 *Horm Mol Biol Clin Investig* 30:1–9. <https://doi.org/10.1515/hmbci-2015-0056>
- 449 16. Antony AC, Tang YS, Khan RA, et al (2004) Translational upregulation of folate
450 receptors is mediated by homocysteine via RNA-heterogeneous nuclear
451 ribonucleoprotein E1 interactions. *J Clin Invest* 113:285–301.
452 <https://doi.org/10.1172/JCI200411548>
- 453 17. Kim H, Lichtenstein AH, Ganz P, et al (2023) Identification of Protein Biomarkers of
454 the Dietary Approaches to Stop Hypertension Diet in Randomized Feeding Studies and
455 Validation in an Observational Study. *J Am Heart Assoc* 12:e028821.
456 <https://doi.org/10.1161/JAHA.122.028821>
- 457 18. Du S, Chen J, Kim H, et al (2023) Plasma Protein Biomarkers of Healthy Dietary
458 Patterns: Results from the Atherosclerosis Risk in Communities Study and the
459 Framingham Heart Study. *J Nutr* 153:34–46.
460 <https://doi.org/10.1016/j.tjnut.2022.11.008>
- 461 19. Walker ME, Song RJ, Xu X, et al (2020) Proteomic and metabolomic correlates of
462 healthy dietary patterns: The framingham heart study. *Nutrients* 12:1–20.
463 <https://doi.org/10.3390/nu12051476>
- 464 20. Warensjö Lemming E, Byberg L, Stattin K, et al (2019) Dietary Pattern Specific
465 Protein Biomarkers for Cardiovascular Disease: A Cross-Sectional Study in 2
466 Independent Cohorts. *J Am Heart Assoc* 8:. <https://doi.org/10.1161/JAHA.118.011860>

- 467 21. Castaño-Martínez T, Schumacher F, Schumacher S, et al (2019) Methionine restriction
468 prevents onset of type 2 diabetes in NZO mice. *FASEB J* 33:7092–7102.
469 <https://doi.org/10.1096/fj.201900150R>
- 470 22. Geng L, Lam KSL, Xu A (2020) The therapeutic potential of FGF21 in metabolic
471 diseases: from bench to clinic. *Nat Rev Endocrinol* 16:654–667.
472 <https://doi.org/10.1038/s41574-020-0386-0>
- 473 23. Bhatt DL, Bays HE, Miller M, et al (2023) The FGF21 analog pegozafermin in severe
474 hypertriglyceridemia: a randomized phase 2 trial. *Nat Med* 29:.
475 <https://doi.org/10.1038/s41591-023-02427-z>
- 476 24. Loomba R, Sanyal AJ, Kowdley K V., et al (2023) Randomized, Controlled Trial of
477 the FGF21 Analogue Pegozafermin in NASH. *N Engl J Med* 1–11.
478 <https://doi.org/10.1056/nejmoa2304286>
- 479 25. Tong TY, Key TJ, Sobiecki JG, Bradbury KE (2018) Anthropometric and physiologic
480 characteristics in white and British Indian vegetarians and nonvegetarians in the UK
481 Biobank. *Am J Clin Nutr* 107:909–920. <https://doi.org/10.1093/ajcn/nqy042>
- 482 26. Tong TYN, Perez-Cornago A, Bradbury KE, Key TJ (2021) Biomarker Concentrations
483 in White and British Indian Vegetarians and Nonvegetarians in the UK Biobank. *J*
484 *Nutr* 151:3168–3179. <https://doi.org/10.1093/jn/nxab192>
- 485 27. Tong TYN, Appleby PN, Bradbury KE, et al (2019) Risks of ischaemic heart disease
486 and stroke in meat eaters, fish eaters, and vegetarians over 18 years of follow-up:
487 results from the prospective EPIC-Oxford study. *BMJ* 366:l4897.
488 <https://doi.org/10.1136/bmj.l4897>
- 489 28. Tong TYN, Appleby PN, Armstrong MEG, et al (2020) Vegetarian and vegan diets

- 490 and risks of total and site-specific fractures: results from the prospective EPIC-Oxford
491 study. BMC Med 18:353. <https://doi.org/10.1186/s12916-020-01815-3>
- 492 29. Chiu T, Lin M-N, Pan W-H, et al (2018) Vegetarian diet, food substitution, and
493 nonalcoholic fatty liver. Tzu Chi Med J 30:102.
494 https://doi.org/10.4103/tcmj.tcmj_109_17
- 495 30. Dye FS, Larraufie P, Kay R, et al (2019) Characterisation of proguanylin expressing
496 cells in the intestine – evidence for constitutive luminal secretion. Sci Rep 9:1–20.
497 <https://doi.org/10.1038/s41598-019-52049-0>
- 498 31. von Volkmann HL, Brønstad I, Tronstad RR, et al (2020) Plasma levels of guanylin
499 are reduced in patients with Crohn’s disease. Scand J Gastroenterol 55:449–453.
500 <https://doi.org/10.1080/00365521.2020.1748224>
- 501 32. Wilson C, Lin JE, Li P, et al (2014) The paracrine hormone for the GUCY2C tumor
502 suppressor, guanylin, is universally lost in colorectal cancer. Cancer Epidemiol
503 Biomarkers Prev 23:2328–2337. <https://doi.org/10.1158/1055-9965.EPI-14-0440>
- 504 33. Sanjoaquin MA, Appleby PN, Spencer EA, Key TJ (2004) Nutrition and lifestyle in
505 relation to bowel movement frequency: a cross-sectional study of 20 630 men and
506 women in EPIC–Oxford. Public Health Nutr 7:77–83.
507 <https://doi.org/10.1079/phn2003522>
- 508 34. Gilsing AMJ, Crowe FL, Lloyd-Wright Z, et al (2010) Serum concentrations of
509 vitamin B12 and folate in British male omnivores, vegetarians and vegans: results
510 from a cross-sectional analysis of the EPIC-Oxford cohort study. Eur J Clin Nutr
511 64:933–939. <https://doi.org/10.1038/ejcn.2010.142>
- 512 35. Hung CJ, Huang PC, Lu SC, et al (2002) Plasma homocysteine levels in Taiwanese

- 513 vegetarians are higher than those of omnivores. J Nutr 132:152–158.
- 514 <https://doi.org/10.1093/jn/132.2.152>
- 515 36. Kaaks R, Bellati C, Venturelli E, et al (2003) Effects of dietary intervention on IGF-I
- 516 and IGF-binding proteins, and related alterations in sex steroid metabolism: The Diet
- 517 and Androgens (DIANA) Randomised Trial. Eur J Clin Nutr 57:1079–1088.
- 518 <https://doi.org/10.1038/sj.ejcn.1601647>
- 519 37. Sobiecki JG, Appleby PN, Bradbury KE, Key TJ (2016) High compliance with dietary
- 520 recommendations in a cohort of meat eaters, fish eaters, vegetarians, and vegans:
- 521 Results from the European Prospective Investigation into Cancer and Nutrition-Oxford
- 522 study. Nutr Res 36:464–477. <https://doi.org/10.1016/j.nutres.2015.12.016>
- 523 38. Gross A, Pack LAP, Schacht GM, et al (2018) Desmoglein 2, but not desmocollin 2,
- 524 protects intestinal epithelia from injury. Mucosal Immunol 11:1630–1639.
- 525 <https://doi.org/10.1038/s41385-018-0062-z>
- 526 39. Yu Y, Mo H, Zhuo H, et al (2022) High Fat Diet Induces Kidney Injury via
- 527 Stimulating Wnt/ β -Catenin Signaling. Front Med 9:1–9.
- 528 <https://doi.org/10.3389/fmed.2022.851618>
- 529 40. Zhao X, Chen X, Zhang Y, et al (2019) Kidney injury molecule-1 is upregulated in
- 530 renal lipotoxicity and mediates palmitate-induced tubular cell injury and inflammatory
- 531 response. Int J Mol Sci 20:. <https://doi.org/10.3390/ijms20143406>
- 532 41. Mori Y, Ajay AK, Chang JH, et al (2021) KIM-1 mediates fatty acid uptake by renal
- 533 tubular cells to promote progressive diabetic kidney disease. Cell Metab 33:1042-
- 534 1061.e7. <https://doi.org/10.1016/j.cmet.2021.04.004>
- 535 42. Ko GJ, Rhee CM, Kalantar-Zadeh K, Joshi S (2020) The effects of high-protein diets

536 on kidney health and longevity. J Am Soc Nephrol 31:1667–1679.
537 <https://doi.org/10.1681/ASN.2020010028>

538 43. Liu H-W, Tsai W-H, Liu J-S, Kuo K-L (2019) Association of Vegetarian Diet with
539 Chronic Kidney Disease. Nutrients 11:279. <https://doi.org/10.3390/nu11020279>

540 44. Schmidt AF, Holmes M V, Preiss D, et al (2019) Phenome-wide association analysis
541 of LDL- cholesterol lowering genetic variants in. 1–10

542 45. Horton JD, Cohen JC, Hobbs HH (2007) Molecular biology of PCSK9: its role in LDL
543 metabolism. Trends Biochem Sci 32:71–77. <https://doi.org/10.1016/j.tibs.2006.12.008>

544 46. Bradbury KE, Crowe FL, Appleby PN, et al (2014) Serum concentrations of
545 cholesterol, apolipoprotein A-I and apolipoprotein B in a total of 1694 meat-eaters,
546 fish-eaters, vegetarians and vegans. Eur J Clin Nutr 68:178–183.
547 <https://doi.org/10.1038/ejcn.2013.248>

548 47. Mensink RP, Zock PL, Kester AD, Katan MB (2003) Effects of dietary fatty acids and
549 carbohydrates on the ratio of serum total to HDL cholesterol and on serum lipids and
550 apolipoproteins: a meta- - analysis of 60 controlled trials. Am J Clin Nutr 77:1146–
551 1155. [https://doi.org/11778166](https://doi.org/10.1177/8166)

552 48. Franzke B, Bileck A, Unterberger S, et al (2022) The plasma proteome is favorably
553 modified by a high protein diet but not by additional resistance training in older adults:
554 A 17-week randomized controlled trial. Front Nutr 9:.
555 <https://doi.org/10.3389/fnut.2022.925450>

556 49. Tan L, Song X, Sun X, et al (2016) ART3 regulates triple-negative breast cancer cell
557 function via activation of Akt and ERK pathways. Oncotarget 7:46589–46602.
558 <https://doi.org/10.18632/oncotarget.10306>

559 50. Tanaka M, Siemann DW (2020) Gas6/Axl signaling pathway in the tumor immune
560 microenvironment. *Cancers (Basel)* 12:1–14. <https://doi.org/10.3390/cancers12071850>

561 51. Hagood JS (2019) Thy-1 as an Integrator of Diverse Extracellular Signals. *Front Cell*
562 *Dev Biol* 7:1–5. <https://doi.org/10.3389/fcell.2019.00026>

563 52. Wee P, Wang Z (2017) Epidermal growth factor receptor cell proliferation signaling
564 pathways. *Cancers (Basel)* 9:1–45. <https://doi.org/10.3390/cancers9050052>

565 53. Pan J hua, Zhou H, Cooper L, et al (2019) LAYN is a prognostic biomarker and
566 correlated with immune infiltrates in gastric and colon cancers. *Front Immunol* 10:1–
567 12. <https://doi.org/10.3389/fimmu.2019.00006>

568 54. Pinette JA, Mao S, Millis BA, et al (2019) Brush border protocadherin CDHR2
569 promotes the elongation and maximized packing of microvilli in vivo. *Mol Biol Cell*
570 30:108–118. <https://doi.org/10.1091/mbc.E18-09-0558>

571 55. Yu X, Kong Y, Dore LC, et al (2007) An erythroid chaperone that facilitates folding of
572 α -globin subunits for hemoglobin synthesis. *J Clin Invest* 117:1856–1865.
573 <https://doi.org/10.1172/JCI31664>

574

575 Funding: The work is supported by the UK Medical Research Council (MR/M012190/1),
576 Wellcome Trust Our Planet Our Health (Livestock, Environment and People, LEAP
577 205212/Z/16/Z) and Cancer Research UK (C8221/A29017 and C8221/A29186). TYNT is
578 support by a Nuffield Departmental of Population Health Fellowship. The funders had no role
579 in study design, data collection, analysis, decision to publish, or preparation of the
580 manuscript.

581 Acknowledgements: This research has been conducted using UK Biobank Resource under
582 application 67506.

583 Competing interests: The authors had no conflicts of interest.

584 Author's contributions: TYNT, TJK and RCT conceived and designed the research question.
585 KSB conducted quality control of the proteomics data. TYNT analyzed the data and wrote
586 the first draft of the manuscript. All authors provided input on data analysis and interpretation
587 of results, revised the manuscript critically for important intellectual content, and read and
588 approved the final manuscript.

Table 1: Baseline characteristics of white British and British Indian participants by diet groups in UK Biobank.

	White British participants						British Indian participants	
	Regular meat eaters N=23,116	Low meat eaters N=23,323	Poultry eaters N=484	Fish eaters N=1,074	Vegetarians N=722	Vegans N=54	Meat eaters N=390	Vegetarians N=163
Age at recruitment	57.2 (8.2)	57.2 (8.0)	56.6 (8.1)	54.4 (8.2)	52.9 (8.1)	55.0 (7.8)	53.9 (8.6)	54.7 (7.7)
Sex								
Women	9,810 (42.4%)	14,861 (63.7%)	389 (80.4%)	756 (70.4%)	483 (66.9%)	28 (51.9%)	170 (43.6%)	102 (62.6%)
Men	13,306 (57.6%)	8,462 (36.3%)	95 (19.6%)	318 (29.6%)	239 (33.1%)	26 (48.1%)	220 (56.4%)	61 (37.4%)
Region of assessment centre								
London	2,558 (11.1%)	2,797 (12.0%)	80 (16.5%)	191 (17.8%)	121 (16.8%)	8 (14.8%)	208 (53.3%)	88 (54.0%)
North-West England	3,957 (17.1%)	3,577 (15.3%)	77 (15.9%)	136 (12.7%)	108 (15.0%)	13 (24.1%)	24 (6.2%)	8 (4.9%)
North-East England	2,830 (12.2%)	3,011 (12.9%)	58 (12.0%)	93 (8.7%)	67 (9.3%)	3 (5.6%)	12 (3.1%)	4 (2.5%)
Yorkshire	3,792 (16.4%)	3,764 (16.1%)	68 (14.0%)	193 (18.0%)	111 (15.4%)	10 (18.5%)	35 (9.0%)	10 (6.1%)
West Midlands	1,957 (8.5%)	1,929 (8.3%)	38 (7.9%)	83 (7.7%)	60 (8.3%)	6 (11.1%)	59 (15.1%)	22 (13.5%)
East Midlands	1,863 (8.1%)	1,889 (8.1%)	38 (7.9%)	84 (7.8%)	60 (8.3%)	4 (7.4%)	15 (3.8%)	15 (9.2%)
South-East England	1,764 (7.6%)	1,790 (7.7%)	28 (5.8%)	69 (6.4%)	57 (7.9%)	5 (9.3%)	17 (4.4%)	9 (5.5%)
South-West England	1,698 (7.3%)	1,871 (8.0%)	38 (7.9%)	103 (9.6%)	60 (8.3%)	4 (7.4%)	10 (2.6%)	3 (1.8%)
Wales	882 (3.8%)	1,084 (4.6%)	23 (4.8%)	53 (4.9%)	35 (4.8%)	0 (0.0%)	5 (1.3%)	3 (1.8%)
Scotland	1,815 (7.9%)	1,611 (6.9%)	36 (7.4%)	69 (6.4%)	43 (6.0%)	1 (1.9%)	5 (1.3%)	1 (0.6%)
Fasting time (hours)	3.8 (2.5)	3.7 (2.3)	3.8 (2.2)	3.6 (2.3)	3.7 (2.4)	3.8 (3.0)	4.1 (2.4)	4.1 (2.5)
Body mass index (kg/m²)								
<20	358 (1.5%)	548 (2.3%)	35 (7.2%)	65 (6.1%)	45 (6.2%)	5 (9.3%)	6 (1.5%)	3 (1.8%)
20.0–22.4	5,911 (25.6%)	7,659 (32.8%)	219 (45.2%)	503 (46.8%)	338 (46.8%)	29 (53.7%)	121 (31.0%)	52 (31.9%)
22.5–24.9	10,239	9,936	159 (32.9%)	379 (35.3%)	251 (34.8%)	13 (24.1%)	180 (46.2%)	71 (43.6%)

	(44.3%)	(42.6%)						
≥25	6,513 (28.2%)	5,105 (21.9%)	68 (14.0%)	123 (11.5%)	87 (12.0%)	7 (13.0%)	80 (20.5%)	35 (21.5%)
Unknown	95 (0.4%)	75 (0.3%)	3 (0.6%)	4 (0.4%)	1 (0.1%)	0 (0.0%)	3 (0.8%)	2 (1.2%)
Alcohol consumption								
<1g/d	3,600 (15.6%)	4,540 (19.5%)	151 (31.2%)	255 (23.7%)	169 (23.4%)	27 (50.0%)	156 (40.0%)	139 (85.3%)
1-7g/d	5,075 (22.0%)	6,721 (28.8%)	152 (31.4%)	293 (27.3%)	200 (27.7%)	9 (16.7%)	96 (24.6%)	18 (11.0%)
8-15g/d	4,986 (21.6%)	5,521 (23.7%)	76 (15.7%)	235 (21.9%)	166 (23.0%)	7 (13.0%)	52 (13.3%)	6 (3.7%)
≥16g/d	9,439 (40.8%)	6,525 (28.0%)	105 (21.7%)	291 (27.1%)	186 (25.8%)	11 (20.4%)	84 (21.5%)	0 (0.0%)
Unknown	16 (0.1%)	16 (0.1%)	0 (0.0%)	0 (0.0%)	1 (0.1%)	0 (0.0%)	2 (0.5%)	0 (0.0%)
Smoking status								
Never	11,729 (50.7%)	12,933 (55.5%)	268 (55.4%)	597 (55.6%)	422 (58.4%)	29 (53.7%)	291 (74.6%)	148 (90.8%)
Previous	8,557 (37.0%)	8,178 (35.1%)	179 (37.0%)	389 (36.2%)	237 (32.8%)	20 (37.0%)	42 (10.8%)	10 (6.1%)
Current < 15 cigs/d	707 (3.1%)	690 (3.0%)	14 (2.9%)	29 (2.7%)	21 (2.9%)	1 (1.9%)	21 (5.4%)	2 (1.2%)
Current 15 or more cigs/d	2,038 (8.8%)	1,438 (6.2%)	21 (4.3%)	55 (5.1%)	37 (5.1%)	3 (5.6%)	31 (7.9%)	3 (1.8%)
Unknown	85 (0.4%)	84 (0.4%)	2 (0.4%)	4 (0.4%)	5 (0.7%)	1 (1.9%)	5 (1.3%)	0 (0.0%)
Physical activity (MET hrs/wk)								
<10	4,735 (20.5%)	4,352 (18.7%)	78 (16.1%)	163 (15.2%)	128 (17.7%)	8 (14.8%)	95 (24.4%)	45 (27.6%)
10-49.9	9,066 (39.2%)	9,883 (42.4%)	183 (37.8%)	482 (44.9%)	324 (44.9%)	29 (53.7%)	127 (32.6%)	51 (31.3%)
≥50	4,089 (17.7%)	3,963 (17.0%)	124 (25.6%)	214 (19.9%)	137 (19.0%)	10 (18.5%)	53 (13.6%)	10 (6.1%)
Unknown	5,226 (22.6%)	5,125 (22.0%)	99 (20.5%)	215 (20.0%)	133 (18.4%)	7 (13.0%)	115 (29.5%)	57 (35.0%)

Numbers shown are mean (SD) or number (%).




Figure 1: Volcano plots of proteins in white British vegetarians (top) and vegans (bottom) compared with regular meat eaters.

The red dotted line signifies p -value threshold for statistical significance. Each dot represents one protein, which were colour-coded by whether the protein is majority expressed (>50%) in one tissue type. Results were based on the multivariable model adjusted for age at recruitment, sex, region, fasting status, body mass index, alcohol consumption, smoking status and physical activity.




Figure 2: Top 10 proteins in vegetarians by diet group and ethnicity.

The top 10 proteins were selected by ranking the *p*-values of proteins comparing white British vegetarians with regular meat eaters, and sorted by betas in white British vegetarians, where the betas represent SD differences. Results were based on the multivariable model adjusted for age at recruitment, sex, region, fasting status, body mass index, alcohol consumption, smoking status and physical activity.




Figure 3: Top 10 proteins in vegans by diet group and ethnicity.

The top 10 proteins were selected by ranking the *p*-values of proteins comparing white British vegans with regular meat eaters, and sorted by betas in white British vegans, where the betas represent SD differences. Results were based on the multivariable model adjusted for age at recruitment, sex, region, fasting status, body mass index, alcohol consumption, smoking status and physical activity.