

## **Overweight and its Associated Risk Factors among Students in Tertiary Institutions of Learning in Mongu District of Western Province, Zambia; A cross-sectional Survey.**

Larry L. Mooka<sup>1,2\*</sup>, Priscilla Funduluka<sup>1</sup> and Mukumbuta Nawa<sup>3</sup>

<sup>1</sup>Department of Disease Prevention and Health Promotion, School of Public Health and Environmental Studies. Levy Mwanawasa Medical University, P. O. Box 33991, Lusaka, Zambia.

<sup>2</sup>Lewanika College of Nursing, Midwifery and Public Health, P. O. Box 910147, Mongu, Zambia

<sup>3</sup>Department of Biostatistics and Epidemiology, School of Public Health and Environmental Studies. Levy Mwanawasa Medical University, P. O. Box 33991, Lusaka, Zambia.

\*Corresponding author: Larry L. Mooka, Lewanika College of Nursing, Midwifery and Public Health, P. O. Box 910147, Mongu, Zambia. Email: [larrymooka@gmail.com](mailto:larrymooka@gmail.com), phone: +260-978-177576

## ABSTRACT

**Introduction:** Overweight and obesity are key public health issues in many high-income countries and have become a major public health concern affecting all ages, including adolescents in low-income and middle-income countries. This study determined the prevalence of overweight and obesity and their associated factors among undergraduate students at colleges and universities in the Mongu district of the Western Province of Zambia.

**Methods:** This was a cross-sectional study. Undergraduate students aged 18 to 26 years were sampled from two colleges and one university in Mongu. Data was collected electronically by trained research assistants using a structured questionnaire using Kobo Collect software. Data was analysed using Stata version 14 software. The analysis included descriptive statistics, using counts, frequencies, percentages, means and medians depending on the distribution of the data. Chi-squared tests of association, bivariate, and multivariate logistic regression were done to test for association between overweight and independent variables. A p-value less than 0.05 was considered significant.

**Results:** A total of 330 students were captured in the survey; over half of the respondents were female students 214 (64.85%), while 116 (35.15%) were male respondents. Body Mass Index (BMI) showed that 21 (6.8%) were underweight, 208 (63.0%) had normal weight, 76 (23.0%) were overweight, and 25 (7.6%) were obese. Factors associated with overweight and obesity included female sex aOR 1.68 (95%CI 1.02 – 2.76), age and alcohol intake.

**Conclusions:** A third of the students were either overweight or obese. Sex, age and alcohol intake were significantly associated with overweight and obesity among students. The prevalence of overweight and obesity is high among young adults, and this calls for early interventions in learning institutions to combat obesity.

**Keywords:** Overweight, obesity, body mass index, prevalence, diet, sex.

## BACKGROUND

Obesity is a key public health issue in many high-income countries and is a steadily rising public health problem in low-income and middle-income countries (Almutairi *et al.*, 2021). Obesity and overweight have grown to epidemic proportions, with over 4 million people dying each year due to being overweight or obese in 2017, according to the global burden of disease (WHO, 2022).

Overweight and obesity are defined as abnormal or excessive fat accumulation that may impair health (WHO, 2022). Body Mass Index (BMI) is a simple weight-for-height index commonly used to classify overweight and obesity in adults. It is defined as a person's weight in kilograms divided by the square of his height in meters ( $\text{kg/m}^2$ ) (WHO, 2022). For adults, a BMI of under 18.5 is considered underweight, a BMI of 18.5 – 24.9 is considered normal weight, a BMI over 25 is considered overweight, and over 30 is obese (WHO, 2022).

The prevalence of overweight and obesity among adults has doubled since 1980 worldwide, and it has increased in many low and middle-income countries (LMICs). Cardiovascular diseases (CVDs), diabetes, musculoskeletal disorders, and some types of cancers associated with overweight and obesity are the leading causes of mortality worldwide (Tateyama *et al.*, 2018). Current projections show that the burden of chronic diseases will become comparable to acute infectious diseases in many LMICs (Tateyama *et al.*, 2018; Nawa *et al.*, 2020).

Sub-Saharan Africa, including Zambia, has experienced an increase in overweight and obesity due to rapid lifestyle changes associated with recent economic growth (Tateyama *et al.*, 2018). Obesity-related diseases currently represent 3.8% of disability-adjusted life years (Tateyama *et al.*, 2019). The estimated national prevalence of overweight and obesity in Zambia was 29.2% in 2014, compared with 26.4% in 2010 (Bwalya *et al.*, 2017; Nawa, 2020).

The increase in overweight and obesity can pose a huge burden on healthcare systems with potential effects on the social and economic well-being of an individual and community. Poor health conditions resulting from overweight and obesity may likewise act as a barrier to developing a good mind, thus paving the way for poor learning behaviour and outcomes among students (Kafyulilo, 2008;).

The study sought to establish the prevalence and to determine the risk factors associated with overweight and obesity among college students in the Mongu District of Western Province, Zambia. While there are few studies on overweight and obesity in Zambia, previous studies have sampled the general population, whilst this study sampled young adults. It will fill the knowledge gap and inform policymakers for this demographic.

## **METHODOLOGY**

### **Study Setting**

Mongu is the capital of the Western Province of Zambia and is geographically located 580 kilometres west of Lusaka, the capital city of Zambia. Whilst Western Province is mostly rural, Mongu is the most urbanised and populous area. The people are involved in trade, civil service and agricultural practices. The study was conducted at three (3) tertiary learning institutions: Mongu College of Education, Lewanika College of Nursing and Midwifery and University of Barotseland.

## **STUDY DESIGN**

This study was a cross-sectional survey conducted between April 2023 and June 2023.

### **Study Respondents**

The study targeted young adults pursuing tertiary education; this demographic represents young educated people in the province with at least some tertiary education. Therefore, this study will indicate the prevalence and risk factors in the young educated class.

### **Inclusion Criteria**

All full-time students enrolled at the three selected colleges and universities in the Mongu district were eligible for the study.

### **Exclusion Criteria**

Pregnant female students, including bodybuilders or weight lifters, were excluded from the study. Students on any form of medication, acutely ill or with known chronic diseases, were also excluded from the study.

## SAMPLE SIZE AND POWER CONSIDERATION

Since the sample for this study was drawn from a finite population, the sample size was determined using the 30% prevalence based on a study done by (Bwalya *et al.*, 2017). A precision of 5% (expressed as 0.05) for a 95% confidence level, with a Z-alpha value of 1.96, was used.

$$n = \frac{Z^2 * P * (1-P)}{E^2}$$

Where:

n = Desired sample

Z = Z-score corresponding to 95% CI

P = Expected prevalence of obesity obtained from previous studies

E = Desired precision, which is 5% (expressed as 0.05).

$$n = \frac{1.96^2 * 0.3 * (1-0.3)}{(0.05)^2} = \underline{324}$$

For non-response and refusals, non-response rate was calculated and compensated using the formulae:

$$\text{Final sample size} = \frac{\text{Effective sample size}}{1 - \text{non response rate anticipated}} \text{ (Bujang, 2021).}$$

## Sampling Technique

Two-stage stratified random sampling technique was applied in this study because the sample obtained was not homogenous (WHO, 2005). The first-tier sample comprised colleges/universities selected randomly from all the learning institutions in the Mongu district. From the three selected learning institutions, all students who met the eligibility criteria constituted the sampling frame of the study. The second tier was a simple random sampling of the students from the selected learning institutions.

## DATA COLLECTION

Five (5) research assistants were trained to take anthropometric measurements and administer the questionnaire using Kobo Toolbox Collect installed on smartphones (<https://ee.kobotoolbox.org/x/hOHLVeP6>).

## **DATA MANAGEMENT AND ANALYSIS**

Data was exported from Kobo Toolbox Collect into Stata version 14.0. Data analysis included computing BMI from the weights and heights ( $w/h^2$ ) and categorising BMI into underweight, normal weight, and overweight/obese. The prevalence of the outcome variables (overweight and obesity) was then estimated. Descriptive statistics were presented as frequency, percentages, median and interquartile range were used. Chi-square tests of association were used to analyse associations between categorical variables. Bivariate and multivariate logistic regression analysis was used to assess the associations between the student's demographics, alcohol intake, food consumption pattern, physical exercise, and eating fruits and vegetables with the outcome. For this study,  $P \leq 0.05$  was considered statistically significant.

## **Ethical Considerations**

The study protocol was submitted to the ERES Converge IRB Committee for ethical review and approval (No. 2023-April-007). Permission to conduct the study was sought in writing from each of the three (3) higher learning institutions from which the study was conducted. Written consent was obtained from respondents after explaining to them the purpose of the study and how the results were to be utilised. Data obtained from respondents was kept in a password-protected computer to guarantee respondents' confidentiality and anonymity. The respondents retained the absolute right and freedom to decline to participate or withdraw from the study at any time without consequence.

## **Dissemination Plan**

Recommendations have been made to the Ministry of Education regarding implementing overweight/obesity college interventions. Findings will be used to advocate for higher institutions-based curriculum and policy change at the national level. The findings will be published as a Master's degree dissertation in peer-reviewed publications. In addition, key findings will be shared with the Ministry of Education and participating colleges and universities through report sharing.

## RESULTS

### Socio-Demographic Profile

As shown in Table 1, over half of the respondents were females, 214 (64.85%), while 116 (35.15%) were male respondents. The median weight was 62kg, and the weight IQR=55-70. The majority of the respondents, 106 (32.12%), were within the 21-23-year age group, while the minority of the respondents, 62 (18.79%), were within the 18-20-year age group. A total of 127 (38.48%) students were recruited from Mongu Catholic College of Education (MOCE). In comparison, 126 (38.18%) students were recruited from Lewanika College of Nursing and Midwifery (LCNM) and 77 (23.33%) students from the University of Barotseland (UBL). A widely held percentage of the respondents, 151 (45.76%), were from a non-health-related field, followed by 126 (38.18%) from a nursing field and 53 (16.06%) from a health-related field. A greater proportion of the respondents, 177 (53.64%), were first-year students.

**Table 1: Socio-demographic profile and continuous variables of respondents**

<b>Variables</b>	<b>Frequency (N=330)</b>	<b>Percent (%)</b>
<b>Sex</b>		
male	116	35.15
female	214	64.85
<b>Age</b>		
18-20	62	18.79
21-23	106	32.12
24-26	70	21.21
>26	92	27.88
<b>Training Institution</b>		
LCNM	126	38.18
MOCE	127	38.48
UBL	77	23.33
<b>Field of Study</b>		
Nursing	126	38.18
Health-related	53	16.06
Non-health related	151	45.76
<b>Level of training</b>		
First-year	177	53.64
Second year	53	16.06
Third year	70	21.21
Fourth-year	30	9.09
<b>Residence</b>		
On-campus	142	43.03
Off-campus	188	56.97
<b>Continuous variables</b>		
<b>Variables</b>	<b>Median</b>	<b>IQR</b>
Weight	62	55-70
Height (m)	1.6	0.1
BMI (kg/m <sup>2</sup> )	23	5

## Overweight and Obesity Prevalence

The overall prevalence of overweight and obesity among respondents was 30.6% (23.0% overweight and 7.6% obese). Female respondents 70 (69.3%) were significantly more overweight and obese than male respondents 31 (30.7%). Concerning the field of study, a higher prevalence of overweight and obesity was found among non-health-related students, 47 (46.5%) compared to nursing students, 29 (28.7%) and health-related students, 25 (24.8%).

## Field of study

Table 2 reveals that students' field of study was significantly associated with overweight and obesity among respondents ( $P=0.003$ ). In addition, the results in Table 5 show that students from a non-health-related field recorded the highest proportion of overweight and obesity 47/101 (46.53%) compared to students from a nursing field 29/101 (28.71%), and a health-related field 25/101 (24.75%) ( $P=0.003$ ).

**Table 2: Association between the field of study and BMI**

BMI	nursing	Health-related	Non-health related	P-value*
<18.5kg/m <sup>2</sup>	14	1	6	0.003
18.5-24.9kg/m <sup>2</sup>	83	27	98	
25-29.9kg/m <sup>2</sup>	25	16	35	
≥30 kg/m <sup>2</sup>	4	9	12	330
Total	126	53	151	

\*Chi-squared Test

## Food Consumption Patterns

Table 3 shows the feeding patterns and food sources of overweight/obese respondents in the study. A high percentage of 33 (40.74%) of the respondents reported consuming snacks once every day. In comparison, 39 (38.61%) of the respondents reported consuming fruits and vegetables once every day and (36.36%) of the respondents consumed fast food once every day (Table 3). A total of 83 (35.62%) of respondents reported consuming junk food once per week, while 234 (71.12%) reported consuming nshima twice per day.



**Table 3: Food consumption pattern of respondents (n=101).**

Variables	Frequency	Percent (%)
<b>Fruits and vegetables</b>		
Once every day	39	38.61
2-3 times a day	33	32.67
Twice in a week	19	18.81
Once in a week	10	9.90
<b>Total</b>	101	100
<b>Snacks</b>		
Once every day	33	40.74
2-3 times a day	11	13.58
Twice in a week	17	20.99
Once in a week	20	24.69
<b>Total</b>	81	100
<b>Fast food</b>		
Once every day	28	36.36
2-3 times a day	9	11.69
Twice in a week	18	23.38
Once in a week	22	28.57
<b>Total</b>	77	100
<b>Junk food</b>		
Once every day	59	25.32
2-3 times a day	38	16.31
Twice in a week	53	22.75
Once in a week	83	35.62
<b>Total</b>	233	100
<b>Nshima</b>		
Once every day	65	19.76
Twice per day	234	71.12
Three times per day	30	9.12
<b>Total</b>	329	100

### Alcohol intake

Cross tabulation and chi-square test revealed a significant association between alcohol consumption and overweight or obesity ( $P=0.003$ ) (Table 4). More students did not take alcohol compared to those who took alcohol, i.e. 245 and 85, respectively.

**Table 4: Association between alcohol intake and BMI**

BMI	Alcohol intake		P-value*
	yes	no	
Underweight ( $<18.5\text{kg/m}^2$ )	3	18	0.003
Normal weight ( $18.5\text{-}24.9\text{kg/m}^2$ )	43	165	
Overweight ( $25\text{-}29.9\text{kg/m}^2$ )	31	45	
Obese ( $\geq 30\text{ kg/m}^2$ )	8	17	

Total	85	245	330
-------	----	-----	-----

\*Chi-squared Test

### Association between independent Variables and dependent variables

The independent variables significantly associated with overweight and obesity among the participants were age group older than 26 years among the students, which had reduced odds 0.37 (95%CI 0.18 – 0.79) compared to students aged between 18 to 20 years. Sex was also statistically significantly associated with being overweight and obese; Female students were less likely to be overweight and obese with an adjusted odds ratio of 0.59 (0.18 – 0.79) compared to their male counterparts after adjusting for age and behavioural factors such as exercises, food and alcohol intake patterns. The other variable associated with overweight and obesity was alcohol intake, which increased the odds by 1.51 aOR 2.51 (1.44 – 4.38). Table 5 shows the associations of selected factors and the outcome variable:

**Table 5: Association between independent and dependent variables**

Variable	Category	Unadjusted OR (95% CI)	Adjusted OR (95% CI)	Best Fit Model AOR (95% CI)
Age	18-20 Yrs.	Ref	Ref	
	21-23 Yrs.	1.15 (0.55 - 2.42)	1.24 (0.58 - 2.66)	1.21 (0.57 - 2.29)
	24 - 26 Yrs.	0.86 (0.39 - 1.88)	0.88 (0.39 - 2.0)	0.84 (0.38 - 1.89)
	Above 26 Yrs.	0.35 (0.17 - 0.71)	<b>0.37 (0.18 - 0.79)</b>	<b>0.36 (0.17 - 0.74)</b>
Sex	Male	Ref	Ref	Ref
	Female	0.75 (0.45 - 1.24)	<b>0.59 (0.18 - 0.79)</b>	0.59 (0.34 - 1.02)
Residence	On-campus	Ref	Ref	
	Off-campus	0.72 (0.45 - 1.17)	0.88 (0.51 - 1.53)	
Field of study	Non-Medical	Ref	Ref	
	Medical Related	0.96 (0.59 - 1.53)	0.84 (0.48 - 1.45)	
	Not at all	Ref	Ref	
Physical exercise	Seldom	1.10 (0.49 - 2.51)	1.03 (0.42 - 2.53)	
	Yes Regularly	1.31 (0.74 - 2.31)	1.06 (0.57 - 1.98)	
Alcohol	No Alcohol	Ref	Ref	Ref
	Takes Alcohol	2.50 (1.49 - 4.19)	<b>2.51 (1.44 - 4.38)</b>	<b>2.56 (1.48 - 4.39)</b>
Snacks	No Snacks	Ref	Ref	
	Takes Snacks	1.07 (.59 - 1.93)	1.32 (0.67-2.58)	
Fast food	No Fast Food	Ref	Ref	
	Takes Fast Foods	0.89 (0.51 - 1.58)	0.87 (0.47 1.61)	
Sugar- sweetened beverages	No Sugar	Ref	Ref	
	Takes Sugar	0.92 0.42 - 2.03)	1.15 (0.48 - 2.77)	

## Model Selection Criteria

Based on the Akaike and Bayesian information, model fitness was assessed for the lowest and highest levels of the coefficient of determination (Pseudo  $R^2$ ). The model with age, sex and alcohol had the lower AIC and BIC but reduced coefficient of determination compared to the full model as shown in table 6. We, therefore, adopted the full model as our best-fit model as it was better able to explain the variation in the outcome despite having higher values of AIC and BIC.

**Table 6: Model Selection Criteria**

Model	Pseudo R-Squared	AIC	BIC	P-Value
Full model (Age, sex, Residence, field of study, physical exercise, alcohol, fruits and vegetables, fast food, junk food, and sugar-sweetened beverages).	0.082	401.30	454.49	0.002
Reduced Model(Age, sex, and alcohol).	0.078	386.92	409.71	<0.001

\*The Akaike Information Criteria (AIC) and Bayesian (BIC) Information Criteria

## DISCUSSION

The current study was conducted to determine the prevalence of overweight and obesity among college students and the associated risk factors. The study has shown that over a fifth of the students were overweight, and at least one-third were overweight or obese. While this is not representative of the general population, it is concerning that a third of young adults in colleges are either overweight or obese. A study that used nationwide data on women in 2014 found a prevalence of 26.4%, which is comparable to this study's findings of 30.6%, although the former only included women (Bwalya *et al.*, 2017). Furthermore, these results correlate with findings from the study carried out on food consumption patterns, physical activity and overweight and obesity among undergraduates of a private university in Nigeria, which showed that 31% of respondents were overweight and 9.3% were obese (Kayode *et al.*, 2020).

This study revealed that female sex reduced the odds of being overweight or obese (AOR=0.59; 95% CI=0.18 - 0.79). This is consistent with studies done by Jiang *et al.* (2018) on overweight and obesity

among Chinese college students and a study done in Brazil among nursing students by Urbanetto *et al.* (2019), which indicated that males were more susceptible to being overweight and obese than females.

Our study found that older age groups above 26 years had reduced odds of obesity compared to the younger age group of 18 to 20 years old. This may be due to the fact that older age groups are more assertive and able to take care of their habits, such as alcohol and junk food. A study done among health sciences college students in Saudi Arabia found that age was not associated with obesity (Makkawy *et al.*, 2021). Our study found that regular alcohol intake was associated with overweight and obesity. This is consistent with findings from a systematic review conducted by Traversy *et al.* (2015), which showed a positive association between alcohol consumption and measures of abdominal adiposity (also known as “beer belly”).

This study did not find significant associations between overweight and obesity and other variables such as physical exercises, taking of snacks, fast foods and sugar-sweetened beverages. Our study may not have been powered to detect differences for all the variables; regular exercise has been shown to reduce weight (Tateyama, 2019). Other studies have also shown that taking snacks, fast foods and sugar-sweetened beverages is associated with overweight and obesity (Moise, 2019).

### **Study limitations**

Anthropometric measurements only included height, weight and body mass index (BMI). Since BMI is not the only accurate measure for adiposity, waist circumference and waist-to-hip ratios should be incorporated in future research. Secondly, this is a cross-sectional study, so it does not adduce causality but associations.

### **Conclusion**

This study has found a high prevalence of overweight and obesity among young adults attending college education. Sex, age, and alcohol intake were significantly associated with overweight and obesity among students.

## References

- Agarwal, K. *et al.* (2021) 'Relationship between BMI and alcohol consumption levels in decision making', *International Journal of Obesity*, 45(11), pp. 2455–2463.
- Almutairi, N.S. *et al.* (2021) 'Identifying factors associated with overweight and obesity among intermediate school students aged 12-15 years in school settings: Mixed-methodology protocol', *BMJ Open*, 11(5).
- Ang, Y.N. *et al.* (2012) 'Multifactorial Influences of Childhood Obesity Multifactorial Influences of Childhood Obesity', (March).
- Al Bashtawy, M. *et al.* (2015) 'Prevalence of Obesity and Physical Inactivity Behaviors among Nursing Students in Mutah University, Jordan', *World Journal of Medical Sciences*, 12(2), pp. 95–102.
- Batiha, A.-M. *et al.* (2018) 'The relationship between physical activity and diet, and overweight and obesity, in young people', *Nursing Children and Young People* [Preprint].
- Bray, G.A. *et al.* (2017) Obesity: a chronic relapsing progressive disease process. A position statement of the World Obesity Federation, *Obesity reviews*: an official journal of the International Association for the Study of Obesity, 18(7), pp. 715–723.
- Bujang, M.A. (2021) 'A step-by-step process on sample size determination for medical research', *Malaysian Journal of Medical Sciences*, 28(2), pp. 15–27.
- Bwalya, B.B. *et al.* (2017) 'Rural-urban differentials in the prevalence of overweight and obesity among women of childbearing age in Zambia', *Epidemiology Biostatistics and Public Health*, 14(2).
- Chaput, J.P. *et al.* (2010) 'Risk factors for adult overweight and obesity: The importance of looking beyond the "Big Two"', *Obesity Facts*, 3(5), pp. 320–327.
- Chaput, J.P. *et al.* (2012) 'Short sleep duration is associated with greater alcohol consumption in adults', *Appetite*, 59(3), pp. 650–655.
- Chen, Y. *et al.* (2020) 'Higher academic stress was associated with increased risk of overweight and obesity among college students in China', *International Journal of Environmental Research and Public Health*, 17(15), pp. 1–12.
- D'Anna, C. *et al.* (2019) 'Physical education status in European school's curriculum, extension of educational offer and planning', in: *Universidad de Alicante Servicio de Publicaciones*.
- Fazzino, T.L. *et al.* (2019) 'A systematic review of the methodology used to study weight change among young adults attending college', *Eating Behaviors*.

French, M.T. *et al.* (2010) 'Alcohol consumption and body weight', *Health Economics*, 19(7), pp. 814–832.

Gaviria, H.A.M. *et al.* (2017) 'Overweight and obesity conditions: Prevalence and associated risk factors in nursing students in a public university in Medellín, Colombia', *Investigacion y Educacion en Enfermeria*, 35(2), pp. 191–198.

Gutiérrez-pliego, L.E. *et al.* (2016) 'Dietary patterns associated with body mass index ( BMI ) and lifestyle in Mexican adolescents', *BMC Public Health*, pp. 1–7.

Harris, J. *et al.* (2019) 'Nutrition transition in Zambia: Changing food supply, food prices, household consumption, diet and nutrition outcomes', *Food Security*, 11(2), pp. 371–387.

Hingorjo, M.R. *et al.* (2009) 'Overweight and obesity in students of a dental college of Karachi: lifestyle influence and measurement by an appropriate anthropometric index.', *JPMA. The Journal of the Pakistan Medical Association*, 59(8), pp. 528–532.

Jiang, S. *et al.* (2018) 'Overweight and Obesity Among Chinese College Students: An Exploration of Gender as Related to External Environmental Influences', *American Journal of Men's Health*, 12(4), pp. 926–934.

Kafyulilo, A.C. (2008) PREVALENCE AND IMPLICATIONS OF OVERWEIGHT AND OBESITY IN CHILDREN'S HEALTH AND LEARNING BEHAVIOUR The Case of Kinondoni and Njombe Districts in Tanzania.

Kayode, O.O. *et al.* (2020) 'Food consumption patterns, physical activity and overweight and obesity among undergraduates of a private university in Nigeria', *Clinical Nutrition Experimental*, 31, pp. 28–34.

Kim, B.Y. *et al.* (2021) 'Association between alcohol consumption status and obesity-related comorbidities in men: data from the 2016 Korean community health survey', *BMC Public Health*, 21(1).

Lobstein, T. *et al.* (2017) 'Comment: obesity as a disease – some implications for the World Obesity Federation's advocacy and public health activities', *Obesity Reviews*, 18(7), pp. 724–726.

Makkawy, E. *et al.* (2021) 'Prevalence of overweight and obesity and their associated factors among health sciences college students, Saudi Arabia', *Journal of Family Medicine and Primary Care*, 10(2), p. 961.

Nawa, M. (2020) 'Investigating the Effect of Prompt Treatment on Malaria Prevalence in Children Aged below Five Years in Zambia□: A Nested Case-Control Study in a Cross-Sectional Survey', *Hindawi, Advances in Public Health* 2020(4289420).

Nawa, M. *et al.* (2020) ‘Modelling malaria reduction in a highly endemic country: Evidence from a Household Survey, Climate and Program Data in Zambia. *Journal of Public Health Research in Africa* 11(1096)

Marcus, K.S. *et al.* (2020) ‘Journal of Obesity and Medical Complications Body Mass Index of Freshman University Students’, pp. 1–5.

McArthur, L. *et al.* (2017) PERCEPTIONS ABOUT OVERWEIGHT AND OBESITY AMONG COLLEGE STUDENTS: APPLICATION OF THE HEALTH BELIEF MODEL APPROVED BY.

Organisation, W.H. (2005) ‘WHO STEPS surveillance manual: the WHO STEPwise approach to chronic disease risk factor surveillance’.

Peltzer, K. *et al.* (2014) ‘Prevalence of overweight/obesity and its associated factors among university students from 22 countries’, *International Journal of Environmental Research and Public Health*, 11(7), pp. 7425–7441.

Raymond, D. *et al.* (2017) ‘Issue 04 | 05 May 2017 The Health Press-Zambia is published by Zambia National Public Health Institute, Ministry of Health Zambia since’, prevention and control, 01(4).

Rodrigues, V.M. *et al.* (2019) ‘Vegetable consumption and factors associated with increased intake among college students: A scoping review of the last ten years’, *Nutrients*.

Di Rosa, C. *et al.* (2020) ‘Very low-calorie ketogenic diets in overweight and obesity treatment: Effects on anthropometric parameters, body composition, satiety, lipid profile and microbiota’, *Obesity Research and Clinical Practice*, 14(6), pp. 491–503.

Sa, J. *et al.* (2021) ‘Sex and racial/ethnic differences in the prevalence of overweight and obesity among U.S. college students, 2011–2015’, *Journal of American College Health*, 69(4), pp. 413–421.

Shmyhol, I. *et al.* (2021) ‘Problems of overweight and obesity among students at general secondary educational institutions’, *Journal of Physical Education and Sport*, 21, pp. 2901–2907.

Tateyama, Y. *et al.* (2018) ‘Obesity matters but is not perceived: A cross-sectional study on cardiovascular disease risk factors among a population-based probability sample in rural Zambia’, *PLoS ONE*, 13(11).

Tateyama, Y. *et al.* (2019) ‘Dietary habits, body image, and health service access related to cardiovascular diseases in rural Zambia: A qualitative study’, *PLoS ONE*, 14(2).

Tomiyama, A.J. (2018) ‘Annual Review of Psychology Stress and Obesity’, *Annu. Rev. Psychol.* 2019, 70, pp. 5–6.

- Tong, T.J. *et al.* (2022) ‘Perception of students on factors contributing to overweight and obesity among high school students in Kiribati: A qualitative study’, PLoS ONE, 17(1 January).
- Traversy, G. *et al.* (2015) ‘Alcohol Consumption and Obesity: An Update’, Current obesity reports, pp. 122–130.
- Urbanetto, J. de S. *et al.* (2019) ‘Stress and overweight/obesity among nursing students’, Revista Latino-Americana de Enfermagem, 27.
- Vandevijvere, S. *et al.* (2015) ‘Increased food energy supply as a major driver of the obesity epidemic: a global analysis’, Bulletin of the World Health Organization, 93(7), pp. 446–456.
- WHO (2016) ‘Adolescents’ Dietary Habits’, Fact Sheet, (March), pp. 2014–2017.
- WHO (2022) ‘World Obesity Atlas 2022’, (March), p. 289.
- Yousif, M.M. *et al.* (2019) ‘Correlation between physical activity, eating behaviour and obesity among Sudanese medical students Sudan’, BMC Nutrition, 5(1).
- ZamStats (2022) ‘2022 CENSUS OF POPULATION AND HOUSING’.