

Evaluating the feasibility, sensitivity, and specificity of next-generation molecular methods for pleural infection diagnosis

3

4 Peter T. Bell^{1,2,3 #}, Timothy Baird^{1,2,4 #}, John Goddard^{1,7}, Olusola S. Olagoke^{2,4}, Andrew Burke^{3,5},
5 Shradha Subedi^{2,6}, Tiana R. Davey^{2,4}, James Anderson^{1,7}, Derek S. Sarovich^{2,4 ^}, and Erin P. Price^{2,4 ^*}

6 ^{#,^} Authors contributed equally

7

8 ¹Department of Respiratory Medicine, Sunshine Coast University Hospital, Birtinya, Queensland,
9 Australia

10 ²Sunshine Coast Health Institute, Birtinya, Queensland, Australia

11 ³School of Medicine, University of Queensland, Herston, Queensland, Australia

12 ⁴Centre for Bioinnovation, University of the Sunshine Coast, Sippy Downs, Queensland, Australia

13 ⁵Department of Respiratory and Thoracic Medicine, The Prince Charles Hospital, Chermside,
14 Queensland, Australia

15 ⁶Department of Infectious Diseases, Sunshine Coast University Hospital, Birtinya, Queensland,
16 Australia

17 ⁷School of Medicine and Dentistry, Griffith University, Sunshine Coast, Queensland, Australia

18

19 ***Corresponding Author**

20 Erin P. Price

21 University of Sunshine Coast

22 eprice@usc.edu.au

23 +61 7 5456 5568

24

25 **Running Title:** Molecular Methods for Pleural Infection Diagnosis

26 **Total Manuscript Word Count:** 3499

27 **Abstract Word Count:** 250

28 **Keywords:** empyema; metagenomics; high-throughput sequencing; parapneumonic effusion; pleural
29 infection; microbiome

30 Abstract

31

32 **Rationale.** Pleural infections are common and associated with substantial healthcare cost, morbidity,
33 and mortality. Accurate pleural infection diagnosis remains challenging due to low culture positivity
34 rates, frequent polymicrobial involvement, and non-specific diagnostic biomarkers.

35 **Objective.** To undertake a prospective pilot study examining the feasibility and challenges associated
36 with molecular methods for diagnosing suspected pleural infection.

37 **Methods.** We prospectively characterised 26 consecutive, clinically suspected pleural infections, and
38 10 consecutive control patients with suspected non-infective pleural effusions, using shotgun
39 metagenomics, bacterial metataxonomics, quantitative PCR, and conventional culture.

40 **Results.** We demonstrate the feasibility of culture-independent molecular techniques for diagnosing
41 suspected pleural infection. Molecular methods exhibited excellent diagnostic performance, with each
42 method identifying 54% (14/26) positive cases among the pleural infection cohort, versus 38% (10/26)
43 with culture. Meta-omics methods unveiled complex polymicrobial infections largely missed by culture.
44 Dominant infecting microbes included streptococci (*S. intermedius*, *S. pyogenes*, *S. mitis*), *Prevotella*
45 spp. (*P. oris*, *P. pleuritidis*), staphylococci (*S. aureus*, *S. saprophyticus*), and *Klebsiella pneumoniae*.
46 However, we encountered challenges that complicated pleural infection interpretation, including: i)
47 uncertainties regarding microbial pathogenicity and the impact of prior antibiotic therapy on diagnostic
48 performance; ii) lack of a clinical diagnostic gold-standard for molecular performance comparisons; iii)
49 potential accidental microbial contamination during specimen collection and processing; and iv)
50 difficulties distinguishing background microbial noise from true microbial signal, particularly in low-
51 biomass specimens.

52 **Conclusions.** Our pilot study demonstrates the potential utility and value of molecular methods in
53 diagnosing pleural infection and highlights key concepts and challenges that should be addressed
54 when designing larger prospective trials.

55 **Key messages**

56

57 *What is already known on this topic*

58 Confident pleural infection diagnosis is often challenging due to low culture positivity rates, frequent
59 polymicrobial involvement, and non-specific diagnostic biomarkers. Limitations of conventional
60 diagnostic tests result in prolonged and inappropriately broad-spectrum antimicrobial use, leading to
61 potentially poorer patient outcomes and avoidable adverse effects.

62

63 *What this study adds*

64 We demonstrate the feasibility, utility, and challenges associated with the use of culture-independent
65 molecular techniques for more accurate pleural infection diagnosis in a real-world clinical setting.

66

67 *How this study might affect research, practice, or policy*

68 These data will help to inform the design of larger prospective clinical trials and identify potential
69 obstacles to be overcome as next-generation sequencing technologies become integrated into routine
70 clinical practice.

71 Introduction

72 Pleural infection is a complex, heterogeneous, and challenging clinical entity associated with
 73 prolonged hospitalization, significant healthcare costs, and high morbidity and mortality¹⁻⁴. Clinical
 74 management frustrations arise due to insensitivities in conventional microbiologic diagnostics (culture-
 75 positive rates of only 20-60%⁵⁻⁹), resulting in prolonged and potentially inappropriate broad-spectrum
 76 antimicrobial use. Culture-based diagnostic sensitivity is poor due to multiple factors, including empiric
 77 antibiotic therapy prior to pleural sampling, low pathogen concentration in pleural fluid, and difficulties
 78 in cultivating fastidious microorganisms^{4,10}. These limitations have obscured characterization of the
 79 pleural 'microbiome' (i.e. the entire microbial community), hampering clinical progress¹¹.

80 Molecular diagnostics enable culture-independent microbial identification from clinical specimens¹².
 81 Such techniques include shotgun metagenomics, metataxonomics, and quantitative PCR (qPCR),
 82 each having unique capabilities and drawbacks. Shotgun metagenomics involves total microbiome
 83 DNA sequencing of a sample, allowing strain-level detection of bacteria, archaea, fungi, parasites,
 84 and DNA viruses, along with their putative functions (e.g. antimicrobial resistance, virulence).
 85 However, metagenomics overlooks RNA viruses, and cannot readily differentiate living from dead
 86 microbes. Metataxonomics involves next-generation amplicon sequencing of conserved bacterial
 87 (16S ribosomal RNA [rRNA]) or fungal internal transcriber spacer (ITS) regions to characterize
 88 bacterial or fungal consortia, respectively; however, these methods typically only resolve microbiota to
 89 the genus level, and cannot be used for relative taxon quantification due to highly variable 16S
 90 rRNA/ITS copy numbers amongst species. Finally, qPCR is an inexpensive, same-day diagnostic
 91 method, with broad resolution ranging from strain to domain level; however, this method is typically
 92 limited to *a priori* target taxa^{13,14}.

93 Several recent studies have applied culture-independent techniques to characterize pleural infections.
 94 This collective work has revealed a pleural microbiome that is often polymicrobial and characterized
 95 by a diverse range of classical pathogens and bystander microbes^{7,8,15-21}. Earlier investigations
 96 implemented bacterial metataxonomics to profile the bacterial microbiome^{7,8,15,21}; however, three
 97 recent studies have implemented metagenomics to more comprehensively profile pleural
 98 microbiota^{16,19,20}. Xu *et al.* 2022 found bacterial dominance in 80 adult Chinese patients with pleural
 99 infections, most commonly *Streptococcus* and *Prevotella* spp., although fungi were occasionally
 100 identified¹⁶. However, the authors only reported taxonomic findings to the genus level, limiting in-

depth understanding of microbiome composition and diversity¹⁶. Liu *et al.* 2023 reported on a single case study of viral pleurisy caused by Epstein-Barr virus, which was successfully treated with acyclovir²⁰. Finally, Liu *et al.* 2023 examined microbiomes of pleural and ascitic fluid in 92 specimens retrieved from 32 Chinese children, identifying *Klebsiella pneumoniae* as the dominant pathogen ($n=17$), followed by *Escherichia coli* ($n=9$), and *Acinetobacter baumannii* ($n=7$); yeasts ($n=6$) and several viruses ($n=33$) were also identified¹⁹. In all three studies, metagenomics yielded a higher pathogen positive rate than conventional culture^{16,19,20}.

To expand knowledge of the pleural microbiome in other cohorts, we undertook a prospective, observational, single-center pilot study of 26 consecutive patients presenting with suspected pleural infection at an Australian hospital, and 10 consecutive control patients lacking clinical suspicion of pleural infection. Next, we compared the diagnostic performance of microbe-enriched shotgun metagenomics, bacterial metataxonomics, panbacterial qPCR, and panfungal qPCR with conventional culture.

Methods

Ethics Statement. The study was approved by The Prince Charles Hospital Human Research Ethics Committee (HREC/2022/QPCH/81858) and the Sunshine Coast Hospital and Health Service Research Governance Office. All participants provided written informed consent.

Study Participants. All patients ≥ 18 years of age presenting with suspected pleural infection (denoted with a '-P' suffix) admitted between March and November 2022 to the Sunshine Coast University Hospital (SCUH), QLD, Australia, and undergoing a diagnostic and/or therapeutic procedure for suspected pleural infection, were prospectively invited.

Due to the lack of a 'gold-standard' pleural infection confirmation test, 'clinical likelihood of pleural infection' was objectively stratified by a specialist respiratory physician using predefined clinical criteria: 1. *Probable Pleural Infection*: (i) clinical evidence of pleural infection based on fever, elevated white cell count with left shift, pleurisy presence, raised inflammatory markers, and suitable clinical presentation; AND (ii) either direct evidence of pleural infection defined as the presence of Gram stain and/or culture positivity on pleural fluid; OR indirect evidence of pleural infection based upon pleural fluid pH < 7.2 as measured on a blood gas analyser and/or glucose < 3.0 mmol/L AND (iii) accompanied by at least two of the following criteria: serum C-reactive protein (CRP) > 100 mg/L; complex pleural fluid and evidence of infection on imaging (pleural fluid loculation/septation on ultrasound and/or computed tomography); fever ($> 38.0^{\circ}\text{C}$). 2. *Possible Pleural Infection*: (i) clinical evidence of pleural infection (as defined in (i) above), AND (ii) negative Gram stain and culture negative on pleural fluid AND pH ≥ 7.2 and glucose > 3.0 mmol/L) AND (iii) accompanied by at least 2 of the following criteria: CRP > 100 mg/L; complex (i.e. loculated) pleural fluid collection on imaging; fever ($\geq 38.0^{\circ}\text{C}$). 3. *Unlikely pleural infection*: (i) No clinical evidence of pleural infection, AND (ii) all of the following criteria are met: pleural fluid Gram stain and culture negative with pH ≥ 7.2 , glucose > 3.0 mmol/L, serum CRP < 100 mg/L, uncomplicated effusion on CT and or ultrasonographic imaging, and an alternate diagnosis confirmed (i.e. heart failure, malignancy). Based on these criteria, 26 consecutive patients with clinically suspected pleural infections were recruited.

For the control group, patients with pleural effusions without clinical suspicion for pleural infection at the time of sampling (denoted with a '-C' suffix), were enrolled. These individuals were approached if they had an alternative effusion cause (e.g., malignancy or heart failure), absence of clinical history to

support active infection, and simple effusion appearance on ultrasound and/or computed tomography. Patients were excluded if they had prior sampling of the pleural space, any clinical concern for current or recent infection, concomitant abdominal ascites, hepatic hydrothorax, or antibiotic administration within 3 weeks of pleural sampling. Ten participants meeting these criteria were enrolled as non-infected control subjects.

Specimen Collection, Storage, and Processing. The study scientific and pathology teams were provided participant categorisation by the study clinicians ('-P' vs. '-C') but no other information. For the pleural infection cohort, specimens were collected as soon as practicable using aseptic technique. Antibiotic commencement ranged from 21 days prior to pleural sampling to immediately post-pleural aspiration (**Table S1**). Following 10mL 1% lidocaine subcutaneous administration, between 10 and 60mL pleural fluid was collected into two sterile specimen pots. One pot was immediately mixed with 7mL stabilization agent²² for total DNA extraction (bacterial metataxonomics and qPCR); the second pot contained only pleural fluid for culture and host DNA depletion and extraction (metagenomics). Specimens were immediately stored at 4°C until processed. In parallel, specimens were also collected for routine microscopy (cell count, white blood cell differential, Gram stain), and biochemistry analyses (pleural fluid pH, protein, glucose, lactate dehydrogenase; serum CRP).

Microbial culturing was undertaken by both SCUH's onsite Pathology laboratory and our Research laboratory, with VITEK 2 (bioMérieux, NSW, Australia) or Illumina NovaSeq 6000 whole-genome sequencing (WGS; Australian Centre for Ecogenomics, QLD, Australia) used respectively to confirm species identity. Bacterial metataxonomics was carried out using Illumina MiSeq 300bp V3-V4 16S rRNA amplicon paired-end sequencing²³ (Ramaciotti Centre for Genomics, NSW, Australia). Metagenomic sequencing of host-depleted DNA²⁴ was performed using Illumina NovaSeq 6000 150bp paired-end reads (17-27 million reads/sample; Azenta Life Sciences, China). For qPCR analyses, total genomic DNA was tested against panbacterial (16S rRNA)²⁵, panfungal (28S rRNA)²⁶, and human (β -globin)²⁷ qPCR assays to quantify microbial-to-host load. For the panbacterial qPCR assay, a cycles-to-threshold (C_T) cut-off value of 31 was used to delineate pleural infection-positive versus -negative cases. Further methodological details are described in Supplementary Methods.

Statistical Analyses. To determine diagnostic test performance, '*Probable Pleural Infection*' was used as the infection-positive reference. Specificity, sensitivity, positive predictive value (PPV), and negative predictive value (NPV) were calculated for microbial culture, qPCR, bacterial

173 metataxonomics, and metagenomics. Spearman's rank correlation (r_s) was used to determine pleural
174 infection/no-infection diagnostic correlations between the four testing methods.

Results

Study participants and clinical outcomes. Twenty-six participants ($n=7$ female; median age=67 years [range 20 – 95]) with suspected pleural infection had an exudative pleural effusion as defined by Light's criteria²⁸. Twenty-four (92%) had loculated (complex) pleural effusions on ultrasound and/or computed tomography imaging, with 20 (77%) also having associated lung consolidation at the time of cross-sectional imaging. Twenty-five (96%) underwent intercostal catheter insertion, whereas one underwent diagnostic/therapeutic needle aspiration. All 26 patients received broad-spectrum antimicrobial therapy (**Table S1**). Intrapleural enzyme therapy (5mg DNase and 10 mg tissue plasminogen activator) was administered in 17 participants (65%), and four (15%) required surgical decortication (**Table 1 & Table S2**). Thirty-day mortality was 12% ($n=3$).

Among the 10 control subjects, 9 were diagnosed with malignant pleural effusion (MPE), and one with decompensated congestive heart failure (**Table 1 & Table S2**). Thirty-day mortality was not recorded for this cohort.

Conventional culture performance. Pathology lab testing returned 10 (38%) microbial culture-positive results from the pleural infection cohort (**Table 2**), whereas only six (23%) were culture-positive from Research lab testing; however, there was 100% overlap with the Pathology results. Unexpectedly, pleural fluid from two control participants, SCHI0152-C and SCHI0175-C, cultured *Candida parapsilosis* and *Moraxella osloensis*, respectively, in the Research lab. No microbes were cultured in the control cohort by the Pathology lab.

qPCR performance. Positive bacterial detection was achieved with panbacterial qPCR in 14/26 (54%) suspect pleural infection cases (**Table 2**). All panbacterial qPCR-positive results were also bacteria-positive according to metagenomics and/or bacterial metataxonomics (**Table 2**). One of the 10 control cases was panbacterial qPCR-positive, albeit at a borderline C_T value of 30.9; this result was not supported by any of the other tested methods. Panfungal qPCR did not identify fungi in any pleural infection or control cohort specimens.

Bacterial metataxonomic and shotgun metagenomic performance. Together, bacterial metataxonomics and metagenomics identified 16/26 (62%) suspect pleural infection cases as infection-positive, with 14/26 (54%) cases determined by each individual method. Metagenomics identified 7/14 (50%) cases as polymicrobial, vs. 5/14 (36%) polymicrobial cases using bacterial

metataxonomics (**Table 2 & Figure 1**). Despite both methods identifying 14 pleural infection cases each, only 12 overlapped, and only three of these cases were identified as polymicrobial by both methods (**Table 2**). Far fewer taxa were identified by bacterial metataxonomics in two of these three cases (SCHI0131-P, SCHI0172-P) (**Table 2**).

Molecular diagnostics versus microbial culture performance. Compared with our 'pleural infection-positive' clinical reference classification, all molecular diagnostic methods demonstrated superior sensitivity and NPVs compared with microbial culture (**Table 3**). As expected, specificity and PPVs remained high across all diagnostic methods.

Conventional culture somewhat correlated with molecular methods (metagenomics: $r_s=0.56$ [$p<0.001$]; bacterial metataxonomics: $r_s=0.39$ [$p=0.05$]; panbacterial qPCR: $r_s=0.39$ [$p=0.05$]) (**Figure 3**). Metagenomics demonstrated the strongest correlation across methods (bacterial metataxonomics: $r_s=0.69$ [$p<0.001$]; panbacterial qPCR: $r_s=0.69$ [$p<0.001$]; conventional culture: $r_s=0.56$ [$p<0.01$]). The strongest individual correlation was between panbacterial qPCR and bacterial metataxonomics ($r_s=0.85$; $p<0.001$; **Figure 3**); this result was expected as both methods target the 16S rRNA gene.

Pleural infection aetiology. A diverse array of oro-naso-hypopharyngeal-, oral-, dental-, respiratory tract-, and gut-associated microbes were detected in the 14 metagenomic infection-positive specimens (**Table 2 & Figure 1**). Streptococci ($n=5$; 2 x *S. intermedius*, 2 x *S. pyogenes*, 1 x *S. mitis*) dominated, followed by *Prevotella* spp. ($n=2$; *P. oris* and *P. pleuritidis*), *Staphylococcus* spp. ($n=2$; *S. aureus* and *S. saprophyticus*), and *Klebsiella pneumoniae* ($n=2$) (**Table 2**). Seven (50%) pleural infection cases were monomicrobial according to metagenomics, of which two harboured *S. intermedius*, along with one each of *K. pneumoniae*, *S. saprophyticus*, *S. aureus*, *Enterococcus faecium*, and *Escherichia coli* (**Table 2 & Figure 2**). In contrast, bacterial metataxonomics identified nine (35%) monomicrobial pleural infection cases, of which four were *Streptococcus* sp., along with one each of *Escherichia-Shigella* sp., *Fusobacterium* sp., *Haemophilus* sp., *Klebsiella* sp., and *Prevotella* sp. (**Table 2 & Figure 2**). Five monomicrobial cases overlapped with the two meta-omics methods; however, bacterial metataxonomics erroneously assigned two monomicrobial pleural infection cases, SCHI0132-P and SCHI0143-P, as '*Streptococcus*', despite the SCHI0132-P isolate being confirmed as coagulase-negative *Staphylococcus* sp. by VITEK 2 (**Table 2**), indicating a probable bacterial metataxonomic database error for this genus.

As expected, bacterial metataxonomics failed to identify fungi or viruses in the pleural infection cases, whereas conventional culture and metagenomics both identified the yeast, *Pichia kudriavzevii*, in SCHI0131-P. Metagenomics further identified the fungi *Rhizoctonia solani*, *Wallemia mellicola*, *Meyerozyma guilliermondii*, *Truncatella angustata*, *Cokeromyces recurvatus*, *Bipolaris maydis*, *Candida orthopsilosis*, and *Gilbertella persicaria* in SCHI0131-P (subject suffered from confirmed Boorhaaves syndrome), *Candida dubliniensis* in SCHI0140-P, and *Candida tropicalis*, *Fonsecaea pedrosoi*, and *Pyricularia pennisetigena* in SCHI0125-P. In all instances, fungi were in relatively low abundance. Metagenomics also identified bacteriophages in 5/26 infection-positive cases, with *Staphylococcus* phage Andhra seen in SCHI0123-P, SCHI0125-P, SCHI0127-P, and SCHI0128-P, and *Streptococcus* phage SM1 seen in SCHI0131-P. *Staphylococcus* phage Andhra was among the most dominant taxa seen in SCHI0125-P and SCHI0128-P, indicating very high phage levels in pleural fluid; both specimens harboured *S. aureus* according to metagenomics, and SCHI0125-P also had *S. saprophyticus* according to culture and WGS (Table 2 & Figure 2).

Metagenomics unexpectedly found microbes in 2/10 control cases (Table 2). The non-pathogenic DNA virus, adeno-associated dependoparvovirus A, was identified in SCHI0173-C – although no helper virus was observed – and *M. osloensis* was confirmed in SCHI0175-C. *C. parapsilosis* cultured from SCHI0152-C was not detected, suggesting possible operator contamination during specimen collection or processing. Bacterial metataxonomics failed to identify the culture-confirmed *M. osloensis* in SCHI0175-C, but identified *Escherichia-Shigella* in SCHI0152-C (1453 reads) and *Corynebacterium* in SCHI0173-C (190 reads), findings that were not supported by the other methods, again likely due to contamination.

Microbiology and clinical outcomes. Of the three pleural infection cases with a <30-day mortality, two (SCHI0128-P and SCHI0134-P) were infected with *K. pneumoniae*, and two (SCHI0131-P and SCHI0128-P) had polymicrobial infections according to metagenomics. In contrast, no patient surviving beyond 30 days ($n=23$) harbored *Klebsiella* spp. according to any testing method.

Antimicrobial resistance gene detection. *In silico* antimicrobial resistance gene detection only identified innate drug resistance genes, with no evidence of acquired resistance development in any case (Table S1).

Discussion

Clinical decision-making in suspected pleural infection is challenging due to low culture positivity rates⁵⁻⁸, resulting in frequent diagnostic uncertainty and antimicrobial overuse. Recent work^{7,8,15,16,19} has highlighted potential advantages of molecular methods for diagnosing and determining the aetiology of pleural infection. Our pilot feasibility study adds to this growing body of work, being one of the first studies to apply shotgun metagenomics on pleural infections, and the first to prospectively collect and profile consecutive pleural fluid specimens. Our findings suggest that all three culture-independent molecular techniques tested in our study – metagenomics, bacterial metataxonomics, and panbacterial qPCR – have excellent performance characteristics for pleural infection diagnosis, and may be superior to conventional culture for diagnosis and aetiological confirmation. Panbacterial qPCR stands out as a cost-effective, simple, and same-day diagnostic that could be readily implemented as a routine test for pleural fluid specimens; such a test may assist with both antimicrobial stewardship efforts and rapid treatment decision-making²⁹. In support of its clinical utility, panbacterial qPCR showed excellent diagnostic performance (sensitivity 82%, specificity 95%, PPV 93%, NPV 86%) and strong correlation with shotgun metagenomics, bacterial metataxonomics, and conventional culture.

In line with recent molecular studies^{7,8,15-17}, we confirm that pleural infections can be either monomicrobial or polymicrobial, with predominant microbes most likely originating from oral, dental, or upper and lower respiratory tract sources, and occasional incursion from gut-borne pathogens (e.g. *E. coli*, *Enterococcus faecium*, *K. pneumoniae*) (Table 2). Furthermore, metagenomics and bacterial metataxonomics unveiled greater microbial heterogeneity than could be detected by conventional culture. Similar to recent work^{16,19}, we observed that polymicrobial bacterial infection was also occasionally associated with fungal coinfection (albeit in very low abundance) using metagenomics, confirming the challenging nature of diagnosing and treating complex polymicrobial pleural infections across microbial domains. Streptococcal and staphylococcal phages were also seen in five cases of pleural infection, with their host species not always detected, suggesting possible lytic activity towards their host within the pleural space, or incidental phage transit from other body sites³⁰. Although not examined here, others have shown that phage presence in clinical specimens can negatively impact culture rates³¹. Despite low sample numbers, we found that *K. pneumoniae* presence was associated

with greater risk of mortality, consistent with recent bacterial metataxonomic study that showed an association between Enterobacteriaceae and mortality⁸.

Immediate antimicrobial administration upon suspicion of pleural infection is important for preventing sepsis³²; however, this practice may negatively affect pleural infection diagnostic rates. Despite this, many prior pleural microbiome studies have identified 100% infection positivity using molecular testing^{7,8,15}. In contrast, only 62% of suspected pleural infections in our study had detectable microbes according to molecular and/or culture assessment. Our lower positivity rate likely reflects our prospective, consecutive recruitment of participants without prior knowledge of culture results or pleural biochemistry, an approach that captures the diagnostic uncertainty facing clinicians in real-world practice. Only two other studies have not used prior knowledge of pleural fluid biochemistry or microbiology for molecular pleural infection identification and characterisation; their infection positivity rate was 70%^{16,17}. Our finding that 38% of suspected pleural infection patients did not have detectable microbes according to any method suggests that a subset of patients with clinically suspected pleural infection may instead have sterile pleural effusions at the time of sampling. In support of this conclusion, these patients had more benign pleural biochemistry using recognised cut-off values (i.e. pH <7.2; glucose <3.0 mmol/L)¹ (**Table 1**). A second possibility is that some infections were missed due to prior antibiotic exposure; however, others have shown that molecular approaches are less impacted by prior antibiotics than conventional culture^{7,16}. A third possibility is that human DNA depletion prior to metagenomic sequencing may have limited our ability to detect extracellular microbial DNA²⁴. To overcome this issue, deeper metagenomic sequencing on total DNA can be conducted; however, such an approach is costly, and risks missing low-abundance microbes due to overwhelming human DNA signal^{24,33}. Prospective randomized trials are needed to understand whether a truncated antibiotic course can be safely used in people with likely sterile effusions as identified with molecular diagnostics, and to quantify the impact of prior antibiotic administration on culture or molecular detection sensitivity.

Sensitive molecular assays come with the trade-off that microorganism presence might represent either a contaminant or a bona fide pathogen. To assess this issue, we investigated whether the control group, predominantly comprised of MPEs, had a pleural microbiome. Although most controls had a sterile pleural space, one MPE patient was positive for *Moraxella osloensis* on culture, WGS, and metagenomics. *M. osloensis*, a commensal of human skin and the upper respiratory tract, most

commonly causes infection in cancer patients³⁴; it is thus feasible that this patient had an undiagnosed infection alongside their MPE. Three other MPE patients also unexpectedly yielded microbes: *C. parapsilosis* was cultured from SCHI0152-C, and bacterial metataxonomics identified *Corynebacterium* and *Escherichia-Shigella* in SCHI0173-C and SCHI0152-C, respectively; however, these findings were not supported by the other methods. Although specimen or sampling contamination represents the most plausible explanation for these results, a recent bacterial metataxonomic study suggested that non-infective pleural disease may contain unique microbial signatures^{7,35}. Strict aseptic practice during specimen collection and processing can mitigate some of these false-positive cases. However, microbial contamination may still occur at many stages of sample handling and analysis, including procedural contamination (e.g. remnant microbial DNA in the specimen jars or local anaesthetics), or residual DNA from transient microbial seeding of the pleura. In addition, index hopping and 'kitome' contamination are known drawbacks of next-generation sequencing, especially in low-biomass specimens like pleural fluid³³. Careful consideration of these challenges will be important in the design of larger prospective studies. Further studies examining the pleural microbiome across a spectrum of pleural diseases, including in people with MPE and congestive heart failure, will also be illuminating.

Other challenges also complicated interpretation of pleural diagnosis and aetiology, including difficulties distinguishing background noise from true microbial signal, lack of appropriate methodology for detecting RNA viruses, and uncertainty regarding microbial causality and pathogenicity (e.g. the role of bystander microbes in pleural disease development and prognosis), particularly in polymicrobial infections. Shotgun metatranscriptomics, which involves total RNA sequencing of microbiome transcripts, may address these difficulties. However, to our knowledge, this technique has not yet been undertaken to investigate pleural infections, so its value in this cohort remains unclear. Perhaps most challenging of all is the current lack of gold-standard clinical diagnostic criteria for pleural infection. In clinical practice, an integrated approach is taken to weigh multiple factors to make a probabilistic diagnosis, including pleural biochemistry, host response, and imaging. In this study, we attempted to standardize integrated probabilistic diagnostic classification by formulating a pragmatic reference standard to compare diagnostic performance between modalities. However, the issue of a validated diagnostic gold-standard remains an unresolved challenge. It is also important to consider the impact of incorporation bias³⁶ when interpreting the performance of

350 conventional culture results, given that such results contribute to the diagnostic 'reference standard',
 351 thereby inflating culture performance estimates.

352 In conclusion, this prospective study demonstrates the feasibility of, and challenges associated with,
 353 molecular tests for pleural infection diagnosis. Whilst such methods hold great clinical utility and
 354 potential superiority in routine pleural infection diagnostics, questions and challenges remain. A large,
 355 rigorous study comparing molecular and culture-based diagnostic techniques in pleural infection is
 356 now required to further address these opportunities for diagnostic advancement, along with the
 357 associated challenges in eventual clinical translation.

Author Contributions: The authors confirm contribution to the paper as follows: study conception and design, PTB, TB, JG, OSS, AB, DSS, EPP; data collection, PTB, TB, JG, OSS, AB, JA, DSS, EPP; analysis and interpretation of results, PTB, TB, JG, OSS, AB, SS, TRD, DSS, EPP; draft manuscript, PTB, TB, DSS, EPP; preparation of manuscript, PTB, JG, OSS, AB, SS, TRD, DSS, EPP. All authors reviewed the results and approved the final version of the manuscript.

Funding: This work was funded by Wishlist Sunshine Coast Hospital Foundation (award 2021-04-CRG to TB, OO, JG, DSS, and EPP) and Advance Queensland (awards AQIRF0362018 to EPP and AQIRF095-2020-CV to OSO). PTB has no funding to declare.

Conflicts of Interest Statement: The authors declare no conflict of interest. The funders had no role in study design, in the collection, analyses, or interpretation of data, in manuscript writing, or in the decision to publish the results.

Acknowledgements: We thank Emma Seaniger and Louise McIntosh for assistance with sample collection and study coordination, Michael Caffery and the Pathology Queensland laboratory scientists for their assistance with culture, and the Australian BioCommons team (led by Ashley Dungan at University of Melbourne, and Melissa Burke at University of Queensland) for providing QIIME 2 training.

Figure Legends

Figure 1. Metagenomic profiles identified from 26 suspected pleural infection specimens and 10 control cohort specimens. *Conventional culture positive; ^16S ribosomal RNA (panbacterial) quantitative PCR positive; #Whole-genome sequencing-confirmed (from culture). Dominant taxa are labelled on each bar.

Figure 2. Bacterial 16S ribosomal RNA (rRNA) metataxonomic profiles identified from 26 suspected pleural infection specimens and 10 control cohort specimens. *Conventional culture positive; ^16S ribosomal RNA (panbacterial) quantitative PCR positive; #Whole-genome sequencing result from culture; ‡Likely incorrect assignment of *Staphylococcus* as *Streptococcus* (1 case confirmed as a coagulase-negative *Staphylococcus* by MALDI-TOF analysis of culture). Dominant taxa are labelled on each bar.

Figure 3. Correlation matrix of the pleural infection diagnostic methods used in this study. 1=perfect correlation; -1=no correlation. Conventional culture showed the weakest correlation when compared with each of the three molecular methods.

394 **Table 1. Clinical, Demographic and Outcome Data**

	Suspected Pleural Infection	Non-infectious Pleural Effusion (Control)
<i>Clinical / Demographics</i>		
Number of participants in group, <i>n</i>	26	10
Age in years (median, range)	67 (20-95)	78 (41-87)
Female sex, <i>n</i> (%)	7 (27)	2 (20)
Immunosuppression, <i>n</i> (%)	3 (12)	1 (10)
Fever, <i>n</i> (%)	23 (90)	–
ICU admission, <i>n</i> (%)	6 (23)	–
<i>Biochemical</i>		
Serum White Cell count, cells x10 ⁹ /L (median, range)	14.5 (8.4-33.6)	8.3 (4.3-15.2)
Serum C-reactive protein, mg/L (median, range)	193 (39-547)	56 (5-124)
Serum Procalcitonin, µg/L (median, range)	0.6 (0.1-9.6)	0.2 (0.1-0.3)
Serum LDH, unit/L (median, range)	236 (108-477)	208 (160-391)
Serum Protein, g/L (median, range)	61 (49-76)	62 (45-72)
Fluid LDH, unit/L (median, range)	865 (173-17400)	239 (81-672)
Fluid Protein, g/L (median, range)	46 (23-58)	43 (37-46)
Fluid pH (median, range)	7.0 (6.6-7.5)	7.4 (7.3-7.5)
Fluid glucose, mmol/L (median, range)	3.9 (2.4-11.9)	6.3 (4.6-10.0)
<i>Imaging</i>		
Consolidation, <i>n</i> (%)	20 (80)	–
Complex appearance of effusion, <i>n</i> (%)	24 (92%)	–
<i>Outcome Data</i>		
IPE, <i>n</i> (%)	17 (65)	–
CTS, <i>n</i> (%)	4 (15)	–
30-day mortality, <i>n</i> (%)	3 (12)	–
Pleural infection Likelihood* (%)	1=0%, 2=38.5%, 3=61.5%	1=100%, 2=0%, 3= 0%

395

396 *Abbreviations:* Cardiothoracic Surgical decortication; IPE, Intrapleural Enzyme therapy; LDH, Lactate
397 dehydrogenase. * 1 = *Unlikely pleural infection*, 2 = *Possible pleural infection*: 3 = *Probable pleural*
398 *infection*.

Table 2. Pleural space diagnostic performance across 26 pleural infection diagnosed participants and 10 controls with a diagnosed non-pleural infection aetiology

Participant ID	Metagenomic result [#]	Bacterial metataxonomic result [†]	Culture result, Research Lab (WGS result)	Culture result, Pathology Lab (MALDI-TOF result)	Panbacterial qPCR [‡]
Diagnosed with Pleural Infection					
SCH10123-P	Polymicrobial (<i>Fusobacterium nucleatum</i> , <i>Fusobacterium hwasookii</i> , <i>Staphylococcus</i> phage Andhra, <i>Prevotella loescheii</i> , <i>Rothia mucilaginosa</i> , <i>Enterococcus faecium</i> , <i>Fusobacterium</i> sp. oral taxon 203)	Monomicrobial (<i>Fusobacterium</i>)	NG	NG	Yes (C _T =20.8)
SCH10125-P	Polymicrobial (<i>Staphylococcus</i> phage Andhra, <i>Prevotella oris</i> , <i>Enterococcus faecium</i> , <i>Streptococcus mitis</i> , <i>Streptococcus anginosus</i> , <i>Escherichia coli</i>^a , <i>Streptococcus pneumoniae</i> , <i>Fonsecaea pedrosoi</i> , <i>Candida tropicalis</i> , <i>Pyricularia pennisetigena</i> , <i>Staphylococcus aureus</i>^b)	Monomicrobial (<i>Prevotella</i>)	<i>Escherichia coli</i>[*] , <i>Staphylococcus saprophyticus</i>[*]	<i>Escherichia coli</i>	Yes (C _T =25.7)
SCH10126-P	ND	ND	NG	NG	No (C _T =33.9)
SCH10127-P	Polymicrobial (<i>Streptococcus pyogenes</i> , <i>Staphylococcus</i> phage Andhra, <i>Streptococcus dysgalactiae</i> , <i>Streptococcus agalactiae</i> , <i>Enterococcus faecium</i> , <i>Rothia mucilaginosa</i>)	Monomicrobial (<i>Streptococcus</i>)	Positive (not characterised)	<i>Streptococcus pyogenes</i>	Yes (C _T =27.5)
SCH10128-P	Polymicrobial (<i>Klebsiella pneumoniae</i> , <i>Staphylococcus</i> phage Andhra, <i>Mycoplasma hyorhinis</i> , <i>Carnobacterium divergens</i> , <i>Actinomyces odontolyticus</i> , <i>Rothia mucilaginosa</i> , <i>Streptococcus mitis</i> , <i>Haemophilus parainfluenzae</i> , <i>Enterococcus faecium</i> , <i>Streptococcus parasanguinis</i> , <i>Staphylococcus aureus</i>)	ND	NG	NG	No (C _T =35.3)

SCHI0131-P ^e	Polymicrobial (<i>Streptococcus mitis</i>, <i>Streptococcus anginosus</i>, <i>Streptococcus pneumoniae</i>, <i>Streptococcus</i> sp. oral taxon 431, <i>Streptococcus oralis</i>, <i>Streptococcus salivarius</i>, <i>Gemella haemolysans</i>, <i>Streptococcus pseudopneumoniae</i>, <i>Streptococcus</i> sp. ChDC B345, <i>Streptococcus gordonii</i>, <i>Streptococcus parasanguinis</i>, <i>Streptococcus cristatus</i>, <i>Streptococcus</i> sp. HSiSM1, <i>Streptococcus australis</i>, <i>Streptococcus</i> sp. oral taxon 064, <i>Streptococcus</i> sp. FDAARGOS_192, <i>Streptococcus intermedius</i>, <i>Streptococcus</i> sp. A12, <i>Veillonella atypica</i>, <i>Actinomyces odontolyticus</i>, <i>Streptococcus sanguinis</i>, <i>Gemella sanguinis</i>, <i>Streptococcus thermophilus</i>, <i>Capnocytophaga leadbetteri</i>, <i>Streptococcus equinus</i>, <i>Streptococcus viridans</i>, <i>Rothia dentocariosa</i>, <i>Capnocytophaga gingivalis</i>, <i>Veillonella parvula</i>, <i>Rothia mucilaginosa</i>, <i>Streptococcus vestibularis</i>, <i>Streptococcus</i> sp. HSiSS2, <i>Prevotella jejuni</i>, <i>Streptococcus constellatus</i>, <i>Streptococcus</i> sp. I-P16, <i>Streptococcus</i> sp. I-G2, <i>Veillonella dispar</i>, <i>Bifidobacterium longum</i>, <i>Streptococcus</i> sp. HSiSS3, <i>Gemella morbillorum</i>, <i>Haemophilus influenzae</i>, <i>Streptococcus</i> sp. HSiSS1, <i>Atopobium parvulum</i>, <i>Streptococcus dysgalactiae</i>, <i>Streptococcus pyogenes</i>, <i>Streptococcus agalactiae</i>, <i>Staphylococcus aureus</i>, <i>Prevotella salivae</i>, <i>Prevotella melaninogenica</i>, <i>Tannerella</i> sp. oral taxon HOT-286, <i>Actinomyces</i> sp. oral taxon 171, <i>Actinomyces oris</i>, <i>Capnocytophaga sputigena</i>, <i>Haemophilus parainfluenzae</i>, <i>Neisseria subflava</i>, <i>Campylobacter showae</i>, <i>Streptococcus</i> phage SM1, <i>Streptococcus mutans</i>, <i>Streptococcus sobrinus</i>, <i>Streptococcus acidominimus</i>, <i>Lactococcus lactis</i>, <i>Lachnospiraceae</i> bacterium oral taxon 500, <i>Actinomyces naeslundii</i>, <i>Actinomyces viscosus</i>, <i>Capnocytophaga</i> sp. oral taxon 878, <i>Neisseria elongata</i>, <i>Lautropia mirabilis</i>, <i>Fusobacterium periodonticum</i>, <i>Fusobacterium nucleatum</i>, <i>Streptococcus</i>	Polymicrobial (<i>Streptococcus</i>, <i>Streptococcus anginosus</i>, <i>Granulicatella</i>, <i>Streptococcus salivarius</i>, <i>Gemella</i>, <i>Prevotella jejuni</i>, <i>Veillonella</i>, <i>Streptococcus constellatus</i>, <i>Prevotella melaninogenica</i>; <i>Neisseria perflava</i>, <i>Neisseria</i>, <i>Fusobacterium periodonticum</i>, <i>Prevotella salivae</i>, <i>Haemophilus influenzae</i>, <i>Leptotrichia</i>, <i>Capnocytophaga gingivalis</i>, <i>Capnocytophaga leadbetteri</i>, <i>Schaalia odontolytica</i>, <i>Neisseria elongata</i>, <i>Atopobium</i>,	<i>Pichia kudriavzevii</i>*	<i>Pichia kudriavzevii</i> ; enteric & mixed anaerobic bacteria	Yes (C _T =17.7)
-------------------------	--	--	------------------------------------	---	----------------------------

	<i>ratti</i> , >70 additional taxa ⁶)	<i>Fusobacterium</i> , <i>Porphyromonas</i> , <i>Bergeyella</i> , <i>Leptotrichia</i> <i>hongkongensis</i> , <i>Oribacterium</i> , <i>Lautropia</i> , <i>Oribacterium</i> , <i>Selenomonas artemidis</i> , <i>Stomatobaculum</i> , <i>Haemophilus</i> , <i>Prevotella</i> <i>shahii</i> , <i>Rothia</i>)			
SCHI0132-P	Monomicrobial (<i>Staphylococcus aureus</i> ^b)	Monomicrobial (<i>Streptococcus</i> ^d)	Positive (not characterised)	Coagulase-negative <i>Staphylococcus</i> sp.	No ($C_T=33.5$)
SCHI0133-P	ND	ND	NG	NG	No ($C_T=33.9$)
SCHI0134-P	Monomicrobial (<i>Klebsiella pneumoniae</i>)	Monomicrobial (<i>Klebsiella</i>)	NG	<i>Klebsiella pneumoniae</i>	Yes ($C_T=23.5$)
SCHI0135-P	ND	ND	NG	NG	No ($C_T=34.5$)
SCHI0136-P	Monomicrobial (<i>Streptococcus intermedius</i>)	ND	NG	<i>Streptococcus</i> <i>intermedius</i>	Yes ($C_T=27.5$)
SCHI0140-P	Polymicrobial (<i>Streptococcus pyogenes</i> , <i>Streptococcus dysgalactiae</i> , <i>Candida dubliniensis</i>)	Polymicrobial (<i>Streptococcus</i> , <i>Streptococcus</i>)	NG	<i>Streptococcus</i> <i>pyogenes</i> , <i>Staphylococcus aureus</i>	Yes ($C_T=25.2$)

		<i>pyogenes</i>)			
SCHi0141-P	ND	ND	NG	NG	No ($C_T=33.5$)
SCHi0143-P	Monomicrobial (<i>Staphylococcus saprophyticus</i>)	Monomicrobial (<i>Streptococcus</i> ^d)	NG	NG	Yes ($C_T=30.1$)
SCHi0148-P	ND	ND	NG	NG	No ($C_T=34.6$)
SCHi0149-P	ND	ND	NG	NG	No ($C_T=31.3$)
SCHi0150-P	ND	Monomicrobial (<i>Haemophilus</i>)	NG	NG	Yes ($C_T=28.1$)
SCHi0157-P	Monomicrobial (<i>Streptococcus intermedius</i>)	Monomicrobial (<i>Streptococcus</i>)	NG	<i>Streptococcus intermedius</i>	Yes ($C_T=24.2$)
SCHi0158-P	ND	Polymicrobial (<i>Fusobacterium</i> , <i>Parvimonas</i> , <i>Eubacterium</i>)	NG	NG	Yes ($C_T=29.2$)
SCHi0159-P	ND	ND	NG	NG	No ($C_T=32.6$)
SCHi0160-P	ND	ND	NG	NG	No ($C_T=32.3$)
SCHi0162-P	ND	ND	NG	NG	No ($C_T=31.2$)
SCHi0163-P	Monomicrobial (<i>Haemophilus parainfluenzae</i>)	Polymicrobial (<i>Haemophilus</i> , <i>Novosphingobium</i>)	NG	NG	Yes ($C_T=23.5$)
SCHi0166-P	Monomicrobial (<i>Escherichia coli</i>)	Monomicrobial (<i>Escherichia-Shigella</i>)	Positive (not characterised)	<i>Escherichia coli</i>	Yes ($C_T=21.5$)
SCHi0169-P	ND	ND	NG	NG	No ($C_T=33.2$)

SCH10172-P	Polymicrobial (<i>Prevotella pleuritidis</i> , <i>Fusobacterium nucleatum</i> , <i>Prevotella enoeca</i> , <i>Prevotella intermedia</i> , <i>Hallela seregens</i> , <i>Porphyromonas gingivalis</i> , <i>Tannerella forsythia</i> , <i>Prevotella buccae</i> , <i>Prevotella oris</i> , <i>Fusobacterium</i> sp. oral taxon 203, <i>Prevotella nigrescens</i> , <i>Prevotella denticola</i> , <i>Streptococcus pyogenes</i> , <i>Prevotella veroralis</i> , <i>Fusobacterium hwasookii</i> , <i>Prevotella aurantiaca</i> , <i>Bacteroides fragilis</i> , <i>Bacteroides heparinolyticus</i> , <i>Prevotella salivae</i> , <i>Prevotella fusca</i> , <i>Prevotella pallens</i> , <i>Prevotella nanceiensis</i> , <i>Prevotella jejuni</i> , <i>Prevotella oulorum</i> , <i>Prevotella maculosa</i> , <i>Prevotella bivia</i> , <i>Prevotella melaninogenica</i> , <i>Prevotella micans</i> , <i>Prevotella amnii</i> , <i>Prevotella conceptionensis</i> , <i>Enterococcus faecium</i>)	Polymicrobial (<i>Enterococcus</i> , <i>Prevotella pleuritidis</i>)	Positive (not characterised)	<i>Fusobacterium nucleatum</i>	Yes ($C_T=20.3$)
Diagnosed with non-Pleural infection (Negative Controls)					
SCH10151-C	ND	ND	NG	NG	No ($C_T=32.7$)
SCH10152-C	ND	Monomicrobial (<i>Escherichia-Shigella</i>)	<i>Candida parapsilosis</i>*	NG	No ($C_T=35.7$)
SCH10161-C	ND	ND	NG	NG	No ($C_T=36.7$)
SCH10164-C	ND	ND	NG	NG	No ($C_T=35.9$)
SCH10165-C	ND	ND	NG	NG	No ($C_T=32.3$)
SCH10168-C	ND	ND	NG	NG	No ($C_T=32.7$)
SCH10170-C	ND	ND	NG	NG	No ($C_T=38.7$)
SCH10173-C	Monomicrobial (Adeno-associated dependoparvovirus A)	Monomicrobial (<i>Corynebacterium</i>)	NG	NG	No ($C_T=31.3$)
SCH10174-C	ND	ND	NG	NG	Yes ($C_T=30.9$)
SCH10175-C	Monomicrobial (<i>Moraxella osloensis</i>)	ND	<i>Moraxella osloensis</i>*	NG	No ($C_T=31.9$)

Abbreviations: C_T, cycles-to-threshold; MALDI-TOF, matrix-assisted laser desorption/ionisation time-of-flight mass spectrometry; ND, not determined; NG, no growth; PE, pleural effusion. Salmon-coloured cells denote possible false-negative results in the Pleural infection cohort. Bolded text denotes footnoted item; gray-shaded fields indicate discordant negative pleural infection diagnosis. [#]Minimum read threshold of ≥400 reads. [†]Minimum read threshold of ≥40 reads. [‡]Based on TRIzol interphase DNA extraction. *Cultures subjected to whole-genome sequencing. ^a*Escherichia coli* excluded from the metagenomic output due to its read count being lower than the background (contaminant) signal. ^bNo *S. saprophyticus* or other coagulase-negative staphylococci were identified, suggesting potential taxonomic misassignment or a detection limit issue. ^c*Pichia kudriavzevii* identified in the metagenome data (1963 reads). ^dProbable taxonomic misassignment of certain '*Staphylococcus*' as '*Streptococcus*' using bacterial metataxonomics. ^eSCH10131-P, yielded exceptionally high and diverse microbial burden across all testing methods; this case differed clinically from others in that the participant had suffered a ruptured oesophagus (Boerhaave syndrome).

Table 3. Comparative diagnostic accuracy of conventional culture and molecular methods for detecting pleural infection when compared to a surrogate gold-standard pleural infection classification of ‘probable pleural infection’*.



	CONVENTIONAL CULTURE	PANBACTERIAL QPCR	BACTERIAL METATAXONOMICS	METAGENOMICS	COMBINED MOLECULAR
SENSITIVITY (95% CI)	70.6 (44.0-89.7)	82.4 (56.6-96.2)	88.2 (63.6-98.5)	82.4 (56.6-96.2)	100 (80.5-100)
SPECIFICITY (95% CI)	100 (82.3-100)	94.7 (74.0-99.9)	94.7 (74.0-99.9)	89.5 (66.9-98.7)	84.2 (60.4-96.6)
PPV (95% CI)	100 (73.5-100)	93.3 (68.1-99.8)	93.8 (69.8-99.8)	87.5 (61.7-98.4)	85 (62.1-96.8)
NPV (95% CI)	79.2 (57.8-92.9)	85.7 (63.7-97.0)	90 (68.3-98.8)	85 (62.1-96.8)	100 (79.4-100)

Abbreviations: CI, confidence interval; NPV, negative predictive value; PPV, positive predictive value; qPCR, quantitative PCR; Combined Molecular, all molecular methods (qPCR, bacterial metataxonomics, metagenomics). **Probable Pleural Infection:* Gram stain and/or culture positive on pleural fluid; or pleural fluid pH < 7.2 and 2 or more of the following: serum C-reactive protein > 100 mg/L; complex pleural fluid on imaging; fever (> 38°C)

References


1. Bedawi EO, Guinde J, Rahman NM, Astoul P. Advances in pleural infection and malignancy. *European Respiratory Review* 2021;30:200002.
2. Cargill TN, Hassan M, Corcoran JP, et al. A systematic review of comorbidities and outcomes of adult patients with pleural infection. *European Respiratory Journal* 2019;54:1900541.
3. Brims F, Popowicz N, Rosenstengel A, et al. Bacteriology and clinical outcomes of patients with culture-positive pleural infection in Western Australia: A 6-year analysis. *Respirology* 2019;24:171-8.
4. Bedawi EO, Ricciardi S, Hassan M, et al. ERS/ESTS statement on the management of pleural infection in adults. *European Respiratory Journal* 2022;2201062.
5. Maskell NA, Batt S, Hedley EL, Davies CW, Gillespie SH, Davies RJ. The bacteriology of pleural infection by genetic and standard methods and its mortality significance. *Am J Respir Crit Care Med* 2006;174:817-23.
6. Rahman NM, Maskell NA, West A, et al. Intrapleural use of tissue plasminogen activator and DNase in pleural infection. *New England Journal of Medicine* 2011;365:518-26.
7. Shiraishi Y, Kryukov K, Tomomatsu K, et al. Diagnosis of pleural empyema/parapneumonic effusion by next-generation sequencing. *Infect Dis (Lond)* 2021;53:450-9.
8. Kanellakis NI, Wrightson JM, Gerry S, et al. The bacteriology of pleural infection (TORPIDS): an exploratory metagenomics analysis through next generation sequencing. *The Lancet Microbe* 2022;3:e294-e302.
9. Menzies SM, Rahman NM, Wrightson JM, et al. Blood culture bottle culture of pleural fluid in pleural infection. *Thorax* 2011;66:658-62.
10. Boyanova L, Vladimir D, Gergova G, et al. Anaerobic microbiology in 198 cases of pleural empyema: a Bulgarian study. *Anaerobe* 2004;10:261-7.
11. Corcoran JP, Wrightson JM, Belcher E, DeCamp MM, Feller-Kopman D, Rahman NM. Pleural infection: past, present, and future directions. *Lancet Respir Med* 2015;3:563-77.
12. Schlaberg R, Chiu CY, Miller S, Procop GW, Weinstock G. Validation of Metagenomic Next-Generation Sequencing Tests for Universal Pathogen Detection. *Arch Pathol Lab Med* 2017;141:776-86.
13. Marchesi JR, Ravel J. The vocabulary of microbiome research: a proposal. *Microbiome* 2015;3:31.
14. Hilton SK, Castro-Nallar E, Pérez-Losada M, et al. Metataxonomic and metagenomic approaches vs. culture-based techniques for clinical pathology. *Front Microbiol* 2016;7:484.
15. Dyrhovden R, Nygaard RM, Patel R, Ulvestad E, Kommedal Ø. The bacterial aetiology of pleural empyema. A descriptive and comparative metagenomic study. *Clin Microbiol Infect* 2019;25:981-6.
16. Xu H, Hu X, Wang W, et al. Clinical application and evaluation of metagenomic next-generation sequencing in pulmonary infection with pleural effusion. *Infect Drug Resist* 2022;15:2813-24.
17. Gimenez-Miranda L, Samhouri BF, Wolf MJ, et al. Diagnostic Yield of 16S Ribosomal Ribonucleic Acid Gene-Based Targeted Metagenomic Sequencing for Evaluation of Pleural Space Infection: A Prospective Study. *Mayo Clin Proc Innov Qual Outcomes* 2023;7:373-81.
18. Elsheikh A, Bhatnagar M, Rahman NM. Diagnosis and management of pleural infection. *Breathe (Sheff)* 2023;19:230146.
19. Liu G, Wang L, Li X, et al. The value of next-generation metagenomic sequencing in pathogen detection of pleural effusions and ascites from children with sepsis. *Front Cell Infect Microbiol* 2023;13:1130483.
20. Liu XP, Mao CX, Wang GS, Zhang MZ. Metagenomic next-generation sequencing for pleural effusions induced by viral pleurisy: A case report. *World J Clin Cases* 2023;11:844-51.
21. Psallidas I, Kanellakis NI, Bhatnagar R, et al. A pilot feasibility study in establishing the role of ultrasound-guided pleural biopsies in pleural infection (The AUDIO Study). *Chest* 2018;154:766-72.
22. Madden DE, Olagoke O, Baird T, et al. Express yourself: quantitative real-time PCR assays for rapid chromosomal antimicrobial resistance detection in *Pseudomonas aeruginosa*. *Antimicrob Agents Chemother* 2022;66:e0020422.
23. Herlemann DP, Labrenz M, Jurgens K, Bertilsson S, Waniek JJ, Andersson AF. Transitions in bacterial communities along the 2000 km salinity gradient of the Baltic Sea. *ISME J* 2011;5:1571-9.
24. Nelson MT, Pope CE, Marsh RL, et al. Human and extracellular DNA depletion for metagenomic analysis of complex clinical infection samples yields optimized viable microbiome profiles. *Cell Rep* 2019;26:2227-40.e5.

25. Nadkarni MA, Martin FE, Jacques NA, Hunter N. Determination of bacterial load by real-time PCR using a broad-range (universal) probe and primers set. *Microbiology (Reading)* 2002;148:257-66.
26. Ahmadi B, Hashemi SJ, Zaini F, et al. A case of onychomycosis caused by *Aspergillus candidus*. *Med Mycol Case Rep* 2012;1:45-8.
27. Klaassen CH, Jeunink MA, Prinsen CF, et al. Quantification of human DNA in feces as a diagnostic test for the presence of colorectal cancer. *Clin Chem* 2003;49:1185-7.
28. Light RW, Macgregor MI, Luchsinger PC, Ball WC, Jr. Pleural effusions: the diagnostic separation of transudates and exudates. *Ann Intern Med* 1972;77:507-13.
29. Insa R, Marin M, Martin A, et al. Systematic use of universal 16S rRNA gene polymerase chain reaction (PCR) and sequencing for processing pleural effusions improves conventional culture techniques. *Medicine (Baltimore)* 2012;91:103-10.
30. Nguyen S, Baker K, Padman BS, et al. Bacteriophage transcytosis provides a mechanism to cross epithelial cell layers. *mBio* 2017;8.
31. Brown-Jaque M, Muniesa M, Navarro F. Bacteriophages in clinical samples can interfere with microbiological diagnostic tools. *Sci Rep* 2016;6:33000.
32. Lim T. Recent advances in the management of pleural sepsis: What is the evidence? *Respirology* 1999;4:207-12.
33. Salter SJ, Cox MJ, Turek EM, et al. Reagent and laboratory contamination can critically impact sequence-based microbiome analyses. *BMC Biol* 2014;12:87.
34. Hadano Y, Ito K, Suzuki J, Kawamura I, Kurai H, Ohkusu K. *Moraxella osloensis*: an unusual cause of central venous catheter infection in a cancer patient. *Int J Gen Med* 2012;5:875-7.
35. Kwok B, Wu BG, Kocak IF, et al. Pleural fluid microbiota as a biomarker for malignancy and prognosis. *Sci Rep* 2023;13:2229.
36. Kea B, Hall MK, Wang R. Recognising bias in studies of diagnostic tests part 2: interpreting and verifying the index test. *Emerg Med J* 2019;36:501-5.



PSI cohort

Control cohort



PSI cohort

Control cohort

